

# ORBITS OF PLANE PARTITIONS OF EXCEPTIONAL LIE TYPE

HOLLY MANDEL AND OLIVER PECHENIK

ABSTRACT. For each minuscule flag variety  $X$ , there is a corresponding minuscule poset, describing its Schubert decomposition. We study an action on plane partitions over such posets, introduced by P. Cameron and D. Fon-der-Flaass (1995). For plane partitions of height at most 2, D. Rush and X. Shi (2013) proved an instance of the cyclic sieving phenomenon, completely describing the orbit structure of this action. They noted their result does not extend to greater heights in general; however, when  $X$  is one of the two minuscule flag varieties of exceptional Lie type  $E$ , they conjectured explicit instances of cyclic sieving for all heights.

We prove their conjecture in the case that  $X$  is the Cayley-Moufang plane of type  $E_6$ . For the other exceptional minuscule flag variety, the Freudenthal variety of type  $E_7$ , we establish their conjecture for heights at most 4, but show that it fails generally. We further give a new proof of a unpublished cyclic sieving of D. Rush and X. Shi (2011) for plane partitions of any height in the case  $X$  is an even-dimensional quadric hypersurface. Our argument uses ideas of K. Dilks, O. Pechenik, and J. Striker (2017) to relate the action on plane partitions to combinatorics derived from  $K$ -theoretic Schubert calculus.

## 1. INTRODUCTION

The *minuscule posets* are a remarkable collection of partially-ordered sets that arise naturally from the representation theory of Lie algebras or alternatively from the Schubert calculus of certain generalized flag varieties. We study dynamic enumerative properties of plane partitions over these posets.

A special case of this situation is that of ordinary plane partitions fitting inside a fixed rectangular box. In this context, P. Cameron and D. Fon-der-Flaass [CFDF95] initiated the study of a combinatorially-natural operator  $\Psi$ . This operator is now generally known as *rowmotion* and has become a subject of intense study (cf., e.g., [Pan09, SW12, AST13, RS13, EP14, PR15, GR15, GR16, DPS17, Vor17]). We will define and describe in detail both minuscule posets and the operation of rowmotion in Sections 2 and 3, respectively.

For any poset  $P$ , let  $\mathbf{PP}^k(P)$  denote the set of plane partitions of height at most  $k$  over  $P$ , that is to say the set of order ideals in the product  $P \times \mathbf{k}$  of  $P$  with a chain poset of size  $k$ . Let  $f_P^k$  denote the generating function that enumerates the elements of  $\mathbf{PP}^k(P)$  by cardinality. In the special case  $k \leq 2$  and  $P$  minuscule, D. Rush and

---

*Date:* October 24, 2017.

*2010 Mathematics Subject Classification.* Primary 05A15; Secondary 05E18, 06A07, 17B25.

*Key words and phrases.* plane partition, increasing tableau, promotion, rowmotion, exceptional Lie algebra, minuscule poset, cyclic sieving phenomenon.

X. Shi [RS13] showed that  $f_P^k$  also encodes the orbit structure of rowmotion via an instance of the **cyclic sieving phenomenon** (introduced by V. Reiner, D. Stanton, and D. White [RSW04]). That is to say that for  $k \leq 2$ , the number of minuscule plane partitions fixed by the  $d$ -fold application of rowmotion  $\Psi^{\circ d}$  is the evaluation of the polynomial  $f_P^k$  at  $\zeta^d$ , where  $\zeta$  is any primitive  $n$ th root-of-unity and  $n$  is the order of  $\Psi$  on  $\mathbf{PP}^k(P)$ . (Since some of our other operators have superscripts in their names, we will consistently denote  $d$ -fold composition of any operator  $\tau$  by  $\tau^{\circ d}$ , instead of the more usual notation  $\tau^d$ , which could otherwise be ambiguous.)

It was noted, however, in [RS13] that this instance of cyclic sieving does not extend to the case  $k \geq 3$  for general minuscule posets. Nonetheless, D. Rush and X. Shi conjectured the following. (The minuscule posets in question are illustrated in Figure 1.)

**Conjecture 1.1** ([RS13, Conjecture 11.1]). *Let  $P$  be one of the two minuscule posets associated to an exceptional Lie algebra of type  $E$  and let  $k \in \mathbb{Z}_{\geq 0}$ . Then  $f_P^k$  is a cyclic sieving polynomial for the action of  $\Psi$  on  $\mathbf{PP}^k(P)$ .*

Our main result is to completely resolve Conjecture 1.1.

**Theorem 1.2.** *Conjecture 1.1 holds for the  $E_6$  minuscule poset (cf. Figure 1A) and all  $k$ , but holds for the  $E_7$  minuscule poset (cf. Figure 1B) only when  $k \leq 4$ .*

Verification of Conjecture 1.1 in the the cases  $k \leq 4$  for the  $E_6$  minuscule poset and  $k \leq 3$  for the  $E_7$  minuscule poset was previously reported in [RS13]. Our new results are therefore:

- cyclic sieving for the  $E_6$  minuscule poset when  $k > 4$ ,
- cyclic sieving for the  $E_7$  minuscule poset when  $k = 4$ , and
- failure of the conjectured cyclic sieving for the  $E_7$  minuscule poset for  $k > 4$ .

Our approach to proving Theorem 1.2 is to extend the ideas of K. Dilks, O. Pechenik, and J. Striker [DPS17] to this minuscule setting, thereby relating the action of  $\Psi$  to the action of  $K$ -promotion on *increasing tableaux*.  $K$ -promotion was first studied in [Pec14] building on combinatorial tools for  $K$ -theoretic Schubert calculus due to H. Thomas and A. Yong [TY09]. We show that the action of  $K$ -promotion is controlled by its behavior on a finite subset of increasing tableaux. By understanding the orbit structure of this subset, we are able to determine the complete orbit structure, thereby establishing Theorem 1.2.

Having developed these methods, it becomes straightforward to prove the following additional result.

**Theorem 1.3.** *Let  $P$  be a minuscule poset associated to an even-dimensional quadric of type  $D$  (cf. Figure 3C) and let  $k \in \mathbb{Z}_{\geq 0}$ . Then  $f_P^k$  is a cyclic sieving polynomial for the action of  $\Psi$  on  $\mathbf{PP}^k(P)$ .*

Theorem 1.3 was previously announced by D. Rush and X. Shi [RS13, Theorem 10.1]; however, they omitted their lengthy proof [RS11, §10] from the published



(A) Cayley-Moufang poset



(B) Freudenthal poset

FIGURE 1. The two minuscule posets associated to exceptional Lie algebras of type  $E$ . Here, we have drawn the posets to resemble Young diagrams in Cartesian (“French”) orientation; the boxes are the elements of the poset and each box covers the box immediately below it and the box immediately to its left (if such boxes exist). Hence the minimal element of each poset is the box at the far left of the bottom row. The Cayley-Moufang poset is associated to  $E_6$ , while the Freudenthal poset is associated to  $E_7$ . (There is no minuscule poset associated to  $E_8$ .)

paper. We believe that our alternative proof of Theorem 1.3 via  $K$ -theoretic combinatorics provides a different understanding.

Often instances of the cyclic sieving phenomenon can be proven using representation-theoretic techniques [RSW04, Rho10], and when instances are established in a more direct fashion (as we do here), it may be a clue toward as yet undeveloped algebra (cf. [Rho17]). The results in Theorems 1.2 and 1.3 perhaps suggest the existence of new symmetric group module structures on the sets  $\mathcal{PP}^k(P)$ ; however, we do not currently understand how to build such structures with the requisite properties in these cases.

This paper is organized as follows. In Section 2, we define the minuscule posets, recalling their classification and certain other properties we will use. In Section 3, we collect the facts we need about the rowmotion operator  $\Psi$ . Section 4 concerns the operation of  $K$ -promotion on increasing tableaux, the key tool in our proofs of

Theorems 1.2 and 1.3. In Section 5, we demonstrate the equivariant bijection between  $\text{PP}^k(P)$  under the action of  $\Psi$  and certain increasing tableaux under  $K$ -promotion, extending the main observation of [DPS17]. Finally, Section 6 is devoted to study of these increasing tableaux; our results in this last section prove Theorems 1.2 and 1.3 via the equivariant bijection of Section 5.

## 2. MINUSCULE POSETS

Let  $\mathbf{G}$  be a complex connected reductive Lie group with maximal torus  $\mathbf{T}$ . Denote by  $W$  the Weyl group  $N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ . The root system  $\Phi$  of  $\mathbf{G}$  may be partitioned  $\Phi^+ \sqcup \Phi^-$  into positive and negative roots according to a choice  $\Delta$  of simple roots. There is a natural poset structure on  $\Phi^+$  obtained as the transitive closure of the covering relation  $\alpha < \beta$  if and only if  $\beta - \alpha \in \Delta$ . The choice of bipartition of  $\Phi$  into positive and negative roots further specifies a choice of a Borel subgroup  $\mathbf{B}_+ \subset \mathbf{G}$  and an opposite Borel subgroup  $\mathbf{B}_- \subset \mathbf{G}$  with  $\mathbf{B}_+ \cap \mathbf{B}_- = \mathbf{T}$ .

We say  $\delta \in \Delta$  is **minuscule** if for every  $\alpha \in \Phi^+$ ,  $\delta^\vee$  appears with multiplicity at most 1 in the simple coroot expansion of  $\alpha^\vee$ . The classification of minuscule roots is well known and is illustrated in Figure 2 in terms of Dynkin diagrams.



FIGURE 2. In each of the finite-type Dynkin diagrams above, each minuscule root is marked as a pale blue circle, while the non-minuscule simple roots are marked in black. In type  $A_n$ , every node is minuscule, while in the other types only the indicated leaves are minuscule. The remaining finite-type Dynkin diagrams are omitted as they have no minuscule nodes.

For each minuscule simple root  $\delta$ , there is an associated **minuscule poset**  $P_\delta$  obtained as the subposet of  $\Phi^+$  induced on those positive roots  $\alpha$  where  $\delta$  appears with nonzero coefficient in the simple root expansion of  $\alpha$ .

Alternatively, one may obtain the minuscule posets via the geometry of certain generalized flag varieties. If  $\mathbf{P}_\delta \supset \mathbf{B}_+$  denotes the maximal parabolic subgroup of

$\mathbf{G}$  associated to the minuscule simple root  $\delta$ , then the space  $X = \mathbf{G}/\mathbf{P}_\delta$  is called a **minuscule variety**. The minuscule varieties are smooth projective varieties with various nice geometric properties (cf., e.g., [BL00] for details). The natural action of the Borel subgroup  $\mathbf{B}_+$  on  $X$  has finitely-many orbits, whose Zariski closures are the **Schubert varieties**. Given two Schubert varieties in  $X$ , it is known that they are either disjoint or else one is a subset of the other. Indeed the poset  $Y_X$  of Schubert varieties of  $X$  with respect to inclusion is a distributive lattice and its corresponding poset of join irreducibles is isomorphic to the minuscule poset  $P_\delta$  for the minuscule simple root  $\delta$ , as constructed above.

The minuscule posets are completely classified. We illustrate them here in Figures 1 and 3. Our focus will be on the **propellers**, the **Cayley-Moufang poset**, and the **Freudenthal poset** as shown in Figures 3C, 1A, and 1B, respectively. The propeller with  $k$  elements ( $k \geq 6$ ) is associated to the minuscule node that is not adjacent to the trivalent node in the  $D_{\frac{k+2}{2}}$  Dynkin diagram, the Cayley-Moufang poset is associated to the unique minuscule simple root for  $E_6$ , and the Freudenthal poset is associated to the unique minuscule simple root for  $E_7$  (cf. Figure 2). The corresponding minuscule varieties are, respectively, even-dimensional quadric hypersurfaces, the octonionic projective plane (or Cayley-Moufang plane), and the Freudenthal variety. The remaining minuscule roots yield minuscule posets that are *rectangles* or *shifted staircases* (illustrated in Figure 3A and B). These correspond respectively to type  $A$  Grassmannians and to maximal orthogonal Grassmannians; we will not study these posets further in this paper.

A poset that is linearly ordered is called a **chain**; we denote the chain of size  $k$  by  $\mathbf{k}$ . By a **plane partition** of height at most  $k$  over a poset  $P$ , we mean an order ideal of the product poset  $P \times \mathbf{k}$ . Clearly, a such a plane partition may be identified with a weakly order-reversing map  $\pi : P \rightarrow \mathbf{k}$  (i.e., a map  $\pi$  such that  $p \leq p'$  implies  $\pi(p) \geq \pi(p')$ ). The poset  $P$  is called **Gaussian** if the generating function  $f_P^k$  enumerating plane partitions of height at most  $k$  over  $P$  according to cardinality may be expressed in the form

$$\frac{(1 - x^{h_1+m})(1 - x^{h_2+m}) \cdots (1 - x^{h_t+m})}{(1 - x^{h_1})(1 - x^{h_2}) \cdots (1 - x^{h_t})},$$

for some nonnegative integers  $t, h_1, h_2, \dots, h_t \in \mathbb{Z}_{\geq 0}$  independent of  $k$ . R. Proctor showed that every minuscule poset is Gaussian [Pro84]. Indeed, it is conjectured that there are no other Gaussian posets. Given the beautiful form of these generating functions, it is perhaps not too surprising that  $f_P^k$  should in some instances play a role in cyclic sieving phenomena, as in Theorems 1.2 and 1.3.

### 3. THE ROWMOTION OPERATOR $\Psi$

If  $P$  is any poset, let  $J(P)$  denote the set of all its order ideals. For  $\mathcal{I} \in J(P)$  an order ideal of  $P$ , we define  $\Psi(\mathcal{I}) \in J(P)$  to be the order ideal generated by the minimal elements of the complement  $P - \mathcal{I}$ . This operator  $\Psi$  is closely related to that described in [BS74, Duc74, CFDF95] in slightly different contexts. It was first



FIGURE 3. Together with the two posets shown in Figure 1, these are exemplars of all five families of minuscule poset, shown in Cartesian orientation. The elements of each poset are the boxes, and each box is covered by any box immediately above it or immediately to its right. Rectangles may have arbitrary height and width. Shifted staircases have arbitrary width, and height equal to their width. Propellers consist of two rows of arbitrary but equal length, overlapping by two boxes in the center. The Cayley-Moufang and Freudenthal posets of Figure 1 are exceptional, forming singleton families.

considered as an action on order ideals by J. Striker and N. Williams [SW12]; we follow them in referring to  $\Psi$  as **rowmotion**.

Here, we recall the facts that we will need about rowmotion. For  $x \in P$ , define the **toggle at  $x$**  to be the operator  $t_x : J(P) \rightarrow J(P)$  defined by

$$t_x(\mathcal{I}) = \begin{cases} \mathcal{I} \cup \{x\}, & \text{for } x \notin \mathcal{I} \text{ and } \mathcal{I} \cup \{x\} \in J(P); \\ \mathcal{I} - \{x\}, & \text{for } x \in \mathcal{I} \text{ and } \mathcal{I} - \{x\} \in J(P); \\ \mathcal{I}, & \text{otherwise.} \end{cases}$$

Following the terminology of [DPS17], a **3-dimensional lattice projection** of a finite ranked poset  $P$  is an order- and rank-preserving map  $\pi : P \rightarrow \mathbb{Z}^3$ , where the partial order on  $\mathbb{Z}^3$  is given by componentwise comparison and  $\mathbb{Z}^3$  is ranked by the sum of the coordinates. Considering the classification of minuscule posets in Figures 1 and 3, observe that for every minuscule  $P$  and any  $k > 0$ , the poset  $P \times \mathbf{k}$  has an injective 3-dimensional lattice projection induced from the obvious embeddings of those pictures in the planar lattice. For  $v = \{\pm 1\}^3$  and  $i \in \mathbb{Z}$ , let

$$(3.1) \quad \tau_{\pi, v}^i = \prod_{\langle v, \pi(x) \rangle = i} t_x.$$

That is,  $\tau_{\pi, v}^i$  is the composition of the toggles  $t_x$  for all  $x$  with  $\pi(x)$  lying on the affine plane defined by  $\langle v, \cdot \rangle = i$ . Since the toggles  $t_{x_1}$  and  $t_{x_2}$  clearly commute whenever neither  $x_1$  nor  $x_2$  covers the other, this composition is well-defined. Then

$(\pi, v)$ -**motion** on  $P$  is the operator

$$\mathbf{Mot}_{\pi, v} := \prod_{i \in \mathbb{Z}} \tau_{\pi, v}^i,$$

where the factors are taken in reverse of the natural order. Note that in this product, only finitely many factors act nontrivially, since  $\pi(P)$  intersects only a finite number of the affine planes defined by  $\langle v, \cdot \rangle = i$ . We recall the following pair of results, which are fundamental to our further analysis of rowmotion on minuscule plane partitions.

**Proposition 3.1** ([CFDF95, Lemma 1]; [DPS17, Proposition 3.18]). *For any finite ranked poset  $P$  with a 3-dimensional lattice projection  $\pi$ , we have*

$$\Psi = \mathbf{Mot}_{\pi, (1,1,1)}$$

as operators on  $J(P)$ . □

**Proposition 3.2** ([DPS17, Theorem 3.25]). *For any finite ranked poset  $P$  with a 3-dimensional lattice projection  $\pi$  and any vectors  $v, w \in \{\pm 1\}^3$ , there is an equivariant bijection between  $J(P)$  under  $\mathbf{Mot}_{\pi, v}$  and  $J(P)$  under  $\mathbf{Mot}_{\pi, w}$ . In particular,  $\mathbf{Mot}_{\pi, v}$  and  $\mathbf{Mot}_{\pi, w}$  induce the same multiset of orbit cardinalities on  $J(P)$ . □*

#### 4. $K$ -PROMOTION OF INCREASING TABLEAUX

**4.1. Definitions.** A classical approach to the cohomological Schubert calculus of complex type  $A$  Grassmannians is to use the jeu de taquin on standard Young tableaux introduced by M.-P. Schützenberger [Sch77]. (See [Ful97, Man01] for modern expositions of the classical jeu de taquin theory.) A theme of modern Schubert calculus has been extending such theories to richer generalized cohomologies and in particular into the  $K$ -theory ring of algebraic vector bundles. For a partial survey of recent work related to  $K$ -theoretic Schubert calculus and the associated combinatorics, see [PY17].

The first  $K$ -theoretic Littlewood-Richardson rule was discovered by A. Buch [Buc02]; this rule, however, was not based on jeu de taquin. H. Thomas and A. Yong [TY09] later found a different Littlewood-Richardson rule for the same structure coefficients by directly extending M.-P. Schützenberger’s jeu de taquin. This latter rule was conjectured in [TY09] to extend to the  $K$ -theoretic Schubert structure coefficients of all minuscule varieties, as was proven by a combination of [BR12, CTY14, BS16]. In this Thomas-Yong theory, the role of standard Young tableaux is filled by *increasing tableaux*.

**Definition 4.1.** Let  $\lambda$  be an order ideal in a minuscule poset, considered as an generalized Young diagram in Cartesian orientation as in Figures 1 and 3. An **increasing tableau** of shape  $\lambda$  is an assignment of a positive integer to each box of  $\lambda$ , such that entries strictly increase from left to right along rows and strictly increase from bottom to top going up columns. Let  $\text{Inc}^q(\lambda)$  denote the set of all increasing tableaux of shape  $\lambda$  with all entries at most  $q$ . We identify  $\text{Inc}^q(\lambda)$  with the set of all strictly order-preserving maps from the poset  $\lambda$  to the chain poset  $\mathbf{q}$  of size  $q$ . For examples of increasing tableaux, see Figures 4 and 5.

Using the  $K$ -jeu de taquin of H. Thomas and A. Yong [TY09], one has a  $K$ -promotion operator  $\mathbf{pro}^q$  on  $\text{Inc}^q(\lambda)$ , directly extending M.-P. Schützenberger’s classical definition of promotion [Sch72].  $K$ -promotion was first studied in [Pec14], and has been further investigated in [BPS16, PSV16, Rho17, DPS17, Pec17, Vor17]. Since studying the action of  $\mathbf{pro}^q$  will be fundamental to our proof of Theorems 1.2 and 1.3, we must recall the definition of  $\mathbf{pro}^q$ . We will indeed use both the original formulation from [Pec14], as well as an equivalent formulation from [DPS17, Proposition 2.4].

Let  $T \in \text{Inc}^q(\lambda)$ . The  **$K$ -Bender-Knuth operator**  $\rho_i$  acts on  $T$  by swapping the letters  $i$  and  $i + 1$  everywhere in  $T$  *where doing so would not violate the increasingness conditions*. More precisely,  $\rho_i$  looks at the set of boxes labeled  $i$  or  $i + 1$ , decomposes that set into edge-connected components, swaps the labels  $i$  and  $i + 1$  in each component that is a single box, and acts trivially on each non-trivial connected component. Notice that each  $\rho_i$  is an involution. Using the characterization of [DPS17, Proposition 2.4],  **$K$ -promotion** may then be defined by

$$\mathbf{pro}^q(T) := \rho_{q-1} \circ \dots \circ \rho_1(T).$$

Examples of the action of  $K$ -Bender-Knuth operators are shown in Figure 4, while an example of the full process of  $K$ -promotion is shown in Figure 5. We will relate  $K$ -Bender-Knuth operators to poset toggles in Section 5.



FIGURE 4. The three increasing tableaux on the right are obtained from the increasing tableau at left by applying the indicated  $K$ -Bender-Knuth operators. The arrows point in both directions since each  $K$ -Bender-Knuth operator is involutive.





FIGURE 5. The calculation of the  $K$ -promotion  $\mathbf{pro}^{13}(T)$  of the increasing tableau  $T$  (on the Cayley-Moufang poset) through the action of the sequence of  $K$ -Bender-Knuth operators  $\rho_1, \dots, \rho_{12}$ .

**4.2. Inflation and deflation.** Suppose  $T \in \text{Inc}^q(\lambda)$  and consider  $T$  as a strictly order-preserving map from  $\lambda$  to the chain poset  $\mathbf{q}$ . We say that  $T$  is **gapless** if this map is surjective; otherwise,  $T$  is **gappy**. We write  $\text{Inc}_{\text{gl}}^q(\lambda)$  for the subset of all gapless tableaux in  $\text{Inc}^q(\lambda)$ . Notice that the set

$$\text{Inc}_{\text{gl}}(\lambda) := \bigcup_q \text{Inc}_{\text{gl}}^q(\lambda)$$

is necessarily finite; this fact will be critical for our ability to prove Theorems 1.2 and 1.3.

For  $T \in \text{Inc}^q(\lambda)$ , let  $N_T := |\text{range}(T)|$  be the number of distinct labels in  $T$ . For each  $q$ , we define the **deflation** map

$$\text{Defl}^q : \text{Inc}^q(\lambda) \rightarrow \coprod_{0 \leq k \leq q} \text{Inc}_{\text{gl}}^k(\lambda)$$

by

$$[\text{Defl}^q(T)](\mathbf{x}) = \#\{h \in \text{range}(T) : h \leq T(\mathbf{x})\},$$

for  $T \in \text{Inc}^q(\lambda)$  and  $\mathbf{x} \in \lambda$ . Note that  $\text{Defl}^q(T) \in \text{Inc}_{\text{gl}}^{N_T}(\lambda)$  and that  $N_T \leq q$ .

For positive integers  $k \leq j$ , let  $\begin{bmatrix} j \\ k \end{bmatrix}$  denote the collection of binary vectors of length  $j$  with  $k$  1's. We now define the **content vector** function

$$\text{Con}^q : \text{Inc}^q(\lambda) \rightarrow \{0, 1\}^q = \coprod_{0 \leq k \leq q} \begin{bmatrix} q \\ k \end{bmatrix}$$

by

$$\text{Con}^q(T) = (c_1, \dots, c_q),$$

where  $c_k = 1$  if  $k \in \text{range}(T)$  and  $c_k = 0$  if  $k \notin \text{range}(T)$ . (Warning: This definition of content differs from the usual notion of content for semistandard tableaux in that here we do not care about the multiplicity of a label, but only its presence or absence.)

**Example 4.2.** If  $T = \begin{array}{|c|c|c|} \hline 4 & 5 & 6 \\ \hline 1 & 2 & 5 \\ \hline \end{array} \in \text{Inc}^7(2 \times 3)$ , then the deflation of  $T$  is

$$\text{Defl}^7(T) = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 1 & 2 & 4 \\ \hline \end{array} \in \text{Inc}_{\text{gl}}^5(2 \times 3).$$

The content vector of  $T$  is  $\text{Con}^7(T) = (1, 1, 0, 1, 1, 1, 0) \in \begin{bmatrix} 7 \\ 5 \end{bmatrix} \subset \{0, 1\}^7$ .  $\diamond$

Note that if  $\text{Defl}^q(T) \in \text{Inc}_{\text{gl}}^k(\lambda)$ , then  $\text{Con}^q(T) \in \begin{bmatrix} q \\ k \end{bmatrix}$ . We denote by  $\text{DeflCon}^q$  the product map

$$\text{DeflCon}^q := (\text{Defl}^q, \text{Con}^q).$$

**Theorem 4.3.** *The map*

$$\text{DeflCon}^q : \text{Inc}^q(\lambda) \rightarrow \coprod_{0 \leq k \leq q} \left( \text{Inc}_{\text{gl}}^k(\lambda) \times \begin{bmatrix} q \\ k \end{bmatrix} \right)$$

*is bijective.*

*Proof.* A two-sided inverse for  $\text{DeflCon}^q$  is given as follows. Let  $[j]$  denote the set  $\{1, 2, \dots, j\}$ . For a binary vector  $v \in \{0, 1\}^q$ , let  $N_v$  be the number of 1's in  $v$ , and define a **vector inflation** map

$$\text{VecInfl}_v^q : [N_v] \rightarrow [q]$$

by

$$\text{VecInfl}_v^q(k) = \min_{n \in [q]} \left( \sum_{\ell=1}^n v\{\ell\} = k \right).$$

(We use curly braces “ $\{ \}$ ” throughout to denote vector components.) An integer  $j \in [q]$  is in the range of  $\text{VecInfl}_v^q$  if and only if  $v\{j\} = 1$ . From this fact and the definitions, it follows that  $\text{VecInfl}_{\text{Con}^q(T)}^q \circ \text{Defl}^q(T) = T$ . Now define the **tableau inflation** map

$$\text{Infl}^q : \coprod_{0 \leq k \leq q} \left( \text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[q]}{k} \right) \rightarrow \text{Inc}^q(\lambda)$$

by

$$\text{Infl}^q(S, v) = \text{VecInfl}_v^q \circ S.$$

Say  $(S, v) \in \text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[q]}{k}$ . Since  $S$  is surjective and  $\text{Infl}^q(S, v)$  maps onto the indices of nonzero components in  $v$ ,  $\text{Con}^q \circ \text{Infl}^q(S, v) = v$ . Also,

$$\begin{aligned} (\text{Defl}^q(\text{VecInfl}_v^q \circ S))(x) &= \#\{h \in \text{range}(\text{VecInfl}_v^q \circ S) : h \leq (\text{VecInfl}_v^q \circ S)(x)\} \\ &= \#\{h : v\{h\} \neq 0 \text{ and } \sum_{\ell=1}^h v\{\ell\} \leq S(x)\} \\ &= S(x). \end{aligned}$$

Therefore,  $\text{Infl}^q$  is a two-sided inverse for  $\text{DeflCon}^q$ .  $\square$

**Example 4.4.** Let  $v = (1, 1, 0, 1, 1, 1, 0) \in \{0, 1\}^7$ . Then,  $N_v = 5$  and  $\text{VecInfl}_v^7 : [5] \rightarrow [7]$  is given by

$$\begin{aligned} \text{VecInfl}_v^7(1) &= 1, \\ \text{VecInfl}_v^7(2) &= 2, \\ \text{VecInfl}_v^7(3) &= 4, \\ \text{VecInfl}_v^7(4) &= 5, \\ \text{VecInfl}_v^7(5) &= 6. \end{aligned}$$

Now, for  $S = \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline 1 & 2 & 4 \\ \hline \end{array}$ , we have  $\text{Infl}^7(S, v) = \text{VecInfl}_v^7 \circ S = \begin{array}{|c|c|c|} \hline 4 & 5 & 6 \\ \hline 1 & 2 & 5 \\ \hline \end{array}$ . Note

that this process has recovered the tableau  $T$  of Example 4.2 from  $S = \text{Defl}^7(T)$  and  $v = \text{Con}^7(T)$ .  $\diamond$

**4.3. Interaction between  $K$ -promotion and deflation.** By determining the relation between the maps  $\text{DeflCon}^q$  and  $\text{pro}^q$ , we will show that the behavior of  $\text{pro}^q$  is controlled by its action on the subset of gapless increasing tableaux. Much of the power of this observation comes from the fact that there are only finitely-many gapless increasing tableaux of any fixed shape  $\lambda$ , and that therefore the action of  $\text{pro}^q$  on the set of all increasing tableaux of shape  $\lambda$  is governed by a finite amount of data.

To establish this relation, we first need to introduce notation for the steps in a different characterization of  $\text{pro}^q$ . Let  $\text{Rep}_{1 \rightarrow \bullet}$  be the operator on  $\text{Inc}^q(\lambda)$  that replaces each instance of 1 by  $\bullet$ . Let  $\text{swap}_n$  be the operator that finds all edge-connected

components containing  $\bullet$  and  $n$ , leaves trivial components unchanged, and switches  $n$  and  $\bullet$  in nontrivial connected components. Let  $\text{Rep}_{\bullet \rightarrow q+1}$  replace each instance of  $\bullet$  with  $q+1$ . Finally, let  $\text{Decr}$  be the operator that decrements each label by 1. Then, [DPS17, Proposition 2.4] gives the following alternative characterization of the operation of  $K$ -promotion on  $\text{Inc}^q(\lambda)$ :

$$(4.1) \quad \text{pro}^q = \text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \cdots \circ \text{swap}_2 \circ \text{Rep}_{1 \rightarrow \bullet}.$$

(This was, in fact, the original definition of  $K$ -promotion introduced in [Pec14].) An example of computing  $K$ -promotion via this alternative but equivalent process is shown in Figure 6.



FIGURE 6. The calculation of the  $K$ -promotion  $\text{pro}^5(T)$  of the increasing tableau  $T$  through the action of a sequence of swaps.

Using this characterization of  $\text{pro}^q$ , we give names to the intermediate products of this process. For  $T \in \text{Inc}^q(\lambda)$ , we define the tableau  $T^{(N)} : \lambda \rightarrow \{2, \dots, q, \bullet\}$  by

$$T^{(N)} := \text{swap}_N \circ \cdots \circ \text{swap}_2 \circ \text{Rep}_{1 \rightarrow \bullet}(T)$$

for  $N > 1$  and  $T^{(1)} := \text{Rep}_{1 \rightarrow \bullet}(T)$ . We define the associated order ideal

$$\lambda(T^{(N)}) := (T^{(N)})^{-1}(\{2, \dots, N\})$$

to be the set of boxes of  $T^{(N)}$  containing entries from  $\{2, \dots, N\}$ . Finally, define

$$T_N := T^{(N)}|_{\lambda(T^{(N)})}.$$

Intuitively,  $T^{(N)}$  shows  $T$  at the point in  $K$ -promotion at which the boxes labeled 2 through  $N$  have been acted upon by  $\text{swap}$  operators, while  $T_N$  is the restriction

of  $T^{(N)}$  to this subset of boxes. Examples of these intermediate tableaux appear in Figure 7.

The following lemmas use these intermediate products to show that  $K$ -promotion acts on a gappy tableau in essentially the same way and as on its deflation, provided that the gappy tableau contains at least one 1. The following simple, but important, result shows why this condition on content is necessary.

**Lemma 4.5.** *If  $\text{DeflCon}^q(T)\{1\} = 0$ , then  $\text{pro}^q(T) = \text{Decr}(T)$ .*

*Proof.* If  $\text{DeflCon}^q(T)\{1\} = 0$ , then  $\text{Rep}_{1 \rightarrow \bullet}(T) = T$ . But then  $\text{swap}_\ell(T) = T$  for all  $\ell$ , since there are no nontrivial short ribbons containing the labels  $\bullet$  and  $\ell$ . Also,  $\text{Rep}_{\bullet \rightarrow 1}(T) = T$ , since there is not  $\bullet$  label in  $T$ . Therefore,  $\text{Decr}$  is the only operator in Equation (4.1) that acts nontrivially on  $T$ .  $\square$



FIGURE 7. The tableau  $T^{(3)}$  is obtained from  $T$  by applying in sequence the operations  $\text{Rep}_{1 \rightarrow \bullet}$ ,  $\text{swap}_2$ , and  $\text{swap}_3$ . We then obtain  $T_3$  from  $T^{(3)}$  by restricting to those boxes with labels less than or equal to 3. The order ideal  $\lambda(T^{(3)})$  is the shape of the tableau  $T_3$ .

**Lemma 4.6.** *Let  $T \in \text{Inc}^q(\lambda)$  with  $\text{Con}^q(T)\{1\} = 1$ . Let  $i_k = \text{VecInfl}_{\text{Con}^q(T)}^q(k)$  for  $k = 1, \dots, N_T$ . Then, for all  $\mathbf{x} \in \lambda$  and  $1 \leq k \leq N_T$ , we have*

$$(4.2) \quad \text{Defl}^q(T)^{(k)}(\mathbf{x}) = \bullet \text{ if and only if } T^{(i_k)}(\mathbf{x}) = \bullet.$$

*Proof.* We proceed by induction on  $k$ . Fix  $T \in \text{Inc}^q(\lambda)$  with  $\text{Con}^q(T)\{1\} = 1$ . In the base case  $k = 1$ ,  $T^{(i_1)} = T^{(1)} = \text{Rep}_{1 \rightarrow \bullet}(T)$  has  $\bullet$  in exactly those boxes  $\mathbf{x} \in \lambda$  such that  $T(\mathbf{x}) = 1$ . On the other hand,  $\text{Defl}^q(T)^{(1)} = \text{Rep}_{1 \rightarrow \bullet}(\text{Defl}^q(T))$  has  $\bullet$  in exactly those boxes  $\mathbf{x} \in \lambda$  such that  $\text{Defl}^q(T)(\mathbf{x}) = 1$ . But since  $\text{Con}^q(T)\{1\} = 1$ ,  $\text{Defl}^q(T)(\mathbf{x}) = 1$  if and only if  $T(\mathbf{x}) = 1$ .

Now, say the lemma holds for all tableaux  $T$  in the case  $k < t$ . First consider the case where  $T^{(i_t)}(\mathbf{x}) = \bullet$  and  $T^{(i_{t-1})}(\mathbf{x}) \neq \bullet$ . This is equivalent to the statement that  $T^{(i_{t-1})}(\mathbf{x}) = i_t$  and  $T^{(i_{t-1})}(\mathbf{x}') = \bullet$  for some  $\mathbf{x}'$  adjacent to  $\mathbf{x}$ . But since  $\text{Rep}_{1 \rightarrow \bullet}$  and  $\text{swap}_j$  for  $j \leq i_{t-1}$  only act on elements of  $T$  labeled with  $\bullet$ 's or values less than or equal to  $i_{t-1}$ ,  $T^{(i_{t-1})}(\mathbf{x}) = i_t$  implies that  $T^{(i_{t-1})}(\mathbf{x}) = T(\mathbf{x}) = i_t$ . This is,

in turn, equivalent by Theorem 4.3 to the statement  $\text{Defl}^q(T)(x) = t$ , implying that  $\text{Defl}^q(T)^{(t-1)}(x) = t$ . Additionally, by the inductive hypothesis,  $T^{(i_{t-1})}(x') = \bullet$  if and only if  $\text{Defl}^q(T)^{(t-1)}(x') = \bullet$ . Therefore, the conditions of this case are equivalent to the conditions  $\text{Defl}^q(T)^{(t)}(x) = \bullet$  and  $\text{Defl}^q(T)^{(t-1)}(x) \neq \bullet$ , as desired.

Consider the other possible case, where  $T^{(i_t)}(x) = \bullet$  and  $T^{(i_{t-1})}(x) = \bullet$ . This scenario is equivalent to the statement that  $T^{(i_{t-1})}(x) = \bullet$  and there is no  $x'$  adjacent to  $x$  such that  $T^{(i_{t-1})}(x') = i_t$ . By similar reasoning as in the previous case, the second condition is equivalent to the statement that there is no  $x'$  adjacent to  $x$  with  $T(x') = i_t$ . This statement is, in turn, equivalent to saying that there is no  $x'$  adjacent to  $x$  with  $\text{Defl}^q(T)(x') = \text{Defl}^q(T)^{(t-1)}(x') = t$ . Since  $\text{Defl}^q(T)^{(t-1)}(x) = \bullet$  by the inductive hypothesis, the conditions of this case are finally equivalent to the conditions that  $\text{Defl}^q(T)^{(t)}(x) = \bullet$  and  $\text{Defl}^q(T)^{(t-1)}(x) = \bullet$ , and we are done.  $\square$

**Lemma 4.7.** *Take  $T$  and  $i_1, \dots, i_{N_T}$  as in Lemma 4.6. Then for  $1 \leq k \leq N_T$ ,*

$$(4.3) \quad \text{Infl}^q\left(\text{Defl}^q(T)_k, \text{Con}^q(T)\right) = T_{i_k}.$$

*That is, the two sides are identical functions on the same domain.*

*Proof.* We proceed again by induction on  $k$ . The base case  $k = 1$  is trivial, since if  $T \in \text{Inc}^q(\lambda)$  is any tableau, then  $\lambda(\text{Defl}^q(T)^{(1)})$  and  $\lambda(T^{(1)})$  are both the empty order ideal.

Now, assume that the lemma holds for all increasing tableaux  $T$  in the case  $k < t$ . We first analyze the left side of Equation (4.3). Take  $x \in \lambda(\text{Defl}^q(T)^{(t)})$ . If  $x \in \lambda(\text{Defl}^q(T)^{(t-1)})$ , then  $[\text{Defl}^q(T)^{(t-1)}](x) \leq t-1$  and hence

$$[\text{swap}_t(\text{Defl}^q(T)^{(t-1)})](x) = \text{Defl}^q(T)^{(t-1)}(x).$$

If instead

$$x \in \lambda(\text{Defl}^q(T)^{(t)}) \setminus \lambda(\text{Defl}^q(T)^{(t-1)}),$$

then either  $[\text{Defl}^q(T)^{(t-1)}](x) = \bullet$  or  $[\text{Defl}^q(T)^{(t-1)}](x) = t$ . In the former case, there must be a box  $x'$  adjacent to  $x$  such that  $[\text{Defl}^q(T)^{(t-1)}](x') = t$ . In the latter case, we must have that for all boxes  $x'$  adjacent to  $x$ ,  $[\text{Defl}^q(T)^{(t-1)}](x') \neq \bullet$ .

Now, we analyze the right side of Equation (4.3). Take  $x \in \lambda(T^{(i_t)})$ . If  $x \in \lambda(T^{(i_{t-1})})$ , then  $x \in \lambda(\text{Defl}^q(T)^{(t-1)}) \subseteq \lambda(\text{Defl}^q(T)^{(t)})$  by the inductive hypothesis. Further, this implies that  $T^{(i_{t-1})}(x) \leq i_{t-1}$ , and so we have

$$[\text{swap}_{\ell+1}(T^{(\ell)})](x) = T^{(\ell)}(x)$$

for all  $i_{t-1} \leq \ell$ . Therefore by the previous paragraph and the inductive hypothesis,

$$T^{(i_t)}(x) = T^{(i_{t-1})}(x) = \text{VecInfl}_{\text{Con}^q(T)}^q \circ \text{Defl}^q(T)^{(t-1)}(x) = \text{VecInfl}_{\text{Con}^q(T)}^q \circ \text{Defl}^q(T)^{(t)}(x).$$

This shows that Equation (4.3) holds at  $x$ . If  $x \in \lambda(T^{(i_t)}) \setminus \lambda(T^{(i_{t-1})})$ , then  $i_{t-1} < T^{(i_t)}(x) \leq i_t$ . Thus, we must have that  $T^{(i_t)}(x) = i_t$ , since the  $\text{swap}$  operators do not change the content of a tableau. Therefore, either  $T^{(i_{t-1})}(x) = \bullet$  or  $T^{(i_{t-1})}(x) = T(x) = i_t$ . In the former case, there must be a box  $x'$  adjacent to  $x$  such that

$T^{(i_{t-1})}(\mathbf{x}') = T(\mathbf{x}') = i_t$ . This occurs exactly when  $\text{Defl}^q(T)^{(t-1)}(\mathbf{x}') = \text{Defl}^q(T)(\mathbf{x}') = t$ . In the latter case, there is no  $\mathbf{x}'$  adjacent to  $\mathbf{x}$  such that  $T^{(i_{t-1})}(\mathbf{x}') = \bullet$ . Since we have assumed that  $\text{Con}^q(T)\{1\} = 1$ , Lemma 4.6 implies that this occurs exactly when  $\text{Defl}^q(T)^{(t-1)}(\mathbf{x}') \neq \bullet$  for every box  $\mathbf{x}'$  adjacent to  $\mathbf{x}$ . Therefore each of these conditions is equivalent to the corresponding condition from the previous paragraph.

Putting these facts together, we have shown that  $\lambda(T^{(i_k)}) = \lambda(\text{Defl}^q(T)^{(k)})$  for all  $k$ , and that moreover, if  $\text{Defl}^q(T)_k(\mathbf{x}) = h$ , then  $T_k(\mathbf{x}) = \text{VecInfl}_{\text{Con}^q(T)}^q(h)$ .  $\square$

*Remark 4.8.* As a corollary to Lemma 4.7, one can show that each  $T_k$  is, in fact, in  $\text{Inc}^q(\lambda(T^{(k)}))$ . However, we do not need this observation here.

**Lemma 4.9.** *Take  $T$  and  $i_1, \dots, i_{N_T}$  as in Lemma 4.6. Then*

$$(4.4) \quad \text{Defl}^q \circ \text{pro}^q(T) = \text{pro}^{N_T} \circ \text{Defl}^q(T).$$

*Proof.* By Equation (4.1), it is sufficient to show that

$$\begin{aligned} \text{Defl}^q \circ \text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet}(T) = \\ \text{Decr} \circ \text{Rep}_{\bullet \rightarrow N_T+1} \circ \text{swap}_{N_T} \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet} \circ \text{Defl}^q(T). \end{aligned}$$

Say  $\mathbf{x} \notin \lambda(T^{(q)})$ ; that is,  $T^{(q)}(\mathbf{x}) = \bullet$ . Then,

$$[\text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1(T)](\mathbf{x}) = q.$$

Now by [DPS17, Lemma 2.1], the content vector of

$$\text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1(T) = \text{pro}^q(T)$$

is a cyclic rotation by one of the content vector of  $T$ . This implies that there are the same number of distinct labels in the content vector of  $\text{pro}^q(T)$ , namely  $N_T$ , and, since  $q$  is included, it must be the greatest such label, so

$$[\text{Defl}^q \circ \text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1(T)](\mathbf{x}) = N_T.$$

On the other hand, since Lemma 4.6 implies that  $\mathbf{x} \notin \lambda(\text{Defl}^q(T)^{(N_T)})$ , we have

$$[\text{Decr} \circ \text{Rep}_{\bullet \rightarrow N_T+1} \circ \text{swap}_{N_T} \circ \dots \circ \text{swap}_1 \circ \text{Defl}(T)](\mathbf{x}) = N_T.$$

Now say  $\mathbf{x} \in \lambda(T^{(q)})$ . We observe that since  $\text{swap}_\ell$  does not affect the tableau  $\text{swap}_{N_T} \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet}(T)$  for  $\ell > i_{N_T}$ , we have that

$$(\text{swap}_q \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet}(T))_q = (\text{swap}_{i_{N_T}} \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet}(T))_{i_{N_T}}.$$

Therefore, it follows from Lemma 4.7, taking  $k = N_T$ , that

$$(\text{swap}_q \circ \dots \circ \text{swap}_1 \circ \text{Rep}_{1 \rightarrow \bullet}(T))_q = \text{VecInfl}_{\text{Con}(T)}^q \circ (\text{swap}_{N_T} \circ \dots \circ \text{Rep}_{1 \rightarrow \bullet} \circ \text{Defl}^q(T))_{N_T}.$$

Now, since  $\mathbf{x} \in \lambda(T^{(q)})$ , we have that  $T^{(q)}(\mathbf{x}) = \ell$  for  $\ell \in \{2, 3, \dots, q\}$ . This implies that  $\ell = i_k$  for some  $2 \leq k \leq N_T$ . Then,

$$\text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1(T)(\mathbf{x}) = i_k - 1.$$

Because the content vector shifts cyclically, this implies that

$$[\text{Defl}^q \circ \text{Decr} \circ \text{Rep}_{\bullet \rightarrow q+1} \circ \text{swap}_q \circ \dots \circ \text{swap}_1(T)](\mathbf{x}) = k - 1.$$

On the other hand,

$$[\text{swap}_q \circ \cdots \circ \text{swap}_1 \circ \text{Defl}(T)](x) = k,$$

so

$$[\text{Decr} \circ \text{Rep}_{\bullet \rightarrow N_T+1} \circ \text{swap}_{N_T} \circ \cdots \circ \text{swap}_1 \circ \text{Defl}(T)](x) = k - 1. \quad \square$$

**4.4. Computation of period.** We use the results of the previous subsection to relate the period of  $T \in \text{Inc}^q(\lambda)$  under  $\text{pro}^q$  to data concerning  $\text{Defl}^q(T)$ . Let

$$\Sigma^q : \{0, 1\}^q \rightarrow \{0, 1\}^q$$

be the cyclic rotation defined by

$$\Sigma^q(v_1, \dots, v_q) = (v_2, \dots, v_q, v_1)$$

for  $(v_1, \dots, v_q) \in \{0, 1\}^q$ .

**Theorem 4.10.** *Fix  $T \in \text{Inc}^q(\lambda)$ . Let  $\tau$  be the period of  $\text{pro}^{N_T}$  on  $\text{Defl}^q(T)$  and  $\ell$  be the period of  $\Sigma^q$  on  $\text{Con}^q(T)$ . If  $q - N_T$  (the number of 0's in  $\text{Con}^q(T)$ ) is  $r \cdot \frac{q}{\ell}$ , then the period of  $\text{pro}^q$  on  $T$  is*

$$\frac{\ell\tau}{\gcd(\ell - r, \tau)}.$$

*Proof.* Define

$$K^q : \coprod_{k \leq q} \text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[q]}{k} \rightarrow \coprod_{k \leq q} \text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[q]}{k}$$

by

$$K^q(S, v) = \begin{cases} (\text{pro}^{N_S}(S), \Sigma^q(v)), & \text{if } v\{1\} = 1; \\ (S, \Sigma^q(v)), & \text{otherwise.} \end{cases}$$

We show that, for  $T \in \text{Inc}^q(\lambda)$ ,

$$(4.5) \quad \text{DeflCon}^q \circ (\text{pro}^q)^{\circ n}(T) = (K^q)^{\circ n} \circ \text{DeflCon}^q(T).$$

Let

$$\pi_1 : \coprod_{k \leq q} \text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[q]}{k} \rightarrow \coprod_{k \leq q} \text{Inc}_{\text{gl}}^k(\lambda)$$

be the projection onto the first factor and define  $\pi_2$  similarly. It is immediate from the definitions that

$$\pi_2 \circ (K^q)^{\circ n} \circ \text{DeflCon}^q(T) = (\Sigma^q)^{\circ n} \circ \text{Con}^q(T)$$

for all  $n$ . But by [DPS17, Lemma 2.1],

$$(\Sigma^q)^{\circ n} \circ \text{Con}^q(T) = \text{Con}^q \circ (\text{pro}^q)^{\circ n}(T).$$

This shows that the two sides of Equation (4.5) are equal in the second factor.

To show that Equation (4.5) holds in the first factor, we proceed by induction on  $n$ . The base case  $n = 0$  is tautological. Assume that Equation (4.5) holds for  $n - 1$  and let

$$w := \text{Con}^q \circ (\text{pro}^q)^{\circ(n-1)}(T) = \pi_2 \circ (K^q)^{\circ(n-1)} \circ \text{DeflCon}^q(T).$$



By Theorem 4.3, we can write

$$(\mathbf{pro}^q)^{\circ(n-1)}(T) = \mathbf{VecInfl}_w^q \circ (K^q)^{\circ(n-1)} \circ \mathbf{DeflCon}^q(T).$$

If  $w\{1\} = 1$ , then

$$\begin{aligned} \pi_1 \circ \mathbf{DeflCon}^q \circ (\mathbf{pro}^q)^{\circ(n)}(T) &= \mathbf{Defl}^q \circ (\mathbf{pro}^q)^{\circ(n)}(T) \\ &= \mathbf{pro}^{N_T} \circ \mathbf{Defl}^q \circ (\mathbf{pro}^q)^{\circ(n-1)}(T) && \text{Lemma 4.9, } w\{1\} = 1 \\ &= \mathbf{pro}^{N_T} \circ \mathbf{Defl}^q \circ \mathbf{VecInfl}_w^q \circ (K^q)^{\circ(n-1)} \circ \mathbf{DeflCon}^q(T) \\ &= \mathbf{pro}^{N_T} \circ \pi_1 \circ (K^q)^{\circ(n-1)} \circ \mathbf{DeflCon}^q(T) && \text{by Theorem 4.3} \\ &= \pi_1 \circ (K^q)^{\circ(n)} \circ \mathbf{DeflCon}^q(T) && \text{because } w\{1\} = 1. \end{aligned}$$

On the other hand, if  $w\{0\} = 0$ , then

$$\begin{aligned} \pi_1 \circ \mathbf{DeflCon}^q \circ (\mathbf{pro}^q)^{\circ(n)}(T) &= \mathbf{Defl}^q \circ (\mathbf{pro}^q)^{\circ(n)}(T) \\ &= \mathbf{Defl}^q \circ (\mathbf{pro}^q)^{\circ(n-1)}(T) && \text{by Lemma 4.5, } w\{1\} = 0 \\ &= \pi_1 \circ (K^q)^{\circ(n-1)} \circ \mathbf{DeflCon}^q(T) \\ &= \pi_1 \circ (K^q)^{\circ(n)} \circ \mathbf{DeflCon}^q(T) && \text{because } w\{1\} = 0. \end{aligned}$$

This completes the proof of Equation 4.5.

Now, since  $\mathbf{DeflCon}^q$  is an injection,  $(\mathbf{pro}^q)^{\circ n}(T) = T$  exactly when  $\mathbf{DeflCon}^q \circ (\mathbf{pro}^q)^{\circ n}(T) = \mathbf{DeflCon}^q \circ T$ . So we are reduced to looking at what powers  $n$  of  $K^q$  stabilize  $\mathbf{DeflCon}^q(T)$ .

Say  $n$  satisfies this property. Projecting onto the second factor, we must have that

$$(\Sigma^q)^{\circ n}(\mathbf{Con}^q(T)) = \mathbf{Con}^q(T),$$

so  $n = t\ell$  for some integer  $t$ , where  $\ell$  is the period of  $\mathbf{Con}^q(T)$  under  $\Sigma^q$ . Projecting onto the first factor, we must have that

$$\mathbf{pro}^{n'}(\mathbf{Defl}^q(T)) = \mathbf{Defl}^q(T),$$

where  $n'$  is  $n$  minus the number of times that  $\mathbf{Con}^q \circ (K^q)^{\circ j} \circ \mathbf{DeflCon}(T)\{1\} = 0$  for  $j \in \{0, \dots, n-1\}$ . If there are  $r \cdot (q/\ell)$  zeros in  $\mathbf{Con}^q(T)$  and  $n = t\ell$ , then  $n' = t \cdot (\ell - r)$ . By our assumption on  $n$ , if  $\tau$  is the order of  $\mathbf{pro}^{N_T}$  on  $\mathbf{Defl}^q(T)$ , then  $s\tau = n' = t(\ell - r)$  for some  $s$ .

Therefore,  $(K^q)^{\circ n} \circ \mathbf{DeflCon}^q(T) = \mathbf{DeflCon}^q(T)$  if and only if  $n = t\ell = s\tau + tr$  for positive integers  $s$  and  $t$ . Hence, the order of  $\mathbf{pro}^q$  on  $T$  is equal to the least positive integer that can be simultaneously written in the forms  $t\ell$  and  $s\tau + tr$  for the same value of  $t$ . But  $t\ell = s\tau + tr$  implies that  $t(\ell - r) = s\tau$ . Since  $0 \leq r < \ell$ , the least such  $t\ell$  is clearly achieved when  $t$  is minimal, which occurs when  $t(\ell - r)$  is minimal, namely the least common multiple of  $\ell - r$  and  $\tau$ . Thus, the least such  $n$  can be expressed as

$$(4.6) \quad n = t\ell = \frac{\ell \operatorname{lcm}(\ell - r, \tau)}{\ell - r} = \frac{\ell\tau}{\gcd(\ell - r, \tau)},$$

as desired. □

## 5. THE EQUIVARIANT BIJECTION

**5.1. Bijection between plane partitions and increasing tableaux.** Let  $\{e_1, e_2\}$  be the standard basis for  $\mathbb{Z}^2$  and let  $\pi_i : \mathbb{Z}^2 \rightarrow \mathbb{Z}$  be the projection map onto the  $i$ th coordinate. We say that a subset  $\lambda \subset \mathbb{Z}^2$  is **connected** if for all  $x, x' \in \lambda$ , there exist  $r \in \mathbb{Z}_{\geq 0}$ ,  $d_1, \dots, d_r \in \{1, 2\}$ , and signs  $n_1, \dots, n_r \in \{\pm 1\}$  such that

$$\left\{ x + \sum_{k=1}^j n_k e_{d_k} : 1 \leq j \leq r \right\} \subseteq \lambda$$

and  $x + \sum_{k=1}^r n_k e_{d_k} = x'$ . Given these data, define  $x_0 = x$  and  $x_\ell = x + \sum_{k=1}^\ell n_k e_{d_k}$  for  $1 \leq \ell \leq r$ . Then the sequence  $(x_\ell)_{0 \leq \ell \leq r}$  is called a **path of length**  $r$ . If  $n_k = 1$  for all  $k$ , then the path is called a **chain**. We call a finite nonempty connected subset of  $\mathbb{Z}_{\geq 0}^2$  a **shape**. Shapes inherit a partial order from the natural componentwise partial order on  $\mathbb{Z}^2$ , and we will conflate a shape  $\lambda$  with the poset thereby induced on it. Note that the restriction of this partial order to a chain in  $\lambda$  is a chain poset, justifying our double use of terminology. A shape poset  $\lambda$  is ranked by restricting the standard rank function  $\text{rk}(x_1, x_2) = x_1 + x_2$  on  $\mathbb{Z}^2$  to the subset  $\lambda$ . Each minuscule poset  $P$  can be realized as a shape by considering its diagram from Figure 1 or 3 and then embedding the diagram in  $\mathbb{Z}^2$  in the obvious way, so that the bottom-leftmost box sits at the origin of  $\mathbb{Z}^2$  and the cover relations in  $P$  correspond to cover relations in  $\mathbb{Z}^2$ . When we consider a minuscule poset  $P$  as a shape, it will always be with respect to this particular embedding in  $\mathbb{Z}^2$ .

We say that the shape  $\lambda$  is **convex** if  $x \in \lambda$  and  $x + k e_d \in \lambda$  for some  $d \in \{1, 2\}$  and  $k \in \mathbb{Z}_{>0}$  together imply that  $\{x + j e_d : 1 \leq j \leq k - 1\} \subseteq \lambda$ .

For any shape  $\lambda$ , let

$$a_i(\lambda) = \max\{\pi_i(x) : x \in \lambda\}$$

and

$$a_\lambda = (a_1(\lambda), a_2(\lambda)) \in \mathbb{Z}^2.$$

We say the shape  $\lambda$  is **justified** if  $(0, 0) \in \lambda$  and  $a_\lambda \in \lambda$ . Note that the minuscule posets are convex and justified, when considered as shapes as above. For any shape  $\lambda$ , define

$$\lambda^* := \{a_\lambda - x : x \in \lambda\}.$$

It is clear that  $\lambda^*$  is a shape and that if  $\lambda$  is convex (resp. justified), then  $\lambda^*$  is convex (resp. justified). Define the **rank** of  $\lambda$  to be

$$\text{rk}(\lambda) := \max_{x \in \lambda} \text{rk}(x).$$

The following elementary lemmas determine useful characteristics of the shapes considered in this paper.

**Lemma 5.1.** *Let  $\lambda$  be a convex shape and let  $x, x' \in \lambda$  be distinct. If  $x < x'$ , then  $\lambda$  contains a chain from  $x$  to  $x'$ .*

*Proof.* Let  $\mathbf{x} = (i, j)$  and  $\mathbf{x}' = (i', j')$ . Because  $\mathbf{x} < \mathbf{x}'$ , we have  $i \leq i'$  and  $j \leq j'$ . Since  $\lambda$  is a shape, there is at least one path from  $\mathbf{x}$  to  $\mathbf{x}'$ . Take such a path  $\mathcal{P} = (\mathbf{x}_\ell)_{0 \leq \ell \leq r}$  of minimal length, characterized by directions  $d_1, \dots, d_r \in \{1, 2\}$  and signs  $n_1, \dots, n_r \in \{\pm 1\}$ . We claim that  $\mathcal{P}$  is a chain.

If not, then  $n_t = -1$  for some  $1 \leq t \leq r$ . Let  $t$  be the least such index. For  $0 \leq \ell \leq r$ , define  $(i_\ell, j_\ell) = \mathbf{x}_\ell$ , so that in particular  $(i_0, j_0) = (i, j)$  and  $(i_r, j_r) = (i', j')$ .

Since  $n_t = -1$ , either  $i_t = i_{t-1} - 1$  or  $j_t = j_{t-1} - 1$ . Without loss of generality, we may assume  $i_t = i_{t-1} - 1$ .

Suppose there is no  $h < t$  with  $d_h = d_t$  and  $n_h = 1$ . Then  $i_t = i_0 - 1$ . Furthermore, since  $i' = i_r \geq i_0$ , there must be a least  $k > t$  such that  $i_k = i_0 = i_{t-1}$ . Let  $q = j_k - j_{t-1}$ . Then clearly the subpath  $(\mathbf{x}_\ell)_{t-1 \leq \ell \leq k}$  of  $\mathcal{P}$  has length at least  $|q| + 2$ . However by convexity, the path

$$\left( (i_{t-1}, j_{t-1}), (i_{t-1}, j_{t-1} + \frac{q}{|q|}), (i_{t-1}, j_{t-1} + 2\frac{q}{|q|}), \dots, (i_{t-1}, j_{t-1} + |q|\frac{q}{|q|}) \right)$$

of length  $|q|$  from  $(i_{t-1}, j_{t-1})$  to  $(i_k, j_k)$  lies entirely inside  $\lambda$ , contradicting the minimality of  $\mathcal{P}$ .

Otherwise, consider the greatest  $h < t$  with  $d_h = d_t$  and  $n_h = 1$ . Then  $i_{h-1} = i_t$ . By minimality of  $t$ , we have  $j_t \geq j_{h-1}$ . Let  $q = j_t - j_{h-1}$ . Now the subpath  $(\mathbf{x}_\ell)_{h-1 \leq \ell \leq t}$  of  $\mathcal{P}$  has length  $q + 2$ . However by convexity, the path

$$((i_{h-1}, j_{h-1}), (i_{h-1}, j_{h-1} + 1), (i_{h-1}, j_{h-1} + 2), \dots, (i_{h-1}, j_{h-1} + q))$$

of length  $q$  from  $(i_{h-1}, j_{h-1})$  to  $(i_t, j_t)$  lies entirely inside  $\lambda$ , contradicting the minimality of  $\mathcal{P}$ .  $\square$

**Lemma 5.2.** *Let  $\lambda$  be a convex justified shape. Then every maximal chain has length  $a_1(\lambda) + a_2(\lambda)$ .*

*Proof.* Since for all  $\mathbf{x} \in \lambda$  we have  $(0, 0) \leq \mathbf{x}$  and  $\mathbf{x} \leq (a_1(\lambda), a_2(\lambda))$ , all maximal chains must start at  $(0, 0)$  and end at  $(a_1(\lambda), a_2(\lambda))$  by Lemma 5.1. But clearly any chain between these two elements has the desired length.  $\square$

For a convex shape  $\lambda$ , an increasing tableau on the poset  $\lambda$  is equivalent to a map  $T : \lambda \rightarrow \mathbb{Z}_{>0}$  such that whenever  $\mathbf{x} \in \lambda$  and  $\mathbf{x} + \mathbf{e}_i \in \lambda$  for  $i \in \{1, 2\}$ , then  $T(\mathbf{x}) < T(\mathbf{x} + \mathbf{e}_i)$ . The **height** of an increasing tableau  $T$  is the maximum value in its range. Recall that  $\text{Inc}^q(\lambda)$  is the set of all increasing tableaux of height at most  $q$  and base  $\lambda$ .

Fix a convex justified shape  $\lambda$  and  $m \in \mathbb{Z}_{>0}$ . Now, we define a poset

$$\lambda \times \mathbf{m} := \{(\mathbf{x}, y) \in \mathbb{Z}^3 : \mathbf{x} \in \lambda, 0 \leq y \leq m - 1\},$$

with the partial order given by restricting the natural partial order on  $\mathbb{Z}^3$ , that is

$$\mathbf{p} \geq \mathbf{q} \text{ if and only if } \mathbf{p} - \mathbf{q} \in \mathbb{Z}_{\geq 0}^3.$$

Note that, as a abstract poset,  $\lambda \times \mathbf{m}$  is, in fact, isomorphic to the product of the posets  $\lambda$  and  $\mathbf{m}$ ; it is useful to us, however, to consider  $\lambda \times \mathbf{m}$  as embedded inside  $\mathbb{Z}^3$  in this particular fashion.

Recall that for any poset  $P$ , we write  $J(P)$  for the collection of all order ideals of  $P$ . Define the map  $\Omega : J(\lambda \times \mathbf{m}) \rightarrow \text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$  as follows. First, define the function

$$\text{ht} : \mathbb{Z}_{\geq 0}^2 \times 2^{(\mathbb{Z}_{\geq 0}^3)} \rightarrow \mathbb{Z}_{\geq 0}$$

(where  $2^{(\mathbb{Z}_{\geq 0}^3)}$  denotes the power set of  $\mathbb{Z}_{\geq 0}^3$ ) by

$$\text{ht}(\mathbf{x}, S) := \#\{y \in \mathbb{Z}_{\geq 0} : (\mathbf{x}, y) \in S\}.$$

Intuitively, if an order ideal  $I$  of  $\lambda \times \mathbf{m}$  is viewed as a pile of blocks stacked on  $\lambda \subset \mathbb{Z}^2$ , then  $\text{ht}(\mathbf{x}, I)$  counts the number of blocks stacked in the column over  $\mathbf{x} \in \lambda$ . In this notation,  $J(\lambda \times \mathbf{m})$  is exactly the set of all plane partitions with  $\text{ht}$  at most  $m$  over the shape poset  $\lambda$ . Next, define  $\Omega : J(\lambda \times \mathbf{m}) \rightarrow \text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$  such that

$$[\Omega(I)](\mathbf{x}) = \text{ht}(\mathbf{a}_\lambda - \mathbf{x}, I) + \text{rk}(\mathbf{x}) + 1.$$

**Proposition 5.3.** *If  $\lambda$  is a convex justified shape, then  $\Omega$  is a bijection.*

*Proof.* Since  $\lambda$  is justified and convex,  $\lambda^*$  has an element of rank 0 and, by Lemma 5.2, all maximal chains in  $\lambda^*$  have equal length. Hence the function  $\Omega$  coincides with the map from [HPPW16, Proposition 8.2], and so by [HPPW16, Proposition 8.2] we have that  $\Omega$  is a bijection between plane partitions over  $\lambda$  and increasing tableaux over  $\lambda^*$ .  $\square$

Note, in particular, that Proposition 5.3 applies to all of the minuscule posets, realized as convex justified shapes as in Figures 1 and 3.

**5.2. Equivariance of  $\Omega$ .** In this section we prove Proposition 5.5, our key to proving Theorems 1.2 and 1.3. Proposition 5.5 is a mild extension of [DPS17, Lemma 4.2] and our proof follows the same outline. We, however, still give significant details of the proof, as there are a few extra subtleties in our case.

The following gives an easy and useful way to determine whether a subset of  $\lambda \times \mathbf{m}$  is an order ideal.

**Lemma 5.4.** *Let  $\lambda$  be any shape. The subset  $Q \subseteq \lambda \times \mathbf{m}$  is an order ideal if and only if both*

- (a)  $(i, j, k) \in Q$  implies that  $(i, j, k') \in Q$ , for  $0 \leq k' \leq k$ ; and
- (b)  $(i, j) \geq (i', j')$  implies that  $\text{ht}((i, j), Q) \leq \text{ht}((i', j'), Q)$ .

*Proof.* The “only if” direction is clear.

Conversely, assume  $Q$  satisfies (a) and (b). Consider  $(i, j, k) \in Q$  and  $(i', j', k') \in \lambda \times \mathbf{m}$  with  $(i', j', k') \leq (i, j, k)$ . We must show  $(i', j', k') \in Q$ .

By assumption,  $(i', j') \leq (i, j)$ . Hence, by condition (b),  $\text{ht}((i, j), Q) \leq \text{ht}((i', j'), Q)$ . Now, condition (a) implies that, for all  $\mathbf{x} \in \lambda$ , we have

$$\text{ht}(\mathbf{x}, Q) = 1 + \max\{k'' \in \mathbb{Z}_{\geq 0} : (\mathbf{x}, k'') \in Q\}.$$

Applying this equality for  $\mathbf{x} = (i, j)$  and  $\mathbf{x} = (i', j')$ , we conclude there is a  $\hat{k} \geq k$  with  $(i', j', \hat{k}) \in Q$ . But since  $k' \leq k \leq \hat{k}$ , condition (a) then implies  $(i', j', k') \in Q$ .  $\square$

**Proposition 5.5.** *Let  $\lambda$  be a convex justified shape. Then  $\Omega$  intertwines  $\text{Mot}_{\text{id},(1,1,-1)}$  and  $\text{pro}^{\text{rk}(\lambda)+m+1}$ . That is, for  $I \in J(\lambda \times \mathbf{m})$ , we have*

$$\Omega \circ \text{Mot}_{\text{id},(1,1,-1)}(I) = \text{pro}^{\text{rk}(\lambda)+m+1} \circ \Omega(I).$$

*Proof.* Let  $a := a_1(\lambda)$  and let  $b := a_2(\lambda)$ ; that is, let  $(a, b) := \mathbf{a}_\lambda$ . Since  $\lambda$  is justified,  $a + b = \text{rk}(\lambda)$ . Fix  $I \in J(\lambda \times \mathbf{m})$ .

Recall that for  $T \in \text{Inc}^{a+b+m+1}(\lambda)$ , we have by definition that

$$\text{pro}^{a+b+m+1}(T) = \rho_{a+b+m} \circ \cdots \circ \rho_1(T).$$

Since  $-(m-1) \leq \langle (1, 1, -1), (x, y, z) \rangle \leq a + b$  for  $(x, y, z) \in \lambda \times \mathbf{m}$ , we have

$$\text{Mot}_{\text{id},(1,1,-1)}(I) = \tau_{\text{id},(1,1,-1)}^{a+b-(a+b+m-1)} \circ \cdots \circ \tau_{\text{id},(1,1,-1)}^{a+b}(I).$$

Therefore, it suffices to show

$$\Omega \left( \tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}(I) \right) = \rho_\ell(\Omega(I))$$

for all  $\ell \in \{1, 2, \dots, a + b + m\}$ .

Fix such an index  $\ell$  and a box  $\mathbf{y} = (i, j) \in \lambda$ . Let  $Y := I \cap (\mathbf{y} \times \mathbf{m})$  be the column of  $\mathbf{y}$  in  $I$ . We consider the effect on  $Y$  of applying  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  to the order ideal  $I \in J(\lambda \times \mathbf{m})$  and the effect on the box  $\mathbf{a}_\lambda - \mathbf{y}$  of applying  $\rho_\ell$  to the increasing tableau  $\Omega(I) \in \text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$ . Considering the definition of  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  from Equation (3.1), note that  $t_{(i,j,i+j-(a+b-(\ell-1)))}$  is the unique toggle in that definition that acts on an element of  $Y$ .

Define the function  $H : \lambda \rightarrow [m+1]$  by  $H(\mathbf{x}) = \text{ht}(\mathbf{x}, I)$  for  $\mathbf{x} \in \lambda$ . By Lemma 5.4, observe that the function  $H$  must be nonincreasing along any chain in  $\lambda$ . There are now four cases to consider depending on the value of  $H(\mathbf{y})$ .

**Case 1:**  $H(\mathbf{y}) > i + j - (a + b - (\ell - 1)) + 1$ :

Here,  $(\mathbf{y}, i + j - (a + b - (\ell - 1))) \in Y$ , so  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  either removes this element from  $Y$  or leaves  $Y$  unchanged. Since, by the case assumption,  $(\mathbf{y}, i + j - (a + b - (\ell - 1)) + 1) \in Y$ , removing  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  from  $I$  does not yield an order ideal. Therefore, the column  $Y$  is unchanged by  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$ . On the other hand, the case assumption implies that  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y}) > \ell + 1$ , so  $\rho_\ell$  does not affect this entry of the increasing tableau  $\Omega(I)$ .

**Case 2:**  $H(\mathbf{y}) = i + j - (a + b - (\ell - 1)) + 1$ :

Here,  $(\mathbf{y}, i + j - (a + b - (\ell - 1))) \in Y$ , while  $(\mathbf{y}, i + j - (a + b - (\ell - 1)) + 1) \notin Y$ . We claim that  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  removes  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  from  $I$  exactly if  $H(\mathbf{y} + \mathbf{e}_d) \leq i + j - (a + b - (\ell - 1))$  for all  $d \in \{1, 2\}$  with  $\mathbf{y} + \mathbf{e}_d \in \lambda$ .

For, on the one hand, if  $\mathbf{y} + \mathbf{e}_d \in \lambda$  and  $H(\mathbf{y} + \mathbf{e}_d) > i + j - (a + b - (\ell - 1))$ , then by Lemma 5.4, removing  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  from  $I$  would not leave an order ideal. On the other hand, say that  $H(\mathbf{y} + \mathbf{e}_d) \leq i + j - (a + b - (\ell - 1))$  for all such  $d$ . To verify that removing  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  from  $I$  results in an order ideal, consider any fixed  $\mathbf{y}' = (i', j') \in \lambda$  with  $\mathbf{y}' > \mathbf{y}$ . By Lemma 5.4, it is enough to check

that  $H(\mathbf{y}') \leq i + j - (a + b - (\ell - 1))$ . Now, by Lemma 5.1, since  $\mathbf{y}' > \mathbf{y}$ , there is a chain from  $\mathbf{y}$  to  $\mathbf{y}'$ . Clearly, this chain contains an element of the form  $\mathbf{y} + \mathbf{e}_d \in \lambda$ . But the function  $H$  is nonincreasing along this chain, so  $H(\mathbf{y}') \leq i + j - (a + b - (\ell - 1))$ , as desired.

On the tableaux side, the case assumption implies that  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y}) = \ell + 1$ . The  $\mathbf{a}_\lambda - \mathbf{y}$  entry of  $\Omega(I)$  is decreased to  $\ell$  by  $\rho_\ell$  exactly when, for all  $d$  with  $\mathbf{a}_\lambda - \mathbf{y} - \mathbf{e}_d \in \lambda^*$ ,  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y} - \mathbf{e}_d) \leq \ell - 1$ . This is easily equivalent to the stated condition on when  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  acts nontrivially on  $Y$ .

**Case 3:**  $H(\mathbf{y}) = i + j - (a + b - (\ell - 1))$ :

Here,  $(\mathbf{y}, i + j - (a + b - (\ell - 1)) - 1) \in Y$  but  $(\mathbf{y}, i + j - (a + b - (\ell - 1))) \notin Y$ . We claim that  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  adds  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  to  $I$  exactly when  $H(\mathbf{y} - \mathbf{e}_d) \geq i + j - (a + b - (\ell - 1)) + 1$  for all  $d \in \{1, 2\}$  with  $\mathbf{y} - \mathbf{e}_d \in \lambda$ .

For, on the one hand, if  $\mathbf{y} - \mathbf{e}_d \in \lambda$  and  $H(\mathbf{y} - \mathbf{e}_d) < i + j - (a + b - (\ell - 1)) + 1$ , then by Lemma 5.4, adding  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  to  $I$  does not yield an order ideal. On the other hand, say that  $H(\mathbf{y} - \mathbf{e}_d) \geq i + j - (a + b - (\ell - 1)) + 1$  for all such  $d$ . To verify that adding  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  to  $I$  results in an order ideal, consider any fixed  $\mathbf{y}' = (i', j') \in \lambda$  with  $\mathbf{y}' < \mathbf{y}$ . By Lemma 5.4, it is enough to check that  $H(\mathbf{y}') \geq i + j - (a + b - (\ell - 1)) + 1$ . Now, by Lemma 5.1, since  $\mathbf{y}' < \mathbf{y}$ , there is a chain from  $\mathbf{y}'$  to  $\mathbf{y}$ . Clearly, this chain contains an element of the form  $\mathbf{y} - \mathbf{e}_d \in \lambda$ . But the function  $H$  is nonincreasing along this chain, so  $H(\mathbf{y}') \geq i + j - (a + b - (\ell - 1)) + 1$  as desired.

On the tableaux side, this case assumption implies that  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y}) = \ell$ . The  $\mathbf{a}_\lambda - \mathbf{y}$  entry of  $\Omega(I)$  is increased to  $\ell + 1$  by  $\rho_\ell$  exactly when, for all  $d$  with  $\mathbf{a}_\lambda - \mathbf{y} + \mathbf{e}_d \in \lambda^*$ ,  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y} + \mathbf{e}_d) \geq \ell + 2$ . This is easily equivalent to the stated condition on when  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  acts nontrivially on  $Y$ .

**Case 4:**  $H(\mathbf{y}) < i + j - (a + b - (\ell - 1))$ :

Here,  $(\mathbf{y}, i + j - (a + b - (\ell - 1))) \notin Y$ , so  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$  either adds this element to  $Y$  or else leaves  $Y$  unchanged. Since, by the case assumption,  $(\mathbf{y}, i + j - (a + b - (\ell - 1)) - 1) \notin I$ , by Lemma 5.4, adding  $(\mathbf{y}, i + j - (a + b - (\ell - 1)))$  to  $I$  does not yield an order ideal. Therefore, the column  $Y$  is unchanged by  $\tau_{\text{id},(1,1,-1)}^{a+b-(\ell-1)}$ . On the other hand, the case assumption implies that  $[\Omega(I)](\mathbf{a}_\lambda - \mathbf{y}) < \ell$ , so  $\rho_\ell$  does not affect this entry of  $\Omega(I)$ .

This completes the proof of Proposition 5.5 □

**Corollary 5.6.** *Let  $\lambda$  be a convex justified shape. Then the multiset of cardinalities of  $\Psi$ -orbits on  $J(\lambda \times \mathbf{m})$  and the multiset of cardinalities of  $\mathbf{pro}^{\text{rk}(\lambda)+m+1}$ -orbits on  $\text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$  are equal.*

*Proof.* By Proposition 5.3,  $\Omega$  is a bijection between  $J(\lambda \times \mathbf{m})$  and  $\text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$ . By Proposition 5.5, the orbit structure of  $\mathbf{pro}^{\text{rk}(\lambda)+m+1}$  on  $\text{Inc}^{\text{rk}(\lambda)+m+1}(\lambda^*)$  is the same as the orbit structure of  $\mathbf{Mot}_{\text{id},(1,1,-1)}$  on  $J(\lambda \times \mathbf{m})$ . By Proposition 3.1, the action of  $\Psi$  on  $J(\lambda \times \mathbf{m})$  is the same as the action of  $\mathbf{Mot}_{\text{id},(1,1,1)}$ . By Proposition 3.2,  $\mathbf{Mot}_{\text{id},(1,1,1)}$  and  $\mathbf{Mot}_{\text{id},(1,1,-1)}$  induce the same multiset of orbit cardinalities on  $J(\lambda \times \mathbf{m})$ . □

## 6. PROOF OF THEOREMS 1.2 AND 1.3

In this final section, we collect the above results into proofs of Theorems 1.2 and 1.3. Consider a poset  $P$  that is either the Cayley-Moufang poset, the Freudenthal poset, or one of the propellers, and realize  $P$  as a convex justified shape  $\lambda$ . By Corollary 5.6, the multiset of  $\Psi$ -orbit cardinalities for  $J(\lambda \times \mathbf{m})$  equals the multiset of  $\mathbf{pro}^{\mathrm{rk}(\lambda)+m+1}$ -orbit cardinalities for  $\mathrm{Inc}^{\mathrm{rk}(\lambda)+m+1}(\lambda^*)$ . Therefore, we may prove Theorems 1.2 and 1.3 by studying increasing tableaux instead of plane partitions. Denote the Cayley-Moufang shape by  $\lambda_{CM}$ , the Freudenthal shape by  $\lambda_F$ , and the propeller shape with  $2k$  boxes ( $k \geq 3$ ) by  $\lambda_k$ . Note that for each of these shapes  $\lambda$ , we have  $\lambda = \lambda^*$ .

By Theorem 4.3, the  $\mathbf{pro}$ -orbit structure of increasing tableaux is controlled by the data of gapless increasing tableaux and binary vectors. But for any fixed  $\lambda$ ,  $\mathrm{Inc}_{\mathrm{gl}}(\lambda)$  is a finite set. Using a computer, we found all 549 gapless increasing tableaux of shape  $\lambda_{CM}$ , as well as all 624 493 gapless increasing tableaux of shape  $\lambda_F$ . For each such tableau  $T \in \mathrm{Inc}_{\mathrm{gl}}^q(\lambda_{CM})$  or  $\mathrm{Inc}_{\mathrm{gl}}^q(\lambda_F)$ , we then determined the size of its orbit under  $\mathbf{pro}^q$  by direct calculation. These data are given in Table 1 for  $\lambda_{CM}$  and in Table 2 for  $\lambda_F$ .

$q$	orbit size	number of orbits
11	1	1
12	3	1
	12	1
13	13	6
14	7	2
	14	12
15	15	13
16	2	1
	4	1
	8	1
	16	4

TABLE 1. The distribution of  $\mathbf{pro}^q$ -orbits of gapless increasing tableaux in  $\mathrm{Inc}_{\mathrm{gl}}^q(\lambda_{CM})$  for each  $q$ .

It is essentially trivial to observe that  $\mathrm{Inc}_{\mathrm{gl}}(\lambda_k)$  consists of exactly three increasing tableaux for any  $k$ . There are two tableaux in  $\mathrm{Inc}_{\mathrm{gl}}^{2k}(\lambda_k)$ , forming a single  $\mathbf{pro}^{2k}$ -orbit, and a single tableau in  $\mathrm{Inc}_{\mathrm{gl}}^{2k-1}(\lambda_k)$ , which is necessarily fixed by  $\mathbf{pro}^{2k-1}$ . These three tableaux and their orbits are illustrated in Figure 8 in the case  $k = 4$  (which is, in fact, completely generic). Table 3 records the observations of this paragraph in form analogous to Tables 1 and 2.

In addition, the number of elements of  $\binom{[M]}{k}$  whose order under  $\Sigma^M$  divides a given divisor of  $M$  is predicted by NAME THIS THING (CITE THIS FACT). Therefore, we can use Tables 1, 2, and 3 to completely determine the multiset of  $\mathbf{pro}^M$ -orbit cardinalities on any set  $\mathrm{Inc}^M(\lambda)$  for  $\lambda \in \{\lambda_{CM}, \lambda_F, \lambda_k\}$ .

$q$	orbit size	number of orbits
17	1	1
18	2	1
	18	2
19	19	30
20	20	228
21	7	3
	21	1044
22	22	3053
	66	2
23	23	5813
	69	13
24	8	7
	24	7195
	48	4
	72	26
25	25	5602
	50	8
	75	21
26	2	2
	26	2495
	52	4
	78	6
27	3	2
	9	4
	27	484

TABLE 2. The distribution of  $\mathbf{pro}^q$ -orbits of gapless increasing tableaux in  $\text{Inc}_{\text{gl}}^q(\lambda_F)$  for each  $q$ .

$q$	orbit size	number of orbits
$2k - 1$	1	1
$2k$	2	1

TABLE 3. The distribution of  $\mathbf{pro}^q$ -orbits of gapless increasing tableaux in  $\text{Inc}_{\text{gl}}^q(\lambda_k)$  for each  $q$ .

**Algorithm 6.1.**

**Input:**  $M, L \in \mathbb{Z}_{>0}$  such that  $L$  divides  $M$ , convex justified shape  $\lambda \in \{\lambda_{CM}, \lambda_F, \lambda_k\}$ , and the corresponding Table 1, 2, or 3.

**Output:**  $x = \#\{T \in \text{Inc}^M(\lambda) : (\mathbf{pro}^M)^{\circ L}(T) = T\}$ , the number of tableaux whose period divides  $L$ .






FIGURE 8. The set  $\text{Inc}_{\text{gl}}(\lambda_4)$  consists of the three illustrated gapless increasing tableaux. The unique element of  $\text{Inc}_{\text{gl}}^7(\lambda_4)$  forms a singleton  $\text{pro}^7$ -orbit, while  $\text{pro}^8$  switches the two elements of  $\text{Inc}_{\text{gl}}^8(\lambda_4)$ , as shown. The situation for  $k \neq 4$  is exactly analogous.

**Initialization:** Set  $x = 0$ .

**Procedure:** Iterate over rows of the table. Say the current row is  $(q, \tau, N)$ , expressing that  $\text{Inc}_{\text{gl}}^q(\lambda)$  has  $N$   $\text{pro}^q$ -orbits of size  $\tau$ .

- For each row, iterate over the positive divisors  $d$  of  $L$ .

(1) Compute the number

$$(6.1) \quad i := \# \left\{ v \in \binom{[M]}{q} : \text{the } \Sigma^m\text{-orbit of } v \text{ has size exactly } L/d \right\}.$$

(2) Compute

$$(6.2) \quad p := \frac{L/d \cdot \tau}{\gcd(qL/(Md), \tau)},$$

the period associated with a tableaux of period  $\tau$ , a content vector of period  $L/d$  with  $q$  ones.

(3) If  $p$  divides  $L$ , add  $i\tau N$  to  $x$ .

#### CHECK AGAINST ALGORITHM

**Theorem 6.2.** *The output of Algorithm 6.1 is equal to the number of order ideals in  $J(\lambda \times \mathbf{m})$  whose order under  $\Psi$  divides  $L$ , where  $m = M - \text{rk}(\lambda) - 1$ .*

*Proof.* By Corollary 5.6, it suffices to show that Algorithm 1 outputs the number of elements of  $\text{Inc}^M(\lambda)$  whose period under  $\text{pro}^M$  divides  $L$ , where  $M = \text{rk}(\lambda) + m + 1$ . By Theorem 4.3, an element  $T \in \text{Inc}^M(\lambda)$  corresponds to a unique pair  $(S, v) \in \coprod_{k \leq M} (\text{Inc}_{\text{gl}}^k(\lambda) \times \binom{[M]}{k})$ . Therefore, we may instead count pairs  $(S, v)$  such that the period of the tableaux associated to  $(S, v)$  divides  $L$ . Now, select a row  $(q, \tau, N)$

from Table 1, Table 2, or Table 3. These data correspond uniquely to  $\tau N$  full-content tableaux in  $\text{Inc}_{\text{gl}}^q(\lambda)$ . Conversely, by Theorem 4.3, each full-content tableaux  $S \in \coprod_{k \leq M} \text{Inc}_{\text{gl}}^k(\lambda)$  must come from a row in this table.

If  $i$  is given by Equation 6.1, then this row gives  $i\tau N$  tableaux. In the notation of Theorem 4.10,  $\ell = L/d$ ,  $\tau = \tau$ , and

$$\ell - r = L/d - \frac{L(M - q)}{dM} = \frac{Lq}{dM}.$$

Therefore the period associated to each of these tableaux is given by Equation 6.2.  $\square$

We have assumed in Algorithm 6.1 that we can execute Step 1. This is performed by the following algorithm.

### Algorithm 2

**Input:** Integer  $L$ , decreasing list  $W$  of all divisors of both  $L$  and  $f$ .

**Output:** Array  $V$ , where  $V[\ell]$  is the number of vectors of length  $T$  with  $f$  ones and with period exactly  $T/d_\ell$

**Initialization:** Let  $k = \text{length}(W)$ . Create a vector  $V$  of  $k$  zeros.

**Procedure:** Loop over  $\ell = 1, \dots, k$ .

$\Rightarrow$  Let  $d_\ell = W[\ell]$ .

$\Rightarrow V[\ell] +$  = the number of content vectors of length  $L$  and  $f$  ones whose period divides  $\ell$ .

$\Rightarrow$  For  $m = 1, \dots, \ell - 1$ , if  $d_m$  divides  $d_\ell$  then  $V[\ell] + = -V[m]$ .

**Theorem 6.3.** *Algorithm ?? computes the number of vectors of length  $T$  with  $f$  ones and period exactly  $T/d_\ell$  CHANGE THESE VARIABLES.*

*Proof.* Only a subset of divisors of  $L$  will yield a nonzero number of content vectors. Since the number of ones in such a content vector is equal to  $q$ , the full-content height of the tableau, if the content vector divides into  $d$  identical segments, then  $L - q$  must be divisible by  $d$ . Therefore,  $d$  must be a divisor of both  $L$  and  $q$ . While  $L$  is arbitrary,  $q$  takes only finitely many values.

Now for a given  $d$ , the known generating function for straight tableaux [RSW04, Theorem 1.1(b)] tells us the number of content vectors of length  $L$  with  $q$  ones whose period is a multiple of  $L/d$ . Since this number will be zero for all but finitely many integers  $d$ , we can then recover the number of tableaux with period exactly  $L/d$  by subtracting off those whose periods strictly divide  $L/d$ .  $\square$

To summarize, we outline the code in `symbolic_Cayley-Moufang_plane.py`. We fix a tableau  $\lambda$  and define a symbolic integer variable  $\mathbf{m}$ . The predicted number of tableaux in  $\text{Inc}^q(\lambda)$  will be a function of  $\mathbf{m}$  which is interpreted as  $m = q - (\text{rk}(\lambda) + 1)$ . Let  $B(\lambda)$  be the number of boxes in  $\lambda$ .

**Algorithm 3:** `symbolic_Cayley-Moufang_plane.py`

**Input:** Shape  $\lambda$ , full content tableaux for  $\lambda$ .

**Output:** Array  $V$ , where  $V[\ell]$  is the number of vectors of length  $T$  with  $f$  ones and with period exactly  $T/d_\ell$

**Initialization:** Let  $\mathfrak{m}$  be a symbolic variable.

**Procedure:** Loop over all combinations  $C$  of prime numbers (with multiplicity) that can divide a number between  $\text{rk}(\lambda) + 1$  to  $B(\lambda)$  inclusive.

- (1) Initialize empty array **dVals**. For  $d \in [2, B(\lambda)]$ , determine whether  $d$  divides a  $q$  value with the combination  $C$  of divisors. If so, append to **dVals**.
- (2) Loop over the elements of **dVals** and compute the predicted period by the Rush-Shi formula. Save as **RushShiPeriods**[ $L/d$ ].
- (3) Loop over full content orbit data.
  - (a) For  $d \in \mathbf{dVals}$ , if  $d$  divides **fcHt**
    - (i) Get exact number of vectors of length  $q$  with **fcHt** ones.
    - (ii) Compute period
    - (iii) number of tableaux to all periods divided by this one.
- (4) Check if Rush-Shi and computed match.

#### ACKNOWLEDGEMENTS

The authors thank Julianna Tymoczko for introducing them to each other. OP was partially supported by a Mathematical Sciences Postdoctoral Research Fellowship from the National Science Foundation. HM was supported by the Graduate Research Fellowship from the National Science Foundation.

#### REFERENCES

- [AST13] Drew Armstrong, Christian Stump, and Hugh Thomas, *A uniform bijection between nonnesting and noncrossing partitions*, Trans. Amer. Math. Soc. **365** (2013), no. 8, 4121–4151. 1
- [BL00] Sara Billey and V. Lakshmibai, *Singular loci of Schubert varieties*, Progress in Mathematics, vol. 182, Birkhäuser Boston, Inc., Boston, MA, 2000. 5
- [BPS16] Jonathan Bloom, Oliver Pechenik, and Dan Saracino, *Proofs and generalizations of a homomesy conjecture of Propp and Roby*, Discrete Math. **339** (2016), no. 1, 194–206. 8
- [BR12] Anders Skovsted Buch and Vijay Ravikumar, *Pieri rules for the  $K$ -theory of cominuscule Grassmannians*, J. Reine Angew. Math. **668** (2012), 109–132. 7
- [BS74] A. E. Brouwer and A. Schrijver, *On the period of an operator, defined on antichains*, Mathematisch Centrum, Amsterdam, 1974, Mathematisch Centrum Afdeling Zuivere Wiskunde ZW 24/74. 5
- [BS16] Anders Skovsted Buch and Matthew J. Samuel,  *$K$ -theory of minuscule varieties*, J. Reine Angew. Math. **719** (2016), 133–171. 7
- [Buc02] Anders Skovsted Buch, *A Littlewood-Richardson rule for the  $K$ -theory of Grassmannians*, Acta Math. **189** (2002), no. 1, 37–78. 7
- [CFDF95] P. J. Cameron and D. G. Fon-Der-Flaass, *Orbits of antichains revisited*, European J. Combin. **16** (1995), no. 6, 545–554. 1, 5, 7
- [CTY14] Edward Clifford, Hugh Thomas, and Alexander Yong,  *$K$ -theoretic Schubert calculus for  $\text{OG}(n, 2n+1)$  and jeu de taquin for shifted increasing tableaux*, J. Reine Angew. Math. **690** (2014), 51–63. 7

- [DPS17] Kevin Dilks, Oliver Pechenik, and Jessica Striker, *Resonance in orbits of plane partitions and increasing tableaux*, J. Combin. Theory Ser. A **148** (2017), 244–274. 1, 2, 4, 6, 7, 8, 12, 15, 16, 20
- [Duc74] P. Duchet, *Sur les hypergraphes invariantes*, Discrete Math. **8** (1974), 269–280. 5
- [EP14] David Einstein and James Propp, *Piecewise-linear and birational toggling*, 26th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2014), Discrete Math. Theor. Comput. Sci. Proc., AT, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2014, pp. 513–524. 1
- [Ful97] William Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997, With applications to representation theory and geometry. 7
- [GR15] Darij Grinberg and Tom Roby, *Iterative properties of birational rowmotion II: rectangles and triangles*, Electron. J. Combin. **22** (2015), no. 3, Paper 3.40, 49. 1
- [GR16] ———, *Iterative properties of birational rowmotion I: generalities and skeletal posets*, Electron. J. Combin. **23** (2016), no. 1, Paper 1.33, 40. 1
- [HPPW16] Zachary Hamaker, Rebecca Patrias, Oliver Pechenik, and Nathan Williams, *Doppelgänger: Bijections of plane partitions*, preprint (2016), 38 pages, arXiv:1602.05535. 20
- [Man01] Laurent Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, SMF/AMS Texts and Monographs, vol. 6, American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2001, Translated from the 1998 French original by John R. Swallow, Cours Spécialisés [Specialized Courses], 3. 7
- [Pan09] Dmitri I. Panyushev, *On orbits of antichains of positive roots*, European J. Combin. **30** (2009), no. 2, 586–594. 1
- [Pec14] Oliver Pechenik, *Cyclic sieving of increasing tableaux and small Schröder paths*, J. Combin. Theory Ser. A **125** (2014), 357–378. 2, 8, 12
- [Pec17] ———, *Promotion of increasing tableaux: frames and homomesies*, Electron. J. Combin. **24** (2017), no. 3, Paper 3.50, 1–14. 8
- [PR15] James Propp and Tom Roby, *Homomesy in products of two chains*, Electron. J. Combin. **22** (2015), no. 3, Paper 3.4, 1–29. 1
- [Pro84] Robert A. Proctor, *Bruhat lattices, plane partition generating functions, and minuscule representations*, European J. Combin. **5** (1984), no. 4, 331–350. 5
- [PSV16] Timothy Pressey, Anna Stokke, and Terry Visentin, *Increasing tableaux, Narayana numbers and an instance of the cyclic sieving phenomenon*, Ann. Comb. **20** (2016), no. 3, 609–621. 8
- [PY17] Oliver Pechenik and Alexander Yong, *Genomic tableaux*, J. Algebraic Combin. **45** (2017), no. 3, 649–685. 7
- [Rho10] Brendon Rhoades, *Cyclic sieving, promotion, and representation theory*, J. Combin. Theory Ser. A **117** (2010), no. 1, 38–76. 3
- [Rho17] ———, *A skein action of the symmetric group on noncrossing partitions*, J. Algebraic Combin. **45** (2017), no. 1, 81–127. 3, 8
- [RS11] David B. Rush and XiaoLin Shi, *On orbits of order ideals of minuscule posets*, unpublished manuscript, available at <http://www-users.math.umn.edu/~reiner/REU/RushShi2011.pdf>, 2011. 2
- [RS13] ———, *On orbits of order ideals of minuscule posets*, J. Algebraic Combin. **37** (2013), no. 3, 545–569. 1, 2
- [RSW04] V. Reiner, D. Stanton, and D. White, *The cyclic sieving phenomenon*, J. Combin. Theory Ser. A **108** (2004), no. 1, 17–50. 2, 3, 26
- [Sch72] M.-P. Schützenberger, *Promotion des morphismes d’ensembles ordonnés*, Discrete Math. **2** (1972), 73–94. 8

- [Sch77] ———, *Combinatoire et représentation du groupe symétrique (Actes Table Ronde CNRS, Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976)*, ch. La correspondance de Robinson, pp. 59–113. Lecture Notes in Math., Vol. 579, Springer, Berlin, 1977. 7
- [SW12] Jessica Striker and Nathan Williams, *Promotion and rowmotion*, European J. Combin. **33** (2012), no. 8, 1919–1942. 1, 6
- [TY09] Hugh Thomas and Alexander Yong, *A jeu de taquin theory for increasing tableaux, with applications to  $K$ -theoretic Schubert calculus*, Algebra Number Theory **3** (2009), no. 2, 121–148. 2, 7, 8
- [Vor17] Corey Vorland, *Homomesy in products of three chains and multidimensional recombination*, preprint (2017), 17 pages, [arXiv:1705.02665](https://arxiv.org/abs/1705.02665). 1, 8

(HM) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720

*E-mail address:* [holly.mandel@berkeley.edu](mailto:holly.mandel@berkeley.edu)

(OP) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109

*E-mail address:* [pechenik@umich.edu](mailto:pechenik@umich.edu)