
Electrodynamic properties of the vacuum near boundaries

Santiago Sanz Wuhl

Institut für Theoretische Physik

Universität Leipzig

Abstract

In my M.Sc. project, I study the effects of the backreaction of a charged Klein-Gordon field coupled to an external electric field, in $1+1$ dimensional spacetime.

In this talk, I motivate and state the problem I intend to solve, discuss existing results from the literature, and present preliminary results from my project.

Table of Contents

► Introduction

► Methods

► Preliminary results

► Bibliography

Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].

Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].
- New interests in studying the effect of **boundary conditions** of QFT in the context of advanced materials [FKV19].

Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].
- New interests in studying the effect of **boundary conditions** of QFT in the context of advanced materials [FKV19].
- **Toy model** for semi classical gravity. [Wro11]

Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].
- New interests in studying the effect of **boundary conditions** of QFT in the context of advanced materials [FKV19].
- **Toy model** for semi classical gravity. [Wro11]
- These calculations were **already done** [AW83], but present counter intuitive results, such as anti-screening behavior for Neumann boundary conditions. These calculations were corrected by [WZ20].

Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].
- New interests in studying the effect of **boundary conditions** of QFT in the context of advanced materials [FKV19].
- **Toy model** for semi classical gravity. [Wro11]
- These calculations were **already done** [AW83], but present counter intuitive results, such as anti-screening behavior for Neumann boundary conditions. These calculations were corrected by [WZ20].
 1. Use of a mode sum formula for the charge density


Motivation

1 Introduction

- **Historically** Vacuum polarization was one of the first quantum electrodynamical effects theoretically studied [Ueh35; HE36].
- New interests in studying the effect of **boundary conditions** of QFT in the context of advanced materials [FKV19].
- **Toy model** for semi classical gravity. [Wro11]
- These calculations were **already done** [AW83], but present counter intuitive results, such as anti-screening behavior for Neumann boundary conditions. These calculations were corrected by [WZ20].
 1. Use of a mode sum formula for the charge density
 2. Neglecting broken down modes for Neumann boundary conditions


The problem

1 Introduction




The problem

1 Introduction



The problem

1 Introduction



Semi classical Klein-Gordon-Maxwell equations

The goal is to solve the coupled system of equations^a

$$\begin{cases} [D_\mu D^\mu + m^2] \phi(x) = 0 \\ \partial_\mu F^{\mu\nu} = \langle : j^\nu : \rangle_\phi \end{cases} \quad (1)$$

^aNotation: The metric signature is $(+, -)$ and the coordinates are $x = (t, z)$

Semi classical Klein-Gordon-Maxwell equations

The goal is to solve the coupled system of equations^a

$$\begin{cases} [D_\mu D^\mu + m^2] \phi(x) = 0 \\ \partial_\mu F^{\mu\nu} = \langle : j^\nu : \rangle_\phi \end{cases} \quad (1)$$

With the covariant derivative

$$D_\mu = \partial_\mu + ieA_\mu,$$

^aNotation: The metric signature is $(+, -)$ and the coordinates are $x = (t, z)$

Semi classical Klein-Gordon-Maxwell equations

The goal is to solve the coupled system of equations^a

$$\begin{cases} [D_\mu D^\mu + m^2] \phi(x) = 0 \\ \partial_\mu F^{\mu\nu} = \langle : j^\nu : \rangle_\phi \end{cases} \quad (1)$$

With the covariant derivative

$$D_\mu = \partial_\mu + ieA_\mu,$$

and the boundary conditions

$$\begin{cases} \phi|_{z=0,1} = 0 \\ E^\phi|_{z=0,1} = 0 \end{cases} \quad (2)$$

^aNotation: The metric signature is $(+, -)$ and the coordinates are $x = (t, z)$

The problem

1 Introduction




Table of Contents

- ▶ Introduction
- ▶ **Methods**
- ▶ Preliminary results
- ▶ Bibliography

Fixing the gauge

2 Methods

1. Constant electric field of strength λ pointing towards positive z .





Figure: The background electric field

Fixing the gauge

2 Methods



1. Constant electric field of strength λ pointing towards positive z .
2. Under the Coulomb gauge

$$A_0(t, z) = -\lambda z, \quad A_1(t, z) = 0$$

up to additive constant

Figure: The background electric field

Fixing the gauge

2 Methods




Figure: The background electric field

1. Constant electric field of strength λ pointing towards positive z .
2. Under the Coulomb gauge

$$A_0(t, z) = -\lambda z, \quad A_1(t, z) = 0$$

up to additive constant

3. To ensure anti-symmetric solutions

$$A_0(t, z) = -\lambda\left(z - \frac{1}{2}\right)$$

Electromagnetic potential

2 Methods




Figure: The background classical electric field and potential

1. Constant electric field of strength λ pointing towards positive z .
2. Under the Coulomb gauge

$$A_0(t, z) = -\lambda z, \quad A_1(t, z) = 0$$

up to additive constant

3. To ensure anti-symmetric solutions

$$A_0(t, z) = -\lambda(z - \frac{1}{2})$$

Electromagnetic potential

2 Methods




Figure: The background electric field and potential, with an additional (antisymmetric) potential.

1. Constant electric field of strength λ pointing towards positive z .
2. Under the Coulomb gauge

$$A_0(t, z) = -\lambda z, \quad A_1(t, z) = 0$$

up to additive constant

3. To ensure antisymmetric solutions

$$A_0(t, z) = -\lambda(z - \frac{1}{2}) + A_0^\phi(z)$$

The set up

2 Methods

Klein-Gordon equation

With the chosen gauge the KGM equations turn into

$$((\partial_t + ieA_0)^2 - \partial_1^2 + m^2)\phi(x) = 0, \quad (3)$$

$$\partial_1^2 A_0^\phi = -\langle \rho(z) \rangle_\phi, \quad (4)$$

with

$$A_0(z) = -\lambda \left(z - \frac{1}{2} \right) + A_0^\phi(z), \quad (5)$$

$$x = (t, z) \in \mathbb{R} \times [0, 1] \quad (6)$$

Time independent Klein-Gordon equation

2 Methods

With the variable separation ansatz $\phi(x) = \phi_n(z)e^{-i\omega_n t}$,

Time independent Klein-Gordon equation

2 Methods

With the variable separation ansatz $\phi(x) = \phi_n(z)e^{-i\omega_n t}$,

Mode equation

For some potential $A_0(z)$

$$\left([\omega_n - eA_0(z)]^2 + \frac{d^2}{dz^2} + m^2 \right) \phi_n = 0 \quad (7)$$

Without backreaction

Time independent Klein-Gordon equation

2 Methods

With the variable separation ansatz $\phi(x) = \phi_n(z)e^{-i\omega_n t}$,

Mode equation

For some potential $A_0(z)$

$$\left([\omega_n - eA_0(z)]^2 + \frac{d^2}{dz^2} + m^2 \right) \phi_n = 0 \quad (7)$$

Without backreaction

$$\left(\left[\omega_n + e\lambda \left(z - \frac{1}{2} \right) \right]^2 + \frac{d^2}{dz^2} + m^2 \right) \phi_n = 0 \quad (8)$$

The external field approximation

2 Methods

Without taking backreaction into account, the KG equation can be solved by

$$\begin{aligned}\phi_n(z) = & a_n D_{i\frac{m^2}{2\lambda} - \frac{1}{2}} \left(\frac{1+i}{\sqrt{\lambda}} \left(\omega_n + \lambda \left(z - \frac{1}{2} \right) \right) \right) \\ & + b_n D_{-i\frac{m^2}{2\lambda} - \frac{1}{2}} \left(\frac{i-1}{\sqrt{\lambda}} \left(\omega_n + \lambda \left(z - \frac{1}{2} \right) \right) \right)\end{aligned}$$

with $D_\nu(z)$ the parabolic cylinder functions.

The external field approximation limitations

2 Methods

Results by [AW83]:




Figure: Positive energy levels for increasing λ **without** considering the backreaction of the massless Klein-Gordon field.

The external field approximation limitations

2 Methods

Results by [AW83]:




Figure: Positive energy levels for increasing λ **without** considering the backreaction of the massless Klein-Gordon field.

The external field approximation limitations

2 Methods

Results by [AW83]:




Figure: Positive energy levels for increasing λ **without** considering the backreaction of the massless Klein-Gordon field.




Figure: Positive energy levels for increasing λ **with** backreaction.

Quantization and renormalization

2 Methods

We quantize the field based on the mode solutions

$$\phi(t, z) = \sum_{n>0} a_n \phi_n(z) e^{-i\omega_n t} + \sum_{n<0} b_n^\dagger \phi_n(z) e^{-i\omega_n t}, \quad (9)$$

with a_n, b_n the operators fulfilling

$$[a_n, a_m^\dagger] = \delta_{nm}, \quad [b_n, b_m^\dagger] = \delta_{nm}, \quad (10)$$

$$[a_n, a_m] = [a_n, b_m] = [a_n, b_m^\dagger] = [b_n, b_m] = 0. \quad (11)$$

This also defines the vacuum by

$$a_n |0\rangle = b_n |0\rangle = 0 \quad (12)$$

Charge density

2 Methods

We calculate the **charge density** as the 0th component of the charge current

$$\rho(z) = ie \langle 0 | \phi^*(z) D_0 \phi(z) - \phi(z) D_0 \phi^*(z) | 0 \rangle \quad (13)$$

Charge density

2 Methods

We calculate the **charge density** as the 0th component of the charge current

$$\rho(z) = ie \langle 0 | \phi^*(z) D_0 \phi(z) - \phi(z) D_0 \phi^*(z) | 0 \rangle \quad (13)$$

Ambjørn and Wolfram calculate it as one naïvely would,

$$\rho(z) = e \left(\sum_{n>0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 + \sum_{n<0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 \right) \quad (14)$$

Hadamard states and point-splitting

2 Methods

Assume that the state is of Hadamard form,

$$w_{\Omega}^{\phi\phi^*}(x, x') = \langle \Omega | \phi(x) \phi^*(x') | \Omega \rangle = H^{\phi\phi^*}(x, x') + R_{\Omega}^{\phi\phi^*}(x, x'), \quad (15)$$

Hadamard states and point-splitting

2 Methods

Assume that the state is of Hadamard form,

$$w_{\Omega}^{\phi\phi^*}(x, x') = \langle \Omega | \phi(x) \phi^*(x') | \Omega \rangle = H^{\phi\phi^*}(x, x') + R_{\Omega}^{\phi\phi^*}(x, x'), \quad (15)$$

with

$$H^{\phi\phi^*}(x, x') = -\frac{1}{4\pi} U(x, x') \log(-(x - x')^2 + i\epsilon(x - x')^0) \quad (16)$$

and

$$U(x, x') = \exp \left(-i \int_0^1 A_{\mu}(x' + s(x - x'))(x - x')^{\mu} ds \right).$$

Hadamard states and point-splitting

2 Methods

Assume that the state is of Hadamard form,

$$w_{\Omega}^{\phi\phi^*}(x, x') = \langle \Omega | \phi(x) \phi^*(x') | \Omega \rangle = H^{\phi\phi^*}(x, x') + R_{\Omega}^{\phi\phi^*}(x, x'), \quad (15)$$

with

$$H^{\phi\phi^*}(x, x') = -\frac{1}{4\pi} U(x, x') \log(-(x - x')^2 + i\epsilon(x - x')^0) \quad (16)$$

and

$$U(x, x') = \exp \left(-i \int_0^1 A_{\mu}(x' + s(x - x'))(x - x')^{\mu} ds \right).$$

Define the product of derivatives of fields as

$$\langle \Omega | D_{\alpha} \phi(x) (D_{\beta} \phi)^*(x) | \Omega \rangle = \lim_{x' \rightarrow x} \left[D_{\alpha} D'_{\beta}{}^* \left(w_{\Omega}^{\phi\phi^*}(x, x') - H^{\phi\phi^*}(x, x') \right) \right], \quad (17)$$

Renormalizing charge densities

2 Methods

Recall the mode decomposition of the field

$$\phi(t, z) = \sum_{n>0} a_n \phi_n(z) e^{-i\omega_n t} + \sum_{n<0} b_n^\dagger \phi_n(z) e^{-i\omega_n t}, \quad (18)$$

the expression for charge density

$$\rho(x) = ie \langle 0 | \phi^*(x) D_0 \phi(x) - \phi(x) D_0 \phi^*(x) | 0 \rangle, \quad D_0 = \partial_0 + ieA_0, \quad (19)$$

Renormalizing charge densities

2 Methods

Recall the mode decomposition of the field

$$\phi(t, z) = \sum_{n>0} a_n \phi_n(z) e^{-i\omega_n t} + \sum_{n<0} b_n^\dagger \phi_n(z) e^{-i\omega_n t}, \quad (18)$$

the expression for charge density

$$\rho(x) = ie \langle 0 | \phi^*(x) D_0 \phi(x) - \phi(x) D_0 \phi^*(x) | 0 \rangle, \quad D_0 = \partial_0 + ieA_0, \quad (19)$$

Realizing the point splitting in the time direction $x' = (t + \tau, z)$, we calculate

$$\langle 0 | \phi^*(x) D_0 \phi(x') | 0 \rangle = -i \sum_{n<0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau+i\epsilon)} \quad (20)$$

Renormalizing charge densities

2 Methods

$$\langle 0 | \phi^*(x) D_0 \phi(x') | 0 \rangle = -i \sum_{n < 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau + i\epsilon)} \quad (21)$$

Renormalizing charge densities

2 Methods

$$\langle 0 | \phi^*(x) D_0 \phi(x') | 0 \rangle = -i \sum_{n < 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau + i\epsilon)} \quad (21)$$

$$\langle 0 | \phi(x) D_0^* \phi^*(x') | 0 \rangle = i \sum_{n > 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{i\omega_n(\tau + i\epsilon)} \quad (22)$$

Renormalizing charge densities

2 Methods

$$\langle 0 | \phi^*(x) D_0 \phi(x') | 0 \rangle = -i \sum_{n < 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau + i\epsilon)} \quad (21)$$

$$\langle 0 | \phi(x) D_0^* \phi^*(x') | 0 \rangle = i \sum_{n > 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{i\omega_n(\tau + i\epsilon)} \quad (22)$$

$$D'_0 H^{\phi\phi^*}(x, x') = -\frac{1}{2\pi} \frac{1}{\tau + i\epsilon} U(x', x) + \mathcal{O}(\tau) = -\frac{1}{2\pi} \left(\frac{1}{\tau + i\epsilon} - ieA_0 \right) + \mathcal{O}(\tau) \quad (23)$$

Renormalizing charge densities

2 Methods

$$\langle 0 | \phi^*(x) D_0 \phi(x') | 0 \rangle = -i \sum_{n < 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau + i\epsilon)} \quad (21)$$

$$\langle 0 | \phi(x) D_0^* \phi^*(x') | 0 \rangle = i \sum_{n > 0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{i\omega_n(\tau + i\epsilon)} \quad (22)$$

$$D'_0 H^{\phi\phi^*}(x, x') = -\frac{1}{2\pi} \frac{1}{\tau + i\epsilon} U(x', x) + \mathcal{O}(\tau) = -\frac{1}{2\pi} \left(\frac{1}{\tau + i\epsilon} - ieA_0 \right) + \mathcal{O}(\tau) \quad (23)$$

$$D'^*_0 H^{\phi^*\phi}(x, x') = -\frac{1}{2\pi} \frac{1}{\tau + i\epsilon} U(x, x') + \mathcal{O}(\tau) = -\frac{1}{2\pi} \left(\frac{1}{\tau + i\epsilon} + ieA_0 \right) + \mathcal{O}(\tau) \quad (24)$$

The renormalized charge density

2 Methods

$$\rho(z) = e \lim_{\tau \rightarrow 0} \left(\sum_{n>0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{i\omega_n(\tau+i\epsilon)} + \sum_{n<0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau+i\epsilon)} \right) + \frac{e^2}{\pi} A_0(z)$$

(25)

The renormalized charge density


2 Methods

$$\rho(z) = e \lim_{\tau \rightarrow 0} \left(\sum_{n>0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{i\omega_n(\tau+i\epsilon)} + \sum_{n<0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 e^{-i\omega_n(\tau+i\epsilon)} \right) + \frac{e^2}{\pi} A_0(z) \quad (25)$$

$$\rho(z) = e \left(\sum_{n>0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 + \sum_{n<0} (\omega_n - eA_0(z)) |\phi_n(z)|^2 \right) \quad (26)$$

The renormalized charge density

2 Methods



The induced charge density ρ resulting from the two different renormalization techniques with Dirichlet boundary conditions. $m = 0$, $\lambda = 5$.


- In orange, the charge density ρ calculated through the mode sum formula.
- In blue, the charge density ρ calculated through Hadamard point-splitting.

The induced fields

2 Methods

The induced electric field:

$$E^\phi(z) = \int_0^z \rho(z') dz' \quad (27)$$




The induced fields

2 Methods


The induced electric field:

$$E^\phi(z) = \int_0^z \rho(z') dz' \quad (27)$$




The induced electric potential

$$A_0^\phi(z) = - \int_0^z E^\phi(z') dz' \quad (28)$$



Closing the loop

2 Methods



Closing the loop

2 Methods

$$\left(\left[\omega_n^{(\kappa+1)} + \lambda \left(z - \frac{1}{2} \right) - \left(A_0^\phi \right)^{(\kappa)}(z) \right]^2 + \frac{d^2}{dz^2} + m^2 \right) \phi_n^{(\kappa+1)} = 0 \quad (29)$$

$$\left(A_0^\phi \right)^{(\kappa)}(z) = - \int_0^z \int_0^{z'} \rho^{(\kappa)}(z'') dz'' dz', \quad (30)$$

Visualizing the impact of backreaction

2 Methods




Figure: The convergence of $\rho^\kappa(z)$ as $\kappa \rightarrow \infty$ for $m = 0$, $\lambda = 15.5$.

Table of Contents

- ▶ Introduction
- ▶ Methods
- ▶ Preliminary results
- ▶ Bibliography

ω_n dependence on λ considering backreaction

3 Preliminary results




Figure: ω_n as a function of λ for the massless case.

Charge densities and $E\phi$

3 Preliminary results




Figure: Charge densities ρ for increasing background electric field λ




Figure: Average polarization for different background electric fields.

Outlook

3 Preliminary results

There is still work to be done:

1. For Dirichlet boundary conditions, study deeper the behavior of the critical λ value.
2. Repeat these calculations for Neumann boundary conditions, and finally for more general Robin boundary conditions

$$\left. \frac{\partial \phi}{\partial z} \right|_{z=0} = h_0 \phi(0) \quad (31)$$

$$\left. \frac{\partial \phi}{\partial z} \right|_{z=1} = -h_1 \phi(1) \quad (32)$$

Table of Contents

- ▶ Introduction
- ▶ Methods
- ▶ Preliminary results
- ▶ Bibliography

- [Ueh35] Edwin A Uehling. “Polarization effects in the positron theory”. In: *Physical Review* 48.1 (1935), p. 55.
- [HE36] Werner Karl Heisenberg and H Euler. “Folgerungen aus der Diracschen Theorie des Positrons”. In: *Z. Phys.* 98 (1936), pp. 714–732. URL: <https://cds.cern.ch/record/423057>.
- [FKV19] Ignat Fialkovsky, Maxim Kurkov, and Dmitri V Vassilevich. “Quantum Dirac fermions in a half-space and their interaction with an electromagnetic field”. In: *Physical Review D* (2019). URL: <https://api.semanticscholar.org/CorpusID:189927793>.
- [Wro11] Michał Wrochna. “Quantum field theory in static external potentials and Hadamard states”. In: (Aug. 15, 2011). DOI: [10.1007/s00023-012-0173-0](https://arxiv.org/abs/1108.2982). arXiv: 1108.2982v2 [math-ph].

- [AW83] Jan Ambjørn and Stephen Wolfram. “Properties of the vacuum. 2. Electrodynamic”. In: *Annals of Physics* 147.1 (1983), pp. 33–56. ISSN: 0003-4916. DOI: [https://doi.org/10.1016/0003-4916\(83\)90066-0](https://doi.org/10.1016/0003-4916(83)90066-0). URL: <https://www.sciencedirect.com/science/article/pii/0003491683900660>.
- [WZ20] Jonathan Wernersson and Jochen Zahn. “Vacuum polarization near boundaries”. In: *Phys. Rev. D* 103, 016012 (2021) (Oct. 12, 2020). DOI: 10.1103/PhysRevD.103.016012. arXiv: 2010.05499v2 [hep-th].

**Electrodynamic properties of the
boundaries**

vacuum near

Thank you for listening