# Vacuum Polarization in 1+1 dimensions

*Authors*

**Abstract**

In this article...

## Contents


| | | |
|---|---|---|
| 1 | Introduction | 1 |
| 2 | Set up | 2 |
| 2.1 | Field modes | 2 |
| 2.2 | External Field Approximation and Critical Field Strength | 4 |
| 3 | Self-consistent fields | 4 |
| 4 | Results | 5 |
| 4.1 | Mode sum formula | 6 |
| 5 | Discussion and outlook | 6 |
| A | Mode sum formula | 7 |
| B | Computing the vacuum polarization | 8 |


## 1 Introduction

Vacuum polarization is a phenomenon in quantum electrodynamics (QED) by which the vacuum of a charged field polarizes when subject to external electric fields, effectively behaving like a dielectric with $\epsilon > 0$. This mechanism was one of the first QED effects studied [3], with consequences that followed closely [14; 6]. This mechanism has been already observed, e.g. in corrections to atomic levels [9] (cf. Ref. [10] for a review), while other effects e.g. the Schwinger effect [12] lay beyond current technological advances (cf. Ref. [4]).

Defining vacuum polarization for generic backgrounds[1] is no easy task. Historically, [3] first defined it through the assumption that the short-distance behavior of the physically relevant states in the presence of strong background electric fields is the same as in their absence, only up to smooth coefficients. In modern bibliography, these states are called Hadamard, (cf Ref. [5]), and they are particularly interesting in the context of quantum field theory on curved spacetimes, where the same idea is used to define observables necessary for the study of e.g. perturbative interacting field theory or semi-classical gravity (cf. [7; 8] for detailed discussions on the definition on such observables).

Semi-classical gravity, a possible first step towards a quantum theory of gravity, couples classical gravity to the expectation value of the stress-energy tensor, which similarly as the vacuum polarization needs to be redefined. To our knowledge, only perturbative corrections to the metric have been studied [13]. In this article we study, as a toy model, the effect of the backreaction of a charged scalar field to a background classical electric field in 1+1 dimensional Minkowski spacetime. This setup was already studied in [1], where it was found that ignoring the backreaction of the scalar field leads to instabilities, and it claims that the inclusion thereof avoids them. However, as pointed out in [11], the expression used in [1] for vacuum polarization $\rho$ cannot be derived from a manifestly gauge invariant prescription. [15] derives the correct expression for $\rho$, which has qualitative disagreements with the one used in [1].

This discussion motivates the following question, which we answer in the present paper: Given the difference in the prescriptions for $\rho$, does using the correct one avoid these instabilities? We will use the correct prescription described in [15] to implement an iterative method to study the effect of the backreaction of the scalar field. It has been found that this expression also alleviates the above mentioned instabilities.

The layout of this article is as follows: Section 2 ...

**Maybe add?**: [2] discusses hadamard states in the pressence of time-like boundaries.

[1] And any other observable defined through the evaluation of products of fields or their derivatives at coinciding points.

This article summarizes and extends the results of the M.Sc. thesis of Santiago Sanz-Wuhl, written in 2025 at the Institut für Theoretische Physik, Universität Leipzig, under the supervision of Jochen Zahn.

## 2 Set up

In 1+1 dimensional spacetime, two charges $q, -q$ are set at $x^1 = 0, x^1 = a$, respectively. Ignoring vacuum polarization effects, this yields a constant electric field of strength $q$ pointing towards positive $x^1$ in the region $x^1 \in [0, a]$, 0 everywhere else. We study the solutions to the semiclassical Klein-Gordon-Maxwell equations (KGM), i.e. the system of coupled differential equations

$$\left(D_\mu D^\mu + m^2\right)\phi = 0, \tag{1a}$$

$$\partial_\mu F^{\mu\nu} = \langle j^\nu \rangle, \tag{1b}$$

in the region $x = (x^0, x^1) \in \mathbb{R} \times [0, a]$. Here, $D_\mu = \partial_\mu + ieA_\mu$ is the gauge covariant derivative,

$$j^\nu(x) = \frac{ie}{2}\left((D^\nu\phi^*(x))\phi(x) - \phi^*(x)D^\nu\phi(x)\right) \tag{2}$$

is the charge current density of the Klein-Gordon field, and $\langle \mathcal{O} \rangle$ is the vacuum expectation value of an observable $\mathcal{O}$, and the metric signature is $(+, -)$. We solve for $F_{\mu\nu}$ and (the mode decomposition of) the two-point function of the vacuum state.

In 1+1 dimensions, $A_1$ can be arbitrarily set to 0. This way, Eq. (1b) reduces to the Poisson equation for the electrostatic potential $A_0$

$$\partial_1^2 A_0(x^1) = -\left\langle j^0 \right\rangle \equiv -\rho, \tag{3}$$

with $\rho$ the vacuum polarization, i.e. the vacuum expectation value of the zeroth component of the charge current density. Eq. (3) will be solved with the boundary conditions

$$\partial_1 A_0(x^1)\big|_{x^1=0,a} = -q. \tag{4}$$

This way, measuring the electric field close enough to the boundary is not affected by vacuum polarization. The boundary conditions (4) fix the solution to Eq. (3) up to an addition constant, which is for convenience chosen so that $A_0(\frac{a}{2}) = 0$. With these boundary conditions and for a given $\rho(x^1)$, Eq. (3) is solved by

$$A_0(x^1) = -q\left(x^1 - \frac{a}{2}\right) - \int_{\frac{a}{2}}^{x^1}\int_0^\chi \rho(\xi)d\xi d\chi. \tag{5}$$

### 2.1 Field modes

In this gauge, the Klein-Gordon equation is of the form

$$\left[(\partial_0 + ieA_0)^2 + \partial_1^2 - m^2\right]\phi = 0, \tag{6}$$

which we will study using Dirichlet boundary conditions (DBC) $\phi|_{x^1=0,a} = 0$ or Neumman boundary conditions (NBC) $\partial_1\phi|_{x^1=0,a} = 0$. More general boundary conditions may be considered, though they are outside of the scope of this work. Using the mode ansatz $\phi_n(x) = \phi_n(x^1)e^{-i\Omega_n x^0}$ the PDE is split into ordinary differential equations for each mode $\phi_n(x^1)$ of the scalar field

$$\left[(\Omega_n - eA_0)^2 + \partial_1^2 - m^2\right]\phi_n = 0. \tag{7}$$

The following dimensionless parameters are defined

$$z = a^{-1}x^1 \tag{8a}$$

$$\lambda = eEa^2 \tag{8b}$$

$$\varepsilon = ae \tag{8c}$$

$$\omega_n = a\Omega_{n,\alpha} \tag{8d}$$

Eq. (7) then takes the form

$$\left[(\omega_n - \varepsilon A_0)^2 + \frac{d^2}{dz^2} - a^2m^2\right]\phi_n = 0. \tag{9}$$

The modes $\phi_n$ are normalized with respect to the symplectic norm

$$(\phi_n, \phi_m) = i\int_0^1 (\phi_n^*\pi_m^* - \pi_n\phi_m)dz, \tag{10}$$

with $\pi_n = (D_0\phi_n)^*$ the canonical conjugate of $\phi_n$.

The general solution to Eq. (6) is thus decomposed in modes

$$\phi(t, z) = \sum_{n>0} a_n\phi_n(z)e^{-i\omega_n t} + \sum_{n<0} b_n^\dagger\phi_n(z)e^{-i\omega_n t}. \tag{11}$$

Positive (negative) values of $n$ are identified with modes of positive (negative) norm. $a_n$ and $b_n$ are the ladder operators for the positive and negative norm solutions, respectively, obeying canonical commutation relations $[a_n, a_n^\dagger] = [b_n, b_n^\dagger] = \delta_{nm}$, and all other commutators vanishing. The ladder operators define the vacuum state $|0\rangle$ as that for which $a\,|0\rangle = b\,|0\rangle = 0$. Note that due to the chosen gauge, $\omega_n = -\omega_{-n}$, and the $b_n^\dagger$ are the creation operators of the negative frequency family of solutions.

As mentioned above, the definition of quantities such as $\rho$ —Wick polynomials—is problematic, since it involves the product of local observables evaluated at coinciding spacetime points (see Eq. (2)). To define such polynomials, one assumes that the states of a field $\phi$ are Hadamard, i.e. they have a similar short-distance behavior to the vacuum of $\phi$ in absence of background fields, and are assumed to be the only physically relevant states [5]. Under this assumption, the divergence of such states is state independenent and locally constructed from the background fields. Since $\phi$ is coupled to the background field through the covariant derivative, one may use the setup at hand as a toy model for semi-classical gravity, a possible first step towards an elusive quantum theory of gravity.

Specifically, one says that the two point function $w_\Omega^{\phi\phi^*}(x,x') = \langle\Omega|\phi(x)\phi^*(x')|\Omega\rangle$ of the scalar field $\phi$ in a Hadamard state $\Omega$ is of the form

$$w_\Omega^{\phi\phi^*}(x,x') \equiv H^{\phi\phi^*}(x,x') + R_\Omega^{\phi\phi^*}(x,x'). \tag{12}$$

Here,

$$H^{\phi\phi^*} = -\frac{V(x,x')}{4\pi}\ln[(t-t')^2-(z-z')^2+i\varepsilon\mathrm{sign}(t-t')] \tag{13}$$

is the Hadamard parametrix of the scalar field in 1+1 spacetime dimensions and $R_\Omega^{\phi\phi^*}$ a state-dependent smooth function. The smooth coefficient $V$, up to order relevant for vacuum polarization reads [11]

$$V(x,x') = \exp\left[-i\varepsilon\int_0^1 A_\mu(x'+s(x-x'))(x-x')^\mu ds\right] + \mathcal{O}(x-x')^2.$$

The Hadamard condition allows us to define expectation values of local observables non-linear in the fields




Fig. 1: The vacuum polarization $\rho$ of the scalar field due to a background electrostatic potential $A_0(z) = -0.5(z-0.5)$, with DBC and $m = 0$.

as a limit of coinciding points:

$$\begin{aligned}&\langle D_\alpha\phi(x)D_\beta^*\phi^*(x)\rangle \\ &\quad = \lim_{x'\to x} D_\alpha D_\beta^{*\prime}(w_\Omega^{\phi\phi^*}(x,x') - H^{\phi\phi^*}(x,x')),\end{aligned} \tag{14}$$

with $\alpha, \beta$ symmetrized multiindices. The above prescription is called *point-splitting with respect to a Hadamard parametrix.*

Eqs. (14), (2) and the mode decomposition (11) together result in the expression for $\rho$ calculated in [15]

$$\begin{aligned}\rho = \lim_{\tau\to 0}\sum_{n>0}\Big[&(\omega_n - \varepsilon A_0)|\phi_n|^2 e^{-i\omega_n\tau} \\ &+ \sum_{n<0}(\omega_n - \varepsilon A_0)|\phi_n|^2 e^{i\omega_n\tau}\Big] + \frac{\varepsilon^2}{\pi}A_0.\end{aligned} \tag{15}$$

An example of $\rho$ using this expression is given in Fig. 1, calculated in the external field approximation explained below, i.e. a homogeneous background electric field given by the potential $A_0(z) = -0.5(z-0.5)$.

To measure the *strength* of the vacuum polarization, the *induced charge*

$$Q_I = \int_0^{\frac{1}{2}} \rho(z)dz \tag{16}$$




Fig. 2: Energies of the three first modes of the massless Klein-Gordon field subject to DBC in the external field approximation as functions of $\lambda$. $\omega_1 \to 0$ as $\lambda \to \lambda_c \approx 16$ defines the critical field strength for these mass and boundary conditions configurations.




Fig. 3: Corresponding induced charge $Q_I$ in Fig. 2 as a function of $\lambda$.

is defined. This charge partially screens the background electric field $\lambda$, and Gauß's law yields that the total electric field at $z = \frac{1}{2}$ as $\lambda + Q_I$.

## 2.2 External Field Approximation and Critical Field Strength

The external field approximation ignores the contribution of the backreaction of the charged scalar field to the background electric field. This approximation displays the instabilities of the system, which the inclusion of backreaction aims to avoid. Eq. (9) is solved with $A_0(z) = -\lambda(z - \frac{1}{2})$, and takes the form

$$\left[\left(\omega_n + \lambda\left(z - \frac{1}{2}\right)\right)^2 + \frac{d^2}{dz^2} - a^2 m^2\right]\phi_n = 0. \quad (17)$$

For each value of $\lambda$, the modes $\phi_n$ and their energies $\omega_n$ can be obtained from the boundary value problem. Fig. 2 displays the energies $\omega_1, \omega_2, \omega_3$ of the first three modes of the massless scalar field with DBC as functions of the background field strength $\lambda$. The value at which $\omega_1$ goes to 0 defines the critical field strength $\lambda_c \approx 16.05039$. Fig. 3 displays the divergent behavior of $Q_I(\lambda)$ for this configuration as $\lambda \to \lambda_c$. $\lambda_c$ adopts different values for different boundary conditions and masses.

[1] claims that the backreaction of $\phi$ to the background electric field avoids these instabilities. However, the expression used therein for the vacuum polarization cannot be derived from gauge invariant renormalization schemes; one should rather use the prescription described in [15]. Their qualitative discrepancies are detailed in Appendix A. This article answers the following question: does considering the backreaction using the correct prescription for the vacuum polarization avoid the instabilities that would otherwise be present?

# 3 Self-consistent fields

As discussed above, the polarization of the vacuum induces a charge density $\rho$, which modifies the background field as in Eq. (5). This modified background field can be used in Eq. (18a) to calculate new solutions, that will have a different vacuum polarization. This iterative procedure, schematized in Fig. 4, is performed *ad infinitum*, and we study its convergence.

More precisely, for each value of $\lambda$ and at each iter-




Fig. 4: Schematic of the iterative procedure for fixed $\lambda$.

ation $\kappa$, we compute the following

$$\frac{d^2\phi_{n,\lambda}^\kappa}{dz^2} = -\left[(\omega_{n,\lambda}^\kappa - \varepsilon A_{0,\lambda}^{\kappa-1})^2 - a^2m^2\right]\phi_{n,\lambda}^\kappa, \tag{18a}$$

$$\begin{aligned}\rho_\lambda^\kappa = \lim_{\tau\to 0}\Big[&\sum_{n>0}(\omega_{n,\lambda}^\kappa - \varepsilon A_{0,\lambda}^{\kappa-1})|\phi_{n,\lambda}^\kappa|^2 e^{-i\omega_{n,\lambda}^\kappa\tau} \\ &+ \sum_{n<0}(\omega_{n,\lambda}^\kappa - \varepsilon A_{0,\lambda}^{\kappa-1})|\phi_{n,\lambda}^\kappa|^2 e^{i\omega_{n,\lambda}^\kappa\tau}\Big] \\ &+ \frac{\varepsilon^2}{\pi}A_{0,\lambda}^{\kappa-1},\end{aligned} \tag{18b}$$

$$\begin{aligned}A_{0,\lambda}^\kappa &= cA_{0,\lambda}^{\kappa-1} \\ &+ (1-c)\left[-\lambda\left(z-\frac{1}{2}\right) - \int_{\frac{1}{2}}^z\int_0^\chi \rho_\lambda^\kappa(\xi)d\xi d\chi\right].\end{aligned} \tag{18c}$$

This procedure has two main differences from the one used in [1]: the expression for the vacuum polarization, and the $c < 1$ prescription in Eq. (18c), which is discussed below. The details on how $\rho$ is calculated are explained in Appendix B. Initial conditions of this procedure are discussed below.

Noticing how Eq. (18c) acts as an update function on $A_{0,\lambda}^{\kappa-1}$, we write this scheme as an infinite dimensional fixed point problem. This is, for each $\lambda$, we look for the fixed point $A_{0,\lambda}$ obeying

$$A_{0,\lambda} = f_\lambda(A_{0,\lambda}), \tag{19}$$

with $f_\lambda$ given by the composition of steps (18a), (18b) and (18c). The fixed points of $f_\lambda$, denominated "self-consistent solutions", are looked for by studying the limit $A_{0,\lambda} = \lim_{\kappa\to\infty} A_{0,\lambda}^\kappa$, where

$$A_{0,\lambda}^{\kappa+1} = f_\lambda(A_{0,\lambda}^\kappa) = f_\lambda^\kappa\left(A_{0,\lambda}^0\right), \text{with } f_\lambda^\kappa = \underbrace{f_\lambda \circ ... \circ f_\lambda}_{k \text{ times}}. \tag{20}$$

If the limit $A_{0,\lambda}$ exists, it is given the name self-consistent potential, as well as the corresponding self-consistent vacuum polarization $\rho_\lambda$, field modes $\phi_{n,\lambda}$ and mode energies $\omega_{n,\lambda}$.

For low enough $\lambda \ll \lambda_c$, the initial value of the sequence $A_{0,\lambda}^0$ can be set to the external field approximation $A_{0,\lambda}^0(z) = -\lambda\left(z - \frac{1}{2}\right)$. This is, however, no longer a valid initial potential for stronger background fields, as it leads to non-convergences in the sequence $A_{0,\lambda}^\kappa$. This is discussed in the Appendix ??(to be added). For higher $\lambda$-values, $A_{0,\lambda}^0$ is instead set by a stored previously calculated self-consistent potential $A_{0,\lambda'}$ with $\lambda = \lambda' + \Delta\lambda$, where $\Delta\lambda > 0$. We use this scheme to probe the $\lambda$-parameter space.

Finally, in Eq. (18c), $c \in [0, 1]$ is introduced in order to avoid an overshooting of the potential: For $c = 0$, the procedure is precisely the one used in [1], and it is expected to quickly converge. For $c$ close to 1, updates will be small corrections to the previous potential. Stronger background field strengths increase the instability of the coupled system, thus increasing the relevance of slower converge schemes. Considering this, we adopt a scheme in which, as $\lambda$ increases, $c$ increases and simultaneously $\Delta\lambda$ decreases.

## 4 Results

Using the iterative procedure above laid out we focus on two particular cases: The massless scalar field subject to DBC, and $m = 1$ with the scalar field subject to NBC. We do not study the massless field subject to NBC, since under these boundary conditions, $\lambda_c = 0$ for $m = 0$, as pointed out in Ref. [15].

Figures 5 and 6 display the dynamics of KGM system in the semiclassical approximation. The external field approximation is given for reference in orange. Notice the different $\lambda$ scales. The system is studied through

the energies of the first modes (top panel), the induced charge $Q_I$ (middle panel) and the total electric field $\lambda+Q_I$ at the midpoint $z = 0.5$ (bottom panel). We observe in both cases how considering backreaction avoids the instabilities reflected in $\omega_1 \to 0$ and $Q_I \to -\infty$ as $\lambda \to \lambda_c$ that the external field approximation presents. Indeed, the energy of the lower modes is raised which "prevents" $\omega_1$ from crossing 0. We also point the reader to notice how the rapid growth of $|Q_I|$ as $\lambda$ crosses $\lambda_c$ keeps $\lambda + Q_I$ below $\lambda_c$. We find this behavior surprising, as intuitively one would expect $\lambda+Q_I$ to stabilize. Considering a quantum background electric field would perhaps fulfill this intuition.

One might also notice the almost identical behavior of the self-consistent solutions to the external field approximation in the weak-field regime $\lambda \ll \lambda_c$ in both cases.

## 4.1 Mode sum formula

For completeness, we compare the results obtained in this work with those obtained in [1]. Fig. 7 displays $Q_I$ in the iterative procedure as a function of $\lambda$ for two different prescriptions for $\rho$. In blue, the correct prescription used throughout this paper, and in orange $\rho$ as defined in [1], truncated to the first mode. Notice the different behaviors in the two regimes: although the truncated mode sum formula presents a noticeably stronger vacuum polarization in the weak-field regime, the strong-field regime behavior is very similar. This might be understood as a result of the lower modes being the most sensitive to background fields, and thus those who contribute the most to vacuum polarization.

# 5 Discussion and outlook

In this article, we have studied the semiclassical Klein-Gordon-Maxwell system equations through an iterative procedure to study the effect of backreaction to a background classical electric field with a well-defined prescription of vacuum polarization. This precise set up was already studied in [1], but the mode sum formula used in their work cannot be derived from a manifestly invariant renormalization scheme, as pointed out in [11; 15].




Fig. 5: In blue, the self-consistent solutions to the coupled semi-classical Klein-Gordon-Maxwell system for a massless scalar field subject to DBC. In orange, for reference, the external field approximation is given. Top panel: Energies of the first three modes as functions of $\lambda$. Middle panel: Corresponding $Q_I$ for each $\lambda$. Bottom panel: Total electric field $\lambda + Q_I$ due to the screening of the vacuum at $z = \frac{1}{2}$.




Fig. 6: Following the format in Fig. 5, the results for the unit mass scalar field subject to NBC. Further details in main text.




Fig. 7: $Q_I$ for the self-consistent solutions for a massless field subject to DBC, by using different expressions for $\rho$. In blue, the (correct) vacuum polarization obtained through renormalization through point-splitting w.r.t. a Hadamard parametrix, and in orange the vacuum polarization obtained through a mode sum formula truncated to one mode used in [1].

## A Mode sum formula

Vacuum polarization in [1] is defined through a mode sum formula

$$\begin{aligned}\rho &= \sum_{n>0} \rho_n \\ &= \sum_{n>0} \left[(\omega_n - \varepsilon A_0)|\phi_n|^2 - (\omega_n + \varepsilon A_0)|\phi_{-n}|^2\right],\end{aligned} \tag{21}$$

so called since the total charge density is calculated as the sum of the contribution $\rho_n$ of each mode to the total charge. The above expression cannot be obtained from a manifestly gauge invariant renormalization scheme, and also relies on a specific pairing of the modes in the series to be convergent. Furthermore, in using the iterative procedure discussed in the following section, [1] truncated the mode sum to the first mode. Fig. 8 shows the different $\rho$ resulting from the different prescriptions. At first sight, it might be surprising that a field subject to DBC has non-zero vacuum polarization at the boundaries. This is precisely due to the missing $\frac{\varepsilon^2}{\pi} A_0$ term in Eq. (21). In using the iterative




Fig. 8: Three different $\rho$ for the same massless scalar field subject to DBC: In blue, the correct prescription as explained in [15]. In orange, $\rho$ calculated through the mode sum (21). In red, $\rho$ defined through the mode sum formula, truncated to the first mode.




Fig. 9: $\rho$ of the massless complex scalar field subject to a constant background electric field of strength $\lambda = 1$, for different mode cut-offs, as indicated in the legend. For reference, the perturbative vacuum polarization is given.

procedure [1] truncates the mode sum (21) to the first mode, which leads to the red curve in Fig. 8, leading to a vacuum polarization aproximately an order of magnitude stronger than the one one would obtain when using Eq. (15).

## B Computing the vacuum polarization

To be able to compute Eq. (15), some rearrangements are to be done. By use of $-\omega_n = \omega_{-n}$, $\rho$ can be expressed as a single sum

$$\rho = \lim_{\tau\to 0} \sum_{n>0} \Big[ (\omega_n - \varepsilon A_0)|\phi_n|^2 - (\omega_n + \varepsilon A_0)|\phi_{-n}|^2 \Big] e^{-i\omega_n \tau} + \frac{\varepsilon^2}{\pi} A_0(z) \tag{22}$$

In general, the limit and the sum do not commute. In order for them to commute, the coefficients of the complex exponentials need to decay fast enough, in order for the series to be absolutely convergent. After commuting, the sum is cut off at some finite mode $N$, which induces oscillations of frequency $\Delta z = \frac{1}{N+1}$. These oscillations are an artifact of the cut off, which must be smoothened out by a low-pass filter.

To achieve this, a low-pass filter is applied to the unfiltered vacuum polarization, by convoluting it with a normalized gaussian function, of approximately 3 oscillations of width. The $\rho$ resulting from the filtering is displayed in Fig. 9, where the filtering has been applied to the vacuum polarization resulting from different mode cut-offs (20, 60 and 100, as indicated by the legend). For reference, the vacuum polarization calculated from perturbative theory as per [15] is also given.

## References

[1] J. Ambjorn and S. Wolfram. Properties of the Vacuum. 2. Electrodynamic. *Annals Phys.*, 147: 33, 1983. doi: 10.1016/0003-4916(83)90066-0.

[2] B. Costeri, C. Dappiaggi, B. A. Juárez-Aubry, and R. D. Singh. The Hadamard parametrix on half-Minkowski with Robin boundary conditions: Fundamental solutions and Hadamard states, Sept. 2025. URL `http://arxiv.org/abs/2509.26035`. arXiv:2509.26035 [math-ph].

[3] P. A. M. Dirac. Discussion of the infinite distribution of electrons in the theory of the positron. *Mathematical Proceedings of the Cambridge Philosophical Society*, 30(2):150–163, 1934. doi: 10.1017/S030500410001656X.

[4] A. Fedotov, A. Ilderton, F. Karbstein, B. King, D. Seipt, H. Taya, and G. Torgrimsson. Advances in QED with intense background fields. *Physics Reports*, 1010:1–138, Apr. 2023. ISSN 03701573. doi: 10.1016/j.physrep.2023.01.003. URL `http://arxiv.org/abs/2203.00019`. arXiv:2203.00019 [hep-ph].

[5] C. J. Fewster and R. Verch. The Necessity of the Hadamard Condition. *Class. Quant. Grav.*, 30:235027, 2013. doi: 10.1088/0264-9381/30/23/235027. _eprint: 1307.5242.

[6] W. Heisenberg and H. Euler. Consequences of Dirac's theory of positrons. *Z. Phys.*, 98(11-12): 714–732, 1936. doi: 10.1007/BF01343663. _eprint: physics/0605038.

[7] S. Hollands and R. M. Wald. Local Wick Polynomials and Time Ordered Products of Quantum Fields in Curved Spacetime. *Communications in Mathematical Physics*, 223(2):289–326, Oct. 2001. ISSN 0010-3616, 1432-0916. doi: 10.1007/s002200100540. URL `http://arxiv.org/abs/gr-qc/0103074`. arXiv:gr-qc/0103074.

[8] S. Hollands and R. M. Wald. Quantum fields in curved spacetime. *Phys. Rept.*, 574:1–35, 2015. doi: 10.1016/j.physrep.2015.02.001. _eprint: 1401.2026.

[9] W. E. Lamb and R. C. Retherford. Fine Structure of the Hydrogen Atom by a Microwave Method. *Phys. Rev.*, 72:241–243, 1947. doi: 10.1103/PhysRev.72.241.

[10] P. J. Mohr, G. Plunien, and G. Soff. QED corrections in heavy atoms. *Phys. Rept.*, 293(5-6):227–369, 1998. doi: 10.1016/s0370-1573(97)00046-x.

[11] J. Schlemmer and J. Zahn. The current density in quantum electrodynamics in external potentials. *Annals Phys.*, 359:31–45, 2015. doi: 10.1016/j.aop.2015.04.006. _eprint: 1501.05912.

[12] J. S. Schwinger. On gauge invariance and vacuum polarization. *Phys. Rev.*, 82:664–679, 1951. doi: 10.1103/PhysRev.82.664.

[13] J. C. Thompson and E. Winstanley. Quantum-corrected anti–de Sitter spacetime. *Physical Review D*, 110(12):125003, Dec. 2024. ISSN 2470-0010, 2470-0029. doi: 10.1103/PhysRevD.110.125003. URL `https://link.aps.org/doi/10.1103/PhysRevD.110.125003`.

[14] E. A. Uehling. Polarization effects in the positron theory. *Phys. Rev.*, 48:55–63, 1935. doi: 10.1103/PhysRev.48.55.

[15] J. Wernersson and J. Zahn. Vacuum polarization near boundaries. *Phys. Rev. D*, 103(1):016012, 2021. doi: 10.1103/PhysRevD.103.016012. _eprint: 2010.05499.