

Boundaries in finite gauge theory

Ilka Brunner Nils Carqueville Domenico Fiorenza

Contents

1 Introduction

Let

- G be a finite group,
- $[\omega] \in H^n(G, \mathbb{C}^\times)$,
- and \mathcal{C} a symmetric monoidal (∞, n) -category for some $n \in \mathbb{Z}_+$.

On the one hand, [?] sketches the construction of a fully extended TQFT

$$\mathrm{Bord}_n \xrightarrow[\text{“classical”}]{I} \mathrm{Fam}_n(\mathcal{C}) \xrightarrow[\text{“quantum”}]{\mathrm{Sum}_n} \mathcal{C} \quad (1.1)$$

based on the data (G, ω) , which is viewed as a fully extended version of Dijkgraaf–Witten theory. On the other hand, in [?] a class of boundary conditions for (classical!?) DW theory is proposed in terms of group extensions.

Goal. We want to reformulate the general aspects of the work [?] in the setting of fully extended TQFT. More precisely, we view the bulk theories of [?] as the classical part I in (??) with $\mathcal{C} = \mathbf{B}^n \mathbb{C}^\times$ (the n -fold delooping of the group \mathbb{C}^\times) and then

- (i) identify *classical* boundary conditions as 1-morphisms with source $*$ in $\mathrm{Fam}_n(\mathbf{B}^n \mathbb{C}^\times)$,
- (ii) subsume the boundary conditions of [?] as special cases,
- (iii) map classical boundary conditions to quantum ones via Sum_n , and connect them to the known quantum boundary conditions, following the work of Ostrik and Fuchs–Schweigert–Valentino in the case $n = 3$,

(iv) do the above not only for framed, but also for oriented, spin, ... TQFTs.

Since [?] is very light on details, we first will work out explicitly how (??) recovers the known bulk theory for $n \in \{1, 2\}$ and hopefully also $n = 3$. We also want to explain in detail how (G, ω) gives rise to an object in $\text{Fam}_n(\mathbf{B}^n \mathbb{C}^\times)$, and how group extensions $1 \rightarrow K \rightarrow H \xrightarrow{r} G \rightarrow 1$ give rise to 1-morphisms $\text{Fam}_n(\mathbf{B}^n \mathbb{C}^\times)(*, \mathbf{B}G)$.

(A more conceptual approach to boundary conditions and defects would be to start with a bordism category “with singularities” [?, Sect. 4.3] instead of Bord_n , but we leave that for another project.)

2 Special objects and 1-morphisms in $\text{Fam}_n(\mathbf{B}^n \mathbb{C}^\times)$

TODO:

- spell out definition of $\text{Fam}_n(\mathcal{C})$
- check that n -functor $\mathbf{B}G \rightarrow \mathbf{B}^n \mathbb{C}^\times$ is precisely an n -cocycle
Issue: For $n > 3$, what is a weak n -functor, i.e. precisely what data and constraints are needed?
Idea: Since both n -categories in $\mathbf{B}G \rightarrow \mathbf{B}^n \mathbb{C}^\times$ are close to trivial, most data of the functor will be trivial, and the constraints (whatever they are) will be trivially satisfied. For $n \in \{1, 2, 3\}$ this is true, see Section ??.
- clarify precisely how $\mathbf{B}G \rightarrow \mathbf{B}^n \mathbb{C}^\times$ induces an n -functor $\mathbf{B}G \rightarrow n\text{-Vect}$, at least for $n \in \{1, 2, 3\}$
- check whether natural transformation from trivial n -functor to $\omega \circ r$ is trivialisation of $r^* \omega$
Issue: induced from issue in second item

3 Bulk theory

3.1 $n = 1$

Since we take the action of G on \mathbb{C}^\times to be trivial, the cocycle

$$\omega = [\omega] \in H^1(G, \mathbb{C}^\times) = \text{Hom}_{\text{Grp}}(G, \mathbb{C}^\times) \quad (3.1)$$

is just a group homomorphism. We consider $\mathbf{B}^1 \mathbb{C}^\times$ as a subcategory of $\text{Vect}_{\mathbb{C}}$, and we will use the pair (G, ω) to construct a functor

$$\text{Bord}_1 \xrightarrow{I} \text{Fam}_1(\text{Vect}_{\mathbb{C}}) \xrightarrow{\text{Sum}_1} \text{Vect}_{\mathbb{C}} . \quad (3.2)$$

According to the cobordism hypothesis, the composite $\text{Sum}_1 \circ I$ is determined by its value on the point pt_+ , so we only need to specify $I(\text{pt}_+)$ and then compute $\text{Sum}_1(I(\text{pt}_+))$. Note that $\mathbf{B}^1 G = *//G$ is the classical action groupoid. We set $I(\text{pt}_+)$ to be the functor

$$\chi^\omega: *//G \longrightarrow \text{Vect}_{\mathbb{C}}, \quad * \longmapsto \mathbb{C}, \quad g \longmapsto \omega(g) \in \text{Aut}(\mathbb{C}) \quad (3.3)$$

for all $g \in G$. Then (??) recovers the known result reviewed in [? , Sect. 1]:

Lemma 3.1. $\text{Sum}_1(\chi^\omega) = \mathbb{C}$ if $\omega(g) = 1$ for all $g \in G$, and $\text{Sum}_1(\chi^\omega) = 0$ otherwise.

Proof. The statement is a particular case of this more general one: let (V, ρ) be a linear representation of the group G , seen as the functor

$$\chi^\rho: *//G \longrightarrow \text{Vect}_{\mathbb{C}}, \quad * \longmapsto V, \quad g \longmapsto \rho(g) \in \text{Aut}(V). \quad (3.4)$$

Then the universal cocone of χ^ρ , viewed as a diagram of shape $*//G$ in $\text{Vect}_{\mathbb{C}}$ is the pair (V_G, π) , where V_G is the vector space of coinvariants for the representation ρ , i.e.,

$$V_G = V / \langle v - \rho(g)v \rangle_{g \in G, v \in V}$$

and $\pi: V \rightarrow V_G$ is the projection to the quotient. Namely, let (W, f) be a cocone for χ^ρ , i.e., a pair consisting of a vector space W together with a linear map $f: \chi^\omega(*) = V \rightarrow W$ such that

$$\begin{array}{ccc} & W & \\ f \nearrow & & \nwarrow f \\ \chi^\omega(*) = V & \xrightarrow{\chi^\rho(g) = \rho(g)} & V = \chi^\omega(*) \end{array} \quad (3.5)$$

Then, by definition of V_G , the morphism f uniquely factors through V_G and so we have a commutative diagram

$$\begin{array}{ccc} & W & \\ & \uparrow \exists! & \\ f \nearrow & V_G & \nwarrow f \\ \pi \nearrow & & \nwarrow \pi \\ V & \xrightarrow{\rho(g)} & V \end{array} \quad (3.6)$$

showing that (V_G, π) enjoys the universal property of the universal cocone. \square

Corollary 3.2. The linear dual $\text{Hom}_{\text{Vect}_{\mathbb{C}}}(\text{Sum}_1(\chi^\omega), \mathbb{C})$ of $\text{Sum}_1(\chi^\omega)$ is naturally isomorphic to the vector space of natural transformations $\text{Hom}_{[*//G, \text{Vect}_{\mathbb{C}}]}(\chi^\omega, \chi^1)$, where $\chi^1: *//G \rightarrow \text{Vect}_{\mathbb{C}}$ is the trivial representation of G on the vector space \mathbb{C} . In other words, we have

$$\text{Hom}_{\text{Vect}_{\mathbb{C}}}(\text{Sum}_1(\chi^\omega), \mathbb{C}) \cong \left\{ \begin{array}{ccc} *//G & & \\ \downarrow \chi^\omega & \Rightarrow & \downarrow \chi^1 \\ \text{Vect}_{\mathbb{C}} & & \mathbb{C} \end{array} \right\}.$$

As a finite dimensional vector space is completely determined by its linear dual, this actually defines $\text{Sum}_1(\chi^\omega)$.

Proof. Again, the statement is true for an arbitrary linear representation (V, ρ) of the group G . The natural isomorphism

$$\text{Hom}_{\text{Vect}_{\mathbb{C}}}(V_G, \mathbb{C}) \cong \left\{ \begin{array}{ccc} *//G & & \\ \downarrow \chi^\rho & \Rightarrow & \downarrow \chi^1 \\ \text{Vect}_{\mathbb{C}} & & \mathbb{C} \end{array} \right\}$$

is then nothing but the universal property of V_G . Namely, an element in the right hand side is a morphism $f: V = \chi^\rho(*) \rightarrow \chi^1(*) = \mathbb{C}$ such that all the diagrams

$$\begin{array}{ccc} V & \xrightarrow{f} & \mathbb{C} \\ \rho(g) \downarrow & & \downarrow \text{id}_{\mathbb{C}} \\ V & \xrightarrow{f} & \mathbb{C} \end{array}$$

commute, for any $g \in G$. This is the same as requiring that all the diagrams

$$\begin{array}{ccc} & \mathbb{C} & \\ f \nearrow & & \nwarrow f \\ V & \xrightarrow{\rho(g)} & V \end{array}$$

commute, and so it is precisely the datum of a morphism $V_G \rightarrow \mathbb{C}$ by the argument in the proof of Lemma ??.

3.2 $n = 2$

TODO:

- The only non-trivial part of the 2-functor $\chi^\omega: \mathbf{B}^2G \rightarrow \mathbf{Alg}_{\mathbb{C}}$ is the coherence 2-morphism $\chi_{g,h}^\omega: \chi^\omega(g) \otimes \chi^\omega(h) \rightarrow \chi^\omega(gh)$, and the constraints (see e.g. [?, Sect. 1.1] for the definitions) on the $\chi_{g,h}^\omega$ precisely say that they are the components of a 2-cocycle. So we can take $\chi_{g,h}^\omega = \omega(g, h)$.
- compute 2-colimit $\text{Sum}_2(\chi)$ for χ as above, obtain twisted group algebra $\mathbb{C}^\omega[G]$

Lemma 3.3. $\text{Sum}_2(\chi^\omega) \cong \mathbb{C}^\omega[G]$, the twisted group algebra of G .

Proof. Let (A, M) be a cocone for χ^ω , i.e., a pair consisting of a \mathbb{C} -algebra A together with a left A -module M (representing a linear functor $\mathbb{C}\text{Mod} \rightarrow {}_A\text{Mod}$) and homotopy commutative diagrams

$$\begin{array}{ccc} & A & \\ M \nearrow & \lambda_g & \nwarrow M \\ \mathbb{C} & \xrightarrow{L_g} & \mathbb{C} \end{array}$$

where the λ_g 's are isomorphisms of left A -modules

$$\lambda_g: M \rightarrow M \otimes_{\mathbb{C}} L_g$$

such that ... (here L_g is the line associated to $g \in G$ by the 2-character ω : $L_g \otimes L_h \cong L_{gh}$ etc. ; this has to be written in detail before this proof in doing the first part of the TODO). The isomorphisms

$$\rho_{g,h}: L_g \otimes L_h \cong L_{gh}$$

endow the direct sum

$$\bigoplus_{g \in G} L_g$$


of a \mathbb{C} -algebra structure, and the isomorphisms λ_g make M a right $(\bigoplus_{g \in G} L_g)$ -module. Therefore, we can see M as a linear functor

$$(\bigoplus_{g \in G} L_g)\text{Mod} \rightarrow {}_A\text{Mod}$$

i.e., as a morphism from $(\bigoplus_{g \in G} L_g)$ to A in $2\text{-Vect}_{\mathbb{C}}$. The algebra $\bigoplus_{g \in G} L_g$ seen as a left $(\bigoplus_{g \in G} L_g)$ -module is a morphism from \mathbb{C} to $\bigoplus_{g \in G} L_g$ in $2\text{-Vect}_{\mathbb{C}}$ and the natural isomorphism

$${}_A M_{\mathbb{C}} \cong {}_A M_{(\bigoplus_{g \in G} L_g)} \otimes_{(\bigoplus_{g \in G} L_g)} (\bigoplus_{g \in G} L_g)_{\mathbb{C}}$$

gives a canonical factorization



exhibiting the algebra $\bigoplus_{g \in G} L_g$ together with itself seen as a left module over itself as the universal cocone.

Finally notice that choosing a linear basis x_g for each line L_g , the isomorphisms $\rho_{g,h}: L_g \otimes L_h \cong L_{gh}$ (and so the multiplication in $\bigoplus_{g \in G} L_g$) read

$$x_g \cdot x_h = \omega(g, h)x_{gh},$$

thus identifying the algebra $\bigoplus_{g \in G} L_g$ with the twisted group algebra $\mathbb{C}^\omega[G]$ of G . \square

3.3 $n = 3$

TODO:

- The only non-trivial part of the 3-functor $\chi^\omega: \mathbf{B}^3 G \rightarrow \mathrm{TC}_{\mathbb{C}}$ are the coherence 3-morphisms which assemble into the modification (also) called ω in [? , Def. A.4.3], and the constraints on ω (see [? , Page 219]) seem to be precisely the 3-cocycle condition.
- **Issue:** The notion of a 3-colimit (to really compute Sum_3) is scary. At least we should make some hand-wavy arguments...

4 Boundary conditions

4.1 $n = 2$

TODO

4.2 $n = 3$

TODO

References

- [] D. Freed, M. Hopkins, J. Lurie, and C. Teleman, *Topological Quantum Field Theories from Compact Lie Groups*, [[arXiv:0905.0731](#)].
- [] T. Leinster, *Basic Bicategories*, [[math/9810017](#)].
- [] J. Lurie, *On the Classification of Topological Field Theories*, *Current Developments in Mathematics* **2008** (2009), 129–280, [[arXiv:0905.0465](#)].
- [] G. Schaumann, *Duals in tricategories and in the tricategory of bimodule categories*, PhD thesis, Friedrich-Alexander-Universität Erlangen-Nürnberg (2013), [urn:nbn:de:bvb:29-opus4-37321](#).
- [] E. Witten, *The “Parity” Anomaly On An Unorientable Manifold*, [[arXiv:1605.02391](#)].
- [] J. Wang, X.-G. Wen and E. Witten, *Symmetric Gapped Interfaces of SPT and SET States: Systematic Constructions*, [[arXiv:1705.06728](#)].
- [] J. Baez and D. Wise, Quantum Gravity Seminar at University of California, Riverside, 2005, <http://math.ucr.edu/home/baez/qg-winter2005>.
- [] B. Balsam and A. Kirillov, Jr., *Turaev-Viro invariants as an extended TQFT*, [[arXiv:1004.1533](#)].
- [] M. Barkeshli, P. Bonderson, M. Cheng, and Z. Wang, *Symmetry, Defects, and Gauging of Topological Phases*, [[arXiv:1410.4540](#)].
- [] I. Brunner, N. Carqueville, and D. Plencner, *Orbifolds and topological defects*, *Comm. Math. Phys.* **315** (2012) 739–769, [[arXiv:1307.3141](#)].
- [] I. Brunner, N. Carqueville, and D. Plencner, *Discrete torsion defects*, *Comm. Math. Phys.* **337** (2015), 429–453, [[arXiv:1404.7497](#)].
- [] M. Barkeshli, C.-M. Jian, and X.-L. Qi, *Genons, twist defects, and projective non-Abelian braiding statistics*, *Phys. Rev. B* **87** (2013), 045130, [[arXiv:1208.4834](#)].
- [] J. Barrett, C. Meusburger, and G. Schaumann, *Gray categories with duals and their diagrams*, [[arXiv:1211.0529](#)].
- [] C. Bachas and M. Petropoulos, *Topological Models on the Lattice and a Remark on String Theory Cloning*, *Commun. Math. Phys.* **152** (1993), 191–202, [[hep-th/9205031](#)].

- J. Barrett and B. Westbury, *Spherical Categories*, [Adv. Math.](#) **143** (1999), 357–375, [[hep-th/9310164](#)].
- J. Barrett and B. Westbury, *Invariants of piecewise-linear 3-manifolds*, [Trans. Amer. Math. Soc.](#) **348** (1996), 3997–4022, [[hep-th/9311155](#)].
- N. Carqueville, *Lecture notes on 2-dimensional defect TQFT*, [[arXiv:1607.05747](#)].
- S. X. Cui, C. Galindo, J. Yael Plavnik, and Z. Wang, *On Gauging Symmetry of Modular Categories*, [Communications in Mathematical Physics](#) **348**:3 (2016), 1043–1064, [[arXiv:1510.03475](#)].
- N. Carqueville, C. Meusburger, and G. Schaumann, *3-dimensional defect TQFTs and their tricategories*, [[arXiv:1603.01171](#)].
- N. Carqueville and A. Quintero Vélez, *Calabi-Yau completion and orbifold equivalence*, [[arXiv:1509.00880](#)].
- N. Carqueville, A. Ros Camacho, and I. Runkel, *Orbifold equivalent potentials*, [Journal of Pure and Applied Algebra](#) **220** (2016), 759–781, [[arXiv:1311.3354](#)].
- N. Carqueville and I. Runkel, *Orbifold completion of defect bicategories*, [Quantum Topology](#) **7**:2 (2016) 203–279, [[arXiv:1210.6363](#)].
- N. Carqueville and I. Runkel, *Introductory lectures on topological quantum field theory*, [[arXiv:1705.05734](#)].
- N. Carqueville, I. Runkel, and G. Schaumann, in preparation.
- A. Davydov, L. Kong, and I. Runkel, *Field theories with defects and the centre functor*, [Mathematical Foundations of Quantum Field Theory and Perturbative String Theory](#), Proceedings of Symposia in Pure Mathematics, AMS, 2011, [[arXiv:1107.0495](#)].
- P. Etingof, D. Nikshych, and V. Ostrik, *Fusion categories and homotopy theory*, [Quantum Topology](#) **1** (2010) 209–273, [[arXiv:0909.3140](#)].
- J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, *Defect lines, dualities, and generalised orbifolds*, [Proceedings of the XVI International Congress on Mathematical Physics](#), Prague, August 3–8, 2009, [[arXiv:0909.5013](#)].
- M. Fukuma, S. Hosono, and H. Kawai, *Lattice Topological Field Theory in Two Dimensions*, [Comm. Math. Phys.](#) **161** (1994), 157–176, [[hep-th/9212154](#)].

- J. Fuchs, J. Priel, C. Schweigert, and A. Valentino, *On the Brauer groups of symmetries of abelian Dijkgraaf-Witten theories*, [Communications in Mathematical Physics](#) **339**:2 (2015), 385–405, [[arXiv:1404.6646](#)].
- J. Fuchs, I. Runkel, and C. Schweigert, *TFT construction of RCFT correlators. 3. Simple currents*, [Nucl. Phys. B](#) **694** (2004) 277–353, [[hep-th/0403157](#)].
- J. Fuchs and C. Schweigert, *Category theory for conformal boundary conditions*, [Fields Institute Communications](#) **39** (2003), 25–71, [[math/0106050](#)].
- J. Fuchs and C. Schweigert, *A note on permutation twist defects in topological bilayer phases*, [Letters in Mathematical Physics](#) **104**:11 (2014), 1385–1405, [[arXiv:1310.1329](#)].
- J. Fuchs, C. Schweigert, and A. Valentino, *Bicategories for boundary conditions and for surface defects in 3-d TFT*, [Communications in Mathematical Physics](#) **321**:2 (2013), 543–575, [[arXiv:1203.4568](#)].
- N. Gurski, *Coherence in Three-Dimensional Category Theory*, [Cambridge Tracts in Mathematics](#) **201**, Cambridge University Press, 2013.
- M. W. Hirsch, *Differential topology*, Springer Graduate Texts in Mathematics **33**, Springer, 1976.
- A. Kitaev and L. Kong, *Models for gapped boundaries and domain walls*, [Commun. Math. Phys.](#) **313** (2012) 351–373, [[arXiv:1104.5047](#)].
- A. Kapustin and L. Rozansky, *Three-dimensional topological field theory and symplectic algebraic geometry II*, [Communications of Number Theory and Physics](#) **4** (2010), 463–549, [[arXiv:0909.3643](#)].
- A. Kapustin, L. Rozansky, and N. Saulina, *Three-dimensional topological field theory and symplectic algebraic geometry I*, [Nuclear Physics B](#) **816** (2009), 295–355, [[arXiv:0810.5415](#)].
- A. Kapustin and N. Saulina, *Surface operators in 3d Topological Field Theory and 2d Rational Conformal Field Theory*, [Mathematical Foundations of Quantum Field Theory and Perturbative String Theory, Proceedings of Symposia in Pure Mathematics](#) **83**, 175–198, American Mathematical Society, 2011, [[arXiv:1012.0911](#)].
- R. J. Lawrence, *An Introduction to Topological Field Theory*, [Proc. Symp. Appl. Math.](#), **51** (1996), 89–128.

- S. Morrison and K. Walker, *Blob homology*, [Geometry & Topology](#) **16** (2012), 1481–1607, [[arXiv:1009.5025](#)].
- J. R. Munkres, *Elementary Differential Topology*, Annals of Mathematics Studies **54**, Princeton University Press, 1967.
- S.-H. Ng and P. Schauenburg, *Higher Frobenius-Schur indicators for pivotal categories*, [Contemporary Mathematics](#) **441** (2007), 63–90, [[math.QA/0503167](#)].
- U. Pachner, *P.L. Homeomorphic Manifolds are Equivalent by Elementary Shellings*, [European Journal of Combinatorics](#) **12:2** (1991), 129–145.
- F. Quinn, *Lectures on axiomatic topological quantum field theory*, IAS/Park City Mathematics Series **1** (1995), 325–433.
- G. Schaumann, *Duals in tricategories and in the tricategory of bimodule categories*, PhD thesis, Friedrich-Alexander-Universität Erlangen-Nürnberg (2013), [urn:nbn:de:bvb:29-opus4-37321](#).
- C. Schweigert and L. Woike, *Orbifold Construction for Topological Field Theories*, [[arXiv:1705.05171](#)].
- V. Turaev and O. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, [Topology](#) **31:4** (1992), 865–902.