

Standard analysis

S. A. Polymath

ABSTRACT. Let \mathcal{L} be a first-order two-sorted language. Let I be some fixed structure. A *standard* structure is an \mathcal{L} -structure of the form $\langle M, I \rangle$ where I is fixed. When I is a compact topological space (and \mathcal{L} meets a few requirements) it is possible to adapt a significant part of model theory to the restricted class of standard structures. This has been demonstrated by Henson and Iovino for Banach spaces (see, e.g. [HI]) and has been generalized to arbitrary structures in [CLCL]. The starting point is to prove that every standard structure has a *positive* elementary extension that is standard and realizes all positive types that are finitely consistent. The main tool is the notion of approximation of a positive formula and of its negation. These have been introduced by Henson and Iovino.

We review and elaborate on the properties of positive formulas and their approximations. In parallel, we introduce *continuous* formulas which are convenient to discuss examples and provide a better counterpart to Henson and Iovino theory and/or real-valued model theory. Finally, we discuss criteria for elimination of quantifiers, omitting types, ω -categoricity, stability, and indiscernibles within the setting of standard structures.

1. Standard structures

We refer to the introduction of [CLCL] for motivations. Aside from that, this paper is reasonably self-contained. In the first part we revisit some results of [CLCL]. We rephrase some theorems and add a few remarks and examples¹. Our notation and terminology diverges slightly, only where necessary.

Let I be some fixed first-order structure which is endowed with a Hausdorff compact topology (in particular, a normal topology). The language \mathcal{L}_I contains a relation symbol for each compact subsets $C \subseteq I^n$ and a function symbol for each continuous functions $f : I^n \rightarrow I$. In particular, there is a constant for each element of I . According to the context, C and f denote either the symbols of \mathcal{L}_I or their interpretation in the structure I . Such a language \mathcal{L}_I is much larger than necessary but it is convenient because it uniquely associates a structure to a topological space. The notion of a *dense set of formulas* (Definition 8) helps to reduce the size of the language when required.

The most straightforward examples of structures I are the unit interval $[0, 1]$ with the usual topology, and its homeomorphic copies $\mathbb{R}^+ \cup \{0, \infty\}$ and $\mathbb{R} \cup \{\pm\infty\}$.

We also fix a first-order language \mathcal{L}_H which we call the language of the **home sort**.

¹We will omit Section 4 from the final version of this paper and refer to [CLCL] instead. We will also omit a few proofs from Sections 3, 5, and 7.

Definition 1. Let \mathcal{L} be a two sorted language that expands both \mathcal{L}_H and \mathcal{L}_I . A **standard structure** is a two-sorted \mathcal{L} -structure of the form $\langle M, I \rangle$, where M is any \mathcal{L}_H -structure, while I is the structure above. We write **H** and **I** to denote the two sorts of \mathcal{L} .

The language \mathcal{L} adds a relation symbol $r_\varphi(x)$ for every $\varphi(x) \in \mathcal{L}_H$. All \mathcal{L} -structures are assumed to model $r_\varphi(x) \leftrightarrow \varphi(x)$. In other words, the Morleyzation of \mathcal{L}_H is assumed in the definition of \mathcal{L} -structure.

Finally and most relevantly, \mathcal{L} contains arbitrarily many function symbols of sort $H^n \rightarrow I$.

Standard structures are denoted by the domain of their home sort.

Clearly, saturated \mathcal{L} -structures exist but, with the exception of trivial cases (when I is finite), they are not standard. As a remedy, below we carve out a set of formulas \mathcal{L}^P , the set of positive formulas, such that every model has an positive elementary saturated extension that is also standard.

As usual, \mathcal{L}_I , \mathcal{L}_H , and \mathcal{L} denote both first-order languages and the corresponding set of formulas. We write \mathcal{L}_x when we restrict variables to x . If $\varphi(x) \in \mathcal{L}$ contains only terms of sort $H^n \rightarrow H$ and relations of sort H^n we will improperly say that $\varphi(x) \in \mathcal{L}_H$ (this is only correct up to equivalence).

Note that \mathcal{L} has two types of atomic formulas:

1. atomic formulas in \mathcal{L}_H (hence, up to equivalence, all formulas of \mathcal{L}_H);
2. formulas of the form $\tau(x; \eta) \in C$, where $C \subseteq I^n$ is compact and $\tau(x; \eta)$ is a tuple of terms of sort $H^{|x|} \times I^{|\eta|} \rightarrow I$.

Definition 2. A formula in \mathcal{L} is **positive** if it uses only the Boolean connectives \wedge, \vee ; the quantifiers \forall^H, \exists^H of sort H ; and the quantifiers \forall^I, \exists^I of sort I .

The set of positive formulas is denoted by \mathcal{L}^P .

A **continuous** formula is a positive formula where only atomic formulas as in (2) above occur. The set of continuous formulas is denoted by \mathcal{L}^c .

We will use Latin letters x, y, z for variables of sort H and Greek letters η, ε for variables of sort I . Therefore we can safely drop the superscript from quantifiers if they are followed by variables.

Notation 3. When we state facts that holds both for \mathcal{L}^P and \mathcal{L}^c we write $\mathcal{L}^{P/c}$ for short.

The positive formulas of Henson and Iovino correspond to those formulas in \mathcal{L}^c that do not have quantifiers of sort I .

Working within \mathcal{L}^P simplifies the comparison with classical model theory but comes at a price. Instead, allowing quantifiers of sort I is almost for free. We will see (cf. Propositions 32 and 44) that the formulas in $\mathcal{L}^{P/c}$ are approximated by formulas in $\mathcal{L}^{P/c}$ without quantifiers of sort I .

The main fact to note about \mathcal{L}^c is that it is a language without equality of sort H . The difference between \mathcal{L}^P and \mathcal{L}^c is similar to the difference between the space of absolutely integrable functions (w.r.t. some measure μ) and the Lebesgue space $L^1(\mu)$. For the latter only equality almost everywhere is relevant. However, even when $L^1(\mu)$ is our focus of interest, it is often easier to argue about real valued functions. For a similar reason \mathcal{L}^P and \mathcal{L}^c are better studied in parallel.

The extra expressive power offered by quantifiers of sort I is convenient. For instance, it is easy to see (cf. Example 4) that \mathcal{L}^c has at least the same expressive power as real valued logic.

Example 4. Let $I = [0, 1]$, the unit interval. Let $\tau(x)$ be a term of sort $H^{|x|} \rightarrow I$. Then there is a positive formula that says $\sup_x \tau(x) = \alpha$. Indeed, consider the formula

$$x [\tau(x) \dot{-} \alpha \in \{0\}] \quad \wedge \quad \forall \varepsilon [\varepsilon \in \{0\} \vee \exists x [\alpha \dot{-} (\tau(x) + \varepsilon) \in \{0\}]]$$

which, in a more legible form, becomes

$$x [\tau(x) \leq \alpha] \quad \wedge \quad \forall \varepsilon > 0 \exists x [\alpha \leq \tau(x) + \varepsilon].$$

The following example is inspired by [HPP].

Example 5. Let \mathcal{U} be a saturated structure of signature \mathcal{L}_H . The relevant part of \mathcal{L} contains symbols for the functions $f : \mathcal{U}^n \rightarrow I$ that are continuous in the following sense: the inverse image of every compact $C \subseteq I$ is type-definable in $\mathcal{L}_H(M)$ for every $M \preceq \mathcal{U}$. It is easy to verify that $\langle \mathcal{U}, I \rangle$ is positively saturated as defined in Section 5.

The theory of normed spaces is one of the motivating examples for standard structures. The requirement for I to be compact seems to be a limitation, but is not a real one. Essentially, functional analysis and model theory are only concerned with the unit ball of normed spaces. To formalize the unit ball of a normed space as a standard structure (where I is the unit interval) is straightforward but cumbersome. Alternatively one can view a normed space as a many-sorted structure: a sort for each ball of radius $n \in \mathbb{Z}^+$. Unfortunately, this also results in a bloated formalism.

A neater formalization is possible if one accepts that some positive elementary extensions (to be defined below) of a normed space could contain vectors of infinite norm hence not be themselves proper normed spaces. However these infinities are harmless if one restricts to balls of finite radius. In fact, inside any ball of finite radius these improper normed spaces are completely standard.

Example 6. Let $I = \mathbb{R} \cup \{\pm\infty\}$. Let \mathcal{L}_H be the language of real (or complex) vector spaces. The language \mathcal{L} contains a function symbol $\|\cdot\|$ of sort $H \rightarrow I$. Normed spaces are standard structures with the natural interpretation of the language. It is easy to verify that the unit ball of a standard structure is the unit ball of a normed space. Note that the same remains true if we add to the language any continuous (equivalently, bounded) operator or functional.

We conclude this introduction by advertising a question in [CLCL]. Is it possible to extend the Positive Compactness Theorem (Theorem 20) to a larger class of languages? E.g. can we allow in \mathcal{L} (and in $\mathcal{L}^{p/c}$) function symbols of sort $H^n \times I^m \rightarrow I$ with both n and m positive? These would have natural interpretations, e.g. a group acting on a compact set I .

2. Henson-Iovino approximations

For C, C' compact subsets of I^n , we write $C' > C$ if C' is a neighborhood of C . For φ, φ' (free variables are hidden) positive formulas possibly with parameters we write $\varphi' > \varphi$ if φ' is obtained by replacing in φ each atomic formula of the form $t \in C$ with $t \in C'$, for some $C' > C$. If no such atomic formula occurs in φ , then $\varphi > \varphi$. We call φ' a **weakening** of φ . Note that $>$ is a dense transitive relation and that $\varphi \rightarrow \varphi'$ in every \mathcal{L} -structure.

We write $\tilde{\varphi} \perp \varphi$ when $\tilde{\varphi}$ is obtained by replacing each atomic formula $t \in C$ occurring in φ with $t \in \tilde{C}$ where \tilde{C} is some compact set disjoint from C . The atomic formulas in \mathcal{L}_H are replaced with their negation. Finally each connective is replaced by its dual i.e., $\vee, \wedge, \exists, \forall$ are replaced by $\wedge, \vee, \forall, \exists$, respectively. We say that $\tilde{\varphi}$ is a **strong negation** of φ . It is clear that $\tilde{\varphi} \rightarrow \neg\varphi$ in every \mathcal{L} -structure.

Lemma 7. For all positive formulas φ

1. for every $\varphi' > \varphi$ there is a formula $\tilde{\varphi} \perp \varphi$ such that $\varphi \rightarrow \neg \tilde{\varphi} \rightarrow \varphi'$;
2. for every $\tilde{\varphi} \perp \varphi$ there is a formula $\varphi' > \varphi$ such that $\varphi \rightarrow \varphi' \rightarrow \neg \tilde{\varphi}$.

Proof. If $\varphi \in \mathcal{L}_H$ the claims are obvious. Suppose φ is of the form $t \in C$. Let φ' be $t \in C'$, for some $C' > C$. Let O be an open set such that $C \subseteq O \subseteq C'$. Then $\tilde{\varphi} = (t \in I \setminus O)$ is as required by the lemma. Suppose instead that $\tilde{\varphi}$ is of the form $t \in \tilde{C}$ for some compact \tilde{C} disjoint from C . By the normality of I , there is $C' > C$ disjoint from \tilde{C} . Then $\varphi' = (t \in C')$ is as required. The lemma follows easily by induction. \square

For every type $p(x)$, we write

$$p'(x) = \{\varphi'(x) : \varphi' > \varphi \text{ for some } \varphi(x) \in p\}$$

in particular $\{\varphi(x)\}' = \{\varphi'(x) : \varphi' > \varphi\}$.

Definition 8. A set of formulas $\mathcal{F} \subseteq \mathcal{L}$ is **p/c-dense** modulo T , a theory, if for every positive/continuous $\varphi(x)$ and every $\varphi' > \varphi$, there is $\psi(x) \in \mathcal{F}$ such that $\varphi(x) \rightarrow \psi(x) \rightarrow \varphi'(x)$ holds in every standard structure that models T .

Example 9. By Lemma 7, the set of negations of positive/continuous formulas is p/c-dense.

Example 10. Let $I = [0, 1]$. The set of formulas built inductively from atomic formulas of the form $\tau \in \{0\}$ is c-dense. In fact, as every compact subset of I is the intersection of finite unions of closed intervals, the formulas built using only atomic formulas of the form $\tau \in [\alpha, \beta]$ are c-dense. Finally note that $\tau \in [\alpha, \beta]$ is equivalent to $\tau \div \beta \in \{0\} \wedge \alpha \div \tau \in \{0\}$.

Example 11. In Section 7 we prove that the set of positive/continuous formulas without quantifiers of sort I is p/c-dense.

3. Morphisms

Let M and N be two standard structures. We say that a partial map $f : M \rightarrow N$ is **p-elementary** if for every $\varphi(x) \in \mathcal{L}^p$ and every $a \in (\text{dom } f)^{|x|}$

1. $M \models \varphi(a) \Rightarrow N \models \varphi(fa)$.

In words, we say that f **preserve the truth** of the positive formulas. A p-elementary map that is total is called a **p-elementary embedding**. When the identity map $\text{id}_M : M \rightarrow N$ is an p-elementary embedding, we write $M \preceq^p N$ and say that M is an p-elementary **substructure** of N .

The discussion of c-elementarity needs some extra care because \mathcal{L}^c does not contains equality in the home sort. Therefore we postpone it till Section 10.

On the other hand, as p-elementary maps are in particular \mathcal{L}_H -elementary, they are injective. However, their inverse need not be p-elementary. In other words, the converse of the implication in (1) may not hold. Hence the following notion of elementarity which is more robust. We say that the map $f : M \rightarrow N$ is an **approximate** p-elementary if for every formula $\varphi(x) \in \mathcal{L}^p$, and every $a \in (\text{dom } f)^{|x|}$

2. $M \models \varphi(a) \Rightarrow N \models \{\varphi(fa)\}'$.

One convenient feature of approximate p-elementarity is that it suffices verify (2) for any/some p-dense set of formulas. It is clear that p-elementarity implies its approximate variant. We will see that with a slight amount of saturation also the converse holds (Proposition 22). Moreover, under full saturation, these morphisms becomes \mathcal{L} -elementarity maps (Corollary 31).

The following holds in general.

Fact 12. If $f : M \rightarrow N$ is approximate p-elementary then

$$M \models \{\varphi(a)\}' \Leftrightarrow N \models \{\varphi(fa)\}'.$$

for every $\varphi(x) \in \mathcal{L}^p$, and every $a \in (\text{dom } f)^{|x|}$.

Proof. \Rightarrow Fix $\varphi' > \varphi$ and assume $M \models \varphi'(a)$. Let $\varphi' > \varphi'' > \varphi$. By the definition of approximate elementarity, $N \models \varphi''(fa)$ and $N \models \varphi'(fa)$ follows.

\Leftarrow Assume the r.h.s. of the equivalence. Fix $\varphi' > \varphi$ and prove $M \models \varphi'(a)$. Let $\varphi' > \varphi'' > \varphi$. By Lemma 7 there is some $\tilde{\varphi} \perp \varphi''$ such that $\varphi'' \rightarrow \neg \tilde{\varphi} \rightarrow \varphi'$. Then $N \models \neg \tilde{\varphi}(fa)$ and therefore $M \models \neg \tilde{\varphi}(a)$. Then $M \models \varphi'(a)$. \square

Finally, we consider the notion of partial embeddings. Though the definition seems weaker, we show that the notion is completely classical. We say that the map $f : M \rightarrow N$ is a **partial embedding** if the implication in (1) holds for all atomic formulas $\varphi(x)$. Note that this is the same as requiring that (1) holds for all formulas without quantifiers of sort H.

Fact 13. If $f : M \rightarrow N$ is a partial embedding then

$$M \models \varphi(a) \Leftrightarrow N \models \varphi(fa)$$

for every $a \in (\text{dom } f)^{|x|}$ and every formula $\varphi(x) \in \mathcal{L}^p$ without quantifiers of sort H.

Proof. The equivalence is trivial for formulas in \mathcal{L}_H so we only consider atomic formulas of the form $t \in C$. Implication \Rightarrow holds by definition. Vice versa, if $M \models \tau(a) \notin C$ then, by normality, $M \models \tau(a) \in \tilde{C}$ for some compact \tilde{C} disjoint of C . By the definition of partial embedding, $N \models \tau(a) \in \tilde{C}$. Hence $N \models \tau(a) \notin C$. Induction is immediate. \square

4. The standard part

Our goal is to prove a compactness theorem for positive theories that only requires standard structures. To bypass lengthy routine applications of ultrafilters, ultrapowers, and ultralimits, we assume the Classical Compactness Theorem and build on that.

In this section we recall the notion of standard part of an element of the elementary extension of a compact Hausdorff topological space. Our goal is to prove Lemma 18 which in turn is required for the proof of the Positive Compactness Theorem (Theorem 20). The reader willing to accept it without proof may skip this section.

Let $\langle N, {}^*I \rangle$ be an \mathcal{L} -structure that extends \mathcal{L} -elementarily the standard structure M . Let η be a free variable of sort I. For each $\beta \in I$, we define the type

$$m_\beta(\eta) = \{\eta \in D : D \text{ compact neighborhood of } \beta\}.$$

The set of the realizations of $m_\beta(\eta)$ in *I is known to nonstandard analysts as the monad of β . The following fact is well-known.

Fact 14. For every $\alpha \in {}^*I$ there is a unique $\beta \in I$ such that ${}^*I \models m_\beta(\alpha)$.

Proof. Negate the existence of β . For every $\gamma \in I$ pick some compact neighborhood D_γ of γ , such that $*I \models \alpha \notin D_\gamma$. By compactness there is some finite $\Gamma \subseteq I$ such that D_γ , with $\gamma \in \Gamma$, cover I . By elementarity, the interpretation in $*I$ of these D_γ cover $*I$. A contradiction. The uniqueness of β follows from normality. \square

We denote by $\text{st}(\alpha)$ the unique $\beta \in I$ such that $*I \models m_\beta(\alpha)$.

Fact 15. For every $\alpha \in *I$ and every compact $C \subseteq I$

$$*I \models \alpha \in C \rightarrow \text{st}(\alpha) \in C.$$

Proof. Assume $\text{st}(\alpha) \notin C$. By normality there is a compact set D disjoint from C that is a neighborhood of $\text{st}(\alpha)$. Then $*I \models \alpha \in D \subseteq \neg C$. \square

Fact 16. For every $\alpha \in (*I)^{|\alpha|}$ and every function symbol f of sort $|\alpha| \rightarrow I$

$$*I \models \text{st}(f(\alpha)) = f(\text{st}(\alpha)).$$

Proof. By Fact 14 and the definition of $\text{st}(-)$ it suffices to prove that $*I \models f(\alpha) \in D$ for every compact neighborhood D of $f(\text{st}(\alpha))$.

Fix one such D . Then $\text{st}(\alpha) \in f^{-1}[D]$. By continuity $f^{-1}[D]$ is a compact neighborhood of $\text{st}(\alpha)$. Therefore $*I \models \alpha \in f^{-1}[D]$ and, as $I \leq *I$ we obtain $*I \models f(\alpha) \in D$. \square

The standard part of $\langle N, *I \rangle$ is the standard structure $\langle N, I \rangle$ that interprets the symbols f of sort $H^n \rightarrow I$ as the functions

$$f^N(a) = \text{st}(*f(a)) \quad \text{for all } a \in N^n,$$

where $*f$ is the interpretation of f in $\langle N, *I \rangle$. Symbols in \mathcal{L}_H maintain the same interpretation.

Fact 17. With the notation as above. Let $\tau(x; \eta)$ be a term of sort $H^{|\alpha|} \times I^{|\eta|} \rightarrow I$. Then for every $a \in M^{|\alpha|}$ and $\alpha \in (*I)^{|\eta|}$

$$\tau^N(a; \text{st}(\alpha)) = \text{st}(*\tau(a; \alpha))$$

Proof. When a t is a function symbol of sort $H^{|\alpha|} \rightarrow I$, the claim holds by definition. When t is a function symbol of sort $I^{|\eta|} \rightarrow I$, the claim follows from Fact 16. Now, assume inductively that

$$t_i^N(a; \text{st}(\alpha)) = \text{st}(*t_i(a; \alpha))$$

holds for the terms $t_1(x; \eta), \dots, t_n(x; \eta)$ and let $t = f(t_1, \dots, t_n)$ for some function f of sort $I^n \rightarrow I$. Then the claim follows immediately from the induction hypothesis and Fact 16. \square

Lemma 18. With the notation as above. For every $\varphi(x; \eta) \in \mathcal{L}^P$, $a \in N^{|\alpha|}$ and $\alpha \in (*I)^{|\eta|}$

$$\langle N, *I \rangle \models \varphi(a; \alpha) \Rightarrow N \models \varphi(a; \text{st}(\alpha))$$

Proof. Suppose $\varphi(x; \eta)$ is \mathcal{L}^P -atomic. If $\varphi(x; \eta)$ is a formula of \mathcal{L}_H the claim is trivial. Otherwise $\varphi(x; \eta)$ has the form $\tau(x; \eta) \in C$. Assume that the tuple $\tau(x; \eta)$ consists of a single term. The general case follows easily from this special case. Assume that $\langle N, *I \rangle \models \tau(a; \alpha) \in C$. Then $\text{st}(*\tau(a; \alpha)) \in C$ by Fact 15. Therefore $\tau^N(a; \text{st}(\alpha)) \in C$ follows from Fact 17. This proves the lemma for atomic formulas. Induction is immediate. \square

Corollary 19. Let M be a standard structure. Let $p(\eta) \subseteq \mathcal{L}^P(M)$ be a type that does not contain existential quantifiers of sort H . Then, if $p(\eta)$ is finitely consistent in M , it is realized in M .

The corollary has also a direct proof. This goes through the observation that the formulas in $p(\eta)$ define compact subsets of I .

Proof. Let $\langle {}^*M, {}^*I \rangle$ be an \mathcal{L} -elementary saturated superstructure of $\langle M, I \rangle$. Then $\langle {}^*M, {}^*I \rangle \models p(\alpha)$ for some $\alpha \in {}^*I$. By the lemma above, ${}^*M \models p(\text{st}(\alpha))$. Now observe that the truth of formulas without existential quantifiers of sort H is preserved by substructures. \square

The exclusion of existential quantifiers of sort H is necessary. For a counterexample take $M = I = [0, 1]$. Assume that \mathcal{L} contains a function symbol for the identity map $\iota : M \rightarrow I$. Let $p(\eta)$ contain the formulas $\exists x (x > 0 \wedge \iota x + \eta \in [0, 1/n])$ for all positive integers n .

5. Compactness

It is convenient to distinguish between consistency with respect to standard structures and consistency with respect to \mathcal{L} -structures. We say that a theory T is **\mathcal{L} -consistent** if $\langle M, {}^*I \rangle \models T$ for some \mathcal{L} -structure $\langle M, {}^*I \rangle$ (this is the classical notion of consistency). We say that T is **standardly consistent** if $M \models T$ is for some standard structure M . By Lemma 18 these two notions coincide if T is positive. Therefore we have the following.

Theorem 20 (Positive Compactness Theorem). Let T be a positive theory. Then, if T is finitely \mathcal{L} -consistent, it is also standardly consistent.

An \mathcal{L} -structure N is **positively λ -saturated** if it realizes all types $p(x; \eta) \subseteq \mathcal{L}^P(N)$ with fewer than λ parameters that are finitely consistent in N . When $\lambda = |N|$ we simply say **p-saturated**. The existence of p-saturated standard structures is obtained from the classical case just as for Theorem 20.

Theorem 21. Every standard structure has a p-elementary extension to a p-saturated standard structure (possibly of inaccessible cardinality).

Proof. Let $\langle {}^*M, {}^*I \rangle$ be an \mathcal{L} -saturated \mathcal{L} -elementary extension of the standard structure $\langle M, I \rangle$. Let $\langle {}^*M, I \rangle$ be its standard part as defined in Section 4. By Lemma 18, $\langle {}^*M, I \rangle$ realizes all finitely consistent positive types with fewer than $|{}^*M|$ parameters. \square

The following proposition shows that a slight amount of saturation tames the positive formulas.

Proposition 22. Let N be a positively ω -saturated standard structure. Then

$$\{\varphi(x; \eta)\}' \leftrightarrow \varphi(x; \eta)$$

holds in N for every formula $\varphi(x; \eta) \in \mathcal{L}^P(N)$.

Proof. We prove \rightarrow , the non trivial implication. The claim is clear for atomic formulas. Induction for conjunction, disjunction and the universal quantifiers is immediate. We consider case of the existential quantifiers of sort H . Assume inductively

$$\text{ih.} \quad \{\varphi(x, z; \eta)\}' \rightarrow \varphi(x, z; \eta)$$

We need to prove

$$\{\exists z \varphi(x, z; \eta)\}' \rightarrow \exists z \varphi(x, z; \eta)$$

From (ih) we have

$$\exists z \{\varphi(x, z; \eta)\}' \rightarrow \exists z \varphi(x, z; \eta)$$

Therefore it suffices to prove

$$\{\exists z \varphi(x, z; \eta)\}' \rightarrow \exists z \{\varphi(x, z; \eta)\}'$$

Replace the variables $x; \eta$ with parameters, say $a; \alpha$, and assume that $N \models \exists z \varphi'(a, z; \alpha)$ for every $\varphi' > \varphi$. We need to prove the consistency of the type $\{\varphi'(a, z; \alpha) : \varphi' > \varphi\}$. By saturation, finite consistency suffices. This is clear if we show that the antecedent is closed under conjunction. Indeed it is easy to verify that if $\varphi_1, \varphi_2 > \varphi$ then $\varphi_1 \wedge \varphi_2 > \varphi'$ for some $\varphi' > \varphi$. In words, the set of approximations of φ is a directed set.

For existential quantifiers of sort I we argue similarly. \square

Remark 23. By Corollary 19, when $\varphi(x; \eta)$ does not contains existential quantifiers of sort H, the proposition above does not require the assumption of saturation. In general, some saturation is necessary: consider the model presented after Corollary 19 and the formula $\exists x (x > 0 \wedge \iota(x) \in \{0\})$.

Remark 24. A consequence of Proposition 22 is that the approximate morphisms defined in Section 3 coincide with their unapproximate version when the codomain is positively ω -saturated. In particular they are invertible by Fact 12.

6. The monster model

We denote by \mathcal{U} some large p-saturated standard structure which we call the **positive monster model**. Truth is evaluated in \mathcal{U} unless otherwise is specified. We denote by T the positive theory of \mathcal{U} . The density of a set of formulas is understood modulo T . Below we say **p-model** for p-elementary substructure of \mathcal{U} . We stress once again that the truth of some $\varphi \in \mathcal{L}^p(M)$ in a p-model M implies the truth of φ (in \mathcal{U}) but not vice versa. However, all p-models agree on the approximated truth, see Fact 12 and Remark 24,

$$M \models \{\varphi\}' \Leftrightarrow \{\varphi\}'.$$

A substructure $M \subseteq \mathcal{U}$ is a **c-model** if

$$c. \quad M \models \varphi(a) \Rightarrow \varphi(a) \quad \text{for all formulas } \varphi(x) \in \mathcal{L}^c \text{ and } a \in M^{|x|}.$$

Note that \mathcal{L}^c is closed under weakening and strong negation. Therefore the argument in the proof of Fact 12 proves that (c) is equivalent to

$$M \models \{\varphi(a)\}' \Leftrightarrow \{\varphi(a)\}' \quad \text{for all formulas } \varphi(x) \in \mathcal{L}^c \text{ and } a \in M^{|x|}.$$

The following fact demonstrates how positive compactness applies. There are some subtle differences from the classical setting. Let $A \subseteq \mathcal{U}$ be a small set throughout this section.

Fact 25. Let $p(x) \subseteq \mathcal{L}^p(A)$ be a type. Then for every $\varphi(x) \in \mathcal{L}^p(\mathcal{U})$

- i. if $p(x) \rightarrow \neg \varphi(x)$ then $\psi(x) \rightarrow \neg \varphi(x)$ for some $\psi(x)$ conjunction of formulas in $p(x)$;
- ii. if $p(x) \rightarrow \varphi(x)$ and $\varphi' > \varphi$ then $\psi(x) \rightarrow \varphi'(x)$ for some conjunction of formulas in $p(x)$.

Proof. (i) is immediate by saturation; (ii) follows from (i) by Lemma 7. \square

Fact 26. Let \mathcal{F} be a p/c-dense set of positive formulas. Then \mathcal{F}' is p/c-dense.

Proof. Let $\varphi' > \varphi$. Pick φ'' such that $\varphi' > \varphi'' > \varphi$ and $\psi \in \mathcal{F}$ such that $\varphi(x) \rightarrow \psi(x) \rightarrow \varphi''(x)$. It suffices to prove that $\psi'(x) \rightarrow \varphi'(x)$ for some $\psi' > \psi$. By Proposition 22, $\psi(x) \leftrightarrow \{\psi(x)\}'$. Therefore, $\psi'(x) \rightarrow \varphi'(x)$ follows from Fact 25. \square

We write $S_x^{\text{p/c}}(A)$ for the set of maximally consistent subsets of $\mathcal{L}_x^{\text{p/c}}(A)$. We define $\text{p/c-tp}(a/A)$, the positive/continuous type of a over A , to be the set of formulas $\{\varphi(x) \in \mathcal{L}^{\text{p/c}}(A) : \varphi(a)\}$.

In general, if \mathcal{F} is any set of formulas, we write $\mathcal{F}\text{-tp}(a/A)$, for the type $\{\varphi(x) \in \mathcal{F}(A) : \varphi(a)\}$. The undecorated symbol $\text{tp}(a/A)$ denotes the \mathcal{L} -type.

Fact 27. Let $p(x) \subseteq \mathcal{L}^{\text{p/c}}(A)$. The following are equivalent

1. $p(x)$ is a maximally standardly consistent subsets of $\mathcal{L}_x^{\text{p/c}}(A)$;
2. $p(x) = \text{p/c-tp}(a/A)$ for some $a \in \mathcal{U}^{|x|}$.

Proof. (1 \Rightarrow 2) If $p(x)$ is consistent then $p(x) \subseteq \text{p/c-tp}(a/A)$ for some $a \in \mathcal{U}^{|x|}$. By maximality $p(x) = \text{p/c-tp}(a/A)$.

(2 \Rightarrow 1) By Proposition 22 and Lemma 7

$$\varphi(x) \leftrightarrow \bigwedge_{\tilde{\varphi} \perp \varphi} \neg \tilde{\varphi}(x).$$

Suppose $\varphi(x) \in \mathcal{L}^{\text{p/c}}(A) \setminus p$. Then $\neg \varphi(a)$. Hence $\tilde{\varphi}(a)$ holds for some $\tilde{\varphi} \perp \varphi$ and $p(x) \rightarrow \neg \varphi(x)$ follows. \square

The fact above extends as follows.

Fact 28. Let $p(x) \subseteq \mathcal{F}(A)$, where \mathcal{F} is a p/c-dense set of positive/continuous formulas. Then the following are equivalent

1. $p(x)$ is a maximally standardly consistent subset of $\mathcal{F}_x(A)$;
2. $p(x) = \mathcal{F}\text{-tp}(a/A)$ for some $a \in \mathcal{U}^{|x|}$;
3. $p(x) \leftrightarrow \text{p/c-tp}(a/A)$ for some $a \in \mathcal{U}^{|x|}$.

Proof. (1 \Rightarrow 2) As above.

(2 \Rightarrow 3) As $q(x) \leftrightarrow q'(x)$, then $q(x) \leftrightarrow p(x) = \mathcal{F}\text{-tp}(a/A)$.

(3 \Rightarrow 1) Immediate. \square

Proposition 29. Let \mathcal{F} be a p/c-dense set of positive/continuous formulas. Then for every formula $\varphi(x) \in \mathcal{L}^{\text{p/c}}$

- i. $\neg \varphi(x) \leftrightarrow \bigvee \{ \psi(x) : \psi'(x) \rightarrow \neg \varphi(x) \text{ and } \psi' > \psi \text{ for some } \psi(x) \in \mathcal{F} \};$
- ii. $\neg \varphi(x) \leftrightarrow \bigvee \{ \neg \psi(x) : \neg \psi(x) \rightarrow \neg \varphi(x) \text{ and } \psi(x) \in \mathcal{F} \}.$

Proof. (i) Only \rightarrow requires a proof. Let $a \in \mathcal{U}^{|x|}$ be such that $\neg \varphi(a)$. Let $p(x) = \text{p/c-tp}(a)$. By Fact 27, $p(x) \rightarrow \neg \varphi(x)$. As \mathcal{F} is p/c-dense, $p(x) \leftrightarrow q(x) = \mathcal{F}\text{-tp}(a)$. Then $q'(x) \rightarrow \neg \varphi(x)$ hence, by compactness, $\psi'(x) \rightarrow \neg \varphi(x)$ for some $\psi(x) \in \mathcal{F}$.

(ii) By density

$$\varphi(x) \leftrightarrow \bigwedge \{ \psi(x) \in \mathcal{F} : \varphi(x) \rightarrow \psi(x) \}.$$

Negating both sides of the equivalence we obtain (ii). \square

7. Elimination of quantifiers of sort I for positive formulas

We write $\mathcal{L}_{\text{lqf}}^{\text{p/c}}$ for the set of positive/continuous formulas without quantifiers of sort I.

In this section we show that $\mathcal{L}_{\text{lqf}}^{\text{p}}$ is p-dense modulo T (the theory of \mathcal{U}). In plain words, this amounts to the elimination of the quantifiers of sort I up to some approximation. In Section 10 we prove the analogous result for $\mathcal{L}_{\text{lqf}}^{\text{c}}$.

A partial map is $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -elementary if it preserves the truth of formulas in $\mathcal{L}_{\text{lqf}}^{\text{p}}$.

Proposition 30. Every $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -elementary map $f : \mathcal{U} \rightarrow \mathcal{U}$ of small cardinality extends to an automorphism of \mathcal{U} .

Proof. By Fact 27, the inverse of an $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -elementary map is also $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -elementary. Then we can extend the map by back-and-forth as usual. \square

Let $A \subseteq \mathcal{U}$ be a small set throughout this section. We apply the proposition above to obtain a strengthening of Fact 28. For $a, b \in \mathcal{U}^{|x|}$ we write $a \equiv_A b$ if a and b satisfy the same \mathcal{L} -formulas over A .

Corollary 31. Let $p(x) = \mathcal{L}_{\text{lqf}}^{\text{p}}\text{-tp}(a/A)$ and $q(x) = \text{tp}(a/A)$. Then $p(x) \leftrightarrow q(x)$.

Proof. Only \rightarrow requires a proof. If $b \models p(x)$ then there is an $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -elementary map $f \supseteq \text{id}_A$ such that $f a = b$. Then f extends to an automorphism. As every automorphism is \mathcal{L} -elementary, $a \equiv_A b$, and the corollary follows. \square

The corollary says that \mathcal{U} does not distinguish between \mathcal{L} , \mathcal{L}^{p} , and $\mathcal{L}_{\text{lqf}}^{\text{p}}$ types. In the classical setting, from an equivalence between types one derives an equivalence between formulas. Without negation, this is not true. Still, we can infer an approximate form of equivalence.

Proposition 32. The set $\mathcal{L}_{\text{lqf}}^{\text{p}}$ is p-dense modulo T .

Proof. Let $\varphi(x)$ be a positive formula. We need to prove that for every $\varphi' > \varphi$ there is some formula $\psi(x) \in \mathcal{L}_{\text{lqf}}^{\text{p}}$ such that $\varphi(x) \rightarrow \psi(x) \rightarrow \varphi'(x)$. By Corollary 31 and Proposition 22

$$\neg\varphi(x) \rightarrow \bigvee_{p'(x) \rightarrow \neg\varphi(x)} p'(x)$$

where $p(x)$ ranges over the maximally consistent $\mathcal{L}_{\text{lqf}}^{\text{p}}$ -types. By Fact 25 and Lemma 7

$$\neg\varphi(x) \rightarrow \bigvee_{\neg\tilde{\psi}(x) \rightarrow \neg\varphi(x)} \neg\tilde{\psi}(x),$$

where $\tilde{\psi}(x) \in \mathcal{L}_{\text{lqf}}^{\text{p}}$. Equivalently,

$$\varphi(x) \leftarrow \bigwedge_{\tilde{\psi}(x) \leftarrow \varphi(x)} \tilde{\psi}(x).$$

By compactness, see Fact 25, for every $\varphi' > \varphi$ there are some finitely many $\tilde{\psi}_i(x) \in \mathcal{L}_{\text{lqf}}^{\text{p}}$ such that

$$\varphi'(x) \leftarrow \bigwedge_{i=1, \dots, n} \tilde{\psi}_i(x) \leftarrow \varphi(x)$$

which yields the interpolant required by the proposition. \square

8. Cauchy completions

For $\tau(x, z) = \tau_1(x, z), \dots, \tau_n(x, z)$ a tuple of terms of sort $H^{|x|+|z|} \rightarrow I$ we define the formula

$$x \sim_\tau y = \bigwedge_{i=1}^n \forall z \ \tau_i(x, z) = \tau_i(y, z),$$

where the expression $\alpha = \beta$ is shorthand for $\langle \alpha, \beta \rangle \in \Delta$, where Δ is the diagonal of I^2 . We also define the type

$$x \sim y = \{x \sim_\tau y : \tau(x, z) \text{ as above}\}.$$

The following fact will be used below without mention.

Fact 33. For any $a = \langle a_i : i < \lambda \rangle$ and $b = \langle b_i : i < \lambda \rangle$

$$a \sim b \Leftrightarrow a_i \sim b_i \text{ for every } i < \lambda.$$

Proof. By induction. Only implication \Leftarrow requires a proof. For limit ordinals induction is trivial. Assume $a_i \sim b_i$ for every $i \leq \lambda$. Let $|a| = |b| = \lambda$ and assume inductively that (note that it suffices to consider tuples τ of arity 1)

$$\forall y, z \ \tau(a, y, z) = \tau(b, y, z)$$

holds for every term $\tau(x, y, z)$, with $|x| = \lambda$ and $|y| = 1$ (universal quantification over the free variables is understood throughout the proof). In particular for any a_λ

$$1. \quad \forall z \ \tau(a, a_\lambda, z) = \tau(b, a_\lambda, z).$$

As $a_\lambda \sim b_\lambda$ then

$$\forall x, z \ \tau(x, a_\lambda, z) = \tau(x, b_\lambda, z)$$

and in particular

$$2. \quad \forall z \ \tau(b, a_\lambda, z) = \tau(b, b_\lambda, z).$$

From (1) and (2) we obtain

$$\forall z \ \tau(a, a_\lambda, z) = \tau(b, b_\lambda, z). \quad \square$$

Note that the approximations of the formula $x \sim_\tau y$ have the form

$$x \sim_{\tau, D} y = \bigwedge_{i=1}^n \forall z \ \langle \tau_i(x, z), \tau_i(y, z) \rangle \in D$$

for some compact neighborhood D of Δ . We write \mathcal{E} for the set containing the pairs τ, D as above. The formulas $x \sim_\varepsilon y$ form a prebase for a system of entougages on $\mathcal{U}^{|x|}$. We refer to this uniformity and the topology associated as the topology induced by I . Though not needed in the sequel, it is worth mentioning that the topology induced by I on $\mathcal{U}^{|x|}$ coincides with the product of the topology induced by I on \mathcal{U} . This can be verified by an argument similar to the proof of Fact 33.

Fact 34. For every $\varphi(x) \in \mathcal{L}^c$

$$x \sim y \rightarrow \varphi(x) \leftrightarrow \varphi(y)$$

Proof. By induction. □

We define

$$x \sim' y = \{x \sim_\varepsilon y : \varepsilon \in \mathcal{E}\}.$$

The following corollary corresponds to the Perturbation Lemma [HI, Proposition 5.15].

Corollary 35. For every $\varphi(x) \in \mathcal{L}^c$, every $\varphi' > \varphi$, and every $\tilde{\varphi} \perp \varphi$ there an $\varepsilon \in \mathcal{E}$ such that

- i. $x \sim_\varepsilon y \wedge \varphi(y) \rightarrow \varphi'(x)$
- ii. $x \sim_\varepsilon y \wedge \tilde{\varphi}(y) \rightarrow \neg\varphi(x)$.

Proof. As $x \sim' y \cup \{\varphi(x)\} \rightarrow \varphi(y)$ by the fact above, (i) follows from Fact 25. Similarly, we obtain (ii) from $x \sim' y \cup \{\tilde{\varphi}(x)\} \rightarrow \neg\varphi(y)$. \square

We say what a type $q(x)$ is **finitely satisfiable** in A if every conjunction of formulas in $q(x)$ has a solution in $A^{|x|}$. This definition coincides with the classical one, but in our context, the notion is less robust. We may happen that $p(x)$ is finitely satisfiable while $q(x) \leftrightarrow p(x)$ is not. In particular if M is a p-model and $q(x) = \text{p-tp}(a/M)$ then $q'(x)$ is always finitely satisfiable while $q(x)$ need not.

We say that a set $A \subseteq \mathcal{U}$ is **Cauchy complete** if it contains all those $a \in \mathcal{U}$ such that $a \sim' x$ is finitely satisfied in A . Note that Cauchy complete sets are in particular closed under \sim -equivalence. The **Cauchy completion** of A is the set

$$\text{Ccl}(A) = \{a : a \sim' x \text{ is finitely satisfied in } A\}.$$

Fact 36. $\text{Ccl}(A)$ is Cauchy complete.

Proof. Suppose that $x \sim' a$ is finitely satisfied in $\text{Ccl}(A)$. Let $\varepsilon \in \mathcal{E}$ be given. We prove that $a \sim_\varepsilon x$ is satisfied in A . Let $\eta \in \mathcal{E}$ be such that $x \sim_\eta y \sim_\eta z \rightarrow x \sim_\varepsilon z$. There is some $b \in \text{Ccl}(A)$ such that $b \sim_\eta a$. There is some $c \in A$ such that $c \sim_\eta b$. Then c satisfies $x \sim_\varepsilon a$, as required. \square

We say that $p(x) \in \mathcal{L}^c(\mathcal{U})$ is a **Cauchy type** if it is consistent and $p(x) \wedge p(y) \rightarrow x \sim y$.

Fact 37. Let M be a c-model and let $p(x) \in \mathcal{L}^c(M)$ be a Cauchy type. Then all realizations of $p(x)$ belong to $\text{Ccl}(M)$.

Proof. If $p(x)$ is a Cauchy type, then $p(x) \rightarrow a \sim x$ for some $a \models p(x)$. As M is a c-model, $p'(x)$ is finitely satisfied in M . Then also $a \sim' x$ is finitely satisfied. Hence $a \in \text{Ccl}(M)$. \square

9. Continuous morphisms

The definition of c-elementary maps is delicate due to the lack of equality. A possible approach is to work in the quotient \mathcal{U}/\sim . A second possibility is to replace functions with relations. The two options differs only in the notation. Here we go for the second.

Let M and N be c-models. Let $R \subseteq M \times N$ be a binary relation. If $a = \langle a_i : i < \lambda \rangle$ and $b = \langle b_i : i < \lambda \rangle$ we write $\langle a, b \rangle \in R$ to abbreviate: $\langle a_i, b_i \rangle \in R$ for all $i < \lambda$. We say that R is an **c-elementary relation** between M and N if

$$M \models \varphi(a) \Rightarrow N \models \varphi(b) \quad \text{for every } \varphi(x) \in \mathcal{L}^c \text{ and every } \langle a, b \rangle \in R.$$

Below, for simplicity we only consider c-elementary relations between \mathcal{U} and itself. Then, by Fact 12 and Proposition 22, the condition above is equivalent to

$$\# \quad \varphi(a) \leftrightarrow \varphi(b) \quad \text{for every } \varphi(x) \in \mathcal{L}^c \text{ and every } \langle a, b \rangle \in R.$$

Therefore if R is c-elementary, also R^{-1} is c-elementary.

In the next section we will also need the notion of $\mathcal{L}_{\text{lqf}}^c$ -elementary relation between \mathcal{U} and itself. This is defined similarly but (#) above is required to hold for formulas in $\mathcal{L}_{\text{lqf}}^c$. We ask the reader to verify that the content of this section applies verbatim to $\mathcal{L}_{\text{lqf}}^c$ -elementary relation.

The relation (\sim) is c-elementary by Fact 34. Then, if the relation R is c-elementary, also $(\sim) \circ R \circ (\sim)$ is c-elementary. We identify a c-elementary relation $R = (\sim) \circ R \circ (\sim)$ with the map

$$\begin{aligned} R_{/\sim} : \mathcal{U}/\sim &\rightarrow \mathcal{U}/\sim \\ [a] &\mapsto [b] \end{aligned} \quad \text{for some/any } b \text{ such that } \langle a, b \rangle \in R.$$

It is easy to verify that the definition is well-given and that $R_{/\sim}$ is injective.

We say that R is **reduced** if there is no $R' \subset R \subseteq (\sim) \circ R' \circ (\sim)$. It is easy to verify that every relation R contains a reduced relation R' such that $R \subseteq (\sim) \circ R' \circ (\sim)$.

Fact 38. Let R be reduced. Then R is the graph of an injective map $f : \text{dom}(R) \rightarrow \text{range}(R)$.

Proof. Let $a \in \text{dom}R$ and assume that $\langle a, b \rangle, \langle a, b' \rangle \in R$. To prove functionality suppose for a contradiction that $b \neq b'$. As R is reduced, $b \not\sim b'$. Then $a \not\sim a$, a contradiction. This prove functionality, for injectivity apply the same argument to R^{-1} . \square

Assume $R = (\sim) \circ R \circ (\sim)$ for simplicity. Let f be a reduced relation such that $f \subseteq R = (\sim) \circ f \circ (\sim)$. Then f is a function defined in exactly one representative a of each (\sim) -equivalence class $[a] \subseteq \text{dom}(R)$. Moreover, $f(a)$ the unique representative of $R_{/\sim}([a])$ in the range of f .

Fact 39. If R is a c-elementary relation then $\text{Ccl}(R)$ is also c-elementary. We also have that $\text{Ccl}(R) = (\sim) \circ \text{Ccl}(R) \circ (\sim)$.

Proof. Let $\varphi(x)$ and $\varphi'' > \varphi' > \varphi$ be given. Let $\langle a, b \rangle \in \text{Ccl}(R)$ and assume $\varphi(a)$. Let $\varepsilon \in \mathcal{E}$ be such that

1. $x \sim_\varepsilon y \wedge \varphi(x) \rightarrow \varphi'(y)$
2. $x \sim_\varepsilon y \wedge \varphi'(x) \rightarrow \varphi''(y)$

By the definition of Cauchy completion, there are $a', b' \sim_\varepsilon a, b$ such that $\langle a', b' \rangle \in R$. Then from (1) we infer that $\varphi'(a')$. As R is c-elementary, $\varphi'(b')$. Then $\varphi''(b)$ follows from (2). As $\varphi'' > \varphi$ is arbitrary, the fact follows from Proposition 32.

The second claim is immediate. \square

Let M and N be c-models. A c-elementary relation such that $\text{dom}(R) = M$ and $\text{range}(R) = N$ is called an approximate **c-isomorphism** between M and N . The modifier *approximate* is due because R is c-elementary between \mathcal{U} and itself. We will only consider these c-isomorphisms, therefore we omit the modifier. A c-automorphism a c-isomorphism of a c-model in itself. The following is noted for future reference.

Remark 40. To be a c-automorphism of \mathcal{U} the relation R only needs to satisfy (#) for atomic formulas of the form $\tau(x) \in C$.

Fact 41. Let R be a c-elementary relation. Let $M \subseteq \text{dom}(R)$ and $N \subseteq \text{range}(R)$ be c-models. Then there is a c-isomorphism between $\text{Ccl}(M)$ and $\text{Ccl}(N)$.

Proof. By Fact 39 we can assume that $\text{Ccl}(M) \subseteq \text{dom}(R)$ and $\text{Ccl}(N) \subseteq \text{range}(R)$, hence a restriction of R is the required c-isomorphism. \square

10. Elimination of quantifiers of sort I for continuous formulas

We verify that the argument in Section 7 can be adapted to continuous formulas.

Proposition 42. All $\mathcal{L}_{\text{qf}}^c$ -elementary maps of small cardinality extend to c-automorphisms of \mathcal{U} .

Proof. We construct by back-and-forth a sequence $f_i : \mathcal{U} \rightarrow \mathcal{U}$ of finite c-elementary reduced relations (i.e. injective maps) and let

$$R = \bigcup_{i \in |\mathcal{U}|} (\sim) \circ f_i \circ (\sim).$$

We will ensure that $\mathcal{U} = \text{dom}(R) = \text{range}(R)$. Then R is a c-automorphism of \mathcal{U} by Remark 40.

Let i be even. Let a be an enumeration of $\text{dom}(f_n)$. Let $b \in \mathcal{U}$. Let $p(x, y) = \text{c-tp}(a, b)$. By c-elementarity the type $p(f_i a, y)$ is finitely satisfied. Let $c \in \mathcal{U}$ realize $p(f_i a, y)$. Let f_{i+1} be a reduced relation such that $f_{i+1} \subseteq f_i \cup \{\langle b, c \rangle\} \subseteq (\sim) \circ f_{i+1} \circ (\sim)$.

When i is odd we proceed similarly with f_i^{-1} . Limit stages are obvious. \square

Corollary 43. Let $a \in \mathcal{U}^{[x]}$. Let $p(x) = \mathcal{L}_{\text{qf}}^c\text{-tp}(a)$ and $q(x) = \text{c-tp}(a)$. Then $p(x) \leftrightarrow q(x)$.

Proof. Only \rightarrow requires a proof. If a contains entries $a_i \sim a_j$, replace a_j with a_i . Note that the tuple a' obtained in this manner has the same c-type of a .

Let $b \models p(x)$ and let b' obtained with the same procedure as a' .

Then the function that maps $a' \mapsto b'$ preserves the truth of $\mathcal{L}_{\text{qf}}^c$ -formulas. Then f extends to an c-automorphism. As every c-automorphism is c-elementary and the corollary follows. \square

Proposition 44. The set $\mathcal{L}_{\text{qf}}^c$ is c-dense modulo T .

Proof. The proof of Proposition 32, where Corollary 43 replaces the reference to Corollary 31. \square

11. The Tarski-Vaught test and the Löwenheim-Skolem theorem

The following proposition is our version of the Tarski-Vaught test.

Theorem 45. Let M be a subset of \mathcal{U} . Let \mathcal{F} be a p-dense set of positive formulas. Then the following are equivalent

1. M is a p-model;
2. for every formula $\psi(x) \in \mathcal{F}(M)$

$$\exists x \psi(x) \Rightarrow \text{for every } \psi' > \psi \text{ there is an } a \in M \text{ such that } \psi'(a);$$
3. for every formula $\psi(x) \in \mathcal{F}(M)$

$$\exists x \neg \psi(x) \Rightarrow \text{there is an } a \in M \text{ such that } \neg \psi(a).$$

If \mathcal{F} is a c-dense set of continuous formulas then (2) and (3) above are equivalent to

- 1'. $\text{Ccl}(M)$ is a c-model.

Moreover, if M is a substructure, then (1'), (2) and (3) are also equivalent to

- 1''. M is a c-model.

Proof. (1 \Rightarrow 2) Assume $\exists x \psi(x)$ and let $\psi' > \psi$ be given. By Lemma 7 there is some $\tilde{\psi} \perp \psi$ such that $\psi(x) \rightarrow \neg \tilde{\psi}(x) \rightarrow \psi'(x)$. Then $\neg \forall x \tilde{\psi}(x)$ hence, by (1), $M \models \neg \forall x \tilde{\psi}(x)$. Then $M \models \neg \tilde{\psi}(a)$ for some $a \in M$. Hence $M \models \psi'(a)$ and $\psi'(a)$ follows from (1).

(2 \Rightarrow 3) Assume (2) and let $\psi(x) \in \mathcal{F}(M)$ be such that $\exists x \neg \psi(x)$. By Proposition 29.i, there are a consistent $\varphi(x) \in \mathcal{F}(M)$ and some $\varphi' > \varphi$ such that $\varphi'(x) \rightarrow \neg \psi(x)$. Then (3) follows.

(3 \Rightarrow 2) Let $\psi' > \psi$ for some $\psi(x) \in \mathcal{F}(M)$. Let $\tilde{\psi} \perp \psi$ such that $\psi(x) \rightarrow \neg \tilde{\psi}(x) \rightarrow \psi'(x)$. By Proposition 29.ii, $\neg \varphi(x) \rightarrow \neg \tilde{\psi}(x)$ for some $\varphi(x) \in \mathcal{F}(M)$ such that $\neg \varphi(x)$ is consistent. Then (2) follows from (3).

(2 \Rightarrow 1) Assume (2). By the classical Tarski-Vaught test $M \leq_H \mathcal{U}$. Then M is the domain of a substructure of \mathcal{U} . Then $M \models \varphi(a) \Rightarrow \varphi(a)$ holds for every atomic formula $\varphi(x)$ and for every $a \in M^{|x|}$. Now, assume inductively

$$M \models \varphi(a, b) \Rightarrow \varphi(a, b).$$

Using (2) and the induction hypothesis we prove that

$$M \models \exists y \varphi(a, y) \Rightarrow \forall y \varphi(a, y).$$

Indeed, for any $\varphi' > \varphi$,

$$\begin{aligned} M \models \exists y \varphi(a, y) &\Rightarrow M \models \exists y \psi(a, y) \quad \text{for some } \psi \in \mathcal{F} \text{ such that } \varphi(a, y) \rightarrow \psi(a, y) \rightarrow \varphi'(a, y) \\ &\Rightarrow M \models \psi(a, b) \quad \text{for some } b \in M \text{ by (2)} \\ &\Rightarrow M \models \varphi'(a, b) \\ &\Rightarrow \varphi'(a, b) \quad \text{by induction hypothesis} \\ &\Rightarrow \exists y \varphi'(a, y). \end{aligned}$$

As $\varphi' > \varphi$ is arbitrary, $\exists y \varphi(a, y)$ follows from Proposition 22.

Induction for the connectives \vee , \wedge , \forall^H , \exists , and \forall^I is straightforward.

(1' \Rightarrow 2) Let $\psi' > \psi'' > \psi$. Reasoning as in the proof of (1 \Rightarrow 2) we obtain that $\text{Ccl}(M) \models \psi''(a)$ for some $a \in \text{Ccl}(M)$. By Corollary 35, $a \sim_\varepsilon x \rightarrow \psi'(x)$ for some ε . As $a \sim' x$ is finitely satisfied in M , it follows that $\psi'(c)$ for some $c \in M$.

(2 \Leftrightarrow 3) The proof above applies verbatim when \mathcal{F} is a c-dense set of continuous formulas.

(2 \Rightarrow 1') Assume (2). We claim that $\text{Ccl}(M)$ is a substructure of \mathcal{U} . Let $a \in M^n$ and let f be a function symbol of sort $H^n \rightarrow H$. We prove that $fa \in M$. We show that $fa \sim' x$ is finitely satisfied in M . Consider the formula $fa \sim_\varepsilon x$ where ε is the pair τ, D . By Lemma 7, there is a formula in $\tilde{\varphi}(x) \in \mathcal{L}^p(M)$ such that

$$fa \sim_\tau x \rightarrow \neg \tilde{\varphi}(x) \rightarrow fa \sim_\varepsilon x$$

By Fact 26.ii there is a consistent formula $\neg\psi(x)$, for some $\psi(x) \in \mathcal{F}(M)$, that implies $fa \sim_\varepsilon x$. Then, by (3), $fa \sim_\varepsilon x$ is satisfied in M . This proves our claim.

Now, we claim that (2) holds also for every $\psi(x) \in \mathcal{F}(\text{Ccl}(M))$. Let $\psi(x, z) \in \mathcal{F}(M)$ and $\psi' > \psi$ be given. Let $b \in \text{Ccl}(M)^{|z|}$. Suppose that $\exists x \psi(x, b)$ and let ε be such that $z \sim_\varepsilon b \rightarrow \exists x \psi''(x, z)$ where $\psi' > \psi'' > \psi$. By Corollary 35, we can also assume that $z \sim_\varepsilon b \wedge \psi''(x, z) \rightarrow \psi'(x, b)$. Let $b' \in M^{|z|}$ be such that $b' \sim_\varepsilon b$. By (2) there is an $a \in M^{|x|}$ such that $\psi''(a, b')$. Then $\psi'(a, b)$ follows. This proves the second claim.

By the two claims above, the inductive argument in the proof in (2 \Rightarrow 1) applies to prove that $\text{Ccl}(M)$ is a c-model.

(1'' \Rightarrow 2) By the same argument as in (1 \Rightarrow 2).

(2 \Rightarrow 1'') As M is a substructure by assumption, the inductive argument in the proof in (2 \Rightarrow 1) applies. \square

Remark 46. The theorem above shows in particular that for every substructure M the following are equivalent

1. M is a c-model;
2. $\text{Ccl}(M)$ is a c-model.

Classically, the first application of the Tarski-Vaught test is in the proof of the downward Löwenheim-Skolem Theorem. Note that here the classical downward Löwenheim-Skolem Theorem holds in full for all \mathcal{L} -structures. In particular every $A \subseteq \mathcal{U}$ is contained in a standard structure of cardinality $|\mathcal{L}(A)|$. In this form the Löwenheim-Skolem Theorem is not very informative. In fact, the cardinality of \mathcal{L} is excessively large because of the abundance of symbols in \mathcal{L}_I .

If there is a countable p/c-dense set \mathcal{F} of positive/continuous formulas, we say that $\mathcal{L}^{p/c}$ is **separable**.

Proposition 47. Let $\mathcal{L}^{p/c}$ be separable. Let A be a countable set. Then there is a countable p/c-model M containing A .

Proof. Let \mathcal{F} be a countable p/c-dense set of positive/continuous formulas. As in the classical proof of the Löwenheim-Skolem Theorem, we construct a countable $M \subseteq \mathcal{U}$ that contains a witness of every consistent formula $\neg\psi(x)$ for $\psi(x) \in \mathcal{F}(M)$. In continuous case we also ensure that M is a substructure. Then the proposition follows from Theorem 45. \square

Proposition 48. Assume that \mathcal{L} has at most countably many symbols of sort $H^n \rightarrow I$. Then, if I is a second countable (the topology has a countable base), then \mathcal{L}^c is separable modulo \mathcal{L} . Moreover if \mathcal{L}_H is countable also then \mathcal{L}^p is separable.

Proof. Fix a countable filter of compact neighborhoods of the diagonal of I^2 . \square

12. Positive omitting types

In this section we prove a version of the omitting types theorem.

A type $p(x)$ is isolated by $\varphi(x)$, a consistent formula, if $\varphi(x) \rightarrow p(x)$. If $p(x)$ is isolated by $\neg\varphi(x)$ for some $\varphi(x) \in \mathcal{L}^p(A)$ we say that it is **p-isolated** by A . We omit reference to A when this is clear from the context (e.g. when A is the set of parameters of $p(x)$).

Note that, by Theorem 45, $p(x)$ is p-isolated by A , then $p(x)$ is realized in every p-model containing A . We also have that if $p(x)$ is realized in M , a p-model, then $p(x)$ is p-isolated by M . In fact, if $b \models p(x)$ then $\neg(b \neq x)$ isolates $p(x)$ where $b \neq x$ is positive by Morleyization.

The continuous case will be considered in the next section. In this case a weaker notion of isolation is appropriate.

Fact 49. Let \mathcal{F} be a p-dense set of positive formulas. Then the following are equivalent

1. $p(x)$ is p-isolated by A ;
2. $p(x)$ is isolated by $\neg\varphi(x)$ for some $\varphi(x) \in \mathcal{F}(A)$;
3. $p(x)$ is isolated by some $\varphi'(x)$ such that $\varphi' > \varphi$ for some consistent $\varphi(x) \in \mathcal{F}(A)$.

Proof. $(1 \Rightarrow 2)$ By Proposition 29.ii.

$(1 \Rightarrow 3)$ By Proposition 29.i.

$(3 \Rightarrow 1)$ Let $\varphi' > \varphi$ be as in (3). Let $\tilde{\varphi} \perp \varphi$ be such that $\varphi(x) \rightarrow \neg\tilde{\varphi}(x) \rightarrow \varphi'(x)$. Then $\neg\tilde{\varphi}(x)$ isolates $p(x)$. \square

Lemma 50. Let $\mathcal{L}^p(A)$ be separable. Let $p(x) \subseteq \mathcal{L}^p(A)$, be non p-isolated by A . Then every consistent formula $\neg\psi(z)$, with $\psi(z) \in \mathcal{L}^p(A)$, has a solution a such that A, a does not p-isolate $p(x)$.

Proof. We construct a sequence of $\mathcal{L}^p(A)$ -formulas $\langle \gamma_i(z) : i < \omega \rangle$ such that any realization a of the type $\{\gamma_i(z) : i < \omega\}$ is the required solution of $\neg\psi(z)$.

Let $\langle \xi_i(x, z) : i < \omega \rangle$ enumerate a countable p-dense subset of $\mathcal{L}_{x,z}^p(A)$. Let $\gamma_0(z)$ be a consistent positive formula such that $\gamma'_0(z) \rightarrow \neg\psi(z)$ for some $\gamma'_0 > \gamma_0$. This exists by Proposition 29. Now we define $\gamma_{i+1}(z)$ and $\gamma'_{i+1} > \gamma_{i+1}$. Let $\gamma_i(z)$ and $\gamma'_i > \gamma_i$ be given. Pick $\tilde{\gamma} \perp \gamma_i$ such that $\gamma_i(z) \rightarrow \neg\tilde{\gamma}(z) \rightarrow \gamma'_i(z)$.

1. If $\neg\xi_i(x, z) \wedge \gamma_i(z)$ is inconsistent, let $\gamma_{i+1} = \gamma_i$ and $\gamma'_{i+1} = \gamma'_i$.

2. Otherwise, pick $\varphi(x) \in p$ such that (#) below is consistent

$$(\#) \quad \neg\tilde{\gamma}(z) \wedge \exists x [\neg\xi_i(x, z) \wedge \neg\varphi(x)].$$

Finally, let $\gamma_{i+1}(z)$ and $\gamma'_{i+1} > \gamma_{i+1}$ consistent positive formulas such that $\gamma'_{i+1}(z)$ implies (#). Such formulas exist by Proposition 29.

Let $a \models \{\gamma_i(z) : i < \omega\}$. We claim that that A, a does not p-isolate $p(x)$. Otherwise, by Fact 49, $\neg\xi_i(x, a) \rightarrow p(x)$ for some consistent $\neg\xi_i(x, a)$. This contradicts $a \models \gamma_{i+1}(z)$.

Therefore the proof is complete if we can show that it is always possible to find the formula $\varphi(x)$ required in (2).

Suppose for a contradiction that $\neg\xi_i(x, z) \wedge \gamma_i(z)$ is consistent while (#) is inconsistent for all formulas $\varphi(x) \in p$. This immediately implies that

$$\exists z [\neg\xi_i(x, z) \wedge \neg\tilde{\gamma}(z)] \rightarrow p(x).$$

By Fact 49, $p(x)$ is p-isolated, a contradiction. \square

From the lemma we easily obtain an omitting types theorem for positive types that is very close to the classical one.

Theorem 51 (Positive Omitting Types). Let \mathcal{L}^p be separable. Let A be countable. Assume also that $p(x) \subseteq \mathcal{L}^p(A)$ is not p-isolated. Then there is a p-model M containing A that omits $p(x)$.

Proof. As in the classical proof, apply the lemma above and the Tarski-Vaught test (Theorem 45) to obtain a countable p-model M that does not isolate $p(x)$. For p-models, p-isolating a type is equivalent to realizing it. \square

13. Continuous omitting types

We say that the type $p(x) \subseteq \mathcal{L}^c(\mathcal{U})$ is **c-isolated** by A if for every $\varepsilon \in \mathcal{E}$ there is a formula $\varphi_\varepsilon(x) \in \mathcal{L}^c(A)$ such that $\neg\varphi_\varepsilon(x)$ is consistent with $p(x)$ and isolates $\exists x' \sim_\varepsilon x p(x')$.

Notice that we are not simply requiring that the type $\exists x' \sim_\varepsilon x p(x')$ is isolated: we also require that $\neg\varphi_\varepsilon(x)$ is consistent with $p(x)$. This is necessary in the proof of Lemma 56.

Assume for the rest of this section that \mathcal{L}^c is separable. Then there is a sequence $\langle \varepsilon_n : n \in \omega \rangle$ such that for every τ, D there is an n such that $x \sim_{\varepsilon_n} y \rightarrow x \sim_\tau y$. We abbreviate $x \sim_{\varepsilon_n} y$ by $x \sim_n y$. We may assume that $x \sim_n y \rightarrow x \sim_{n+1} y$. If the type $p(x)$ is c-isolated, say $\neg\varphi_n(x)$ isolates $\exists x' \sim_n x p(x')$, then we may assume that $\neg\varphi_{n+1}(x) \rightarrow \neg\varphi_n(x)$.

Fact 52. Let \mathcal{L}^c be separable. Let $p(x) \subseteq \mathcal{L}^c(A)$ be c-isolated by A . Let M be a c-model containing A . Then $p(x)$ is realized in $\text{Ccl}(M)$.

Proof. Let $a_n \in M$ be a solution of $\neg\varphi_n(x)$. By the assumption above, the type

$$p(x) \cup \{a_n \sim_n x : n \in \omega\}$$

is consistent. Its realizations belong to $\text{Ccl}(M)$. \square

In words, the proof above shows that if $p(x)$ is c-isolated, then in every c-model contains a sequence of elements that converges, in the topology induced by I , to a realization of $p(x)$.

Fact 53. Let \mathcal{L}^c be separable. If $p(x) \subseteq \mathcal{L}^c(A)$ is realized in $\text{Ccl}(M)$ for some c-model M containing A , then $p(x)$ is c-isolated by M .

Proof. Let $b \in \text{Ccl}(M)^{|x|}$ realize $p(x)$. Let $\varepsilon \in \mathcal{E}$ be given. By Fact 49 it suffices to find a continuous formula $\varphi(x)$ and some $\varphi' > \varphi$ such that $\varphi'(x) \rightarrow \exists x' \sim_\varepsilon x p(x')$.

Let $\eta' \in \mathcal{E}$ be such that $x \sim_{\eta'} y \sim_{\eta'} z \rightarrow x \sim_\varepsilon z$. Pick some $a \in M^{|x|}$ that satisfies $x \sim_{\eta'} b$. Then $x \sim_{\eta'} a \rightarrow \exists x' \sim_\varepsilon x p(x')$. Finally, note that $(\sim_{\eta'}) > (\sim_\varepsilon)$ for some $\eta \in \mathcal{E}$. \square

Lemma 54. Let $\mathcal{L}^c(A)$ be separable. Let $p(x) \subseteq \mathcal{L}^c(A)$, be non c-isolated. Then every consistent formula $\neg\psi(z)$, with $\psi(z) \in \mathcal{L}^c(A)$, has a solution a such that A, a does not c-isolate $p(x)$.

Proof. The proof is essentially that of Lemma 50. We go over it again to check that the stronger requirement of consistency does not invalidate the argument.

Let $\varepsilon \in \mathcal{E}$ witness that $p(x)$ is not c-isolated by A . We construct a sequence of $\mathcal{L}^c(A)$ -formulas $\langle \gamma_i(z) : i < \omega \rangle$ such that any realization a of the type $\{\gamma_i(z) : i < \omega\}$ is the required solution of $\neg\psi(z)$, witnessed by the same ε .

Let $\langle \xi_i(x, z) : i < \omega \rangle$ enumerate a countable c-dense subset of $\mathcal{L}_{x,z}^c(A)$. Let $\gamma_0(z)$ be a consistent continuous formulas such that $\gamma'_0(z) \rightarrow \neg\psi(z)$ for some $\gamma'_0 > \gamma_0$. This exists by Proposition 29. Now we define $\gamma_{i+1}(z)$ and $\gamma'_{i+1} > \gamma_{i+1}$. Let $\gamma_i(z)$ and $\gamma'_i > \gamma_i$ be given. Pick $\tilde{\gamma} \perp \gamma_i$ such that $\gamma_i(z) \rightarrow \neg\tilde{\gamma}(z) \rightarrow \gamma'_i(z)$.

1. If $\neg\xi_i(x, z) \wedge \gamma_i(z)$ is inconsistent with $p(x)$, let $\gamma_{i+1}(z) = \gamma_i(z)$ and $\gamma'_{i+1}(z) = \gamma'_i(z)$.

2. Otherwise, pick $\varphi(x) \in p$ such that (#) below is consistent

$$(\#) \quad \neg\tilde{\gamma}(z) \wedge \exists x [\neg\xi_i(x, z) \wedge \neg\exists x' \sim_\varepsilon x \varphi(x')].$$

Finally, let $\gamma_{i+1}(z)$ and $\gamma'_{i+1} > \gamma_{i+1}$ consistent continuous formulas such that $\gamma'_{i+1}(z)$ implies (#). Such formulas exist by Proposition 29.

Let $a \models \{\gamma_i(z) : i < \omega\}$. We claim that that A, a does not c-isolate $p(x)$, witnessed by ε . Otherwise $\neg\xi_i(x, a) \rightarrow \exists x' \sim_\varepsilon x p(x')$ for some $\neg\xi_i(x, a)$ consistent with $p(x)$. This contradicts $a \models \gamma_{i+1}(z)$.

Therefore the proof is complete if we can show that it is always possible to find the formula $\varphi(x)$ required in (2).

Suppose for a contradiction that $\neg\xi_i(x, z) \wedge \gamma_i(z)$ is consistent with $p(x)$ while (#) is inconsistent for all formulas $\varphi(x) \in p$. This immediately implies that

$$\exists z [\neg\xi_i(x, z) \wedge \neg\tilde{\gamma}(z)] \rightarrow \exists x' \sim_\varepsilon x p(x').$$

By Fact 49, this yields the desired contradiction. \square

Theorem 55 (Continuous Omitting Types). Let \mathcal{L}^c be separable. Let A be countable. Assume also that $p(x) \subseteq \mathcal{L}^c(A)$ is not c-isolated. Then there is a c-model M containing A such that $\text{Ccl}(M)$ omits $p(x)$.

Proof. By assumption $\exists y \sim_\varepsilon x p(y)$ is not c-isolated for some $\varepsilon \in \mathcal{E}$. Apply Lemma 50 and the Tarski-Vaught test (Theorem 45) to obtain a countable c-model M that does not isolate $\exists y \sim_\varepsilon x p(y)$. By Fact 53, $\text{Ccl}(M)$ omits $p(x)$. \square

14. Continuous ω -categoricity

In the discussion of countable categoricity we assume that the underlying theory is complete. This assumption is not necessary (it is rather a consequence of categoricity), but it allows to work inside a monster model and simplify the notation.

Let M be a c-model. We say that M is **atomic** if for every finite tuple a of elements of M the type $\text{c-tp}(a)$ is c-isolated.

Lemma 56. Let \mathcal{L}^c be separable. Let $k : M \rightarrow N$ be a finite c-elementary bijection between two atomic c-models. Then for every $c \in M$ there is $d \in \text{Ccl}(N)$ such that $k \cup \{ \langle c, d \rangle \}$ is a c-elementary relation.

Proof. Let a enumerate $\text{dom} k$. Let $p(x, y) = \text{c-tp}(a, c)$, it suffices to prove that $p(x, y)$ is realized by some for some $d \in \text{Ccl}(N)$. Let

$$\neg\varphi_n(x, y) \rightarrow \exists x', y' \sim_n x, y p(x', y').$$

As $\neg\varphi_n(x, y)$ is consistent with $p(x, y)$ then $a, c \models \neg\varphi_n(x, y)$. Then by c-elementarity, $\neg\varphi(ka, y)$ is consistent. Let $d_n \in M$ be such that $ka, d_n \models \neg\varphi_n(x, y)$. Let b, d be a realization of the type $\{ka, d_n \sim_n x, y : n \in \omega\} \cup p(x, y)$. As $ka \sim b$, also ka, d realizes $p(x, y)$. As $d \in \text{Ccl}(N)$, the proof is complete \square

Fact 57. Let M and N be countable atomic c-models. Then $\text{Ccl}(M)$ and $\text{Ccl}(N)$ are c-isomorphic.

Proof. We construct by back-and-forth a sequence $f_n : \text{Ccl}(M) \rightarrow \text{Ccl}(N)$ of finite c-elementary reduced relations (i.e. injective maps) and let

$$R = \bigcup_{n \in \omega} (\sim) \circ f_n \circ (\sim).$$

We will ensure that $M \subseteq \text{dom}(R)$ and $N \subseteq \text{range}(R)$. By Fact 41 this suffices to prove the fact.

Let n be even. Let a be an enumeration of $\text{dom}(f_n)$. Let $c \in M$. Let $p(x, y) = \text{c-tp}(a, b)$. As M is atomic by Lemma 56, for some $d \in \text{Ccl}(N)$ the relation $f_n \cup \{\langle c, d \rangle\}$ is c-elementary. Let f_{n+1} be a reduced relation such that $f_{n+1} \subseteq f_n \cup \{\langle a, c \rangle\} \subseteq (\sim) \circ f_{n+1} \circ (\sim)$.

When n is odd we proceed similarly with f_n^{-1} . \square

Let \mathcal{L}^c be separable. We say that T is **c- ω -categorical** if $\text{Ccl}(M)$ and $\text{Ccl}(N)$ are c-isomorphic for any two countable c-models M, N . The following analogue of Ryll-Nardzewski's Theorem follows as usual.

Theorem 58. Assume that \mathcal{L}^c is separable. Then the following are equivalent

1. T is c- ω -categorical;
2. every consistent type $p(x) \subseteq \mathcal{L}^c$, for x any finite tuple of variables, is c-isolated.

Proof. (1 \Rightarrow 2) This is just the classical argument. If $p(x)$ is consistent it is realized in some countable c-model M . If $p(x)$ is not c-isolated, by the continuous omitting type theorem there is a c-model N such that $\text{Ccl}(N)$ omits $p(x)$. Then $\text{Ccl}(M)$ and $\text{Ccl}(N)$ are not c-isomorphic, hence T is not c- ω -categorical.

(2 \Rightarrow 1) As, every c-model is atomic, c- ω -categoricity follows from Fact 57 as in the classical case. \square

The classical argument proves also the following.

Proposition 59. Fix a finite tuple of variables x . The following are equivalent

1. every consistent $p(x) \subseteq \mathcal{L}^c(A)$ is c-isolated;
2. for every given $\varepsilon \in \mathcal{E}$ there are at most $n_\varepsilon < \omega$ many types $p_i(x) \subseteq \mathcal{L}^c(A)$ such that $\exists y \sim_\varepsilon x$ $p_i(y)$ are pairwise inconsistent.

Proof. (1 \Rightarrow 2) Let $\varepsilon \in \mathcal{E}$ be given. Let $p_i(x) \subseteq \mathcal{L}^c$, for $i < \lambda$, be such that $\exists y \sim_\varepsilon x$ $p_i(y)$ are pairwise inconsistent. Suppose that λ is maximal and, for a contradiction, that it is infinite. Let $\neg\varphi_i(x)$, where $\varphi_i(x) \in \mathcal{L}^c$, be the formulas that isolate $\exists y \sim_\varepsilon x$ $p_i(x)$. As these types are pairwise inconsistent, then the type $q(x) = \{\varphi_i(x) : i < \lambda\}$ is consistent. The type $\exists y \sim_\varepsilon x$ $q(x)$ is inconsistent with all $\exists y \sim_\varepsilon x$ $p_i(x)$. This contradicts the maximality of λ .

(2 \Rightarrow 1?) Let τ, D be given and pick a compact neighborhood of the diagonal E such that $E \circ E \circ E \subseteq D$. Let $p_i(x) \subseteq \mathcal{L}^c(A)$, for $i < n$, be such that $\exists y \sim_{\tau, E} x$ $p_i(x)$ are pairwise inconsistent. By compactness there are some $\varphi_i(x) \in p_i$ such that $\exists y \sim_{\tau, E} x$ $\varphi_i(x)$ are pairwise inconsistent. Assume $n = n_{\tau, E}$ is maximal. Let $F = E \circ E$. We claim that for every a

$$\bigvee_{i=1}^n \exists y \sim_{\tau, F} x \ \varphi_i(y)$$

In fact, if $q(x) = \text{c-tp}(a/A)$, where a does not satisfy the formula above, then $\exists y \sim_{\tau, E} x \ q(y)$ would be inconsistent with every $\exists y \sim_{\tau, E} x \ \varphi_i(x)$. This contradicts the maximality of n .

Let $q(x) \subseteq \mathcal{L}^c(A)$. We prove that $\exists y \sim_{\varepsilon} x \ q(y)$ is c-isolated. By maximality $\exists y \sim_{\tau, E} x \ q(y)$ is consistent with, say, $\exists y \sim_{\tau, E} x \ \varphi_1(x)$. Without loss of generality, assume $q(x) \rightarrow \exists y \sim_{\tau, E} x \ \varphi_1(x)$. Note that

$$\exists y \sim_{\tau, E} x \ \varphi_1(x) \rightarrow \neg \bigvee_{i=2}^n \exists y \sim_{\tau, F} x \ \varphi_i(y)$$

We claim that

$$\neg \bigvee_{i=2}^n \exists y \sim_{\tau, F} x \ \varphi_i(y) \rightarrow$$

Let

$$\mathcal{C} = \left\{ \langle C_1, \dots, C_n \rangle : E \subseteq C_i \subseteq D \text{ compact, } \forall x \bigvee_{i=1}^n \exists y \sim_{\tau, C_i} x \ \varphi_i(y) \right\}$$

, for $i < n$ be such that

$$1. \quad \neg \bigvee_{i=1}^n \varphi_i(x) \rightarrow \exists y \sim_{\varepsilon} x \ q(y)$$

Suppose not for a contradiction. Let a satisfy the antecedent but not the consequence of (1).

□

15. Example: metric spaces

We discuss a simple example. Let I be $\mathbb{R}^+ \cup \{0, \infty\}$, with the topology that makes it homeomorphic to the unit interval. Let M, d is a metric space. Let \mathcal{L}_M contain symbols for functions $M^n \rightarrow M$ that are uniformly continuous. Let \mathcal{L} contain a symbol for d and possibly for some functions of $M^n \rightarrow I$ that are uniformly continuous w.r.t. the metric d .

Let $\langle M, I \rangle$ be the structure that interprets the symbols of the language as natural.

Let \mathcal{U} be a positive monster model that is an \mathcal{L}^p elementary extension of M . Clearly, d does not define a metric on \mathcal{U} as there are pairs of elements at infinite distance. However, when restricted to a ball of finite radius, d defines a pseudometric on \mathcal{U} . Therefore the notion of convergent sequence makes perfectly sense in \mathcal{U} . As all functions have been required to be uniformly continuous, it is immediate that $x \sim_1 y$ is equivalent to $d(x, y) = 0$.

Fact 60. Let \mathcal{U} be as above. Then for every p-model N and $a \in \mathcal{U}$ the following are equivalent

1. $p(x) = \text{p-tp}(a/N)$ is a Cauchy type;
2. there is a sequence $\langle a_i : i \in \omega \rangle$ of elements of N that converges to a .

Proof. (2 \Rightarrow 1) Let $\langle \varepsilon_i : i \in \omega \rangle$ be a sequence of reals that converges to 0 and such that $d(a_i, a) \leq \varepsilon_i$ for every $i \in \omega$. Then the formulas $d(a_i, x) \leq \varepsilon_i$ are in $p(x)$. Then every element realizing $p(x)$ is at distance 0 from a . Therefore $p(x) \rightarrow a \sim x$.

(1 \Rightarrow 2) As $p(x)$ is Cauchy type, $p(x) \rightarrow a \sim x$. Then $p'(x) \rightarrow d(a, x) < 2^{-i}$ for all i . By compactness (see Fact 25) there are formulas $\varphi_i(x) \in p$ such that $\varphi'_i(x) \rightarrow d(a, x) < 2^{-i}$ for some $\varphi'_i > \varphi_i$. By \mathcal{L}^p -elementarity there is an $a_i \in N$ such that $\varphi'_i(a_i)$. As $d(a, a_i) < 2^{-i}$ for all i , the sequence $\langle a_i : i \in \omega \rangle$ converges to a . □

References

- [G] Isaac Goldbring, *Lecture notes on nonstandard analysis*, UCLA Summer School in Logic 2012, available at www.math.uci.edu/~isaac/NSA%20notes.pdf.
- [H] Bradd Hart, *An Introduction To Continuous Model Theory* (2023), arXiv:2303.03969
- [Hr] Ehud Hrushovski, *Stable group theory and approximate subgroups*, J. Amer. Math. Soc. **25** (2012), no. 1, 189–243.
- [HPP] Ehud Hrushovski, Ya'acov Peterzil, and Anand Pillay, *Groups, measures, and the NIP*, J. Amer. Math. Soc. **21** (2008), no. 2, 563–596.
- [HI] C. Ward Henson and José Iovino, *Ultraproducts in analysis*, Analysis and logic (Mons, 1997), London Math. Soc. Lecture Note Ser., vol. 262, Cambridge Univ. Press, Cambridge, 2002, pp. 1–110.
- [CLCL] C.L.C.L. Polymath, *Continuous Logic for the Classical Logician* (202?). T.a.



Danger after this point

16. Elimination of quantifiers

Let T be a positive theory. We say that T has (approximate, positive) elimination of quantifiers if the quantifier-free positive formulas are p-dense modulo T . Equivalently, if every complete type is equivalent to a quantifier-free positive types.

The following fact is routinely proved by back-and-forth.

Fact 61. The following are equivalent

1. T has elimination of quantifiers;
2. every finite partial embedding $k : M \rightarrow N$ between p-models of T is an approximate p-elementary map;
3. for every finite partial embedding $k : M \rightarrow N$ between ω -saturated p-models of T , and for every $b \in M$ there is a $c \in N$ such that $k \cup \{\langle b, c \rangle\} : M \rightarrow N$ is also a partial embedding.

17. Pseudofinite randomizations

Let T be a complete first-order theory of signature L with an infinite model. Let \mathcal{U} be a model of T . Let Ω be a finite set. We denote by \mathcal{U}^Ω the set of functions $\Omega \rightarrow \mathcal{U}$. The elements of \mathcal{U}^Ω are denoted by letters decorated with a circumflex accent \hat{a}, \hat{b} , etc. The value of \hat{a} at $\omega \in \Omega$ is denoted by a_ω . We now define models of the form (\mathcal{U}^Ω, I) , where I is the real interval $[0, 1]$ with the usual topology.

The language \mathcal{L}_H is empty. The language \mathcal{L} contains functions of sort $H^{|x|} \rightarrow I$, one for each formula $\varphi(x) \in L$. These are denoted by $\text{Pr } \varphi(\hat{x})$. Variables of sort H are decorated with a circumflex accent. The interpretation of the function symbol $\text{Pr } \varphi(\hat{x})$ is

$$\text{Pr}(\varphi(\hat{a})) = \frac{\#\llbracket \varphi(\hat{a}) \rrbracket}{\#\Omega},$$

where by # denotes the finite cardinality, and

$$\llbracket \varphi(\hat{a}) \rrbracket = \{ \omega \in \Omega : \mathcal{U} \models \varphi(a_\omega) \}.$$

The pseudofinite randomization of T is the positive theory

$$T^{\text{pfr}} = \left\{ \varphi \in \mathcal{L}^P : \langle \mathcal{U}^\Omega, I \rangle \models \varphi \text{ for all sufficiently large finite set } \Omega \right\}$$

In this section we prove that T^{pfr} is a complete theory with elimination of quantifiers. We prove the lemma that is required for the back-and-forth.

Lemma 62. Let $k : M \rightarrow N$ be a finite partial embedding between positively ω -saturated models of T^{pfr} . Then for every $\hat{b} \in M$ and there is a $\hat{c} \in N$ such that $k \cup \{ \langle \hat{b}, \hat{c} \rangle \} : M \rightarrow N$ is a partial embedding.

Proof. Let \hat{a} be an enumeration of $\text{dom} k$. Let $p(\hat{x}, \hat{z})$ be set of formulas of the form $\gamma = \text{Pr} \varphi(\hat{x}, \hat{z})$, for some $\gamma \in I$, that hold in M, \hat{b}, \hat{a} . Note that $p(\hat{x}, \hat{z})$ implies the atomic type of \hat{b}, \hat{a} in M . Therefore it suffices to show that $p(\hat{x}, k\hat{a})$ is finitely consistent in N .

Pick a finite set of these formulas, say $\gamma_i = \text{Pr} \varphi_i(\hat{x}, \hat{z})$, for $i < n$. We may assume that the formulas $\varphi_i(x, z)$ define a partition of $\mathcal{U}^{[x, z]}$. By the saturation of N , it suffices to prove that for every $\varepsilon > 0$ there is a $\hat{c} \in N$ such that

$$\gamma_i =_\varepsilon \text{Pr} \varphi_i(\hat{c}, k\hat{a}) \quad (\alpha =_\varepsilon \beta \text{ is a shorthand for } |\alpha - \beta| \leq \varepsilon)$$

is true in N for all $i < n$.

Some preliminary work is required. For each $J \subseteq n$ define the following L -formulas

$$\xi_J(z) = \bigwedge_{i \in J} \exists x \varphi_i(x, z) \wedge \bigwedge_{i \in n \setminus J} \neg \exists x \varphi_i(x, z)$$

These formulas $\xi_J(z)$ partition $\mathcal{U}^{[z]}$. For $J \subseteq n$ let α_J be such that

$$2. \quad \alpha_J = \text{Pr}(\xi_J(\hat{z}))$$

is in $p(\hat{x}, \hat{z})$. Let $\mathcal{J} = \{ J \subseteq n : \alpha_J \neq 0 \}$. When $J \in \mathcal{J}$, there are some $\beta_{i,J}$ such that the formulas

$$3. \quad \beta_{i,J} = \text{Pr}(\varphi_i(\hat{x}, \hat{z}) \mid \xi_J(\hat{z})).$$

are among the consequences of $p(\hat{x}, \hat{z})$. As the formulas $\varphi_i(x, z)$ define a partition, the $\beta_{i,J}$ add up to 1 for any fixed J . Note that $\beta_{i,J} = 0$ for $i \notin J$. Clearly, have that

$$4. \quad \gamma_i = \sum_{J \in \mathcal{J}} \beta_{i,J} \alpha_J.$$

It is plain that for $i \in J$ the following is a consequence of $p(\hat{x}, \hat{z})$

$$5. \quad 1 = \text{Pr}(\exists x \varphi_i(x, \hat{z}) \mid \xi_J(\hat{z}))$$

Now we prove that (2) & (5) \Rightarrow (1) holds in \mathcal{U}^Ω when Ω large enough (larger than n/ε suffices). Strictly speaking, this implication is not a positive formula, but the reader can easily verify that a suitable approximation of (2) and (5) suffices.

Let $\hat{a}' \in \mathcal{U}^\Omega$ satisfy (2) and (5) for every $i \in J \subseteq n$. We define $\hat{c}' \in \mathcal{U}^\Omega$ such that

$$\mathcal{U}^\Omega \models \beta_{i,J} =_\varepsilon \text{Pr}(\varphi_i(\hat{c}', \hat{a}') \mid \xi_J(\hat{a}')).$$

We may define \hat{c}' separately in each event $\llbracket \xi_J(\hat{a}') \rrbracket$. Partition $\llbracket \xi_J(\hat{a}') \rrbracket$ into events $E_i \subseteq \Omega$, for $i \in J$, such that $\#E_i / \#\Omega =_\varepsilon \beta_{i,J}$. Then define c'_ω for $\omega \in E_i$ to be any witness of $\exists x \varphi_i(x, a_\omega)$. By (5), we can always find such a witness.

Finally, by (4), we deduce

$$\mathcal{U}^\Omega \models \gamma_i =_\epsilon \Pr(\varphi_i(\hat{c}', \hat{a}')).$$

As $k\hat{a}$ satisfy (2) and (5), we conclude that some $\hat{c} \in N$ satisfy (1). \square

Now, from Fact 61 we obtain that T^{pfr} has elimination of quantifiers. Completeness follows.

Corollary 63. The theory T^{pfr} is complete and has elimination of quantifiers.

18. Stability

A partitioned formula $\varphi(x; z) \in \mathcal{L}^P$ is stable if there is no sequence $\langle a_i; b_i : i < \omega \rangle$ and no $\tilde{\varphi} \perp \varphi$ such that for every $i, j < \omega$

$$i < j \Rightarrow \varphi(a_i; b_j)$$

$$i > j \Rightarrow \tilde{\varphi}(a_i; b_j)$$

Using the terminology of [Hr], $\varphi(x; z)$ is stable if it is stably separated from all $\tilde{\varphi} \perp \varphi$.

Note that by compactness if $\varphi(x; z)$ and $\tilde{\varphi} \perp \varphi$ are stably separated then there is a maximal length m of a sequence $\langle a_i; b_i : i < m \rangle$ such that (1) and (2) above.

19. Elementary relations

Let M and N be models. We say that $R \subseteq M \times N$ is an **p-elementary relation** between M and N if for every $\varphi(x) \in \mathcal{L}$

$$M \models \varphi(a) \Rightarrow N \models \varphi(b) \quad \text{for every } a \text{ and } b \text{ such that } a R b.$$

Note that, when $a = a_1, \dots, a_n$ and $b = b_1, \dots, b_n$ are tuples, $a R b$ stands for $a_i R b_i$ for every $i \in \{1, \dots, n\}$.

Fact 64. Let $R \subseteq \mathcal{U}^2$ be an p-elementary relation of cardinality $< \kappa$. Then there is a total and surjective p-elementary relation $S \subseteq \mathcal{U}^2$ containing R .

Proof. We apply the usual back-and-forth construction with a pinch of extra caution. Let a be an enumeration of the domain of R . Let $\bar{a} = \langle a_i : i < \lambda \rangle$ be an enumeration of all tuples of length $|\bar{a}|$ such that $a R a_i$. As κ is inaccessible, $\lambda < \kappa$. Let $b \in U$. It suffices to prove that there is a c such that $R \cup \{\langle b, c \rangle\}$ is an \mathbb{L} -relation. Let $p(x, z) = \text{tp}(b, a)$ and let

$$q(x, \bar{z}) = \bigcup_{i < \lambda} p(x, z_i).$$

We claim that $q(x, \bar{a})$ is a finitely consistent type. A finite conjunction of formulas in $q(x, \bar{a})$ has the form $\psi(x, a_{i_1}) \wedge \dots \wedge \psi(x, a_{i_n})$. As $\psi(b, a)$ and $a_{i_1}, \dots, a_{i_n} R a, \dots, a$, we conclude that the condition $\psi(x, a_{i_1}) \wedge \dots \wedge \psi(x, a_{i_n})$ is satisfied. The existence of the required element c follows by saturation. \square

20.

Corollary 65. Let $A \subseteq \mathcal{U}$ have cardinality $< \kappa$. Let $p(x) = \text{tp}_{\mathbb{L}}(a/A)$, where $a \in \mathcal{U}^{|x|}$ is a tuple of length $|x| < \kappa$. Then

$$p(\mathcal{U}) = \{b : b R a, R \in \text{Aut}(\mathcal{U}/A)\}$$

Let \mathcal{M} and \mathcal{N} be models. We say that $R \subseteq M \times N$ is an **\mathbb{L} -(elementary) relation** between \mathcal{M} and \mathcal{N} (or on \mathcal{M} if the two coincide) if for every $\varphi(x) \in \mathbb{L}$

$$\mathcal{M} \models \varphi(a) \Leftrightarrow \mathcal{N} \models \varphi(b) \quad \text{for every } a \text{ and } b \text{ such that } a R b.$$

Recall that, when $a = a_1, \dots, a_n$ and $b = b_1, \dots, b_n$ are tuples, $a R b$ stands for $a_i R b_i$ for every $i \in \{1, \dots, n\}$.

We define an equivalence relation $(\sim_{\mathcal{M}})$ on M as follows

$$1. \quad a \sim_{\mathcal{M}} b \Leftrightarrow \left(\mathcal{M} \models \varphi(a) \leftrightarrow \varphi(b) \text{ for every } \varphi(x) \in \mathbb{L}(M) \right),$$

where $|x| = |a| = |b| = 1$. Note that this relation would be trivial had we included in $\mathbb{L}(M)$ equality between elements of M .

The following proposition is easily proved by induction on the syntax.

Proposition 66. The following are equivalent for every $a, b \in M$.

1. $a \equiv_{\mathcal{M}} b$;
2. $\mathcal{M} \models \tau(a) = \tau(b)$ for every $\tau(x) \in \mathbb{T}(M)$, with $|x| = 1$.

Lemma 67. The relation $(\sim_{\mathcal{M}}) \subseteq M^2$ is an \mathbb{L} -relation. Moreover, it is maximal among the \mathbb{L} -relations on \mathcal{M} , i.e. no \mathbb{L} -relation properly contains $(\sim_{\mathcal{M}})$.

Proof. Assume $a \sim_{\mathcal{M}} b$, where $a = a_1, \dots, a_n$ and $b = b_1, \dots, b_n$. Recall that this means that $a_i \sim_{\mathcal{M}} b_i$ for all $i \in \{1, \dots, n\}$. Let Δ denote the diagonal relation on M . Note that $a_i \sim_{\mathcal{M}} b_i$ is equivalent to saying that $\Delta \cup \{(a_i, b_i)\}$ is an \mathbb{L} -relation. As \mathbb{L} -relations are closed under composition $\Delta \cup \{(a_1, b_1), \dots, (a_n, b_n)\}$ is \mathbb{L} -elementary. It follows that for every $\varphi(x) \in \mathbb{L}$

$$2. \quad \mathcal{M} \models \varphi(a) \Leftrightarrow \mathcal{M} \models \varphi(b).$$

This proves that $(\sim_{\mathcal{M}})$ is an \mathbb{L} -relation. Finally, maximality is immediate. \square

Lemma 68. Let $R \subseteq M \times N$ be total and surjective \mathbb{L} -relation. Then there is a unique maximal \mathbb{L} -relation containing R . This maximal \mathbb{L} -relation is equal to both $(\sim_{\mathcal{M}}) R$ and $R (\sim_{\mathcal{N}})$, where juxtaposition of relations stands for composition.

Proof. It is immediate to verify that $(\sim_{\mathcal{M}}) R$ is an \mathbb{L} -relation containing R . Let S be any maximal \mathbb{L} -relation containing R . By maximality, $(\sim_{\mathcal{M}}) S = S$. As S is a total relation $(\sim_{\mathcal{M}}) \subseteq S S^{-1}$. Therefore, by the lemma above, $(\sim_{\mathcal{M}}) = S S^{-1}$. As R is a surjective relation, $S \subseteq S S^{-1} R$. Finally, by maximality, we conclude that $S = (\sim_{\mathcal{M}}) R$. A similar argument proves that $S = R (\sim_{\mathcal{N}})$. \square

We write $\text{Aut}(\mathcal{M})$ for the set of maximal, total and surjective, \mathbb{L} -relations $R \subseteq M^2$. The choice of the symbol Aut is motivated by the lemma above. In fact any such relation R induces a unique automorphism on the (properly defined) quotient structure $\mathcal{M}/\sim_{\mathcal{M}}$.