

Local stability in structures with a standard sort

L. S. Polymath

ABSTRACT.

1. Structures with a standard sort

Let S be some Hausdorff compact topological space. We associate to S a first order structure in a language \mathcal{L}_S that has a symbol for each compact subset $C \subseteq S$ and a function symbol for each continuous functions $f : S^n \rightarrow S$. According to the context, C and f denote either the symbols of \mathcal{L}_S or their interpretation in the structure S .

Throughout these notes the letter C always denotes a compact subset of S , or a tuple of such sets. Finally, note that we could allow in \mathcal{L}_S relation symbols for all compact subsets of S^n , for any n . But this would clutter the notation adding very little to the theory.

We also fix an arbitrary first-order language which we denote by \mathcal{L}_H and call the language of the home sort.

1 Definition Let \mathcal{L} be a two sorted language. The two sorts are denoted by H and S . The language \mathcal{L} expands \mathcal{L}_H and \mathcal{L}_S with symbols sort $H^n \times S^m \rightarrow S$. An \mathcal{L} -structure is a structure of signature \mathcal{L} that interprets these symbols in equicontinuous functions (i.e. uniformly continuous w.r.t. the variables in H). A **standard structure** is a two-sorted \mathcal{L} -structure of the form $\langle M, S \rangle$, where M is any structure of signature \mathcal{L}_H and S is fixed. Standard structures are denoted by the domain of their home sort.

We denote by \mathcal{F} the set of \mathcal{L} -formulas constructed inductively from atomic formulas of the form (i) and (ii) below using Boolean connectives \wedge, \vee ; the quantifiers \forall^H, \exists^H of sort H ; and the quantifiers \forall^S, \exists^S of sort S .

We write $x \in C$ for the predicate associated to a compact set C , and denote by $|\cdot|$ the length of a tuple.

2 Definition We call atomic formulas those of the form

- i. atomic and negated atomic formulas of \mathcal{L}_H but not equalities and inequalities
- ii. $\tau \in C$, where τ is a k -tuple terms of sort $H^n \times S^m \rightarrow S$, and C a compact.

Let $M \subseteq N$ be standard structures. We say that M is an \mathcal{F} -elementary substructure of N if the latter models all $\mathcal{F}(M)$ -sentences that are true in M .

Let x and ξ be variables of sort H , respectively S . A standard structure M is \mathcal{F} -saturated if it realizes every type $p(x; \xi) \subseteq \mathcal{F}(A)$, for any $A \subseteq M$ of cardinality smaller than $|M|$, that is finitely consistent in M . The following theorem is proved in [AAVV] for signatures that do not contain symbols sort $H^n \times S^m \rightarrow S$ with $n \cdot m > 0$. A similar framework has been independently introduced in [PC] – only without quantifiers of sort H and with an approximate notion of satisfaction. The corresponding compactness theorem is proved there with a similar method. The setting in [PC] is a generalization of that used by Henson and Iovino for Banach spaces [HI]. By the elimination of quantifiers of sort S , proved in [AAVV, Proposition 3.6], the two approaches are equivalent – up to approximations.


In [Z] it is observed that, under the assumption of equicontinuity, the proof in [AAVV] extends to the case $n \cdot m > 0$.

3 Theorem (Compactness) Every standard structure has a \mathcal{F} -saturated \mathcal{F} -elementary extension.

To a standard structure $\langle M, S \rangle$ we associate a canonical expansion $\langle M, S, K(S) \rangle$, where $K(S)$ is the set of compact subsets of S endowed with the Vietoris topology. The sort of $K(S)$ is denoted by K and is treated as a second standard sort. To the atomic formulas in Definition 1 we add the atomic formulas of the form

iii. $\tau(x; \xi) \in X$, where X is a variable of sort K .

After closing under the (positive) logic connectives, we obtain the set of formulas that we denote by \mathcal{F}_K . Formulas in \mathcal{F}_K are denoted by $\varphi(x; \xi; X)$, where $X = X_1, \dots, X_n$ be a tuple of variables of sort K . If $C = C_1, \dots, C_n$ is a tuple of compact subsets, then $\varphi(x; \xi; C)$, is a formula in \mathcal{F} . This we call an **instance** of $\varphi(x; \xi; X)$. All formulas in \mathcal{F} are instances of formulas in \mathcal{F}_K .

 In Appendix A we prove that Theorem 3 extends to these expanded standard structures.

4 Definition Let $\varphi \in \mathcal{F}_K$ be a formula – possibly with some (hidden) free variables. The **pseudonegation** of $\varphi \in \mathcal{F}_K$ is the formula obtained by replacing the atomic formulas in \mathcal{L}_H by their negation and every connective $\wedge, \vee, \forall, \exists$ by their respective duals $\vee, \wedge, \exists, \forall$. The atomic formulas of the form $\tau \in X$ remain unchanged.

The pseudonegation of φ is denoted by $\sim\varphi$. Clearly, when X does not occur in φ , we have $\sim\varphi \leftrightarrow \neg\varphi$.

The following fact is immediate and will be used without further mention.

5 Fact The following hold for every $C \subseteq C'$ and $\tilde{C} \cap C = \emptyset$, every standard structure M , and every $\mathcal{F}_K(M)$ -formula $\varphi(X)$

$$M \models \varphi(C) \rightarrow \varphi(C')$$

$$M \models \sim\varphi(\tilde{C}) \rightarrow \neg\varphi(C).$$

6 Fact For every $C \subseteq C'$ there is a $\tilde{C} \subseteq C'$ disjoint from C such that

$$M \models \varphi(C) \rightarrow \neg\sim\varphi(\tilde{C}) \rightarrow \varphi(C')$$

for every M , and every $\mathcal{F}_K(M)$ -formula $\varphi(X)$. Conversely, for every $\tilde{C} \cap C = \emptyset$ there is a $C' \supseteq C$ such that

$$M \models \varphi(C) \rightarrow \varphi(C') \rightarrow \neg\sim\varphi(\tilde{C}).$$

for every M , and every $\mathcal{F}_K(M)$ -formula $\varphi(X)$.

Proof. When $\varphi(X)$ is the atomic formula $\tau \in X$, the first claim holds with $\tilde{C} = C' \setminus O$ where O is any open set $C \subseteq O \subseteq C'$. The second claim holds with as C' any neighborhood of C disjoint from \tilde{C} . Induction on the syntax of $\varphi(X)$ proves the general case. \square

The above fact has the following useful consequence.

7 Fact Let $\varphi(X)$ be a $\mathcal{F}_K(M)$ -formula, where M is a standard structure. Then the following are equivalent for every tuple C

$$M \models \varphi(C) \leftarrow \bigwedge \left\{ \varphi(C') : C' \text{ neighborhood of } C \right\}$$

$$M \models \neg\varphi(C) \rightarrow \bigvee \left\{ \sim\varphi(\tilde{C}) : \tilde{C} \cap C = \emptyset \right\}.$$

(The converse implications are trivial – therefore not displayed.)

Note that the fact also holds if we restrict the above C' and \tilde{C} to range over the subsets of some neighborhood of C .

8 Fact The equivalent conditions in Fact 7 hold in all \mathcal{F} -saturated structures.

Proof. Prove the first of the two implications by induction on the syntax of $\varphi(X)$. The existential quantifier of sort H is the only connective that requires attention. Assume inductively that

$$\bigwedge \left\{ \varphi(a; C') : C' \text{ neighborhood of } C \right\} \rightarrow \varphi(a; C)$$

holds for every $\varphi(a; C)$. Then induction for the existential quantifier follows from

$$\bigwedge \left\{ \exists x \varphi(x; C') : C' \text{ neighborhood of } C \right\} \rightarrow \exists x \bigwedge \left\{ \varphi(x; C') : C' \text{ neighborhood of } C \right\}$$

which is a consequence of saturation. \square

We say that M is \mathcal{F} -maximal if it models all $\mathcal{F}(M)$ -sentences that hold in some of its \mathcal{F} -elementary extensions. By Fact 8 and the following, \mathcal{F} -saturated standard structures are \mathcal{F} -maximal.

9 Fact Let $\varphi(X)$ be an $\mathcal{F}_K(M)$ -formula, where M is a standard model. Then the following are equivalent

1. M is \mathcal{F} -maximal
2. the equivalent conditions in Fact 7 hold for every $\varphi(C)$.

Proof. $1 \Rightarrow 2$. If (2) in Fact 7 fails, $\neg\varphi(C) \wedge \neg\sim\varphi(\tilde{C})$ holds in M for every $\tilde{C} \cap C = \emptyset$. Let \mathcal{U} be an \mathcal{F} -saturated \mathcal{F} -elementary extension of M . As M is \mathcal{F} -maximal, $\neg\varphi(C) \wedge \neg\sim\varphi(\tilde{C})$ holds also in \mathcal{U} . Then (2) in Fact 7 fails in \mathcal{U} , contradicting Fact 8.

$2 \Rightarrow 1$. Let N be any \mathcal{F} -elementary extension of M . Suppose $N \models \varphi(C)$. By (2), it suffices to prove that $M \models \varphi(C')$ for every C' neighborhood of C . Suppose not. Then, by (2) in Fact 7, $M \models \sim\varphi(\tilde{C})$ for some $\tilde{C} \cap C' = \emptyset$. Then $N \models \sim\varphi(\tilde{C})$ by \mathcal{F} -elementarity. As $\tilde{C} \cap C = \emptyset$, this contradicts Fact 8. \square

In what follows we fix a large saturated standard structure which we denote by \mathcal{U} .

2. A duality in $K(S)$

Write $K(S)$ for the set of compact subsets of S . In this section \mathcal{D} and \mathcal{C} range over subsets of $K(S)^n$. We define

$$\sim\mathcal{D} = \{\tilde{C} : \tilde{C} \cap C \neq \emptyset \text{ for every } C \in \mathcal{D}\}.$$

The following fact motivates the definition.

10 Fact For any $\mathcal{F}_K(M)$ -formula $\varphi(X)$. Let $\mathcal{D} = \{C : \varphi(C)\}$. Then $\sim\mathcal{D} = \{\tilde{C} : \sim\varphi(\tilde{C})\}$.

Proof. We need to prove that

$$\sim\varphi(\tilde{C}) \Leftrightarrow \text{for every } C, \text{ if } C \cap \tilde{C} = \emptyset \text{ then } \neg\varphi(C)$$

Implication \Rightarrow follows immediately from Fact 5. To prove \Leftarrow assume the r.h.s. Then $\neg\varphi(S \setminus O)$ holds for every open set $O \supseteq \tilde{C}$. From the second implication in Fact 7 we obtain $\sim\varphi(\tilde{C}')$ for some $\tilde{C} \subseteq \tilde{C}' \subseteq O$. As O is arbitrary, we obtain $\varphi(\tilde{C})$ from the second implication in Fact 7. \square

We prove a couple of straightforward inclusions.

11 Fact For every $\mathcal{D} \subseteq \mathcal{C}$ we have $\sim\mathcal{C} \subseteq \sim\mathcal{D}$. Moreover, $\mathcal{D} \subseteq \sim\sim\mathcal{D}$.

Proof. Let $C \notin \sim\mathcal{D}$. Then $\tilde{C} \cap C = \emptyset$ for some $\tilde{C} \in \mathcal{D}$. As $\tilde{C} \in \mathcal{C}$ we conclude that $C \notin \sim\mathcal{C}$. For the second claim, let $C \notin \sim\sim\mathcal{D}$. Then $\tilde{C} \cap C = \emptyset$ for some $\tilde{C} \in \sim\mathcal{D}$. Then $C \notin \mathcal{D}$ follows. \square

12 Fact For every \mathcal{D} the following are equivalent

1. $\mathcal{D} = \sim\sim\mathcal{D}$.
2. $\mathcal{D} = \sim\mathcal{C}$ for some \mathcal{C} .

Proof. Only $2 \Rightarrow 1$ requires a proof. From Fact 11 we obtain $\sim\sim\mathcal{C} \subseteq \sim\mathcal{C}$. Assume (2) then $\sim\sim\mathcal{D} \subseteq \mathcal{D}$ which, again by Fact 11, suffices to prove (1). \square


We say that \mathcal{D} is **involutive** if $\mathcal{D} = \sim\sim\mathcal{D}$. We say that \mathcal{D} is **closed** if

$$C \in \mathcal{D} \Leftrightarrow C' \in \mathcal{D} \text{ for every neighborhood } C' \text{ of } C.$$

Note that we are requiring closure in two distinct contexts: topological, and Boolean (upward closure by inclusion).

13 Theorem The following are equivalent

1. \mathcal{D} is involutive
2. \mathcal{D} is closed.

 It is not difficult to see that (2) holds if and only if \mathcal{D} is closed in the Vietoris topology and includes all the supersets of its elements.

Proof. $2 \Rightarrow 1$. It suffices to prove $\sim\sim\mathcal{D} \subseteq \mathcal{D}$. Let $C \in \sim\sim\mathcal{D}$. Let $O \supseteq C$ be open. Then $S \setminus O \notin \sim\mathcal{D}$. Then $\tilde{C} \in \mathcal{D}$ for some $\tilde{C} \subseteq O$. Assume (2). Then, by \Rightarrow every $C' \supseteq O$ is in \mathcal{D} . As O is arbitrary, $C \in \mathcal{D}$ by \Leftarrow .

$1 \Rightarrow 2$. It suffices to show that (2) holds with $\sim\mathcal{D}$ for \mathcal{D} . Implication \Rightarrow in the definition of closed is obvious. To prove \Leftarrow , assume $C \notin \sim\mathcal{D}$. Then $C \cap \tilde{C}$ for some $\tilde{C} \in \mathcal{D}$. Let C' be a neighborhood of C disjoint of \tilde{C} . Then the r.h.s. of the equivalence fails for C' . \square

3. Stable formulas – the finitary case

There is more than one way to define stability in our context. In this section we present the strongest possible requirement. It is not the most interesting one, but it is very similar to the classical definition, so it is useful for comparison.

Let $\varphi(x; z; X)$ be a formulas in $\mathcal{F}_K(\mathcal{U})$. The variables of sort H are partitioned in two tuples $x; z$ which may be infinite. The following is the classical definition of stability which we dub *finitary* for the reason explained below.

14 Definition We say that $\varphi(x; z; C)$ is **finitarily stable** if for some $m < \omega$ there is no sequence $\langle a_i; b_i : i < m \rangle$ such that for every $i < n$

$$\varphi(a_n; b_i; C) \wedge \neg \varphi(a_i; b_n; C).$$

From Fact 7 we easily obtain the following equivalent definition which only mentions formulas in \mathcal{F} . The reader may wish to compare it with Definition/Theorem 20.

15 Fact The following are equivalent

1. $\varphi(x; z; C)$ is finitarily stable
2. for some m there is no sequence $\langle a_i; b_i; C_i : i < m \rangle$ such that for every $i < n$

$$\varphi(a_n; b_i; C) \wedge \sim \varphi(a_i; b_n; C_n) \text{ and } C \cap C_n = \emptyset.$$

In a classical context, i.e. relying on \mathcal{L} -saturation, we can replace m in Definition 14 by ω . But this may not be true if only have \mathcal{F} -saturation. Therefore we dedicate a separate section to the non finitary version of stability.

The following definitions are classical, i.e. they rely on the full language \mathcal{L} .

16 Definition A **global $\varphi(x; z; C)$ -type** is a maximally (finitely) consistent set of formulas of the form $\varphi(x; b; C)$ or $\neg \varphi(x; b; C)$ for some $b \in \mathcal{U}^{|z|}$.

Global $\varphi(x; z; C)$ -types should not be confused with $\varphi(x; z; X)$ -types which are introduced in the following section.

In this section the symbol \mathcal{D} always denotes a subset of \mathcal{U}^z .

17 Definition We say that \mathcal{D} is **approximable** by $\varphi(x; z; C)$ if for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that

$$b \in \mathcal{D} \Leftrightarrow \varphi(a; b; C) \quad \text{for every } b \in B.$$

It goes without saying that these definitions describe two faces of the same coin. In fact, it is clear that \mathcal{D} is approximable by $\varphi(x; z; C)$ if and only if

$$p(x) = \{\varphi(x; b; C) : b \in \mathcal{D}\} \cup \{\neg \varphi(x; b; C) : b \notin \mathcal{D}\}$$

is a global $\varphi(x; z; C)$ -type.

The following theorem is often rephrased by saying that, when $\varphi(x; z; C)$ is finitarily stable, then all global $\varphi(x; z; C)$ -types are definable. The proof of the classical case applies verbatim because no saturation is required. We state it here only for comparison with its non-finitary counterpart, see Theorem 26.

18 Theorem Let $\varphi(x; z; C)$ be finitarily stable. Let \mathcal{D} be approximable by $\varphi(x; z; C)$. Then there are some $\langle a_{i,j} : i < k, j < m \rangle$ such that for every $b \in \mathcal{U}^z$

$$b \in \mathcal{D} \Leftrightarrow \bigvee_{i < k} \bigwedge_{j < m} \varphi(a_{i,j}; b; C)$$

4. Stable formulas – the non-finitary case

In this section \mathcal{D} is always a subset of $\mathcal{U}^z \times K(S)^n$, where n has to be inferred from the context (typically, $n = |X|$). We write $\sim \mathcal{D}$ for the set obtained by applying the definition in Section 2 to all fibers of \mathcal{D} . We say that \mathcal{D} is involutive/closed if all its fibers are such.

The letters ε and δ always denote definable compact symmetric neighborhood of the diagonal of S^2 . By $\varepsilon + \delta$ we denote the composition $\varepsilon \circ \delta$, where the two sets are viewed as relations.

If C is a subsets of S , we write C^ε for the set whose elements are ε -close to elements of C . We write C^{ε° for the set of elements that are ε° -close to some element of C , where ε° is the interior of ε . Then C^{ε° is open. This definition extends naturally to when $C = C_1, \dots, C_k$.

19 Definition We say that $\varphi(x; z; X)$ is **stable** if there is no sequence $\langle a_i; b_i : i < \omega \rangle$ such that for some C , some $\tilde{C} \cap C = \emptyset$, and for every $i < n$

$$\varphi(a_n; b_i; C) \wedge \sim \varphi(a_i; b_n; \tilde{C})$$

It is convenient to work with the following weaker notion of stability. Note that ε -stable implies ε' -stable for every $\varepsilon \subseteq \varepsilon'$.

20 Definition / Theorem We say that $\varphi(x; z; X)$ is **ε -stable** if the following equivalent conditions hold

1. there is no sequence $\langle a_i; b_i; C_i; \tilde{C}_i : i < \omega \rangle$ such that for every $i < n < \omega$

$$\varphi(a_n; b_i; C_i) \wedge \sim \varphi(a_i; b_n; \tilde{C}_i) \quad \text{and} \quad C_i^\varepsilon \cap \tilde{C}_i = \emptyset.$$
2. there is a maximal m such that some sequence $\langle a_i; b_i; C_i; \tilde{C}_i : i < m \rangle$ is such that for every $i < n < m$

$$\varphi(a_n; b_i; C_i) \wedge \sim \varphi(a_i; b_n; \tilde{C}_i) \quad \text{and} \quad C_i^\varepsilon \cap \tilde{C}_i = \emptyset.$$

Proof. $2 \Rightarrow 1$. Any counter example to (1) immediately provides a counterexample to (2).

1 \Rightarrow 2. Negate (2). For a given m let $\langle a_i; b_i; C_i; \tilde{C}_i : i < m \rangle$ be a witness of (2). Note that we can find an open neighborhood of the diagonal $\delta_m^\circ \supseteq \varepsilon$ such that for every $i < n < m$

$$\varphi(a_n; b_i; C_i) \wedge \sim \varphi(a_i; b_n; \tilde{C}_i) \quad \text{and} \quad C_i^{\delta_m^\circ} \cap \tilde{C}_i = \emptyset.$$

Without loss of generality we can assume that $\delta_{m+1}^\circ \subseteq \delta_m^\circ$.

We use \mathcal{F}_K -saturation in structures with two standard sorts S and $K(S)$. Consider the following type that has free variables $\langle x_i; z_i; X_i; \tilde{X}_i : i < \omega \rangle$, where X_i and \tilde{X}_i are tuples variables of sort K

$$p(x_i; z_i; X_i; \tilde{X}_i : i < \omega) = \left\{ \varphi(x_n; z_i; X_i) : i < n < \omega \right\} \cup \left\{ \sim \varphi(a_i; b_n; \tilde{X}_n) : i < n < \omega \right\} \cup$$

$$\{X_i^{\delta_m} \cap \tilde{X}_i = \emptyset : i < m < \omega\}.$$

We leave to the reader to verify that the latter set translates into a set of \mathcal{F}_K -formulas (because it requires that $\langle X_i, \tilde{X}_i \rangle$ belongs to some compact subsets of $K(S)^2$).

Let $\langle a_i; b_i; C_i; \tilde{C}_i : i < \omega \rangle$ be a realization of the above type. Clearly, this witnesses the negation of (1). \square

21 Theorem The following are equivalent

1. $\varphi(x; z; X)$ is stable.
2. $\varphi(x; z; X)$ is ε -stable for every ε .

Proof. $2 \Rightarrow 1$. Given any counter example to (1), let ε be such that $C^\varepsilon \cap \tilde{C} = \emptyset$. To obtain a counter example to (2), set $C_i = C$ and $\tilde{C}_i = \tilde{C}$ for every $i < \omega$.

\triangle $1 \Rightarrow 2$. Negate (2). Pick ε and $\langle a_i; b_i; C_i; \tilde{C}_i : i < \omega \rangle$ that are a counter example to (1) in Definition 20. By the compactness of $K(S)$ in the Vietoris topology, for every δ there are C and \tilde{C} such that for infinitely many $i < \omega$ we have

$$C_i \subseteq C^\delta, \quad C \subseteq C_i^\delta \quad \text{and} \quad \tilde{C}_i \subseteq \tilde{C}^\delta, \quad \tilde{C} \subseteq \tilde{C}_i^\delta.$$

Pick δ such that $2\delta \subseteq \varepsilon$. Refine the sequence so that the inclusions above hold for every i . Then, as $C_i^{2\delta} \cap \tilde{C}_i = \emptyset$, for every $i < \omega$. Then $C^\delta \cap \tilde{C}_i = \emptyset$, for every $i < \omega$. And finally $C \cap \tilde{C} = \emptyset$. \square

22 Definition A **global $\varphi(x; z; X)$ -type** is a maximally (finitely) consistent set of formulas of the form $\varphi(x; b; C)$ or $\sim \varphi(x; b; C)$ for some $b \in \mathcal{U}^{|z|}$ and some C .

It is convenient to have a different characterization of global types, therefore the following definition.

23 Definition We say that \mathcal{D} is **approximable** by $\varphi(x; z; X)$ if for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \varphi(a; b; C)$ and
2. $\langle b, C \rangle \in \mathcal{D} \Leftarrow \varphi(a; b; C)$ for every $b \in B$ and every C .

We remark that, by Facts 10 and 11, when \mathcal{D} is involutive (2) can be rephrased as

$$\langle b, C \rangle \in \sim \mathcal{D} \Rightarrow \sim \varphi(a; b; C) \quad \text{for every } b \in B \text{ and every } C.$$

24 Fact For every involutive \mathcal{D} the following are equivalent

1. \mathcal{D} is approximable by $\varphi(x; z; X)$
2. the following is a global $\varphi(x; z; X)$ -type

$$p(x) = \{ \varphi(x; b; C) : \langle b, C \rangle \in \mathcal{D} \} \cup \{ \sim \varphi(x; b; C) : \langle b, C \rangle \in \sim \mathcal{D} \}.$$

Proof. First we prove that $p(x)$ is maximal as soon as it is consistent – in fact, this only depends on \mathcal{D} being involutive. We need to consider two cases. First, assume that $\varphi(x; b; C)$ is consistent with $p(x)$. Then consistency implies that $\langle b, \tilde{C} \rangle \notin \sim \mathcal{D}$ for every $\tilde{C} \cap C = \emptyset$. But this implies that $C \in \sim \sim \mathcal{D} = \mathcal{D}$, therefore $\varphi(x; b; C) \in p$. The second case is when $\sim \varphi(x; b; C)$ is consistent with $p(x)$. This implies that $\langle b, \tilde{C} \rangle \notin \mathcal{D}$ for every $\tilde{C} \cap C = \emptyset$. Then $\langle b, C \rangle \in \sim \mathcal{D}$, therefore $\sim \varphi(x; b; C) \in p$.

$2 \Rightarrow 1$. Let B be a given finite subset of \mathcal{U}^z . The finite consistency of $p(x)$ provides some a such that for every finite B an a such that

$$\begin{aligned} \langle b, C \rangle \in \mathcal{D} &\Rightarrow \varphi(a; b; C) \quad \text{and} \\ \langle b, C \rangle \in \sim \mathcal{D} &\Rightarrow \sim \varphi(a; b; C) \end{aligned} \quad \text{for every } b \in B \text{ and every } C.$$

By what remarked above, these yield (1) and (2) in Definition 30.

$1 \Rightarrow 2$. Let B be given. Let a be as in Definition 30. Then, by what remarked above, $a \models p(x) \upharpoonright B$. \square

The proof of the main theorem of this section requires the following notion of approximation. The definition is inspired by the classical notion of approximation from below in [Z15] where it is used as rephrasing of the notion of honest definability in [CS13]. Here this notion plays a purely technical role, but we submit it as potentially interesting in itself.

25 Definition We say that \mathcal{D} is **approximable** by $\varphi(x; z; X)$ **from ε -below** if for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that for every C

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \varphi(a; b; C)$ for every $b \in B$ and
2. $\langle b, C^\varepsilon \rangle \in \mathcal{D} \Leftarrow \varphi(a; b; C)$ for every $b \in \mathcal{U}^z$.

26 Theorem Let $\varphi(x; z; X)$ be stable. Assume that \mathcal{D} is approximable by $\varphi(x; z; X)$. Then for every ε there are some $\langle a_{i,j} : i < k, j < m \rangle$ such that for every $b \in \mathcal{U}^z$ and every C

$$\begin{aligned} \langle b, C \rangle \in \mathcal{D} &\Rightarrow \bigvee_{i < k} \bigwedge_{j < m} \varphi(a_{i,j}; b; C^\varepsilon) \\ \langle b, C^\varepsilon \rangle \in \mathcal{D} &\Leftarrow \bigvee_{i < k} \bigwedge_{j < m} \varphi(a_{i,j}; b; C) \end{aligned}$$

Proof. Note that we can take ε sufficiently small that $\varphi(x; z; X)$ is ε -stable. Then theorem is an immediate consequence of the following three lemmas. \square

27 Lemma Let $\varphi(x; z; X)$ be ε -stable. Assume that \mathcal{D} is approximable by $\varphi(x; z; X)$ from ε -below. Then there are some $\langle a_i : i < k \rangle$ such that for every $b \in \mathcal{U}^z$ and every C

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \bigvee_{i < k} \varphi(a_i; b; C^\varepsilon)$
2. $\langle b, C^\varepsilon \rangle \in \mathcal{D} \Leftarrow \bigvee_{i < k} \varphi(a_i; b; C)$

Proof. We define recursively the required parameters a_i together with some auxiliary parameters b_i . The element a_n is chosen such that

3. $\langle b, C^\varepsilon \rangle \in \mathcal{D} \Leftarrow \varphi(a_n; b; C)$ for every $b \in \mathcal{U}^z$ and every C

and

4. $\langle b_i, C \rangle \in \mathcal{D} \Rightarrow \varphi(a_n; b_i; C)$ for every $i < n$ and every C .

This is possible because \mathcal{D} is ε -approximated from below. Note that (3) immediately guarantees (2). Now, assume (1) fails for $k = n$, and choose b_n and C_n witnessing this. Then, by Fact 7, for some $\tilde{C}_n \cap C_n^\varepsilon = \emptyset$

5. $\langle b_n, C_n \rangle \in \mathcal{D} \quad \& \quad \bigwedge_{i < n} \sim \varphi(a_i; b_n; \tilde{C}_n)$

Suppose for a contradiction that the construction never ends. Then, as (5) guarantees that $\langle b_i, C_i \rangle \in \mathcal{D}$ for every i , from (4) we obtain $\varphi(a_n; b_i; C_n)$ for every $i < n$. From (5) we also obtain $\sim \varphi(a_i; b_n; \tilde{C}_n)$, for every $i \leq n$. This contradicts ε -stability. \square

28 Lemma Let $\varphi(x; z; X)$ be ε -stable. Assume that \mathcal{D} is approximable by $\varphi(x; z; X)$. Let m be maximal so that a sequence as in (3) of Definition/Theorem 20 exists. Let $\bar{x} = \langle x_i : i \leq m \rangle$ where the x_i are copies of x . Then the formula

$$\sigma(\bar{x}; z; X) = \bigwedge_{i \leq m} \varphi(x_i; z; X)$$

approximate \mathcal{D} from ε -below.

Proof. Negate the claim and let B witness that $\sigma(\bar{x})$ does not approximate \mathcal{D} from ε -below. Suppose that a_0, \dots, a_{n-1} and b_0, \dots, b_{n-1} have been defined. Choose a_n such that for every $b \in B \cup \{b_0, \dots, b_{n-1}\}$ and every C

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \varphi(a_n; b; C)$ and
2. $\langle b, C^\varepsilon \rangle \in \mathcal{D} \Leftarrow \varphi(a_n; b; C^\varepsilon)$

Note that the latter implication is equivalent to: for every C there is some $\tilde{C} \cap C^\varepsilon = \emptyset$ such that

3. $\langle b, C^\varepsilon \rangle \notin \mathcal{D} \Rightarrow \sim \varphi(a_n; b; \tilde{C}).$

Now, as the lemma is assumed to fail, we can choose b_n and C_n such that

4. $\langle b_n, C_n^\varepsilon \rangle \notin \mathcal{D} \quad \& \quad \bigwedge_{i=0}^n \varphi(a_i; b_n; C_n)$

Note that (4) ensure that $\langle b_i, C_i^\varepsilon \rangle \notin \mathcal{D}$ for every i . There is some \tilde{C}_i that witnesses (3) for $\langle b_i, C_i^\varepsilon \rangle \notin \mathcal{D}$. We claim that the procedure has to stop after $\leq m$ steps. In fact, from (3) we obtain $\sim\varphi(a_n; b_i; \tilde{C}_i)$ for every $i < n$. On the other hand, by (4) we have that $\varphi(a_i; b_n; C_n)$ for every $i < n$. Therefore, $\langle a_{m-i}; b_{m-i}; C_{m-i}; \tilde{C}_{m-i} : i \leq m \rangle$ contradicts the maximality of m . \square

29 Lemma If $\varphi(x; z; C)$ is stable then $\sigma(\tilde{x}; z; C)$ in the previous lemma is ε -stable.

Proof. Let ε be given and let m be maximal such that a sequence as in (3) of Theorem 21 exists. Let k be sufficiently large so that every m -coloring of a graph of size k has a monochromatic subgraph of size $> m$. Let $\langle \tilde{a}_i; b_i; C_i; \tilde{C}_i : i < k \rangle$ be a sequence witnessing instability as in (3) of Theorem 21. Then for every pair $i < n$ there is some $j < m$ such that $\sim\varphi(a_{j,n}; b_i; \tilde{C}_i)$. By the choice of k there is a $j < m$ such that $\sim\varphi(a_{j,n}; b_i; \tilde{C}_i)$ obtains for $> m$ many i . Therefore we can extract a subsequence that contradicts the maximality of m . \square

5. An approximate version of approximability

We present a second version of Theorem 26 which uses a weaker variant of δ -approximability. This notion applies naturally to definable functions which we discuss in the next section.

30 Definition We say that \mathcal{D} is δ -approximable by $\varphi(x; z; X)$ if for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \varphi(a; b; C^\delta)$ and
2. $\langle b, C^\delta \rangle \in \mathcal{D} \Leftarrow \varphi(a; b; C)$ for every $b \in B$ and every C .

We say δ -approximable from ε -below if we can also require that

- 2'. $\langle b, C^{\delta+\varepsilon} \rangle \in \mathcal{D} \Leftarrow \varphi(a; b; C)$ for every $b \in \mathcal{U}$ and every C .

It is evident that, when \mathcal{D} is closed as defined in Section 2 then approximable is equivalent to δ -approximable for every δ .

We restate a variant of Theorem 26. The proof follows closely that in the previous section.

31 Theorem Let $\varphi(x; z; X)$ be stable. Assume that \mathcal{D} is δ -approximable by $\varphi(x; z; X)$. Then for every ε there are some $\langle a_{i,j} : i < k, j < m \rangle$ such that for every $b \in \mathcal{U}^z$ and every C

$$\begin{aligned} \langle b, C \rangle \in \mathcal{D} &\Rightarrow \bigvee_{i < k} \bigwedge_{j < m} \varphi(a_{i,j}; b; C^{\delta+\varepsilon}) \\ \langle b, C^{\delta+\varepsilon} \rangle \in \mathcal{D} &\Leftarrow \bigvee_{i < k} \bigwedge_{j < m} \varphi(a_{i,j}; b; C) \end{aligned}$$

Proof. The theorem is an immediate consequence of the following two lemmas and Lemma 29 in the previous section. \square

32 Lemma Let $\varphi(x; z; X)$ be stable. Assume that \mathcal{D} is δ -approximable by $\varphi(x; z; X)$ from ε -below. Then there are some $\langle a_i : i < k \rangle$ such that for every $b \in \mathcal{U}^z$ and every C

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \bigvee_{i < k} \varphi(a_i; b; C^{\delta+\varepsilon})$
2. $\langle b, C^{\delta+\varepsilon} \rangle \in \mathcal{D} \Leftarrow \bigvee_{i < k} \varphi(a_i; b; C)$

Proof. We define recursively the required parameters a_i together with some auxiliary parameters b_i . The element a_n is chosen such that

3. $\langle b, C^{\delta+\varepsilon} \rangle \in \mathcal{D} \Leftarrow \varphi(a_n; b; C)$ for every $b \in \mathcal{U}^z$ and every C

and

4. $\langle b_i, C \rangle \in \mathcal{D} \Rightarrow \varphi(a_n; b_i; C^\delta)$ for every $i < n$ and every C .

This is possible because \mathcal{D} is δ -approximated from ε -below. Note that (3) immediately guarantees (2). Now, assume (1) fails for $k = n$, and choose b_n and C_n witnessing this. Then, by Fact 7, for some $\tilde{C}_n \cap C_n^{\delta+\varepsilon} = \emptyset$

5. $\langle b_n, C_n \rangle \in \mathcal{D} \ \& \ \bigwedge_{i < n} \sim \varphi(a_i; b_n; \tilde{C}_n)$

Suppose for a contradiction that the construction never ends. Then, as (5) guarantees that $\langle b_i, C_i \rangle \in \mathcal{D}$ for every i , from (4) we obtain $\varphi(a_n; b_i; C_n^\delta)$ for every $i < n$. From (5) we also obtain $\sim \varphi(a_i; b_n; \tilde{C}_n)$, for every $i \leq n$. This contradicts ε -stability. \square

33 Lemma Let $\varphi(x; z; X)$ be ε -stable. Assume that \mathcal{D} is δ -approximable by $\varphi(x; z; X)$. Let m be maximal so that a sequence as in (3) of Definition/Theorem 20 exists. Let $\bar{x} = \langle x_i : i \leq m \rangle$ where the x_i are copies of x . Then the formula

$$\sigma(\bar{x}; z; X) = \bigwedge_{i \leq m} \varphi(x_i; z; X)$$

δ -approximate \mathcal{D} from ε -below.

Proof. Negate the claim and let B witness that $\sigma(\bar{x})$ does not δ -approximate \mathcal{D} from ε -below. Suppose that a_0, \dots, a_{n-1} and b_0, \dots, b_{n-1} have been defined. Choose a_n such that for every $b \in B \cup \{b_0, \dots, b_{n-1}\}$ and every C

1. $\langle b, C \rangle \in \mathcal{D} \Rightarrow \varphi(a_n; b; C^\delta)$ and
2. $\langle b, C^{\varepsilon+\delta} \rangle \in \mathcal{D} \Leftarrow \varphi(a_n; b; C^\varepsilon)$

Note that the latter implication is equivalent to: for every C there is some $\tilde{C} \cap C^\varepsilon = \emptyset$ such that

3. $\langle b, C^{\varepsilon+\delta} \rangle \notin \mathcal{D} \Rightarrow \sim \varphi(a_n; b; \tilde{C})$.

Now, as the lemma is assumed to fail, we can choose b_n and C_n such that

$$4. \quad \langle b_n, C_n^{\varepsilon+\delta} \rangle \notin \mathcal{D} \quad \& \quad \bigwedge_{i=0}^n \varphi(a_i; b_n; C_n)$$

Note that (4) ensure that $\langle b_i, C_i^{\delta+\varepsilon} \rangle \notin \mathcal{D}$ for every i . There is some \tilde{C}_i that witnesses (3) for $\langle b_i, C_i^{\delta+\varepsilon} \rangle \notin \mathcal{D}$. We claim that the procedure has to stop after $\leq m$ steps. In fact, from (3) we obtain $\sim\varphi(a_n; b_i; \tilde{C}_i)$ for every $i < n$. On the other hand, by (4) we have that $\varphi(a_i; b_n; C_n)$ for every $i < n$. Therefore, $\langle a_{m-i}; b_{m-i}; C_{m-i}; \tilde{C}_{m-i} : i \leq m \rangle$ contradicts the maximality of m . \square

6. Stable definable functions

In this section we specialize the notions introduced in the previous sections to formulas $\varphi(x; z; X)$ of the form $\tau(x; z) \in X$ where $\tau(x; z)$ is an S -valued \mathcal{F} -definable function. Notice that $\sim(\tau(x; z) \in X) = \tau(x; z) \in X$.

We say that $\tau(x; z)$ is **stable** if so is $\tau(x; z) \in X$. The following fact is a suggestive characterization of the stability of functions.

34 Fact Let $\tau(x; z)$ be as above. Then the following are equivalent

1. the formula $\tau(x; z) \in X$ is unstable
2. there is a sequence $\langle a_i; b_i : i < \omega \rangle$ such that

$$\lim_{i \rightarrow \infty} \lim_{j \rightarrow \infty} \tau(a_i; b_j) \neq \lim_{j \rightarrow \infty} \lim_{i \rightarrow \infty} \tau(a_i; b_j)$$

3. there is a sequence $\langle a_i; b_i : i < \omega \rangle$ and some ε such that for every $i < j < \omega$

$$\langle \tau(a_i; b_j), \tau(a_j; b_i) \rangle \notin \varepsilon.$$

Proof. $2 \Leftrightarrow 3$ Clear.

$1 \Rightarrow 2$. Let $C \cap \tilde{C} = \emptyset$ and $\langle a_i; b_i : i < \omega \rangle$ be as given by (1). That is, $\tau(a_i; b_j) \in C$ and $\tau(a_j; b_i) \in \tilde{C}$ hold for every $i < j < \omega$. We can restrict to a subsequence such that the two limits exist; C contains the limit on the left; and \tilde{C} contains the limit on the right – which therefore are distinct.

$2 \Rightarrow 1$. Let C and \tilde{C} be disjoint neighborhoods of the two limits in (2). Then (1) is witnessed by a tail of the sequence $\langle a_i; b_i : i < \omega \rangle$. \square

Let $f : \mathcal{U}^z \rightarrow S$ be a function. We define

$$\mathcal{D}_f = \{ \langle b, C \rangle : f(b) \in C, b \in \mathcal{U}^z, C \in K(S) \}$$

35 Fact The following are equivalent

- 1 \mathcal{D}_f is approximable by $\tau(x; z) \in X$
- 2 for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that $\tau(a; b) = f(b)$ for every $b \in B$.

36 Fact (conjecture) The following are equivalent

- 1 \mathcal{D}_f is δ -approximable by $\tau(x; z) \in X$
- 2 for every finite $B \subseteq \mathcal{U}^z$ there is an $a \in \mathcal{U}^x$ such that $\langle \tau(a; b); f(b) \rangle \in \delta$ for every $b \in B$.

References

- [AAVV] Claudio Agostini, Stefano Baratella, Silvia Barbina, Luca Motto Ros, and Domenico Zambella, *Continuous logic in a classical setting* (2025). t.a. in Bull. Iranian Math. Soc.
- [A] Josef Auslander, *Topological Dynamics*, Scholarpedia (2008), doi:10.4249/scholarpedia.3449.
- [BY] Itai Ben Yaacov, *Model theoretic stability and definability of types, after A. Grothendieck*, Bull. Symb. Log. **20** (2014), no. 4, 491–496.
- [BU] Itai Ben Yaacov and Alexander Usvyatsov, *Continuous first order logic and local stability*, Trans. Amer. Math. Soc. **362** (2010), 5213–5259.
- [BBHU] Itai Ben Yaacov, Alexander Berenstein, C. Ward Henson, and Alexander Usvyatsov, *Model theory for metric structures*, Model theory with applications to algebra and analysis. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 350, Cambridge Univ. Press, Cambridge, 2008, pp. 315–427.
- [CS13] Artem Chernikov and Pierre Simon, *Externally definable sets and dependent pairs*, Israel J. Math. **194** (2013), no. 1, 409–425, [arXiv:1007.4468](#).
- [HI] C. Ward Henson and José Iovino, *Ultraproducts in analysis*, Analysis and logic (Mons, 1997), London Math. Soc. Lecture Note Ser., vol. 262, Cambridge Univ. Press, Cambridge, 2002, pp. 1–110.
- [K] H. Jerome Keisler, *Model Theory for Real-valued Structures*, in Beyond First Order Model Theory (José Iovino, ed.), Vol. II, Chapman and Hall/CRC, 2023.
- [PC] Nicolas Chavarría and Anand Pillay, *On pp-elimination and stability in a continuous setting*, Ann. Pure Appl. Logic **174** (2023), 1–14.
- [Z15] Domenico Zambella, *Elementary classes of finite VC-dimension*, Arch. Math. Logic (2015), 511–520.
- [Z] ———, *Standard analysis* (2023), [arXiv:2311.15711](#)