

Scratch paper

Anonymous
Università di Torino
April 2019

Abstract Poche idee, ben confuse.

1 Introduction

2 Samples

The **finite (fractional) expansion** of a structure M is a many-sorted expansion that has a sort for M , which we call the home-sort, a sort for \mathbb{R} , the set of real numbers, and, for every n , a sort for the functions $s : M^n \rightarrow \mathbb{R}$ that are almost always 0.

These functions will be interpreted as **(signed and fractional) multisets**. Namely $s(a)$ interpreted as the number of times a occurs in s . We use the following suggestive notation:

1. $a \in s = s(a)$ this is called the **multiplicity** of a in s ;
2. $|s| = \sum_{a \in M} a \in s$ this is called the **size** of s ;
3. $|\{a \in s : \varphi(a; b)\}| = \sum_{a \models \varphi(x; b)} a \in s$.

We call these multisets **samples** and refer to their sorts collectively as **sample-sort**.

We will write M^f for the expansion above. The language of M^f is denoted by L^f . It expands L , the language of rings, and the language of vector spaces over \mathbb{R} . Each interpreted in the obvious sort. Moreover, L^f has a symbols for the functions in 1-3 above. In particular, 3 requires a symbol for every $\varphi(x; z) \in L$.

Let \mathcal{U} be a monster model of cardinality κ , a cardinal larger than the cardinality of L . We introduce two elementary extension of \mathcal{U}^f .

2.1 Definition We denote by \mathcal{U}^{f*} a saturated elementary extension of \mathcal{U}^f of cardinality κ . As all saturated models of cardinality κ are isomorphic, we can assume that \mathcal{U} is the home-sort of \mathcal{U}^{f*} . We denote by \mathcal{U}^{f*+} some elementary extension of \mathcal{U}^{f*} that realizes all types in $S(\mathcal{U}^{f*})$. Hence we have $\mathcal{U}^f \prec \mathcal{U}^{f*} \prec \mathcal{U}^{f*+}$. Elements of sample-sort in these models are called **finite samples**, **internal samples**, and **(external) samples** respectively. \square

The use of \mathcal{U}^{f*+} is not ideal, it is introduced to replace the use of global types of sample-sort which also have drawbacks.

We write $\text{supp}(s)$ for the **support** of s , that is, the set of those a in \mathcal{U}^{f*} such that $a \in s$ is positive.

Let s be an external sample. For every formula $\varphi(x; z) \in L$ and every $b \in \mathcal{U}^{|z|}$ we define

$$\text{Av}_s \varphi(x; b) = \frac{|\{a \in s : \varphi(a; b)\}|}{|s|},$$

Below, ε always ranges over the positive standard reals. If a and b are hyperreals, we write $a \approx_\varepsilon b$ for $|a - b| < \varepsilon$. We write $a \approx b$ if $a \approx_\varepsilon b$ holds for every ε . We write $\text{st}(a)$, for the standard part of the hyperrational number a . That is, the unique real number μ such that $\mu \approx a$.

The standard part of Av_s induces a finite probability measure on the algebra of definable subsets of $\mathcal{U}^{|x|}$. This measure has a unique extension to a Lebesgue probability measure on a σ -algebra. This is known as the **Loeb measure**.

2.2 Definition We say that a sample s is **smooth** for $\varphi(x; z)$ over M if, for every $s' \equiv_M s$ and every $b \in \mathcal{U}^{|z|}$, we have $\text{Av}_{s'} \varphi(x; b) \approx \text{Av}_s \varphi(x; b)$. □

2.3 Definition Let $\varphi(x; z) \in L$ and let M be a model. An external s is

1. **invariant** if, for every $b' \equiv_M b \in \mathcal{U}^{|z|}$, we have $\text{Av}_s \varphi(x; b) \approx \text{Av}_s \varphi(x; b')$.
2. **definable** if, for every ε , the set $\{b \in \mathcal{U}^{|z|} : \text{Av}_s \varphi(x; b) < \varepsilon\}$ is definable over M .
3. **satisfiable** if, for all $b \in \mathcal{U}^{|z|}$, if $\text{Av}_s \varphi(x; b) \not\approx 0$, then $\varphi(a; b)$ for some $a \in M^{|z|}$.
4. **generically stable** if all of the above holds.

When $\varphi(x; z)$ and M are not clear from the context we add: **for $\varphi(x; z)$ over/in M .**

2.4 Fact Smooth samples are generically stable. (For some given $\varphi(x; z)$ and M .)

Proof We prove in turn 1-3 of the definition above.

1. If s is smooth for $\varphi(x; z)$ over M , then there is an internal sample $s' \in \mathcal{U}^{f*}$ such that $\text{Av}_{s'} \varphi(x; b) \approx \text{Av}_s \varphi(x; b)$ for all $b \in \mathcal{U}^{|z|}$. Then 1 follows.
2. The set $\{b \in \mathcal{U}^{|z|} : \text{Av}_s \varphi(x; b) < \varepsilon\}$ is definable in \mathcal{U}^{f*} because, by invariance of s , we may replace s with any internal sample $s' \equiv_M s$. As this set is invariant over M , it is definable over M^f and therefore, over M .
3. Let $b \in \mathcal{U}^{|z|}$ be such that $\text{Av}_s \varphi(x; b) > \varepsilon$. Pick a sample s' such that $s \equiv_M s' \perp_{M^f} b$. By the smoothness of s , we obtain $\text{Av}_{s'} \varphi(x; b) > \varepsilon$. Let $s_0 \in M^f$ such that $\text{Av}_{s_0} \varphi(x; b) > \varepsilon$. Clearly, some element of $\text{supp}(s_0)$ proves 3. □

Smooth samples are invariant. The converse need not be true in general, but we have the following.

2.5 Fact For every internal sample s , the following are equivalent

1. s is invariant for $\varphi(x; z)$ over M ;

2. s is smooth for $\varphi(x; z)$ over M .

Proof By the fact above, implication $2 \Rightarrow 1$ holds for every sample. We prove $1 \Rightarrow 2$. Let $s \in \mathcal{U}^{f*}$ be an invariant internal sample. Assume for a contradiction that $\text{Av}_{s'} \varphi(x; b) \not\approx_\varepsilon \text{Av}_s \varphi(x; b)$ for some $b \in \mathcal{U}^{|x|}$ and some sample $s' \equiv_M s$. Pick a internal sample $s'' \equiv_{M, s} s'$. Then $\text{Av}_{s''} \varphi(x; b) \not\approx_\varepsilon \text{Av}_s \varphi(x; b)$. As $s'' \equiv_M s$ are both in \mathcal{U}^{f*} , this contradicts the invariance of s . \square

The following is a useful characterization of smoothness.

2.6 Lemma *The following are equivalent for every external sample s*

1. s is smooth over M ;
2. for every $b \in \mathcal{U}^{|z|}$ there is a finite sample $s_0 \in M^f$ such that $\text{Av}_s \varphi(x; b) \approx_\varepsilon \text{Av}_{s_0} \varphi(x; b)$.
(The sample s_0 depends on M , epsilon and b .)
3. there are some finite samples $s_1, \dots, s_n \in M^f$ such that for every $b \in \mathcal{U}^{|z|}$ we have $\text{Av}_s \varphi(x; b) \approx_\varepsilon \text{Av}_{s_i} \varphi(x; b)$ for some s_i . (The samples s_1, \dots, s_n as well as the number n depend on ε and M .)

Proof Implication $3 \Rightarrow 1$ is clear. We prove $1 \Rightarrow 2$, so, fix a $b \in \mathcal{U}^{|z|}$. Let μ be the standard part of $\text{Av}_s \varphi(x; b)$. Let s' be a sample such that $s \equiv_M s' \downarrow_{M^f} b$. By smoothness, $\text{Av}_{s'} \varphi(x; b) \approx \mu$. As $s' \downarrow_{M^f} b$ there is an $s_0 \in M^f$ such that $\text{Av}_{s_0} \varphi(x; b) \approx_\varepsilon \mu$.

We prove $2 \Rightarrow 3$ by compactness. Suppose not, then the following global type is dinitely consistent, hence realized in \mathcal{U}^{f*}

$$p(y) = \left\{ \text{Av}_y \varphi(x, b) \not\approx_\varepsilon \text{Av}_{s_0} \varphi(x; b) : s_0 \in M^f, b \in \mathcal{U}^{|z|} \right\}$$

This is a contradiction. \square

2.7 Definition Let $\varphi(x; z) \in L$. Let $\mathcal{B} \subseteq \mathcal{U}^{|z|}$ be arbitrary. We say that s is **uniform** on \mathcal{B} if for every ε there is a finite sample $s_0 \in M^f$ such that $\text{Av}_s \varphi(x, b) \approx_\varepsilon \text{Av}_{s_0} \varphi(x, b)$ for every $b \in \mathcal{B}$. (The sample s_0 depends on ε and M .) We add **for $\varphi(x; z)$ over M** when these are not clear from the context. \square

2.8 Lemma Let s be uniform on \mathcal{B} and such that $\text{Av}_s \varphi(x; b) > \varepsilon$ for every $b \in \mathcal{B}$. Then there is a finite cover of \mathcal{B} , say $\mathcal{B}_1, \dots, \mathcal{B}_n$, such that all types $p_i(x) = \{\varphi(x; b) : b \in \mathcal{B}_i\}$ are consistent.

Proof Let $s_0 \in M^f$ be as in Definition 2.7. As s_0 is finite, we may assume that $\text{supp}(s_0) = \{a_1, \dots, a_n\}$. Let $\mathcal{B}_i = \{b \in \mathcal{B} : \varphi(a_i; b)\}$. As $\text{Av}_{s_0} \varphi(x; b) > 0$ for all $b \in \mathcal{B}$, these \mathcal{B}_i are the required cover of \mathcal{B} . \square

2.9 Corollary If s be uniform on $\langle b_i : i < \omega \rangle$ for $\varphi(x; z)$ over M , where $\langle b_i : i < \omega \rangle$ is a sequence of indiscernibles such that $\text{Av}_s \varphi(x; b_i) > \varepsilon$ for every $i < \omega$, then the type $\{\varphi(x; b_i) : i < \omega\}$ is consistent. \square

2.10 Corollary *Let s be uniform on \mathcal{C} for $\varphi(x; z)$ over M . Then $\text{Av}_s[\psi(x; c) \nleftrightarrow \psi(x; c')] > \varepsilon$ does not hold for all $c, c' \in \mathcal{C}, c \neq c'$.*

Proof Suppose for a contradiction that a set \mathcal{C} as above exists. Apply Lemma 2.8 to the formula $\varphi(x; z, z') = [\psi(x; z) \nleftrightarrow \psi(x; z')]$ and the set $\mathcal{B} = \{\langle c, c' \rangle \in \mathcal{C}^2 : c \neq c'\}$. The sets \mathcal{B}_i obtained from the lemma induce a finite coloring of the complete graph on \mathcal{C} . By the Ramsey theorem there is an infinite monochromatic set $A \subseteq \mathcal{C}$. Hence $\{\psi(x; a) \nleftrightarrow \psi(x; a') : a, a' \in A, a \neq a'\}$ is consistent. As $|A| > 2$ this is impossible. \square

3 The nip formulas

3.1 Theorem *Let $\varphi(x; z) \in L$ be a nip formula. Then every sample is uniform on M over M .*

Proof Questo è Vapnik-Chervonenkis. \square

For smooth samples the Vapnik-Chervonenkis Theorem can be strengthened as follows. (The difference is that below b ranges over $\mathcal{U}^{|z|}$.)

3.2 Corollary *Let $\varphi(x; z) \in L$ be nip. Every smooth sample s is uniform for $\varphi(x; z)$ on \mathcal{U} over M .*

Proof By Theorem 3.1, there is an $s_0 \in M^f$ such that $\text{Av}_s \varphi(x; b) \approx_\varepsilon \text{Av}_{s_0} \varphi(x; b)$ for all $b \in M^{|z|}$.

Suppose for a contradiction that $\text{Av}_s \varphi(x; b') \not\approx_\varepsilon \text{Av}_{s_0} \varphi(x; b')$ for some $b' \in \mathcal{U}^{|z|}$. Pick a $s' \subseteq \mathcal{U}^{|x|}$ such that $b' \downarrow_M s' \equiv_M s$. By the smoothness of s , we obtain $\text{Av}_{s'} \varphi(x; b') \approx_\varepsilon \text{Av}_{s_0} \varphi(x; b')$. As $b' \downarrow_M s'$, the same formula holds for some $b \in M^{|z|}$. Once again by smoothness, $\text{Av}_s \varphi(x; b) \not\approx_\varepsilon \text{Av}_{s_0} \varphi(x; b)$. A contradiction. \square

3.3 Definition *A formula $\varphi(x; z) \in L$ is **distal** if there is a formula $\psi(x; z_1, \dots, z_n) \in L$ such that for every finite set $A \subseteq \mathcal{U}^f$ and every $b \in \mathcal{U}^{|x|}$ there are $a_1, \dots, a_n \in \text{supp}(s)$ such that $\psi(x; a_1, \dots, a_n) \leftrightarrow p(x)$, where $p(x) = \text{tp}_\varphi(b/A)$.* \square

3.4 Theorem (false) *The following are equivalent*

1. $\varphi(x; z) \in L$ is distal;
2. for every sample $s \in \mathcal{U}^{f+}$, if s is generically stable for $\varphi(x; z)$ over M then s is smooth for $\varphi(x; z)$ over M .