

Esercizio 1. Let $\varepsilon(x, y) \subseteq L(A)$ be a type-definable equivalence relation with $< \lambda$ classes. Prove that every λ -saturated model containing A intersects every class of $\varepsilon(x, y)$.

Esercizio 2. Let $L = \{<\}$ and let N be a ω_1 -saturated extension of \mathbb{Q} . Prove that there is an embedding $f: \mathbb{R} \rightarrow N$. Is it elementary? Is it an isomorphism?

Esercizio 3. Let $L = \{<\}$ and let N be a saturated extension of \mathbb{Q} . Prove that there $2^{|N|}$ Dedekind cuts of N .

Esercizio 4. Let M be an arbitrary structure of cardinality larger than $|L|$. Prove that M has an ω -homogeneous elementary extension of the same cardinality.

Esercizio 5. Let M and N be elementarily homogeneous structures of the same cardinality λ . Suppose that $M \models \exists x p(x) \Leftrightarrow N \models \exists x p(x)$ for every $p(x) \subseteq L$ such that $|x| < \lambda$. Prove that the two structures are isomorphic.

Esercizio 6. Let L be a countable language containing L_{gr} and assume \mathcal{U} is a group. Prove that the following are equivalent

1. every model M is a normal subgroup of \mathcal{U}
2. some ω -saturated model M is a normal subgroup of \mathcal{U}
3. \mathcal{U} is a BFC group.

A group G is BFC (has boundedly finite conjugacy classes) if $\{g a g^{-1} : g \in G\}$, as a ranges over G , have at most n elements, for some fixed n .

Esercizio 7. Prove that for every regular $\lambda \geq |L|$ every theory T with an infinite model has a model that is λ -saturated and λ -homogeneous.

Esercizio 8. Let λ be such that every first-order theory T with an infinite model has a saturated model of cardinality λ . Prove that $\lambda = \lambda^{<\lambda}$.