Esercizio 1. (Martina Apicella & Andrea Morello) Let $a \equiv_M b$. Prove that there is a sequence $\bar{c} = \langle c_i : i < \omega \rangle$ such that $a, \bar{c}$ and $b, \bar{c}$ are both sequences of M-indiscernibles. Does the claim hols with an arbitrary set A for M?

Esercizio 2. (Beatrice Calderaro & Gabriele Maiolino) Let $\langle c_i : i < \omega \rangle$ be an indiscernible sequence. Prove that there is an indiscernible sequence $\langle d_i : i < \omega \rangle$ such that $d_0, d_1 = c_1, c_0$.

Esercizio 3. (Mattia Lo Mele & Daniele Lanciotti) Let $\langle \mathcal{D}_i : i < \omega \rangle$ be an A-indiscernible sequence in $\mathcal{U}^{\mathrm{eq}}$. Prove that there is a formula $\varphi(x; z) \in L$ and an A-indiscernible sequence $\langle b_i : i < \omega \rangle$ in $\mathcal{U}^z$ such that $\mathcal{D}_i = \varphi(\mathcal{U}; b_i)$.

Esercizio 4. (Francesco Dam & Filippo Giugie) Let M be a graph with the property that for every finite $A \subseteq M$ there is a $c \in M$ such that $A \subseteq r(c, M)$. The edges of M are colored with infinitely many colors but every clique contains at most k colors, for some fixed $k \in \omega$. Prove that M has an infinite monochromatic complete subgraph.