



## 2020 2021 DEqs S2 Midterm

Electrical Engineering (Trường Đại học Quốc tế, Đại học Quốc gia Thành phố Hồ Chí Minh)



Scan to open on Studocu

## MIDTERM EXAMINATION

Academic year 2020-2021, Semester 2

Duration: 120 minutes

<b>SUBJECT:</b> <b>Differential Equations (MAFE202IU)</b>	
Head of Department of Mathematics	Lecturer:
Signature:	Signature:
Professor Pham Huu Anh Ngoc	Full name: Pham Huu Anh Ngoc

**Instructions:**

- Each student is allowed a scientific calculator and a maximum of two double-sided sheets of reference material (size A4 or similar), stapled together and marked with their name and ID. All other documents and electronic devices are forbidden.

**Question 1.** (20 marks) (Memorization) When forgetfulness is taken into account, the rate of memorization of a subject is given by

$$\frac{dA}{dt} = k_1(M - A) - k_2A, \quad A(0) = 0$$

where  $k_1 > 0, k_2 > 0$ ,  $A(t)$  is the amount memorized in time  $t$ ,  $M$  is the total amount to be memorized and  $M - A$  is the amount remaining to be memorized.

Find the limiting value of  $A(t)$  as  $t \rightarrow \infty$ . Interpret the result.

**Question 2.** (20 marks) Prove that the differential equation

$$(\cos x - xy^2 + y)dx + (y(1 - x^2) + x)dy = 0,$$

is exact. Solve the differential equation.

**Question 3.** (20 marks) Find the solution to the initial value problem

$$xy' + (3x + 1)y = e^{-3x}, \quad y(1) = 0.$$

**Question 4.** (20 marks) Find a particular solution of the following differential equation

$$y'' - 8y' + 16y = 6x^2 + 2020 + 2021e^{4x}.$$

**Question 5.** (20 marks) Find the general solution of the following differential equation

$$x^2y'' - 5xy' + 9y = x^2, \quad x \in (0, \infty).$$

END.

# SOLUTIONS:

**Question 1.** The limiting value of the population is 1,000,000. The population will reach 500,000 in 5.29 months.

**Question 2.** The given differential equation is rewritten as

$$(e^{2y}dx + xde^{2y}) - \cos(xy)(xdy + ydx) + dy^2 = 0.$$

Then, we get

$$d(e^{2y}x) - \cos(xy)d(xy) + dy^2 = d(e^{2y}x) + d(-\sin(xy)) + dy^2 = 0.$$

Therefore,

$$d(e^{2y}x - \sin(xy) + y^2) = 0.$$

Thus the general solution is given by

$$e^{2y}x - \sin(xy) + y^2 = C.$$

**Question 3.** Consider the differential equation

$$y' - (\sin x)y = 2 \sin x.$$

The integrating factor is given by  $I(x) = e^{\cos x}$ . Thus, we get

$$e^{\cos x}y' - e^{\cos x}(\sin x)y = 2e^{\cos x} \sin x.$$

This gives

$$\frac{d}{dx}(e^{\cos x}y) = 2 \int e^{\cos x} \sin x dx = -2e^{\cos x} + C.$$

Therefore, the general solution is

$$y(x) = -2 + \frac{C}{e^{\cos x}}.$$

Since  $y(\frac{\pi}{2}) = 1$ , the particular solution is  $y(x) = -2 + \frac{3}{e^{\cos x}}$ .

**Question 4.** a) The form of a particular solution of the differential equation

$$y'' - 4y' + 3y = e^{2x}(x^3 + 1) + e^x(x + 1)$$

is given by

$$y_p(x) = e^{2x}(Ax^3 + Bx^2 + Cx + D) + e^x(Ex^2 + Fx).$$

The general solution of the differential equation

$$y'' - 4y' + 3y = e^x(x + 1)$$

is given by

$$y(x) = c_1e^x + c_2e^{3x} - e^x(\frac{1}{4}x^2 + \frac{3}{4}x).$$

**Question 5.** a)  $a = b = q, q \in \mathbb{R}$ .

b) Note that  $y_1(x) = x + 1$  is a particular solution of the differential equation

$$(1 - 2x - x^2)y'' + 2(x + 1)y' - 2y = 0.$$

By the Liouville formula,  $y_2(x) = x^2 + x + 2$  is a solution of this equation such that  $y_1, y_2$  are linearly independent. So, the general solution is given by

$$y(x) = c_1(x + 1) + c_2(x^2 + x + 2).$$