

SAMPLE OF FINAL EXAMINATION

Semester XXX • Date: XXX • Duration: 120 minutes

INSTRUCTIONS:

Each student is allowed a scientific calculator and a maximum of TWO double-sided sheets of reference material (size A4 or similar) marked with their name and ID. All other documents and electronic devices are forbidden.

Each question carries 10 marks.

.....

Question 1. Show that the function $U(x, y) = e^{-x} \sin(y)$ satisfies the Laplace equation

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} = 0.$$

Question 2. Find the equation of the plane tangent to the surface

$$z(x, y) = x^2 e^{x-y}$$

at the point $(2, 2, 4)$.

Question 3. Find the critical points of the following function $f(x, y)$, and determine whether each critical point corresponds to a local maximum, local minimum or a saddle point?

$$f(x, y) = x^4 + 2y^2 - 4xy.$$

Question 4. Using Lagrange multipliers, find the absolute maximum and minimum values of the function

$$f(x, y) = x - y \quad \text{subject to} \quad g(x, y) = x^2 + y^2 - 3xy = 20.$$

Question 5. Evaluate the integral $\iint_D y^2 dA$ where D is bounded by: $x = 1, y = 2x+2, y = -x-1$.

Question 6. Find the volume of the solid bounded by the plane $z = 0$ and hyperboloid $z = 3 - \sqrt{1 + x^2 + y^2}$.

Please turn over...

Question 7. Evaluate the line integral $\int_C xy ds$, where C is a portion of the ellipse

$$\frac{x^2}{4} + \frac{y^2}{16} = 1$$

in the first quadrant, oriented counterclockwise.

Question 8. Find $\iiint_E z \, dV$, where E is the solid tetrahedron with vertices $(0, 0, 0)$, $(4, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 5)$.

Question 9. Let D be part of the annulus $\{(x, y) : 1 \leq x^2 + y^2 \leq 9\}$ that lies in the upper half plane. Let C be the boundary of D , positively oriented. Compute

$$\int_C (xy + e^{x^2}) dx + (\sin(\sqrt{y}) + x^2) dy.$$

Question 10. Let $F = \langle 2x + 3y, 3x - 2y \rangle$. Find $f(x, y)$ such that $F = \nabla f$.

—END OF QUESTION PAPER—

Short answer

Q.1

Q.2 $f_x = (2x + x^2)e^{x-y}$, $f_y = -x^2e^{x-y}$. So

$$x = -x - y - 1$$

Q3. $f_x = 4x^3 - 4y$; $f_y = 4y - 4x$; $x_{xx} = 12x^2$, $f_{yy} = 4$, $f_{xy} = -4$, and $D(x, y) = 38x^2 - 16$

$D(0, 0) = -16 < 0 \rightarrow f$ has saddle point at $(0, 0)$

$D(1, 1) = D(-1, -1) = 32 > 0$; $f_{xx}(1, 1) = f_{xx}(-1, -1) = 12 > 0 \Rightarrow f$ has local minimum at $(1, 1)$ and $(-1, -1)$

Q.4 $\nabla f(x, y) = \langle 1, -1 \rangle$; $\nabla g(x, y) = \langle 2x - 3y, 2y - 3x \rangle$. Lagrang multiplier is

$$\begin{cases} 2x - 3y = 1/\lambda \\ 2y - 3x = -1/\lambda \\ x^2 + y^2 - 3xy = 20 \end{cases}$$

$\Rightarrow (x, y) = (\pm 2, \pm 2)$. So maximum $f(x, y) = 4$ at $(2, -2)$, minimum $f(x, y) = -4$ at $(-2, 2)$

Q5. $I = \int_{-1}^1 \int_{-x-1}^{2x+2} y^2 dy dx = 12$

Q6. $V = \int_0^{2\pi} \int_0^{2\sqrt{2}} \int_0^{3-\sqrt{1+r^2}} 1 dz dr d\theta = \frac{20\pi}{3}$

Q7. $r(t) = \langle 2 \cos(t), 4 \sin(t) \rangle$, $0 \leq t \leq \pi/2$, and

$$|r'(t)| = \sqrt{4 \cos^2(t) + 16 \sin^2(t)} = 2\sqrt{1 + 3 \cos^2(t)}$$

$$I = \int_0^{\pi/2} 16 \sin(t) \cos(t) \sqrt{1 + 3 \cos^2(t)} dt = 112/9$$

Q8. The equation of plane passing given vertices is $\frac{x}{4} + y + \frac{z}{5} = 1$, and hence

$$E = \{(x, y, z) : 0 \leq x \leq 4, 0 \leq y \leq 1 - \frac{x}{4}, 0 \leq z \leq 5 - \frac{5x}{4} - 5y\}.$$

Thus,

$$\iiint_E z dV = \int_0^4 \int_0^{1-\frac{x}{4}} \int_0^{5-\frac{5x}{4}-5y} z dx dy dz = \frac{25}{6} \approx 4.1667.$$

Q9. By Green's theorem, it equals $\iint_D x dA = 0$.

Q10. $f(x, y) = x^2 + 3xy - y^2 + C$.