


# Applied Linear Algebra

## Final

Tử Queng Khuông mến soạn tặng



Special thanks : 88 HN      88 TNgan      88 Duy Võ  
88 BND      88 Thuy      88 HNM  
88 Nguyen      88 DThao      88 LO  
88 YI      88 Ming      88 DHien  
88 Wmyu      88 TL  
88 Lkuc      88 Muyu  
88 Kae      88 Dakoe  
88 Moon      88 MThao

**Aladin**

**iem-  
buddy**  
College Tutoring

4 mức độ a - z :

(4) : Chắc chắn sẽ ra thi

(3) : Quan trọng , 70% ra thi

(2) : Dễ , chỉ là kỹ năng

(1) : Mang tính lý thuyết

lum  
88 Quang Khương

### Question 1

1. Find eigenvalues and eigenvectors of the following matrices

a)

$$A = \begin{pmatrix} 4 & -5 \\ 2 & -3 \end{pmatrix}$$

(4)

b)

$$A = \begin{pmatrix} 6 & -3 \\ -2 & 1 \end{pmatrix}$$

lun  
88 Quang Khadong

a)  $A - \lambda I = \begin{pmatrix} 4 - \lambda & -5 \\ 2 & -3 - \lambda \end{pmatrix}$

$\Rightarrow \det(A - \lambda I) = (4 - \lambda)(-3 - \lambda) + 10 = 0$

$\Leftrightarrow -12 - 4\lambda + 3\lambda + \lambda^2 + 10 = 0$

$\Leftrightarrow \begin{cases} \lambda_1 = 2 \\ \lambda_2 = -1 \end{cases} \quad (\text{Eigen value})$

Now we find eigen vectors:

$$\begin{aligned} N(A - \lambda_1 I) &= \left( \begin{array}{cc|c} 2 & -5 & 0 \\ 2 & -5 & 0 \end{array} \right) \xrightarrow{\text{RREF}} \left( \begin{array}{cc|c} 1 & -\frac{5}{2} & 0 \\ 0 & 0 & 0 \end{array} \right) \Rightarrow x_1 - \frac{5}{2}x_2 = 0 \\ &\Rightarrow \begin{cases} x_1 = \frac{5t}{2} \\ x_2 = t \end{cases} \quad \text{Eigen vector: } t \begin{pmatrix} \frac{5}{2} \\ 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} N(A - \lambda_2 I) &= \left( \begin{array}{cc|c} 5 & -5 & 0 \\ 2 & -2 & 0 \end{array} \right) \xrightarrow{\text{RREF}} \left( \begin{array}{cc|c} 1 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right) \quad x_1 - x_2 = 0 \\ &\Rightarrow x_1 = x_2 = t \quad \text{Eigen vector: } t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \end{aligned}$$

$\Rightarrow \begin{cases} x_1 = x_2 = t \\ x_2 = x_2 = t \end{cases} \quad \text{Eigen vector: } t \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

### Question 2

2. Find eigenvalues and eigenvectors of the following matrix

$$A = \begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ 1 & -2 & 0 \end{pmatrix}$$

(2)

$A - \lambda I = \begin{bmatrix} -2 - \lambda & 2 & -3 \\ 2 & 1 - \lambda & -6 \\ 1 & -2 & -\lambda \end{bmatrix} \quad \begin{matrix} (-2 - \lambda) & 2 \\ 2 & (1 - \lambda) \\ 1 & -2 \end{matrix}$

$$\begin{aligned} \det(A - \lambda I) &= (-2 - \lambda)(1 - \lambda)(-\lambda) + (-12) + 12 - (1 - \lambda)(-3) - (-\lambda)(-6)(-2 - \lambda) \\ &\quad - (-\lambda)(2)(2) \end{aligned}$$

$$\begin{aligned}
 &= (-2 + \lambda + \lambda^2)(-\lambda) + 3 - 3\lambda - 12(-2 - \lambda) + 4\lambda \\
 &= -\lambda^3 - \lambda^2 + 2\lambda + 3 - 3\lambda + 24 + 12\lambda + 4\lambda \\
 &= -\lambda^3 - \lambda^2 + 15\lambda + 27
 \end{aligned}$$

. Let  $\det(A - \lambda I) = 0$

$$(E) \quad -\lambda^3 - \lambda^2 + 15\lambda + 27 = 0$$

(E)

$$\begin{cases} \lambda_1 = -3 \\ \lambda_2 = 1 + \sqrt{10} \\ \lambda_3 = 1 - \sqrt{10} \end{cases} \quad \left( \text{Eigen values} \right)$$

lum  
88 Quang Khadong

Now we find eigen vectors:

$$N(A - \lambda_1 I) = \left[ \begin{array}{ccc|c} 1 & 2 & -3 & 0 \\ 2 & 4 & -6 & 0 \\ 1 & -2 & 3 & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & -\frac{3}{2} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\begin{aligned} \Rightarrow x_1 &= 0 \\ x_2 &= \frac{3}{2}x_3 \\ x_3 &= x_3 = t \end{aligned}$$

$$\begin{bmatrix} 0 \\ \frac{3t}{2} \\ t \end{bmatrix} \rightarrow t \begin{bmatrix} 0 \\ \frac{3}{2} \\ 1 \end{bmatrix}$$

Eigen space

$$t \begin{bmatrix} 0 \\ \frac{3}{2} \\ 1 \end{bmatrix}$$

Eigen vector

$$N(A - \lambda_2 I) = \left[ \begin{array}{ccc|c} -3 - \sqrt{10} & 2 & -3 & 0 \\ 2 & -\sqrt{10} & -6 & 0 \\ 1 & -2 & -1 - \sqrt{10} & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{ccc|c} 1 & 0 & \frac{1 + \sqrt{10}}{3} & 0 \\ 0 & 1 & \frac{2(1 + \sqrt{10})}{3} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\begin{aligned} \Rightarrow x_1 &= -\frac{1 + \sqrt{10}}{3}t \\ x_2 &= -\frac{2(1 + \sqrt{10})}{3}t \\ x_3 &= x_3 = t \quad (\text{free variable}) \end{aligned}$$

$$\Rightarrow t \begin{bmatrix} -\frac{1 + \sqrt{10}}{3} \\ -\frac{2(1 + \sqrt{10})}{3} \\ 1 \end{bmatrix} \quad (\text{Eigen vector})$$

$$N(\lambda - \lambda_3 I) = \left[ \begin{array}{ccc|c} -3 + \sqrt{10} & 2 & -3 & 0 \\ 2 & +\sqrt{10} & -6 & 0 \\ 1 & -2 & -1 + \sqrt{10} & 0 \end{array} \right] \xrightarrow{\text{REF}} \left[ \begin{array}{ccc|c} 1 & 0 & -\frac{-1 + \sqrt{10}}{3} & 0 \\ 0 & 1 & -\frac{2(-1 + \sqrt{10})}{3} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\Rightarrow \begin{cases} x_1 = \frac{-1 + \sqrt{10}}{3} t \\ x_2 = \frac{2(-1 + \sqrt{10})}{3} t \\ x_3 = x_3 = t \end{cases} \quad \Rightarrow t \begin{bmatrix} \frac{-1 + \sqrt{10}}{3} \\ \frac{2(-1 + \sqrt{10})}{3} \\ 1 \end{bmatrix} \quad (\text{Eigen vector})$$

88 Quang Khue

Question 1:

1. Let matrix

$$A = \begin{pmatrix} 1 & -4 & 9 & -7 \\ -1 & 2 & -4 & 1 \\ 5 & -6 & 10 & 7 \end{pmatrix}$$

Find rank(A), nullity of A ( $\dim \text{Nul}(A)$ ).

(4)

$$A = \begin{pmatrix} 1 & -4 & 9 & -7 \\ -1 & 2 & -4 & 1 \\ 5 & -6 & 10 & 7 \end{pmatrix} \quad R_2 \leftarrow R_1 + R_2 \\ R_3 \leftarrow 5R_2 + R_3$$

$$A = \begin{pmatrix} 1 & -4 & 9 & -7 \\ 0 & -2 & 5 & -6 \\ 0 & 4 & -10 & 12 \end{pmatrix} \quad R_3 \leftarrow 2R_2 + R_3$$

$$A = \begin{pmatrix} \text{pivot} & 1 & -4 & 9 & -7 \\ 0 & \text{pivot} & -2 & 5 & -6 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

lum  
88 Quang Khê Long

$$2 \text{ pivots} \Rightarrow \text{Rank}(A) = 2 \\ \dim \text{Nul}(A) = 2$$

(Rank là số pivot columns  
 $\dim \text{Nul}$  là số cột ko có pivot)

Question 2:

2. Let  $B = \{(8, 11, 0), (7, 0, 10), (1, 4, 6)\}$  be a basis of  $\mathbb{R}^3$ . Find the coordinate of vector  $x = (3, 19, 2)$ .

$$x = \begin{bmatrix} 3 \\ 19 \\ 2 \end{bmatrix}$$

$$B = \left\{ \begin{bmatrix} 8 \\ 11 \\ 0 \end{bmatrix}, \begin{bmatrix} 7 \\ 0 \\ 10 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 6 \end{bmatrix} \right\}$$

$$A\bar{x} = \bar{b} \quad [x]_B, B = v_1, v_2, v_3$$

$$\alpha v_1 + \beta v_2 + \gamma v_3 = x$$

$$\alpha \begin{bmatrix} 8 \\ 11 \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 7 \\ 0 \\ 10 \end{bmatrix} + \gamma \begin{bmatrix} 1 \\ 4 \\ 6 \end{bmatrix} = \begin{bmatrix} 3 \\ 19 \\ d \end{bmatrix}$$

$$\begin{cases} 8\alpha + 7\beta + \gamma = 3 \\ 11\alpha + 4\beta = 19 \\ 10\beta + 6\gamma = 2 \end{cases}$$

$$\Rightarrow \begin{cases} \alpha = 1 \\ \beta = -1 \\ \gamma = 2 \end{cases}$$

$$\Rightarrow [\alpha]_B = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$$

num  
88 Queng Kholidng

### Question 3

3. Find the dimension and basic of the subspace

$$H = \left\{ \begin{pmatrix} a - 4b - 2c \\ 2a + 5b - 4c \\ -a + 2c \\ -3a + 7b + 6c \end{pmatrix}, a, b, c \in \mathbb{R} \right\}$$

(4)

$$H = \left\{ a \begin{bmatrix} 1 \\ 2 \\ -1 \\ -3 \end{bmatrix} + b \begin{bmatrix} -4 \\ 5 \\ 0 \\ 7 \end{bmatrix} + c \begin{bmatrix} -2 \\ -4 \\ 2 \\ 6 \end{bmatrix} \right\}$$

$$\Rightarrow H = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \\ -3 \end{bmatrix}, \begin{bmatrix} -4 \\ 5 \\ 0 \\ 7 \end{bmatrix}, \begin{bmatrix} -2 \\ -4 \\ 2 \\ 6 \end{bmatrix} \right\}$$

Basis of  $H$  = basis of  $\text{col}(A)$  where

$$A = \begin{pmatrix} 1 & -4 & -2 \\ 2 & 5 & -4 \\ -1 & 0 & 2 \\ -3 & 7 & 6 \end{pmatrix}$$

$R_2 \leftarrow (-2)R_1 + R_2$   
 $R_3 \leftarrow R_1 + R_3$   
 $R_4 \leftarrow 3R_1 + R_4$

$$A = \begin{pmatrix} 1 & -4 & -2 \\ 0 & 13 & 0 \\ 0 & -4 & 0 \\ 0 & -5 & 0 \end{pmatrix}$$

$R_3 \leftarrow 13/4 R_3 + R_2$   
 $R_4 \leftarrow 13/5 R_4 + R_2$

$$A = \begin{pmatrix} 1 & -4 & -2 \\ 0 & 13 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

pivot

lum  
88 Quang Kháng

pivot ở 2 cột đầu  $\Rightarrow$  basis là 2 cột đầu

BASIS

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \\ -3 \end{bmatrix}, \begin{bmatrix} -4 \\ 5 \\ 0 \\ 7 \end{bmatrix} \right\}$$

Since there are 2 vectors (2 rows)

$$\Rightarrow \text{DIM} = 2$$

#### Question 4

4. Let  $S = \{(1, -4), (3, -5)\}$ , and  $T = \{(-9, 1), (-5, -1)\}$  be bases in  $\mathbb{R}^2$ .

a. Find the change of coordinates matrix from  $T$  to  $S$ .

$$[v]_T$$

(4)

b. Let vector  $v = (1, -2)$ . Find the coordinate of  $v$  in the basis  $T$  then use the transition matrix at question a) to find the coordinate of  $v$  in the basis  $S$ .

$$[v]_S$$

$$S = \left\{ \begin{bmatrix} 1 \\ -4 \end{bmatrix}, \begin{bmatrix} 3 \\ -5 \end{bmatrix} \right\}$$

$$T = \left\{ \begin{bmatrix} -9 \\ 1 \end{bmatrix}, \begin{bmatrix} -5 \\ -1 \end{bmatrix} \right\}$$

a) Find  $\mathbf{I}_{T \rightarrow S}$

$$\left( \begin{array}{cc|cc} S & T \\ 1 & 3 & -9 & -5 \\ -4 & -5 & 1 & -1 \end{array} \right)$$

$$\Rightarrow \left( \begin{array}{cc|cc} 1 & 0 & 6 & 4 \\ 0 & 1 & -5 & -3 \end{array} \right)$$

$$\Rightarrow \mathbf{I}_{T \rightarrow S} = \begin{pmatrix} 6 & 4 \\ -5 & -3 \end{pmatrix}$$

lum  
88 Quang Khaldng

b) Find  $[v]_T$ :  $\left( \begin{array}{cc|c} -9 & -5 & 1 \\ 1 & -1 & -2 \end{array} \right)$

$$\left( \begin{array}{cc|c} 1 & 0 & -11/14 \\ 0 & 1 & 13/14 \end{array} \right)$$

$$\Rightarrow [v]_T = \begin{pmatrix} -11/14 \\ 13/14 \end{pmatrix}$$

. Find  $[v]_S$  :  $[v_S] = \mathbf{I}_{T \rightarrow S} [v_T]$

$$= \begin{pmatrix} 6 & 4 \\ -5 & -3 \end{pmatrix} \begin{pmatrix} -11/14 \\ 13/14 \end{pmatrix} = \begin{pmatrix} 1/7 \\ 2/7 \end{pmatrix}$$

# Sample Test |

(4)

Question 1

QUESTION 1 Find the inverse of matrix

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 2 & 2 & 1 \\ 4 & -1 & 3 \end{bmatrix}$$

Nhớ c/m  $\det \neq 0$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 2 & 2 & 1 & 0 & 1 & 0 \\ 4 & -1 & 3 & 0 & 0 & 1 \end{array} \right) \quad \begin{aligned} R_2 &\leftarrow (-2)R_1 + R_2 \\ R_3 &\leftarrow (-4)R_1 + R_3 \end{aligned}$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 2 & -3 & -2 & 1 & 0 \\ 0 & -1 & -5 & -4 & 0 & 1 \end{array} \right) \quad \begin{aligned} R_1 &\leftarrow 2R_3 + R_2 \\ R_3 &\leftarrow (-1)R_3 \end{aligned}$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & -13 & -10 & 1 & 2 \\ 0 & 1 & 5 & 4 & 0 & -1 \end{array} \right) \quad \begin{aligned} R_2 &\leftarrow -\frac{1}{13}R_2 ; R_2 \leftrightarrow R_3 \end{aligned}$$

lumen  
88 Quang Khaldong

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 1 & 5 & 4 & 0 & -1 \\ 0 & 0 & 1 & \frac{10}{13} & -\frac{1}{13} & -\frac{2}{13} \end{array} \right) \quad \begin{aligned} R_1 &\leftarrow R_1 + (-2)R_3 \\ R_2 &\leftarrow R_2 + (-5)R_3 \end{aligned}$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{7}{13} & \frac{2}{13} & \frac{4}{13} \\ 0 & 1 & 0 & \frac{4}{13} & \frac{5}{13} & -\frac{3}{13} \\ 0 & 0 & 1 & \frac{10}{13} & -\frac{1}{13} & -\frac{2}{13} \end{array} \right) \quad \Rightarrow A^{-1} = \left( \begin{array}{ccc} -\frac{7}{13} & \frac{2}{13} & \frac{4}{13} \\ \frac{4}{13} & \frac{5}{13} & -\frac{3}{13} \\ \frac{10}{13} & -\frac{1}{13} & -\frac{2}{13} \end{array} \right)$$

Question 2

QUESTION 2 Find the rank and the base of column space and null space of matrix

$$A = \begin{bmatrix} 1 & 6 & 2 & -4 \\ -3 & 2 & -2 & -8 \\ 4 & -1 & 3 & 9 \end{bmatrix} \quad \begin{aligned} R_1 &\leftarrow R_1 + 6R_3 \\ R_2 &\leftarrow 3R_1 + R_2 \\ R_3 &\leftarrow (-4)R_1 + R_3 \end{aligned}$$

$$\left( \begin{array}{cccc|c} 25 & 0 & 20 & 50 & \\ 0 & 20 & 4 & -20 & \\ 0 & -25 & -5 & 25 & \end{array} \right) \quad \begin{aligned} R_1 &\leftarrow 1/5 R_1 \\ R_2 &\leftarrow 1/4 R_2 \\ R_3 &\leftarrow -1/5 R_3 \end{aligned}$$

$$\left( \begin{array}{cccc} 5 & 0 & 4 & 10 \\ 0 & 5 & 1 & -5 \\ 0 & 5 & 1 & -5 \end{array} \right) \quad \begin{aligned} R_1 &\leftarrow \frac{1}{5}R_1 \\ R_2 &\leftarrow R_2 \\ R_3 &\leftarrow R_3 - R_2 \end{aligned}$$

$$\left( \begin{array}{cc|cc} 1 & 0 & \frac{4}{5} & 2 \\ 0 & 1 & \frac{1}{5} & -1 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad \text{pivot column}$$

$$\Rightarrow \text{Rank}(A) = 2$$

Basis of  $A^\perp$ :

$$\left\{ \begin{pmatrix} 1 \\ -3 \\ 4 \end{pmatrix}, \begin{pmatrix} 6 \\ 2 \\ -1 \end{pmatrix} \right\}$$

Homework  
88 Quang Khê Long

We solve for  $A = 0 \Leftrightarrow$

$$\begin{cases} x_1 + \frac{4}{5}x_3 + 2x_4 = 0 \\ x_2 + \frac{1}{5}x_3 - x_4 = 0 \\ x_3 = t \\ x_4 = s \end{cases} \quad \begin{aligned} x_3 &= t \quad (\text{free variable}) \\ x_4 &= s \end{aligned}$$

$$\begin{cases} x_1 = -\frac{4}{5}t - 2s \\ x_2 = -\frac{1}{5}t + s \\ x_3 = t \\ x_4 = s \end{cases}$$

$$\Rightarrow \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -\frac{4}{5} \\ -\frac{1}{5} \\ 1 \\ 0 \end{pmatrix} t + \begin{pmatrix} -2 \\ 1 \\ 0 \\ 1 \end{pmatrix} s$$

$$\text{null}(A) = \begin{pmatrix} -\frac{4}{5}t - 2s \\ -\frac{1}{5}t + s \\ t \\ s \end{pmatrix}$$

$$\text{A basis of null}(A) = \left\{ \begin{pmatrix} -\frac{4}{5} \\ -\frac{1}{5} \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 1 \\ 0 \\ 1 \end{pmatrix} \right\}$$

### Question 3

**QUESTION 3** Let  $A = \begin{bmatrix} 4 & -3 \\ 2 & -1 \end{bmatrix}$

- (a) Find the eigenvalues and eigenvectors of  $A$
- (b) Find nonsingular matrix  $P$  and diagonal matrix  $D$  such that  $A = PDP^{-1}$

$$a) A - \lambda I = \begin{pmatrix} 4-\lambda & -3 \\ 2 & -1-\lambda \end{pmatrix}$$

$$\Rightarrow \det(A - \lambda I) = (4-\lambda)(-1-\lambda) - 2(-3) = 0 \\ = -4 - 4\lambda + \lambda^2 + 6 = 0 \\ = \lambda^2 - 3\lambda + 2$$

$$\text{Let } \det(A - \lambda I) = 0 \Rightarrow \begin{cases} \lambda_1 = 2 \\ \lambda_2 = 1 \end{cases} \quad (\text{Eigen values})$$

. Now we find eigen vectors:

$$N(A - \lambda_1 I) = \left( \begin{array}{cc|c} 2 & -3 & 0 \\ 2 & -3 & 0 \end{array} \right) R_1 \leftarrow \frac{1}{2}R_1 \\ R_2 \leftarrow R_1 - R_2$$

lum  
88 Quang Khadong

$$\Rightarrow \left( \begin{array}{cc|c} 1 & -3/2 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow \begin{cases} x_1 = \frac{3t}{2} \\ x_2 = t \end{cases} \quad (\text{free variable})$$

$$\Rightarrow \text{Eigen vector : } t \begin{pmatrix} \frac{3}{2} \\ 1 \end{pmatrix} \text{ with } \lambda_1 = 2$$

$$N(A - \lambda_2 I) = \left( \begin{array}{cc|c} 3 & -3 & 0 \\ 2 & -2 & 0 \end{array} \right) R_1 \leftarrow \frac{1}{3}R_1 \\ R_2 \leftarrow R_2 - R_1$$

$$\left( \begin{array}{cc|c} 1 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow \begin{cases} x_1 = x_2 = t \\ x_2 = t \end{cases} \quad (\text{free variable})$$

$$\Rightarrow \text{Eigen vector } t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ with } \lambda_2 = 1$$

Find  $D$  &  $L$  with  $A = L D L^{-1}$

Ans  $D = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$

$$\lambda_1 = 2 \quad \lambda_2 = 1$$
$$L = \begin{pmatrix} \frac{3}{2} & 1 \\ 1 & 1 \end{pmatrix}$$

Eigen value

Verify:

Since  $L$  has inverse  $\Rightarrow \det(L) = \frac{1}{2} \neq 0$

$$\left( \begin{array}{cc|cc} \frac{3}{2} & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{array} \right) \begin{matrix} R_1 \leftarrow \frac{2}{3}R_1 \\ R_2 \leftarrow (-\frac{2}{3})R_2 + R_1 \end{matrix}$$

$$\left( \begin{array}{cc|cc} 1 & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & -\frac{1}{2} & 1 & -\frac{3}{2} \end{array} \right) \begin{matrix} R_2 \leftarrow (-2)R_2 \end{matrix}$$

$$\left( \begin{array}{cc|cc} 1 & \frac{2}{3} & \frac{2}{3} & 0 \\ 0 & 1 & -2 & 3 \end{array} \right) \begin{matrix} -\frac{2}{3}R_2 + R_1 \end{matrix}$$

lum  
88 Quang Khê Long

$$\left( \begin{array}{cc|cc} 1 & 0 & 2 & -2 \\ 0 & 1 & -2 & 3 \end{array} \right) \Rightarrow L^{-1} = \begin{pmatrix} 2 & -2 \\ -2 & 3 \end{pmatrix}$$

$$LDL^{-1} = \begin{pmatrix} \frac{3}{2} & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -2 \\ -2 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 2 & -2 \\ -2 & 3 \end{pmatrix} = \begin{pmatrix} 4 & -3 \\ 2 & -1 \end{pmatrix} = A$$

Find  $A^{1000}$

$$A^{1000} = L^0 D^{1000} L^{-1} = \begin{pmatrix} \frac{3}{2} & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1^{1000} & 0 \\ 0 & 1^{1000} \end{pmatrix} \begin{pmatrix} 2 & -2 \\ -2 & 3 \end{pmatrix}$$

# Sample Test 2 (4)

**Question 1:**

**Question 1. (25 marks)** Solve the linear system of equations

$$\begin{cases} 2x_1 + x_2 + 3x_3 = 7 \\ x_1 - x_2 + x_3 = 0 \\ x_1 + 4x_2 - 2x_3 = 7 \end{cases}$$

by the following steps

(a) Write the system in matrix form  $Ax = b$

(b) Find the inverse of matrix  $A$

(c) Solve  $x$  by  $x = A^{-1}b$

$$A \cdot x = b$$

$$\Rightarrow x = b \cdot A^{-1}$$

$$= b \cdot \cancel{A^{-1}}$$

$$a) \quad \left( \begin{array}{ccc|c} 2 & 1 & 3 & 7 \\ 1 & -1 & 1 & 0 \\ 1 & 4 & -2 & 7 \end{array} \right) \left( \begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right) = \left( \begin{array}{c} 7 \\ 0 \\ 7 \end{array} \right)$$

$$\det \neq 0$$

$$b) \quad A = \left( \begin{array}{ccc|ccc} 2 & 1 & 3 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 & 1 & 0 \\ 1 & 4 & -2 & 0 & 0 & 1 \end{array} \right) \quad R_1 \leftarrow R_1 + R_2$$

$$R_2 \leftarrow (-2)R_2 + R_1$$

$$R_3 \leftarrow R_2 - R_3$$

$$\left( \begin{array}{ccc|ccc} 3 & 0 & 4 & 1 & 1 & 0 \\ 0 & 3 & 1 & 1 & -2 & 0 \\ 0 & -5 & 3 & 0 & 1 & -1 \end{array} \right) \quad R_3 \leftarrow R_3 \times \left( \frac{3}{5} \right) + R_2$$

$$\left( \begin{array}{ccc|ccc} 3 & 0 & 4 & 1 & 1 & 0 \\ 0 & 3 & 1 & 1 & -2 & 0 \\ 0 & 0 & \frac{14}{5} & 1 & -\frac{7}{5} & -\frac{3}{5} \end{array} \right) \quad \frac{1}{3}R_1$$

$$\frac{1}{3}R_2$$

$$\frac{5}{14}R_3$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & \frac{4}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 1 & \frac{1}{3} & \frac{1}{3} & -\frac{2}{3} & 0 \\ 0 & 0 & 1 & \frac{5}{14} & -\frac{1}{2} & -\frac{3}{14} \end{array} \right) \quad R_1 - \frac{4}{3}R_3$$

$$R_2 - \frac{1}{3}R_3$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{4} & 1 & \frac{9}{4} \\ 0 & 1 & 0 & \frac{3}{14} & -\frac{1}{2} & \frac{1}{14} \\ 0 & 0 & 1 & \frac{5}{14} & -\frac{1}{2} & -\frac{3}{14} \end{array} \right)$$

$$\Rightarrow A^{-1} = \begin{pmatrix} -\frac{1}{4} & 1 & \frac{9}{4} \\ \frac{3}{14} & -\frac{1}{2} & \frac{1}{14} \\ \frac{5}{14} & -\frac{1}{2} & -\frac{3}{14} \end{pmatrix}$$

$$c) \quad x = A^{-1}b = \begin{pmatrix} -\frac{1}{4} & 1 & \frac{9}{4} \\ \frac{3}{14} & -\frac{1}{2} & \frac{1}{14} \\ \frac{5}{14} & -\frac{1}{2} & -\frac{3}{14} \end{pmatrix} \begin{pmatrix} 7 \\ 0 \\ 7 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$

lum  
88 Quang Khudong

## Question 2

**Question 2. (25 marks)** Find the base of column space and null space of matrix

$$A = \begin{bmatrix} 1 & 2 & 4 & 1 & 2 \\ 2 & 1 & 2 & 2 & 1 \\ 1 & 4 & 2 & 1 & 1 \\ 3 & 3 & 6 & 3 & 3 \end{bmatrix} \quad \begin{array}{l} R_1 + (-2)R_2 \\ R_2 + (-2)R_1 \\ R_3 + (-1)R_1 \\ R_4 + (-3)R_1 \end{array}$$

$$\left( \begin{array}{ccccc} -3 & 0 & 0 & -3 & 0 \\ 0 & -3 & -6 & 0 & -3 \\ 0 & 2 & -2 & 0 & -1 \\ 0 & -3 & -6 & 0 & -3 \end{array} \right) \quad \begin{array}{l} -1/3 R_1 \\ -1/3 R_2 \\ 1/2 R_3 \\ R_2 - R_4 \end{array}$$

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 1 \\ 0 & 1 & -1 & 0 & -1/2 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad 2R_3 + R_2$$

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 1 \\ 0 & 3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad 1/3 R_3$$

lumen  
88 Quang Khê Long

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad R_2 \leftrightarrow R_3$$

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad R_2 - R_3$$

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad -1/2 R_3$$

$$\left( \begin{array}{ccccc} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1/2 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow \text{Basis } \{ \alpha_1, \alpha_2, \alpha_3 \} = \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ 4 \\ 3 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \\ 2 \\ 6 \end{pmatrix} \right\}$$

To find null space, we solve matrix A

$$\left( \begin{array}{ccccc|c} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1/2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

Homogeneous system  
88 Quang Kholid

$$\left\{ \begin{array}{l} x_1 + x_4 = 0 \\ x_2 = 0 \\ x_3 + 1/2x_5 = 0 \\ x_4 = t \\ x_5 = s \end{array} \right. \quad \begin{array}{l} (\text{free variable}) \end{array}$$

$$\Rightarrow \left\{ \begin{array}{l} x_1 = -t \\ x_2 = 0 \\ x_3 = 1/2s \\ x_4 = t \\ x_5 = s \end{array} \right.$$

$$\Rightarrow \text{null } A = \begin{pmatrix} -t \\ 0 \\ 1/2s \\ t \\ s \end{pmatrix}$$

$$\therefore \text{Basis of null}(A) = \left\{ \begin{pmatrix} -1 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1/2 \\ 0 \\ 1 \end{pmatrix} \right\}$$

### Question 3:

**Question 4. (25 marks)** Let  $A = \begin{bmatrix} 7 & 3 \\ 0 & 4 \end{bmatrix}$

- Find all eigenvalues and eigenvectors of  $A$ .
- Find nonsingular matrix  $P$  and diagonal matrix  $D$  such that  $A = PDP^{-1}$ .
- Use the formula  $A^{20} = PD^{20}P^{-1}$  to compute  $A^{20}$ .

lum  
88 Quang Khadong

$$a) A - \lambda I = \begin{pmatrix} 7-\lambda & 3 \\ 0 & 4-\lambda \end{pmatrix}$$

$$\det(A - \lambda I) = (7-\lambda)(4-\lambda) - 0 \times 3 \\ = 28 - 7\lambda - 4\lambda + \lambda^2$$

$$\text{Let } \det(A - \lambda I) = 0$$

$$\Leftrightarrow \lambda^2 - 11\lambda + 28 = 0$$

$$\Rightarrow \begin{cases} \lambda_1 = 7 \\ \lambda_2 = 4 \end{cases} \quad (\text{Eigen values})$$

. Now we find eigen vectors

$$N(A - \lambda_1 I) = \left( \begin{array}{cc|c} 0 & 3 & 0 \\ 0 & -3 & 0 \end{array} \right) \xrightarrow[1/3R_2]{}$$

$$\left( \begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

$$\Rightarrow \begin{cases} x_1 = t \\ x_2 = 0 \end{cases}$$

$\Rightarrow$  Eigen vector  $t \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  with  $\lambda_1 = 7$

$$N(A - \lambda_2 I) = \left( \begin{array}{cc|c} 3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right) \xrightarrow[1/3R_1]{}$$

$$\Rightarrow \left( \begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right) \Rightarrow \begin{cases} x_1 = -t \\ x_2 = t \end{cases} \quad (\text{free variable})$$

$\Rightarrow$  Eigenvector  $t \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  with  $\lambda_2 = 4$

b) . Find  $D$  &  $L$  with  $A = L D L^{-1}$

Ans  $D = \begin{pmatrix} 7 & 0 \\ 0 & 4 \end{pmatrix}$

Eigen value

$$\lambda_1 = 7 \quad \lambda_2 = 4$$
$$L = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

Eigen vector

Verify:

. Since  $L$  has inverse  $\Rightarrow \det(L) = 1 \neq 0$

$$\left( \begin{array}{cc|cc} 1 & -1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{array} \right) \xrightarrow{R_1 + R_2}$$

$$\left( \begin{array}{cc|cc} 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{array} \right)$$

lun  
88 Quang Khadong

$$\Rightarrow L^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow L D L^{-1} = A$$

. Find  $A^{20} = L D^{20} L^{-1}$

$$= \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 7^{20} & 0 \\ 0 & 4^{20} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 7^{20} & -4^{20} \\ 0 & 4^{20} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 7^{20} & 7^{20} - 4^{20} \\ 0 & 4^{20} \end{pmatrix}$$

(4)

Score on last try: 1 of 1 pts. See Details for more.

[Next question](#)[Get a similar question](#)

$$\vec{v}_1, \vec{v}_2, \vec{v}_3$$

Find an orthogonal basis for the space spanned by  $\begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ -2 \\ 0 \end{bmatrix}$ , and  $\begin{bmatrix} -4 \\ 2 \\ -6 \\ -4 \end{bmatrix}$ .

Orthogonal basis :  $u$ Orthonormal basis :  $e$ 

Using Gram-Schmidt process

Vector projection of  $\vec{v}$  onto  $\vec{u}$ :  $\text{proj}_{\vec{u}}(\vec{v})$ 

$$\frac{\vec{v} \cdot \vec{u}}{|\vec{u}|^2} \cdot \vec{u}$$

$$\left\{ \begin{array}{l} \vec{u}_1 = \vec{v}_1 = (0, 1, 0, -1) \\ \vec{e}_1 = \frac{\vec{u}_1}{|\vec{u}_1|} = \left(0, \frac{\sqrt{2}}{2}, 0, -\frac{\sqrt{2}}{2}\right) \end{array} \right.$$

$$\left\{ \begin{array}{l} \vec{u}_2 = \vec{v}_2 - \text{proj}_{\vec{u}_1}(\vec{v}_2) \\ \Rightarrow \vec{u}_2 = (-2, 0, -2, 0) \\ \vec{e}_2 = \frac{\vec{u}_2}{|\vec{u}_2|} = \left(-\frac{\sqrt{2}}{2}, 0, -\frac{\sqrt{2}}{2}, 0\right) \end{array} \right.$$

$$\left\{ \begin{array}{l} \vec{u}_3 = \vec{v}_3 - \text{proj}_{\vec{u}_1}(\vec{v}_3) - \text{proj}_{\vec{u}_2}(\vec{v}_3) = (1, -1, -1, -1) \\ \vec{e}_3 = \frac{\vec{u}_3}{|\vec{u}_3|} = \left(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\right) \end{array} \right.$$

lum  
88 Quang Khaldong

=> Orthogonal basis / vector :  $\left\{ \begin{pmatrix} u_1 \\ 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} u_2 \\ -2 \\ 0 \\ -2 \\ 0 \end{pmatrix}, \begin{pmatrix} u_3 \\ 1 \\ -1 \\ -1 \\ -1 \end{pmatrix} \right\}$

=> Orthonormal basis / vector  $\{(e_1), (e_2), (e_3)\}$

# Các dạng phu (1)

1) Determine whether  $W = \{(x, 2x) : x \text{ is a real number}\}$  with standard operations in  $\mathbb{R}^2$  is a vector space

1.  $u + v$  is in  $V$  (closure under addition)

$$\begin{aligned}(r, 2r) + (s, 2s) &= (r+s, 2r+2s) \\&= \left[ \frac{r+s}{x}, \frac{2(r+s)}{2x} \right] \quad \checkmark\end{aligned}$$

2.  $u + v = v + u$  (commutativity of addition)

$$\begin{aligned}u + v &= (r, 2r) + (s, 2s) = (r+s, 2r+2s) \\&= (s+r, 2s+2r) \\&= (s, 2s) + (r, 2r) = v + u \quad \checkmark\end{aligned}$$

3.  $u + (v+w) = (u+v) + w$  (Associativity of addition)

$$\begin{aligned}(r, 2r) + ((s, 2s) + (t, 2t)) &= (r, 2r) + (s+t, 2s+2t) \\&= (r+s+t, 2r+2s+2t) \\&= ((r+s)+t, (2r+2s)+2t) \\&= (r+s, 2r+2s) + (t, 2t) \\&= ((r, 2r)+(s, 2s)) + (t, 2t) \quad \checkmark\end{aligned}$$

lumen  
88 Quang Khadong

4.  $V$  has a vector  $0$  such that :  $u + 0 = 0 + u = u$  (additive identity)

$$\begin{aligned}(0, 0) &= (0, 2(0)) \text{ is in } W \\(0, 0) + (r, 2r) &= (r, 2r) = (r, 2r) + (0, 0) \quad \checkmark\end{aligned}$$

5. For every vector  $u$  in  $V$ , there is a vector  $-u$  in  $V$  such that :

$$u + (-u) = 0 \quad (\text{additive inverse})$$

$$\begin{aligned}(r, 2r) + (-r, -2r) &= (r-r, 2r-2r) \\&= (0, 0)\end{aligned}$$

$$\text{So } (-r, -2r) = - (r, 2r) \quad \checkmark$$

6.  $cw$  in in  $V$  (closure under scalar multiplication)

$$\begin{aligned}c(r, 2r) &= (cr, c2r) \\&= \left[ \frac{cr}{x}, \frac{2(c r)}{2x} \right] \Rightarrow \text{Axiom hold}\end{aligned}$$

7.  $c(u+v) = cu + cv$  (distributive properties)

$$\begin{aligned} c[(r, 2r) + (s, 2s)] &= c(r+s, 2r+2s) \\ &= (c(r+s), c(2r+2s)) \\ &= (cr+cs, 2cr+2cs) \\ &= (cr, 2cr) + (cs, 2cs) \\ &= c(r, 2r) + c(s, 2s) \quad \checkmark \\ &\Rightarrow \text{Seventh axiom hold} \end{aligned}$$

8.  $(c+d)u = cu + du$  (distributive properties)

$$\begin{aligned} (c+d)(r, 2r) &= ((c+d)r, (c+d)2r) \\ &= (cr+dr, 2cr+2dr) \\ &= (cr, 2cr) + (dr, 2dr) \\ &= c(r, 2r) + d(r, 2r) \\ &\Rightarrow \text{Eighth axiom hold} \quad \checkmark \end{aligned}$$

9.  $c(dv) = (cd)v$  (associative property)

$$\begin{aligned} c(d(r, 2r)) &= c(dr, 2dr) \\ &= (cdr, 2cdr) \\ &= cd(r, 2r) \quad \checkmark \end{aligned}$$

10)  $1(u) = u$  (multiplication - identity)

$$\begin{aligned} 1(r, 2r) &= (1r, 12r) \\ &= (r, 2r) \quad \checkmark \end{aligned}$$

$\Rightarrow W$  is a vector space

lum  
88 Quang Khê Long

'a' to make the matrix orthogonal

(2)

$$A = \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \\ \frac{\sqrt{2}}{2} & a & 0 \end{bmatrix}$$

To let it be,  $A \times A^T = I$

$$\begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \\ \frac{\sqrt{2}}{2} & a & 0 \end{bmatrix} \times \begin{bmatrix} \frac{\sqrt{2}}{2} & 0 & \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} & 0 & a \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Consider (1,3)

$$\left(\frac{\sqrt{2}}{2}\right) \left(\frac{\sqrt{2}}{2}\right) - \left(\frac{\sqrt{2}}{2}a\right) + 0 = 0$$

$$\Rightarrow a = \frac{\sqrt{2}}{2}$$

Lim  
88 Quang Khadong

Determine whether the set  $\mathbb{R}^2$ , along with standard operations

$$\cdot (x_1, y_1) + (x_2, y_2) = (x_1 x_2, y_1 y_2)$$

$$\cdot c(x_1, y_1) = (cx_1, cy_1)$$

(1)

1.  $u + v$  is in  $V$

$$(x_1, y_1) + (x_2, y_2) = (x_1 x_2, y_1 y_2)$$

is in  $\mathbb{R}^2 \checkmark$

2.  $u + v = v + u$

$$\begin{aligned}(x_1, y_1) + (x_2, y_2) &= (x_1 x_2, y_1 y_2) \\&= (x_2 x_1, y_2 y_1) \\&= (x_2, y_2) + (x_1, y_1) \checkmark\end{aligned}$$

lum  
88 Aneng Khadung

3.  $u + (v + w) = (u + v) + w$

$$\begin{aligned}(x_1, y_1) + ((x_2, y_2) + (x_3, y_3)) &= (x_1, y_1) + (x_2 x_3, y_1 y_2 y_3) \\&= (x_1 x_2 x_3, y_1 y_2 y_3) \\&= (x_1 x_2, y_1 y_2) + (x_3, y_3) \\&= ((x_1, y_1) + (x_2, y_2)) + (x_3, y_3)\end{aligned}$$

$\Rightarrow$  Third axiom holds

4.  $V$  has a vector  $0$  such that

$$u + 0 = 0 + u = u \quad \vec{0} = (1, 1)$$

$$(x_1, y_1) + (1, 1) = (x_1 \cdot 1, y_1 \cdot 1) = (x_1, y_1)$$

$$(1, 1) + (x_1, y_1) = (1x_1, 1y_1) = (x_1, y_1) \checkmark$$

5. For every vector  $u$  in  $V$ , there is a vector  $-u$  in  $V$  such that :

$$u + (-u) = 0 \quad (\text{additive inverse})$$

$$x_1 \neq 0, y_1 \neq 0$$

$$(x_1, y_1) + \left( \frac{1}{x_1}, \frac{1}{y_1} \right) = \left( x_1 \cdot \frac{1}{x_1}, y_1 \cdot \frac{1}{y_1} \right) = (1, 1) = \vec{0}$$

$$-(x_1, y_1) = \left( \frac{1}{x_1}, \frac{1}{y_1} \right)$$

But  $(0, 0)$  is in  $\mathbb{R}^2$  and has no additive inverse  $\Rightarrow$  Fifth axiom failed

$\Rightarrow$  The set is not vector space

Leave  $V = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in \mathbb{R}^3 \mid x_3 = x_1 + x_2 \right\}$  is a subspace of  $\mathbb{R}^3$

Need to show

- (1)  $V \neq \emptyset$
- (2)  $\vec{x}, \vec{y} \in V \Rightarrow \vec{x} + \vec{y} \in V$
- (3)  $\alpha \in \mathbb{R}, \vec{x} \in V \Rightarrow \alpha \vec{x} \in V$

Proof: (1)  $0 = 0 + 0, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \in V$ , so  $V \neq \emptyset$

(2) Suppose  $\vec{x}, \vec{y} \in V$

$$\vec{x} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \vec{y} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$x_1, x_2, x_3, y_1, y_2, y_3 \in \mathbb{R} \text{ and } \begin{cases} x_3 = x_1 + x_2 \\ y_3 = y_1 + y_2 \end{cases}$$

lum  
88 Quang Khê

$$\vec{x} + \vec{y} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix} \text{ and } x_3 + y_3 = (x_1 + x_2) + (y_1 + y_2) = (x_1 + y_1) + (x_2 + y_2)$$

so  $\vec{x} + \vec{y} \in V$

(3) Take any  $\vec{x} \in V$  and  $\alpha \in \mathbb{R}$

$$\vec{x} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$x_1, x_2, x_3 \in \mathbb{R} \text{ and } x_3 = x_1 + x_2$$

$$\alpha \vec{x} = \begin{pmatrix} \alpha x_1 \\ \alpha x_2 \\ \alpha x_3 \end{pmatrix} \text{ and } \alpha x_3 = \alpha (x_1 + x_2) = \alpha x_1 + \alpha x_2, \text{ so } \alpha \vec{x} \in V$$

$\therefore V$  is a subspace

Determine whether  $W = \{(a, b, c) \mid a = b^2\}$  is a subspace of  $\mathbb{R}^3$

Need to show

(1)  $0 \in W$

(2)  $\vec{x}, \vec{y} \in W \Rightarrow \vec{x} + \vec{y} \in W$

(3)  $\alpha \in \mathbb{R}, \vec{x} \in W \Rightarrow \alpha \vec{x} \in W$

lum  
88 Quang Khê

(1)  $(0, 0, 0) \in W$  because  $0 = 0^2$  ( $a = b^2$ )

(2)  $(4, 2, 1) \in W \quad 4 = 2^2$   
 $a \quad b \quad c$

$(9, 3, 1) \in W \quad 9 = 3^2$   
 $a \quad b \quad c$

Then  $(4, 2, 1) + (9, 3, 1) = (13, 5, 2)$

Since  $13 \neq 5^2 \Rightarrow$  Not closed under vector addition

$\therefore$  Not a subspace

Determine whether  $W = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$  is a subspace of  $\mathbb{R}^3$

(1)  $0^2 + 0^2 + 0^2 \leq 1$

$(0, 0, 0) \in W, W \neq \emptyset$

(2)  $(1, 0, 0) \in W$  b/c  $1^2 + 0^2 + 0^2 \leq 1$

$(0, 1, 0) \in W$  b/c  $0^2 + 1^2 + 0^2 \leq 1$

$(1, 0, 0) + (0, 1, 0) = (1, 1, 0)$

But  $1^2 + 1^2 + 0^2 = 2 > 1$ , so  $(1, 1, 0) \notin W \Rightarrow$  Not closed under vector addition  
 $\Rightarrow$  Not subspace

Leave  $W = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2 \mid a = -b \right\}$  is a subspace of  $\mathbb{R}^2$

(1)

Need to show

- (1)  $V \neq \emptyset$
- (2)  $\vec{x}, \vec{y} \in V \Rightarrow \vec{x} + \vec{y} \in V$
- (3)  $\alpha \in \mathbb{R}, \vec{x} \in V \Rightarrow \alpha \vec{x} \in V$

lumen  
88 Quang Khê Long

(1)  $0 = -0$ , so  $\begin{pmatrix} 0 \\ 0 \end{pmatrix} \in W$ , so  $W \neq \emptyset$

(2) Spse,  $x, y \in W$

$$x = \begin{pmatrix} a \\ b \end{pmatrix}, y = \begin{pmatrix} c \\ d \end{pmatrix} \text{ with } a = -b \text{ and } c = -d$$

$$x + y = \begin{pmatrix} a+c \\ b+d \end{pmatrix} = \begin{pmatrix} -b+d \\ b+d \end{pmatrix} = \begin{pmatrix} -(b+d) \\ b+d \end{pmatrix}$$

So  $x + y \in W$

(3) Spse  $\alpha \in \mathbb{R}, \alpha x \in W$

$$\alpha x = \alpha \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} \alpha a \\ \alpha b \end{pmatrix} \text{ with } \alpha a = -\alpha b$$

So  $\alpha x \in W$

$\therefore W$  is a subspace of  $\mathbb{R}^2$

Determine  $W = \{(a, b, c) \mid a \leq b \leq c\}$  is a subspace of  $\mathbb{R}^3$

(1)  $(0, 0, 0) \in W$   $b/c$   $0 \leq 0 \leq 0$

(2)  $(1, 2, 3) \in W$   $b/c$   $1 \leq 2 \leq 3$

and  $-1 \in \mathbb{R}$

Then  $-1(1, 2, 3) = (-1, -2, -3)$

but  $-1 \leq -2 \leq -3$  is false not closed under vector scalar multiplication

$\Rightarrow$  Not subspace

# Linear Combination

Write  $w$  as a linear combination of  $u$  &  $v$

(2)

$$u = \begin{bmatrix} 1 \\ 1 \\ 2 \\ -1 \end{bmatrix}, \quad v = \begin{bmatrix} 3 \\ 0 \\ 2 \\ 1 \end{bmatrix}, \quad w = \begin{bmatrix} -11 \\ 4 \\ -2 \\ -9 \end{bmatrix}$$

$$\left[ \begin{array}{cc|c} u & v & w \\ 1 & 3 & -11 \\ 1 & 0 & 4 \\ 2 & 2 & -2 \\ -1 & 1 & -9 \end{array} \right] \xrightarrow{\substack{R_3 - 2R_2 \\ R_4 + R_2}} \left[ \begin{array}{cc|c} 1 & 3 & -11 \\ 1 & 0 & 4 \\ 0 & 2 & -10 \\ 0 & 1 & -5 \end{array} \right] \xrightarrow{\substack{R_1 - R_2 \\ R_3/2}} \left[ \begin{array}{cc|c} 0 & 3 & -15 \\ 1 & 0 & 4 \\ 0 & 1 & -5 \\ 0 & 1 & -5 \end{array} \right]$$

$$\begin{array}{l} R_1 \leftrightarrow R_2 \\ R_1/3 \rightarrow \end{array} \left[ \begin{array}{cc|c} 1 & 0 & 4 \\ 0 & 1 & -5 \\ 0 & 1 & -5 \\ 0 & 1 & -5 \end{array} \right] \xrightarrow{\substack{R_3 - R_2 \\ R_4 - R_2}} \left[ \begin{array}{cc|c} 1 & 0 & 4 \\ 0 & 1 & -5 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

$$\Rightarrow x = 4$$

$$y = -5$$

$$\Rightarrow 4u + (-5)v = w$$

$$\Rightarrow 4 \begin{bmatrix} 1 \\ 1 \\ 2 \\ -1 \end{bmatrix} + (-5) \begin{bmatrix} 3 \\ 0 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -11 \\ 4 \\ -2 \\ -9 \end{bmatrix}$$

If possible, write  $\vec{v} = (-1, 7, 2)$  as a linear combination of  $\vec{u}_1 = (1, 3, 5)$

$$c_1 \cdot \vec{u}_1 + c_2 \cdot \vec{u}_2 + c_3 \cdot \vec{u}_3 = \vec{v}$$

$$c_1(1, 3, 5) + c_2(2, -1, 3) + c_3(-3, 2, -4) = (-1, 7, 2)$$

$$c_1 + 2c_2 - 3c_3 = -1$$

$$3c_1 - c_2 + 2c_3 = 7$$

$$5c_1 + 3c_2 - 4c_3 = 2$$

lun  
88 Quang Khaling

$$\left[ \begin{array}{ccc|c} \vec{u}_1 & \vec{u}_2 & \vec{u}_3 & \vec{v} \\ 1 & 2 & -3 & -1 \\ 3 & -1 & 2 & 7 \\ 5 & 3 & -4 & 2 \end{array} \right] \xrightarrow{\text{refl}} \left[ \begin{array}{ccc|c} 1 & 0 & 1/7 & 0 \\ 0 & 1 & -4/7 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] \quad 0=1$$

$c_1 \quad c_2 \quad c_3$

$\therefore \vec{v}$  can not be expressed as a linear comb

## Spanning Vector

Let  $v_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $v_2 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$ , does  $\{v_1, v_2\}$  span  $\mathbb{R}^2$

(Let  $\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2$  Can we write  $x_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$  ?)

$$\Rightarrow \left[ \begin{array}{cc} 1 & 1 \\ 1 & -2 \end{array} \right] \left[ \begin{array}{c} x_1 \\ x_2 \end{array} \right] = \left[ \begin{array}{c} a \\ b \end{array} \right] \xrightarrow{\text{Aug}} \left[ \begin{array}{cc|c} 1 & 1 & a \\ 1 & -2 & b \end{array} \right] \quad R_2 \leftarrow R_2 - R_1$$

(2)

$$\left[ \begin{array}{cc|c} 1 & 1 & a \\ 0 & -3 & b-a \end{array} \right] \quad R_2 \leftarrow \frac{1}{3} R_2$$

$$\left[ \begin{array}{cc} 1 & 1 \\ 1 & -2 \end{array} \right]$$

$$\det = \begin{vmatrix} 1 & 1 \\ 1 & -2 \end{vmatrix} = 1 \cdot (-2) - 1 \cdot 1 = -3 \neq 0$$

$\Rightarrow$  span

$$\left[ \begin{array}{cc|c} 1 & 1 & a \\ 0 & 1 & \frac{a-b}{3} \end{array} \right] \quad R_1 \leftarrow R_1 - R_2$$

$$\left[ \begin{array}{cc|c} 1 & 0 & a - \frac{a-b}{3} \\ 0 & 1 & \frac{a-b}{3} \end{array} \right]$$

$$\Rightarrow \boxed{\begin{aligned} x_1 &= a - \frac{a-b}{3} \\ x_2 &= \frac{a-b}{3} \end{aligned}}$$

Yes, these span  $\mathbb{R}^2$

Let  $v_1 = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} -2 \\ 4 \end{pmatrix}$

Does  $v_1, v_2$  span  $\mathbb{R}^2$ ?

$$\left[ \begin{array}{cc|c} 1 & -2 & a \\ -2 & 4 & b \end{array} \right] \quad \ell_2 \leftarrow 2\ell_1 + \ell_2$$

(2)

$$\left[ \begin{array}{cc|c} 1 & -2 & a \\ 0 & 0 & 2a+b \end{array} \right]$$

$$\Rightarrow 0 = 2a + b$$

Not true  $\nexists a, b$

$\Rightarrow$  No solution

$\Rightarrow$  Vectors do not span  $\mathbb{R}^2$

Can  $(-1, 0, 0)$   $(1, 0, 0)$   $(0, 0, 3)$  span  $\mathbb{R}^3$

$$x_1 \begin{pmatrix} -1 \\ 0 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$$

$$\left[ \begin{array}{ccc|c} -1 & 1 & 0 & a \\ 0 & 0 & 0 & b \\ 0 & 0 & 3 & c \end{array} \right]$$

$$0 = b \text{ Not true } \nexists b$$

$\Rightarrow$  No sol

$\Rightarrow$  Vectors do not span  $\mathbb{R}^3$

lum  
88 Quang Khê

Mo:  $\det = 0 \Rightarrow$  ko Span

$\det \neq 0 \Rightarrow$  Span

# Vector independence

$$\begin{pmatrix} v_1 \\ 1 \\ -1 \\ 1 \end{pmatrix} \begin{pmatrix} v_2 \\ 1 \\ 1 \\ 0 \end{pmatrix} \begin{pmatrix} v_3 \\ -1 \\ 1 \\ 2 \end{pmatrix}$$

$$c_1v_1 + c_2v_2 + \dots + c_nv_n = 0$$

$$\left[ \begin{array}{cccc|c} 1 & 1 & -1 & 0 \\ -1 & 1 & 1 & a \\ 1 & 0 & 2 & 0 \end{array} \right] \xrightarrow{\substack{R_2 + R_1 \\ R_3 - R_1}} \left[ \begin{array}{cccc|c} 1 & 1 & -1 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 1 & 3 & 0 \end{array} \right]$$

(2)

$$\xrightarrow{R_2/2} \left[ \begin{array}{cccc|c} 1 & 1 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 3 & 0 \end{array} \right] \xrightarrow{\substack{R_1 - R_2 \\ R_3 + R_2}}$$

$$\longrightarrow \left[ \begin{array}{cccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 3 & 0 \end{array} \right]$$

lum  
88 Quang Khaldong

$$\xrightarrow{R_3/3} \left[ \begin{array}{cccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

$$\left\{ \begin{array}{l} x_1 = 1 \\ x_2 = 2 \\ x_3 = 3 \end{array} \right.$$

$$\left\{ \begin{array}{l} x_1 = 1 \\ x_2 = 5 \\ x_3 = 35 \end{array} \right. \quad \begin{matrix} 1 & 2 \\ 3 & 6 \end{matrix}$$

$$\xrightarrow{R_1 + R_1 + R_3} \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

$\Rightarrow$  Linearly Independent

Bié thang  $\rightarrow \det \neq 0 \rightarrow L$ . In  
 $\rightarrow \det = 0 \rightarrow L$ . De

$$\left| \begin{array}{ccc|c} 1 & 2 & 3 & 7 \\ 0 & 4 & 5 & 8 \\ 0 & 0 & 6 & 9 \end{array} \right| \Rightarrow \left\{ \begin{array}{l} x_1 + 2x_2 + 3x_3 = 7 \\ 4x_2 + 5x_3 = 8 \\ 6x_3 = 9 \end{array} \right.$$

$\det \neq 0$

$$\left| \begin{array}{ccc|c} 1 & 2 & 3 & 7 \\ 0 & 4 & 5 & 8 \\ 0 & 0 & 0 & 0 \end{array} \right|$$

$\det = 0$

7. For which values of  $t$  is each set linearly independent?

$$S = \{(t, 1, 1), (1, t, 1), (1, 1, t)\}$$

$$\begin{bmatrix} t & 1 & 1 \\ 1 & t & 1 \\ 1 & 1 & t \end{bmatrix} \neq 0$$

$$\Rightarrow \det \neq 0$$

$$\Leftrightarrow t(t^2 - 1) - (t - 1) + (1 - t) \neq 0$$

$$\Leftrightarrow t^3 - t - t + 1 - t + 1 \neq 0$$

$$\Leftrightarrow t^3 - 3t + 2 \neq 0$$

$$\Rightarrow \begin{cases} t \neq -2 \\ t \neq 1 \end{cases}$$

lum  
88 Quang Khulóng

(2)

$\{1, 2, 3, \dots\}$

$$0^{>l} = 0$$

Vô số nghiệm 0 0 0 | 0

Vô nghiệm 0 0 0 | ≠0

$$0 \propto = 3$$

$$\Rightarrow \propto = \frac{3}{0}$$

FINAL EXAMINATION

Semester 2, 2022-2023 • Date: June 07, 2023 • Duration: 75 minutes

SUBJECT: Applied Linear Algebra

Department of Mathematics  
Vice Chair:

Dr. Nguyen Minh Quan

Lecturer

Dr. Le Minh Tuan

**INSTRUCTIONS:** Each student is allowed a scientific calculator and a maximum of TWO double-sided sheets of reference material (size A4 or similar) marked with their name and ID. *All other documents and electronic devices are forbidden.*

Solve each of the following problems. Show your work clearly. You must write out all relevant steps. Simply having the correct answer does not give you credit.

Problem 1 (20 points).

Evaluate the following determinant: 
$$\begin{vmatrix} 1 & -3 & 2 & 0 \\ -3 & -1 & 0 & -2 \\ 2 & 1 & 3 & 1 \\ 2 & 0 & -3 & 0 \end{vmatrix}$$

Mid: Thay đổi Row  $\times (-1)$

det

Problem 2 (20 points).

Find all real numbers  $\lambda$  such that the homogeneous system:

$$\left( \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 2 & 3 \\ 2 & -3 \end{bmatrix} \right) \mathbf{x} = \mathbf{0} \quad \left\{ \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{bmatrix} 2 & 3 \\ 2 & -3 \end{bmatrix} \right\} \mathbf{x} = \mathbf{0}$$

has a nontrivial solution.

vô số nghiệm

det = 0

$$\det \left\{ \begin{bmatrix} \lambda-2 & -3 \\ -2 & \lambda+3 \end{bmatrix} \right\} = 0$$

$$\Rightarrow (\lambda-2)(\lambda+3) - (-2)(-3) = 0$$

a) Show that  $S$  is an ordered basis of  $\mathbb{R}^3$ .

span  $\mathbb{R}^3$

b) Find  $[v]_S$ , the coordinates of  $v$  with respect to  $S$ .

$$T(u+v) = T(u) + T(v) \text{ and } T(cu) = cT(u)$$

Problem 4 (30 points).

Let  $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be a mapping defined by  $T \begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}$ , for all  $\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} \in \mathbb{R}^4$ .

a) Show that  $T$  is linear transformation.

b) Does  $T$  map  $\mathbb{R}^4$  onto  $\mathbb{R}^3$ ?

THE END!

RREF

| Nếu có 1 hàng vô số nghiệm, toàn là 0 ( $\det=0$ )

→  $T$  doesn't map  $\mathbb{R}^4$  onto  $\mathbb{R}^3$

Problem 4:

$$(i) T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$$

$$\therefore \vec{u} = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} \quad \vec{v} = \begin{pmatrix} 2 \\ 3 \\ 4 \\ 5 \end{pmatrix} \quad c = 2$$

$$\therefore T(\vec{u} + \vec{v}) = T\left(\begin{pmatrix} 3 & w \\ 5 & x \\ 7 & y \\ 9 & z \end{pmatrix}\right) = \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}, \quad = \begin{pmatrix} 28 \\ -26 \\ -18 \end{pmatrix}$$

$$\therefore T\left(\begin{pmatrix} 1 & u \\ 2 & x \\ 3 & y \\ 4 & z \end{pmatrix}\right) + T\left(\begin{pmatrix} 2 \\ 3 \\ 4 \\ 5 \end{pmatrix}\right) = \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}, \quad + \quad \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}, \quad + \quad \begin{pmatrix} 12 \\ -12 \\ -9 \end{pmatrix} \quad + \quad \begin{pmatrix} 16 \\ -14 \\ -9 \end{pmatrix} = \begin{pmatrix} 28 \\ -26 \\ -18 \end{pmatrix}$$

$$(ii) T(c\vec{u}) = c \cdot T\vec{u}$$

$$\therefore T\left(\begin{pmatrix} 2 \\ 4 \\ 6 \\ 8 \end{pmatrix}\right) = \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}, \quad = \begin{pmatrix} 24 \\ -24 \\ -18 \end{pmatrix}$$

$$2 \cdot T\left(\begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}\right) = 2 \cdot \begin{bmatrix} 2w - 3x + 4y + z \\ w - x + 3y - 5z \\ 3w + x - 2y - 2z \end{bmatrix}, \quad = 2 \begin{pmatrix} 12 \\ -12 \\ -9 \end{pmatrix} \\ = \begin{pmatrix} 24 \\ -24 \\ -18 \end{pmatrix}$$

$\Rightarrow$  Linear Trans

Đặc biệt cảm ơn :

Nguyễn Thành Luân  
Mai Khanh Thuận

Mến chúc tất cả các bạn thi tốt

lun  
Queng Khuong

**Aladin**

**iem-  
buddy**  
College Tutoring

Queng Khuong via Aladin / Tô review:

- Deterministic model in OR
- Linear Algebra
- Engineering Economy
- Cal / Thay các biến
- Probability - Statistics
- Mechanics - Dynamics