



Contents lists available at ScienceDirect

Ticks and Tick-borne Diseases

journal homepage: www.elsevier.com/locate/ttbd

Original article

Regional prevalences of *Borrelia burgdorferi*, *Borrelia bissetiae*, and *Bartonella henselae* in *Ixodes affinis*, *Ixodes pacificus* and *Ixodes scapularis* in the USARicardo G. Maggi^{a,*}, Marcée Toliver^b, Toni Richardson^a, Thomas Mather^c, Edward B. Breitschwerdt^a^a Intracellular Pathogens Research Laboratory, Comparative Medicine Institute, College of Veterinary Medicine, North Carolina State University (NCSU), 1060 William Moore Drive, Raleigh, NC 27607, USA^b Public Health Pest Management Section, NC Department of Environment and Natural Resources, 10005 Waterford Court, Raleigh, NC 27613, USA^c Department of Plant Sciences and Entomology, Center for Vector-Borne Disease, College of the Environment and Life Sciences, University of Rhode Island, 231 Woodward Hall, Kingston, RI 02881, USA

ARTICLE INFO

Keywords:

Borrelia
B. bissetiae
Bartonella
Ixodes
I. affinis
I. scapularis

ABSTRACT

The objective of this work was to determine the prevalence of *Borrelia* and *Bartonella* species in *Ixodes* spp. ticks collected from 16 USA states. Genus PCR amplification and sequence analysis of *Bartonella* and *Borrelia* 16SsRNA-23SsRNA intergenic regions were performed on DNA extracted from 929 questing adult ticks (671 *Ixodes scapularis*, 155 *Ixodes affinis*, and 103 *Ixodes pacificus*). Overall, 129/929 (13.9%) *Ixodes* ticks were PCR positive for *Borrelia burgdorferi* sensu stricto, 48/929 for *B. bissetiae* whereas 23/929 (2.5%) were PCR positive for a *Bartonella henselae*. *Borrelia bissetiae* or *B. burgdorferi* s.s. and *B. henselae* co-infections were found in *I. affinis* from North Carolina at a rate of 4.5%; in a single *I. scapularis* from Minnesota, but not in *I. pacificus*. For both bacterial genera, PCR positive rates were highly variable depending on geographic location and tick species, with *Ixodes affinis* (n = 155) collected from North Carolina, being the tick species with the highest prevalence's for both *Borrelia* spp. (63.2%) and *B. henselae* (10.3%). Based on the results of this and other published studies, improved understanding of the enzootic cycle, transmission dynamics, and vector competence of *Ixodes* species (especially *I. affinis*) for transmission of *Borrelia* spp. and *B. henselae* should be a public health research priority.

1. Introduction

Blacklegged ticks (Acari: Ixodidae) are obligate vertebrate blood feeders and known vectors for a variety of zoonotic viruses, bacteria, and protozoan pathogens worldwide. In the United States, some species, including *I. affinis*, *I. pacificus* and *I. scapularis* are generalist host feeders and are established or potential vectors for 16 zoonotic pathogens (Nelder et al., 2016), including *Borrelia burgdorferi* (causative agent of Lyme disease) (Clark, 2004; Clark et al., 2002; Harrison et al., 2010; Holden et al., 2006; Johnson et al., 2017; Maggi et al., 2010; Nadolny et al., 2011; Nelder et al., 2016; Oliver et al., 2003; Rudenko et al., 2013; Tokarz et al., 2017; Wroblewski et al., 2017), *Bartonella henselae* (bartonellosis/cat scratch disease) (Holden et al., 2006), *B. quintana* (bartonellosis/trench fever) (Chang et al., 2001, 2002), *Anaplasma phagocytophilum* (anaplasmosis) (Adelson et al., 2004; Belongia, 2002; Holden et al., 2006; Johnson et al., 2017; Murphy et al., 2017;

Swanson and Norris, 2007; Tokarz et al., 2017; Wroblewski et al., 2017), *Babesia microti* (babesiosis) (Adelson et al., 2004; Belongia, 2002; Swanson and Norris, 2007), and *B. miyamotoi* (relapsing fever borreliosis/ *B. miyamotoi* disease) (Barbour, 2014; Breuner et al., 2017; Han et al., 2016; Johnson et al., 2017; King et al., 2017; Nelder et al., 2016; Tokarz et al., 2017; Wroblewski et al., 2017), and Ehrlichia muris-like (EMLA) (Murphy et al., 2017).

Lyme disease, caused by *B. burgdorferi* sensu stricto, is estimated to affect 300,000 people a year in the United States (Lyme-CDC, 2018). Lyme disease accounts for almost 75% of VBD cases reported from the highly endemic region of the northeastern United States (Paddock et al., 2016). As vectors for disease transmission, ticks are associated with approximately 95% of human vector-borne disease (VBD) cases reported to the Centers for Disease Control and Prevention (CDC) (Paddock et al., 2016). In recent years, bartonellosis has been recognized as an emerging/re-emerging zoonotic infectious disease

* Corresponding author.

E-mail addresses: rgmaggi@ncsu.edu (R.G. Maggi), marceetoliver@gmail.com (M. Toliver), tgrichar@ncsu.edu (T. Richardson), tmather@uri.edu (T. Mather), ebbreits@ncsu.edu (E.B. Breitschwerdt).

caused by numerous mammalian reservoir-adapted *Bartonella* species. At least 18 *Bartonella* spp. have been implicated as a cause of disease pathology in animals or humans (Breitschwerdt, 2017). Several arthropod vectors, including sand flies, lice, fleas, biting flies and ticks are known to transmit various *Bartonella* spp. to animals and humans in the USA and abroad (Breitschwerdt, 2014, 2017; Breitschwerdt et al., 2010; Chomel and Kasten, 2010; Guptill, 2010; Mada and Joel Chandranesan, 2017; Mayne, 2011). *Bartonella* spp. infections have been associated with a wide range of diseases such as Carrión's disease, trench fever, cat-scratch disease, bacillary angiomatosis, peliosis hepatis, endocarditis, myocarditis, chronic lymphadenopathy, and neurological and rheumatologic disorders. It is estimated that *B. henselae* infections affect between 13,000 to 45,000 people yearly in the USA alone (Nelson et al., 2016).

In 1926, Noguchi reported *Dermacentor andersoni* transmission of *B. bacilliformis* (Noguchi et al., 1926). Subsequently, historical evidence supporting ticks as competent vectors for transmission of *Bartonella* organisms was mostly circumstantial and primarily based on clinical case reports, epidemiological associations, or the detection of *Bartonella* spp. DNA in ticks of the genus *Ixodes* (Adelson et al., 2004; Chang et al., 2002; Dietrich et al., 2010; Halos et al., 2005; Nelder et al., 2016; Vilcins et al., 2009), *Amblyomma* (Billeter et al., 2008), *Dermacentor* (Billeter et al., 2011; Chang et al., 2002; Ebani et al., 2015), *Rhipicephalus* (Billeter et al., 2011; Erekat et al., 2016), and *Hyalomma* (Erekat et al., 2016) among others. The role of ticks as competent vectors for *Bartonella* transmission remains controversial (Harrison et al., 2012; Telford and Wormser, 2010; Tjisse-Klasen et al., 2011). Recent laboratory studies have provided evidence supporting *Ixodes ricinus* transmission of *B. henselae* to cats using an artificial tick feeding model (Cotte et al., 2008; Reis et al., 2011). *In vivo* laboratory transmission of *B. birtlesii* by *I. ricinus* nymphs was confirmed using a rodent model (Cotte et al., 2008; Reis et al., 2011). We are not aware of laboratory confirmation of *B. henselae* by an *Ixodes* sp.

Since a relatively high prevalence of *Bartonella* or *Borrelia* spp. have been reported in *Ixodes* spp. (Adelson et al., 2004; Chang et al., 2001; Dietrich et al., 2010) from the USA and Europe, the present study surveyed the presence of both bacterial genera in three *Ixodes* species: *I. affinis*, *I. pacificus*, and *I. scapularis*.

2. Methods

2.1. Tick collection, processing, and DNA extraction

In total, 816 individual adult questing ticks were tested, including 671 *I. scapularis*, 155 *I. affinis* (from North Carolina), and 103 *I. pacificus* from southern California (Table 1). Questing ticks were collected by dragging. No tick was visibly engorged. Each tick was individually cleaned using three washes of 75% ethanol and processed for DNA extraction using Qiagen DNeasy Blood & Tissue Kits® (Qiagen, Germantown, MD USA 20874). DNA quality and quantity was assessed using NanoDrop® (Thermo Fisher Scientific, Waltham, MA USA 02451) measurement at 260/280 ratio. House keeping DNA testing, targeting mitochondrial 16S rRNA DNA using primers Tick16S1s: 5' CTG CTC AAT GAT TTT TTA AAT TGC TGT GGT 3' and Tick16S460as: 5' CCG GTC TGA ACT CAG ATC AAG TAG GA 3' (developed and validated in our laboratory), was used to assess DNA quality for DNA amplification.

2.2. Molecular testing

Polymerase chain reaction, targeting DNA of *Bartonella* spp. and *Borrelia* spp., was performed as previously reported (Maggi et al., 2011, 2008, 2010; Mascarelli et al., 2016) with minor modifications. Briefly, screening of *Bartonella* ITS region was performed using oligonucleotides 325s: 5' CTT CAG ATG ATG ATC CCA AGC CTT CTG GCG 3' and 1100as: 5' GAA CCG ACG ACC CCC TGC TTG CAA AGC A 3' as forward and reverse primers, respectively. Amplification was performed in a 25-

Table 1
Borrelia burgdorferi s.s. and *Bartonella henselae* PCR results derived from *Ixodes* spp. ticks collected from different regions of the US.

| Location | Number of ticks | <i>B. burgdorferi</i> s.s. positive ticks (%) | <i>B. henselae</i> positive ticks (%) | <i>B. burgdorferi/B. henselae</i> Co-infected ticks (%) |
|----------------|------------------|---|---------------------------------------|---|
| California | 1 ^a | 0 (0) | 0 (0) | 0 (0) |
| California | 103 ^b | 0 (0) | 1 (1.0) | 0 (0) |
| Connecticut | 13 ¹ | 8 (62) | 0 (0) | 0 (0) |
| Florida | 2a ¹ | 0 (0) | 0 (0) | 0 (0) |
| Louisiana | 1 ^a | 0 (0) | 0 (0) | 0 (0) |
| Maine | 55 ^a | 0 (0) | 0 (0) | 0 (0) |
| Maryland | 17 ^a | 7 (41) | 0 (0) | 0 (0) |
| Massachusetts | 3 ^a | 2 (67) | 0 (0) | 0 (0) |
| Minnesota | 62 ^a | 21 (34) | 1 (2) | 1 (2) |
| New Jersey | 9 ^a | 4 (44) | 0 (0) | 0 (0) |
| New York | 8 ^a | 6 (75) | 0 (0) | 0 (0) |
| North Carolina | 348 ^a | 1 (0.3) | 5 (1.4) | 0 (0) |
| North Carolina | 155 ^c | 55 (35.5) | 16 (10.3) | 3 (1.9) ^d |
| Pennsylvania | 21 ^a | 15 (71) | 0 (0) | 0 (0) |
| Rhode Island | 120 ^a | 8 (6.7) | 0 (0) | 0 (0) |
| Virginia | 8 ^a | 0 (0) | 0 (0) | 0 (0) |
| Washington DC | 2 ^a | 1 (50) | 0 (0) | 0 (0) |
| Wisconsin | 1 ^a | 1 (100) | 0 (0) | 0 (0) |

^a *Ixodes scapularis*.

^b *Ixodes pacificus*.

^c *Ixodes affinis*.

^d Four additional *I. affinis* ticks were co-infected with *B. bissetiae* and *B. henselae*.

μl final volume reaction containing 12.5 μl of MyTaq™ Red masterMix Mix (Bioline) 0.2 μl of 100 μM of each forward and reverse primer (IDT® DNA Technology, Coralville, IA USA 52241), 7.5 μl of molecular-grade water, and 5 μl of DNA from each sample tested. PCR negative controls were prepared using 5 μl of molecular grade water. Positive controls for PCR were prepared by using 5 μl of *B. henselae* genomic DNA at a concentration of 0.001 pg/μl (0.5 genome copies per microliter). Conventional PCR was performed in an Eppendorf Mastercycler EP-gradient® (Eppendorf Inc., Westbury, NY USA 11590) under the following conditions: a single hot-start cycle at 95 °C for 2 min followed by 55 cycles of denaturing at 94 °C for 15 s, annealing at 68 °C for 15 s, and extension at 72 °C for 18 s. Amplification was completed by an additional cycle at 72 °C for 1 min, and products were analyzed by 2% agarose gel electrophoresis with detection using ethidium bromide under ultraviolet light. Amplicon products were sequenced (Genewiz®, Research Triangle Park, NC USA 27709) to establish species strain identification.

PCR screening for *Borrelia* spp. DNA targeting the ITS region was performed using IGSA: 5' CACGA CTT TCT TCG CCT TAA AGC 3' and IGSB: 5' GTT AAG CTC TTA TTC CTT GAT GGT A 3' as forward and reverse primers, respectively. Amplification was performed in a 25 μl final volume reaction as described above for *Bartonella* DNA amplification. Positive PCR controls were prepared using 5 μl of DNA of genomic *B. burgdorferi* strain B31 at concentration of 0.005 pg/μl. Conventional PCR was performed in an Eppendorf Mastercycler EPgradient® under the following conditions: a single hot-start cycle at 95 °C for 2 min followed by 55 cycles of denaturing at 94 °C for 15 s, annealing at 64 °C for 15 s, and extension at 72 °C for 18 s. Amplification was completed by an additional cycle at 72 °C for 30 s. As described for *Bartonella* testing, products were analyzed by 2% agarose gel electrophoresis with detection using ethidium bromide under ultraviolet light, and positive samples were sequenced for species and strain identification.

3. Results

Based upon house keeping gene amplification, PCR inhibition was not detected for any tick extracted DNA sample (results not shown). The

Table 2

Prevalence of *Borrelia burgdorferi*, *Borrelia bissettiae*, *Bartonella henselae* and *Borrelia/B. henselae* co-infections in *Ixodes affinis*, *Ixodes pacificus*, and *Ixodes scapularis* ticks from the United States.

| | N = | <i>B. burgdorferi</i> Infected (%) | <i>B. bissettiae</i> Infected (%) | <i>B. henselae</i> Infected (%) | <i>Borrelia</i> spp. + <i>B. henselae</i> Co-Infected (%) |
|--------------------------|-----|---------------------------------------|--------------------------------------|------------------------------------|--|
| <i>Ixodes scapularis</i> | 671 | 74 (11.0) | 0 (0) | 6 (0.9) | 1 (0.1) |
| <i>Ixodes affinis</i> | 155 | 55 (35.5) | 48 (31) | 16 (10.3) | 7 (4.5) |
| <i>Ixodes pacificus</i> | 103 | 0 (0) | 0 (0) | 1 (1.0) | 0 (0) |
| Total <i>Ixodes</i> spp. | 929 | 129 (13.9) | 48 (5.2) | 23 (2.5) | 8 (0.9) |

limit of detection for DNA amplification was 0.005 pg/μl and 0.001 pg/μl for *B. burgdorferi* and *B. henselae* DNA, respectively, as previously established (Lantos et al., 2014; Maggi et al., 2012, 2010). *Borrelia* and *Bartonella* DNA were amplified in all PCR positive controls (0.005 pg/μl and 0.001 pg/μl concentration, respectively), but not from any of the PCR negative controls used during this work.

3.1. *Bartonella* DNA screening

B. henselae, the only *Bartonella* species PCR amplified and sequenced from ticks in this study, was found in California (1/103 *I. pacificus*), Minnesota (1 of 62 *I. scapularis*), and North Carolina (5/348 *I. scapularis*, 16/155 *I. affinis*), for an overall prevalence of 2.6% (Table 2). The highest prevalence of *B. henselae* infection was found in *I. affinis* collected from NC, where *B. henselae* was amplified from 16 tick samples (10%).

3.2. *Borrelia* DNA screening

74 of 671 (11%) *I. scapularis* and 55/155 (36%) *I. affinis* were PCR positive for *B. burgdorferi*. *Borrelia* DNA was not amplified from 103 *I. pacificus* ticks from southern California or from *I. scapularis* collected in Florida, Louisiana or Virginia. The prevalence of *B. burgdorferi* in *I. scapularis* varied from 0.5–71.4% in the remaining states (Table 1). In addition to *B. burgdorferi*, *B. bissettiae* DNA, a member of the *B. burgdorferi* sensu lato complex (Maggi et al., 2010; Rudenko et al., 2014, 2011), was PCR amplified and sequenced from 48/155 (31%) *I. affinis* tick extractions. *Borrelia bissettiae* DNA was not amplified from ticks from other states. No other *Borrelia* spp. were amplified from the ticks in this study. Overall, *B. bissettiae* and/or *B. burgdorferi* DNA was amplified and sequenced from 98/155 (63.2%) *Ixodes affinis* (Table 2).

3.3. Co-infection

B. burgdorferi and *B. henselae* co-infections were infrequently documented among the 929 *Ixodes* ticks tested. In Minnesota, only 1/21 (5%) *B. burgdorferi* positive ticks was co-infected with *B. henselae*. Co-infection was more prevalent for *I. affinis* (Table 2). *B. burgdorferi* and *B. henselae* co-infection was found in 3/155 (2%) *I. affinis*, whereas *B. bissettiae* and *B. henselae* co-infection was found in 4/155 (2.6%) *I. affinis*, with an overall co-infection rate of 7/155 (4.5%) ticks tested.

4. Discussion and conclusions

In the United States, *Ixodes* ticks are the predominant vectors for *B. burgdorferi* transmission among animals and humans. Based upon PCR amplification of *Bartonella* DNA from questing ticks, *Ixodes* spp. are potential vectors for *B. henselae* transmission. In this study, we determined the prevalence of *Bartonella* and *Borrelia* spp. DNA in three *Ixodes* adult species (*I. scapularis*, *I. affinis*, and *I. pacificus*) collected from different regions of the US. As previously reported, our results indicate a higher prevalence for *B. burgdorferi* in the North and Northeast, than for other regions of the US (Table 1), with an average of 11.0% for *I. scapularis*, 35.5% for *I. affinis*, and 0% for *I. pacificus* in southern California (Table 2). Interestingly, other than *B. bissettiae*, no

other *Borrelia* spp. DNA was amplified in this study. In *I. affinis*, *B. bissettiae* or *B. burgdorferi* DNA was amplified from 63.2% of 155 *I. affinis* from North Carolina (Maggi et al., 2010) compared to 0.3% of 348 *I. scapularis* ticks from North Carolina tested in an identical manner. Although *I. affinis* is infrequently reported to attach to humans, even infrequent exposures to this tick species may carry a risk for transmission of *B. burgdorferi* and/or *B. bissettiae*.

Bartonella henselae was the only *Bartonella* species detected in this study. A very low prevalence of *B. henselae* DNA was found in *I. scapularis* from all regions surveyed (Table 1). The highest *B. henselae* prevalence was found in *I. affinis* ticks, with 10% of the 155 surveyed ticks being *B. henselae* PCR/DNA sequence positive. Based upon DNA sequencing, two *B. henselae* 16S-23S ITS strain types (*B. henselae* Houston 1 or San Antonio 2 strain) were identified. Our prevalence results are in contrast to previous studies where *Bartonella* DNA prevalences as high as 34.5% and 37.7% to 38.2% have been reported in New Jersey (USA) for *I. scapularis* and in Europe for *I. ricinus* (Adelson et al., 2004; Corrain et al., 2012; Dietrich et al., 2010; Ebani et al., 2015). This discrepancy could be attributed to regional (even at small geographical scale) differences, as reported from France (Vayssier-Tausat et al., 2016), where *B. henselae* prevalence varied from 0 to 36% in *Ixodes ricinus* collected within relatively close proximities.

Two different (and perhaps co-related) facts may account for differences in *Bartonella* prevalence among Ixodid ticks from the same or from disparate geographic regions: spatial-temporal distribution (clustering) and host association differences between larvae, nymph, and adult ticks (Dietrich et al., 2010; Nelder et al., 2014; Paul et al., 2016). Dietrich et al. reported a 14-fold higher *B. henselae* DNA prevalence in *I. ricinus* nymphal ticks when compared with adult ticks from the same geographic regions (Dietrich et al., 2010). Similarly, PCR positive rates varied from 11.8% to 38.2% for nymphs and from 0% to 12% for adult ticks depending on the region surveyed (Dietrich et al., 2010). Host preference, regional and/or seasonal variances seem to be the most likely reasons for the un-even pathogen prevalences for *Bartonella* and *Borrelia* spp. reported in *I. ricinus* in Europe (Bogumila and Adamska, 2005; Corrain et al., 2012; Cotte et al., 2010; Dietrich et al., 2010; Egyed et al., 2012; Halos et al., 2005; Hercik et al., 2007; La Scola et al., 2004; Raileanu et al., 2017; Sytykiewicz et al., 2012; Tjisse-Klasen et al., 2011) and in *I. scapularis* in the US (Adelson et al., 2004; Becker et al., 2016; Belongia, 2002; Hinrichsen et al., 2001; Lukin and Grabarev, 1999; Nelder et al., 2016). The potential impact/s of environmental, geospatial and tick life stage were not assessed in this study.

In this study, co-detection of *B. henselae* and *B. burgdorferi* DNA within a tick was infrequent, with *I. affinis* ticks being the species with the highest percentage of co-infection (5%).

Our co-infection findings also differ somewhat from previous reports from the US and Europe. Co-infection with *Bartonella* spp. and *B. burgdorferi* was reported in 8.4% of 107 *I. scapularis* tested from New Jersey (Adelson et al., 2004). In contrast, co-infection with *B. henselae* and *B. burgdorferi* was not documented in 419 adults *I. ricinus* from Germany and France (Dietrich et al., 2010). It is important to note that, for most of the states surveyed in this study, a very low number of ticks were available for testing. Therefore, our results are not representative of optimal tick sample size or distributions necessary for accurate

assessment of pathogen prevalence's within the three *Ixodes* spp. throughout the US. However, we were able to confirm *B. henselae* DNA in all three tick species tested. In addition to vectorial differences among the three *Ixodes* spp., host preferences, regional, and/or seasonal variances may be responsible in part for differences in *Bartonella* and *Borrelia* spp. prevalence's.

Based upon the results of this and other studies (Clark et al., 2002; Harrison et al., 2010; Maggi et al., 2010; Rudenko et al., 2013), a thorough understanding of the enzootic cycle as well as the vector competence of *I. affinis* for potential transmission of both *B. bissettiae*, *B. burgdorferi* and *B. henselae* should be emphasized. With the advent of new and improved diagnostic technologies, its seems that bartoneliosis and borreliosis disease awareness continues to increase (Nelder et al., 2016). Nevertheless, due to human population growth into suburban and rural areas, fluctuations in tick's host species densities, and the apparent climate changes that may favor *Ixodes* tick expansions and survival (Nelder et al., 2014; Paul et al., 2016), the full spectrum of animal and human tick-transmitted diseases remains incompletely characterized and funding to investigate the impacts will provide a more thorough risk assessment.

Authors' contribution

EBB and RGM supervised and conducted the study as presented here. MT and TM collected ticks for testing. TR assisted with PCR testing of extracted tick DNA. All authors contributed to, read and approved the final version of the manuscript.

Funding

This research was supported by the State of North Carolina, the Lyme Disease Association, Inc., and by Bayer Animal Health.

Competing interests

The authors declare that they have no competing interest.

Acknowledgments

We gratefully thank all individuals who participated in collecting and providing ticks throughout the study. We particularly want to thank Dr. Matt Eberts, Baxter MN, and to Bayer Animal Health, for funding part of this work.

References

- Adelson, M.E., Rao, R.V., Tilton, R.C., Cabets, K., Eskow, E., Fein, L., Occhi, J.L., Mordechai, E., 2004. Prevalence of *Borrelia burgdorferi*, *Bartonella* spp., *Babesia microti*, and *Anaplasma phagocytophilum* in *Ixodes scapularis* ticks collected in Northern New Jersey. *J. Clin. Microbiol.* 42, 2799–2801.
- Barbour, A.G., 2014. Phylogeny of a relapsing fever *Borrelia* species transmitted by the hard tick *Ixodes scapularis*. *Infect. Genet. Evol.* 27, 551–558.
- Becker, N.S., Margos, G., Blum, H., Krebs, S., Graf, A., Lane, R.S., Castillo-Ramirez, S., Sing, A., Fingerle, V., 2016. Recurrent evolution of host and vector association in bacteria of the *Borrelia burgdorferi* sensu lato species complex. *BMC Genomics* 17, 734.
- Belongia, E.A., 2002. Epidemiology and impact of coinfections acquired from *Ixodes* ticks. *Vector Borne Zoonotic Dis.* 2, 265–273.
- Billeter, S.A., Miller, M.K., Breitschwerdt, E.B., Levy, M.G., 2008. Detection of two *Bartonella* tamiae-like sequences in *Amblyomma americanum* (Acari: Ixodidae) using 16S-23S intergenic spacer region-specific primers. *J. Med. Entomol.* 45, 176–179.
- Billeter, S.A., Caceres, A.G., Gonzales-Hidalgo, J., Luna-Caypo, D., Kosoy, M.Y., 2011. Molecular detection of *Bartonella* species in ticks from Peru. *J. Med. Entomol.* 48, 1257–1260.
- Bogumila, S., Adamska, M., 2005. *Capreolus capreolus* and *Ixodes ricinus* as a reservoir of *Bartonella* in north-western Poland. *Wiad. Parazytol.* 51, 139–143.
- Breitschwerdt, E.B., 2014. Bartonellosis: one health perspectives for an emerging infectious disease. *ILAR J.* 55, 46–58.
- Breitschwerdt, E.B., 2017. Bartonellosis, one health and all creatures great and small. *Vet. Dermatol.* 28, 96–e21.
- Breitschwerdt, E.B., Maggi, R.G., Chomei, B.B., Lappin, M.R., 2010. Bartonellosis: an emerging infectious disease of zoonotic importance to animals and human beings. *J. Vet. Emerg. Crit. Care (San Antonio)* 20, 8–30.
- Breuner, N.E., Dolan, M.C., Repligie, A.J., Sexton, C., Hojgaard, A., Boegler, K.A., Clark, R.J., Eisen, L., 2017. Transmission of *Borrelia miyamotoi* sensu lato relapsing fever group spirochetes in relation to duration of attachment by *Ixodes scapularis* nymphs. *Ticks Tick. Dis.* 8, 677–681.
- Chang, C.C., Chomei, B.B., Kasten, R.W., Romano, V., Tietze, N., 2001. Molecular evidence of *Bartonella* spp. in questing adult *Ixodes pacificus* ticks in California. *J. Clin. Microbiol.* 39, 1221–1226.
- Chang, C.C., Hayashidani, H., Pusterla, N., Kasten, R.W., Madigan, J.E., Chomei, B.B., 2002. Investigation of *Bartonella* infection in Ixodid ticks from California. *Comp. Immunol. Microbiol. Infect. Dis.* 25, 229–236.
- Chomei, B.B., Kasten, R.W., 2010. Bartonellosis, an increasingly recognized zoonosis. *J. Appl. Microbiol.* 109, 743–750.
- Clark, K., 2004. *Borrelia* species in host-seeking ticks and small mammals in northern Florida. *J. Clin. Microbiol.* 42, 5076–5086.
- Clark, K.L., Oliver Jr., J.H., James, A.M., Durden, L.A., Banks, C.W., 2002. Prevalence of *Borrelia burgdorferi* sensu lato infection among rodents and host-seeking ticks in South Carolina. *J. Med. Entomol.* 39, 198–206.
- Corrain, R., Drigo, M., Fenati, M., Menandro, M.L., Mondin, A., Pasotto, D., Martini, M., 2012. Study on ticks and tick-borne zoonoses in public parks in Italy. *Zoonoses Public Health* 59, 468–476.
- Cotte, V., Bonnet, S., Le Rhun, D., Le Naour, E., Chauvin, A., Boulouis, H.J., Lecuelle, B., Lilin, T., Vayssier-Taussat, M., 2008. Transmission of *Bartonella henselae* by *Ixodes ricinus*. *Emerging Infect. Dis.* 14, 1074–1080.
- Cotte, V., Bonnet, S., Cote, M., Vayssier-Taussat, M., 2010. Prevalence of five pathogenic agents in questing *Ixodes ricinus* ticks from western France. *Vector Borne Zoonotic Dis.* 10, 723–730.
- Dietrich, F., Schmidgen, T., Maggi, R.G., Richter, D., Matuschka, F.R., Vonthein, R., Breitschwerdt, E.B., Kempf, V.A., 2010. Prevalence of *Bartonella henselae* and *Borrelia burgdorferi* sensu lato DNA in *Ixodes ricinus* ticks in Europe. *Appl. Environ. Microbiol.* 76, 1395–1398.
- Ebani, V.V., Bertelloni, F., Turchi, B., Filogari, D., Cerri, D., 2015. Molecular survey of tick-borne pathogens in Ixodid ticks collected from hunted wild animals in Tuscany, Italy. *Asian Pac. J. Trop. Med.* 8, 714–717.
- Egyed, L., Elio, P., Sreter-Lancz, Z., Szell, Z., Balogh, Z., Sreter, T., 2012. Seasonal activity and tick-borne pathogen infection rates of *Ixodes ricinus* ticks in Hungary. *Ticks Tick. Dis.* 3, 90–94.
- Ereqat, S., Nasereddin, A., Vayssier-Taussat, M., Abdelkader, A., Al-Jawabreh, A., Zaid, T., Azmi, K., Abdeen, Z., 2016. Molecular evidence of *Bartonella* species in Ixodid ticks and domestic animals in palestine. *Front. Microbiol.* 7, 1217.
- Guptill, L., 2010. Bartonellosis. *Vet. Microbiol.* 140, 347–359.
- Halos, L., Jamal, T., Maillard, R., Beugnet, F., Le Menach, A., Boulouis, H.J., Vayssier-Taussat, M., 2005. Evidence of *Bartonella* sp. in questing adult and nymphal *Ixodes ricinus* ticks from France and co-infection with *Borrelia burgdorferi* sensu lato and *Babesia* sp. *Vet. Res.* 36, 79–87.
- Han, S., Hickling, G.J., Isao, J.I., 2016. High prevalence of *Borrelia miyamotoi* among adult blacklegged ticks from white-tailed deer. *Emerg. Infect. Dis.* 22, 316–318.
- Harrison, B.A., Rayburn Jr., W.H., Toliver, M., Powell, E.E., Engber, B.R., Durden, L.A., Robbins, R.G., Prendergast, B.F., Whitt, P.B., 2010. Recent discovery of widespread *Ixodes affinis* (Acari: Ixodidae) distribution in North Carolina with implications for Lyme disease studies. *J. Vector Ecol.* 35, 174–179.
- Harrison, A., Bown, K.J., Montgomery, W.I., Birtles, R.J., 2012. *Ixodes ricinus* is not an epidemiologically relevant vector of *Bartonella* species in the wood mouse (*Apodemus sylvaticus*). *Vector Borne Zoonotic Dis.* 12, 366–371.
- Hercik, K., Hasova, V., Janacek, J., Branny, P., 2007. Molecular evidence of *Bartonella* DNA in Ixodid ticks in Czechia. *Folia Microbiol.* 52, 503–509.
- Hinrichsen, V.L., Whitworth, U.G., Breitschwerdt, E.B., Hegarty, B.C., Mather, I.N., 2001. Assessing the association between the geographic distribution of deer ticks and seropositivity rates to various tick-transmitted disease organisms in dogs. *J. Am. Vet. Med. Assoc.* 218, 1092–1097.
- Holden, K., Boothby, J.I., Kasten, R.W., Chomei, B.B., 2006. Co-detection of *Bartonella henselae*, *Borrelia burgdorferi*, and *Anaplasma phagocytophilum* in *Ixodes pacificus* ticks from California, USA. *Vector Borne Zoonotic Dis.* 6, 99–102.
- Johnson, T.L., Graham, C.B., Boegler, K.A., Cherry, C.C., Maes, S.E., Pilgard, M.A., Hojgaard, A., Buttke, D.E., Eisen, R.J., 2017. Prevalence and diversity of tick-borne pathogens in nymphal *Ixodes scapularis* (Acari: Ixodidae) in Eastern National Parks. *J. Med. Entomol.* 54, 742–751.
- King, J.L., Smith, A.D., Mitchell, E.A., Allen, M.S., 2017. Validation of droplet digital PCR for the detection and absolute quantification of *Borrelia* DNA in *Ixodes scapularis* ticks. *Parasitology* 144, 359–367.
- La Scola, B., Holmberg, M., Raoult, D., 2004. Lack of *Bartonella* sp. in 167 *Ixodes ricinus* ticks collected in central Sweden. *Scand. J. Infect. Dis.* 36, 305–306.
- Lantos, P.M., Maggi, R.G., Ferguson, B., Varkey, J., Park, L.P., Breitschwerdt, E.B., Woods, C.W., 2014. Detection of *Bartonella* species in the blood of veterinarians and veterinary technicians: a newly recognized occupational hazard? *Vector Borne Zoonotic Dis.* 14, 563–570.
- Lukin, E.P., Grabarev, P.A., 1999. Ecological aspects of human pathogens Ehrlichia and Bartonella. *Med. Parazitol. (Mosk)* 17–22.
- Lyme-CDC, 2018. Lyme Disease Cases-CDC. CDC. <https://www.cdc.gov/lyme/stats/humancases.html>.
- Mada, P.K., Joel Chandranesan, A.S., 2017. Bartonellosis. *StatPearls, Treasure Island (FL)*.
- Maggi, R.G., Raverty, S.A., Lester, S.J., Huff, D.G., Haulena, M., Ford, S.L., Nielsen, O., Robinson, J.H., Breitschwerdt, E.B., 2008. *Bartonella henselae* in captive and hunter-harvested beluga (*Delphinapterus leucas*). *J. Wildl. Dis.* 44, 871–877.
- Maggi, R.G., Reichelt, S., Toliver, M., Engber, B., 2010. *Borrelia* species in *Ixodes affinis*

- and *Ixodes scapularis* ticks collected from the coastal plain of North Carolina. *Ticks Tick. Dis.* 1, 168–171.
- Maggi, R.G., Mascarelli, P.E., Pultorak, E.L., Hegarty, B.C., Bradley, J.M., Mozayani, B.R., Breitschwerdt, E.B., 2011. *Bartonella* spp. bacteremia in high-risk immunocompetent patients. *Diagn. Microbiol. Infect. Dis.* 71, 430–437.
- Maggi, R.G., Mozayani, B.R., Pultorak, E.L., Hegarty, B.C., Bradley, J.M., Correa, M., Breitschwerdt, E.B., 2012. *Bartonella* spp. bacteremia and rheumatic symptoms in patients from Lyme disease-endemic region. *Emerg. Infect. Dis.* 18, 783–791.
- Mascarelli, P.E., Tartara, G., Pereyra, N.B., Maggi, R.G., 2016. Detection of *Mycoplasma haemocanis*, *Mycoplasma haematoparvum*, *Mycoplasma suis* and other vector-borne pathogens in dogs from Córdoba and Santa Fé, Argentina. *Parasit. Vectors* 9, 642.
- Mayne, P.J., 2011. Emerging incidence of Lyme borreliosis, babesiosis, bartonellosis, and granulocytic ehrlichiosis in Australia. *Int. J. Gen. Med.* 4, 845–852.
- Murphy, D.S., Lee, X., Larson, S.R., Johnson, D.K., Loo, T., Paskewitz, S.M., 2017. Prevalence and distribution of human and tick infections with the ehrlichia muris-like agent and *Anaplasma phagocytophilum* in Wisconsin, 2009–2015. *Vector Borne Zoonotic Dis.* 17, 229–236.
- Nadolin, R.M., Wright, C.L., Hynes, W.L., Sonenshine, D.E., Gaff, H.D., 2011. *Ixodes affinis* (Acari: Ixodidae) in southeastern Virginia and implications for the spread of *Borrelia burgdorferi*, the agent of Lyme disease. *J. Vector Ecol.* 36, 464–467.
- Neider, M.P., Russell, C., Lindsay, L.R., Dhar, B., Patel, S.N., Johnson, S., Moore, S., Kristjanson, E., Li, Y., Ralevski, F., 2014. Population-based passive tick surveillance and detection of expanding foci of blacklegged ticks *Ixodes scapularis* and the Lyme disease agent *Borrelia burgdorferi* in Ontario, Canada. *PLoS One* 9, e105358.
- Neider, M.P., Russell, C.B., Sheehan, N.J., Sander, B., Moore, S., Li, Y., Johnson, S., Patel, S.N., Sider, D., 2016. Human pathogens associated with the blacklegged tick *Ixodes scapularis*: a systematic review. *Parasit. Vectors* 9, 265.
- Nelson, C.A., Saha, S., Mead, P.S., 2016. Cat-scratch disease in the United States, 2005–2013. *Emerg. Infect. Dis.* 22, 1741–1746.
- Noguchi, H., 1926. Etiology of oroya fever: V. The experimental transmission of *Bartonella bacilliformis* by ticks (*DERMACENTOR Andersoni*). *J. Exp. Med.* 44, 729–734.
- Oliver Jr., J.H., Lin, T., Gao, L., Clark, K.L., Banks, C.W., Durden, L.A., James, A.M., Chandler Jr., F.W., 2003. An enzootic transmission cycle of Lyme borreliosis spirochetes in the southeastern United States. *Proc. Natl. Acad. Sci. U. S. A.* 100, 11642–11645.
- Paddock, C., Lane, K., Staples, J., Labruna, M., 2016. Global health impacts of vector-borne diseases: changing paradigms for tick-borne diseases in the Americas. Forum on Microbial Threats; Board on Global Health; Health and Medicine Division; National Academies of Sciences, Engineering, and Medicine, vol. A8 National Academies Press, Washington (DC).
- Paul, R.E., Cote, M., Le Naour, E., Bonnet, S.I., 2016. Environmental factors influencing tick densities over seven years in a French suburban forest. *Parasit. Vectors* 9, 309.
- Raileanu, C., Moutailler, S., Pavel, I., Porea, D., Mihalca, A.D., Savuta, G., Vayssier-Tausat, M., 2017. *Borrelia* diversity and co-infection with other tick borne pathogens in ticks. *Front. Cell. Infect. Microbiol.* 7, 36.
- Reis, C., Cote, M., Le Rhun, D., Lecuelle, B., Levin, M.L., Vayssier-Tausat, M., Bonnet, S.I., 2011. Vector competence of the tick *Ixodes ricinus* for transmission of *Bartonella birtlesii*. *PLoS Negl. Trop. Dis.* 5, e1186.
- Rudenko, N., Golovchenko, M., Grubhoffer, L., Oliver Jr., J.H., 2011. Updates on *Borrelia burgdorferi* sensu lato complex with respect to public health. *Ticks Tick. Dis.* 2, 123–128.
- Rudenko, N., Golovchenko, M., Honig, V., Mallatova, N., Krbkova, L., Mikulasek, P., Fedorova, N., Belfiore, N.M., Grubhoffer, L., Lane, R.S., Oliver Jr., J.H., 2013. Detection of *Borrelia burgdorferi* sensu stricto ospC alleles associated with human Lyme borreliosis worldwide in non-human-biting tick *Ixodes affinis* and rodent hosts in Southeastern United States. *Appl. Environ. Microbiol.* 79, 1444–1453.
- Rudenko, N., Golovchenko, M., Belfiore, N.M., Grubhoffer, L., Oliver Jr., J.H., 2014. Divergence of *Borrelia burgdorferi* sensu lato spirochetes could be driven by the host diversity of *Borrelia* strains isolated from ticks feeding on a single bird. *Parasit. Vectors* 7 (4).
- Swanson, K.L., Norris, D.E., 2007. Co-circulating microorganisms in questing *Ixodes scapularis* nymphs in Maryland. *J. Vector Ecol.* 32, 243–251.
- Sytykiewicz, H., Karbowiak, G., Werszko, J., Czerniewicz, P., Sprawka, I., Mitrus, J., 2012. Molecular screening for *Bartonella henselae* and *Borrelia burgdorferi* sensu lato co-existence within *Ixodes ricinus* populations in central and eastern parts of Poland. *Ann. Agric. Environ. Med.* 19, 451–456.
- Telford 3rd, S.R., Wormser, G.P., 2010. *Bartonella* spp. transmission by ticks not established. *Emerg. Infect. Dis.* 16, 379–384.
- Tijssen-Klasen, E., Fonville, M., Gassner, F., Nijhof, A.M., Hovius, E.K., Jongejans, F., Takken, W., Reimerink, J.R., Overgaauw, P.A., Sprong, H., 2011. Absence of zoonotic *Bartonella* species in questing ticks: first detection of *Bartonella clarridgeiae* and *Rickettsia felis* in cat fleas in the Netherlands. *Parasit. Vectors* 4, 61.
- Tokarz, R., Tagliafierro, T., Cucura, D.M., Rochlin, I., Sameroff, S., Lipkin, W.I., 2017. Detection of *Anaplasma phagocytophilum*, *Babesia microti*, *Borrelia burgdorferi*, *Borrelia miyamotoi*, and Powassan virus in ticks by a multiplex real-time reverse transcription-pcr assay. *mSphere* 2.
- Vayssier-Tausat, M., Moutailler, S., Femenia, F., Raymond, P., Croce, O., La Scola, B., Fournier, P.E., Raoult, D., 2016. Identification of novel zoonotic activity of *Bartonella* spp., France. *Emerg. Infect. Dis.* 22, 457–462.
- Vilcins, I.M., Kosoy, M., Old, J.M., Deane, E.M., 2009. *Bartonella*-like DNA detected in *Ixodes tasmani* ticks (Acari: Ixodida) infesting Koalas (*Phascogale cinerea*) in Victoria, Australia. *Vector Borne Zoonotic Dis.* 9, 499–503.
- Wroblewski, D., Gebhardt, L., Prusinski, M.A., Meehan, L.J., Halse, T.A., Musser, K.A., 2017. Detection of *Borrelia miyamotoi* and other tick-borne pathogens in human clinical specimens and *Ixodes scapularis* ticks in New York State, 2012–2015. *Ticks Tick. Dis.* 8, 407–411.