

Summer terms

Dominik Glandorf

2022-08-19

```
terms = get_term_data()
students = get_student_sub()
terms_sub = merge(terms, students[,c('mellon_id', 'admitdate')], by="mellon_id")
```

Where do we have summer terms?

```
table(terms_sub[,c('term_part_code', 'term_part_desc')])
```

```
##           term_part_desc
## term_part_code  Fall Fall Quarter Spring Spring Quarter Summer Summer-1
##           03         0           0      0           0         0         0
##           14         0           0 166662           1         0         0
##           25         0           0      0           0         0         5
##           51         0           0      0           0      5012         0
##           92 176788         4      0           0         0         0
##           term_part_desc
## term_part_code Summer Quarter (Med School) and Summer Academic Term (OIR)
##           03                                           0
##           14                                           0
##           25                                           0
##           51                                           10
##           92                                           0
##           term_part_desc
## term_part_code Winter Winter Quarter
##           03 172894           5
##           14         0           0
##           25         0           0
##           51         0           0
##           92         0           0
```


We are interested in term part code 51, which in general does not occur that much.

```
1-(sum(terms_sub$term_part_code %in% c("03", "14", "92")) / length(terms_sub$term_part_code))
```

```
## [1] 0.009675891
```

In around 1% of the terms it is other than Fall, Winter or Spring.

```
ggplot(terms_sub, aes(fill=admitdate, x=term_part_code)) +  
  geom_bar(position="stack", stat="count")
```



Summer terms were only tracked from 2015 or 2016 on.

Conclusion: do not include summer terms for comparability.