


# Proton Dynamics in Cavities

Dominik Wille

October 23, 2014



# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Analytic calculations</b>	<b>4</b>
2.1	Constructing the single-ion correlation function . . . . .	5
2.1.1	One surface . . . . .	5
2.1.2	Two surfaces . . . . .	6
2.2	Limiting case . . . . .	7
<b>3</b>	<b>Brownian Dynamics</b>	<b>8</b>
3.1	Langevin Symulation . . . . .	8
3.1.1	Priciple . . . . .	8
3.1.2	Physical background . . . . .	8
3.2	Connection to diffusion . . . . .	9
3.2.1	Fixed step size . . . . .	9
3.2.2	Gaussian random walk . . . . .	11
3.3	Mean square distance . . . . .	12
<b>4</b>	<b>Free particle - MSD</b>	<b>13</b>
<b>5</b>	<b>Infinite plates</b>	<b>14</b>
5.1	Simple random walk vs. Gaussian random walk . . . . .	14
5.2	Probability density . . . . .	16
5.3	Correlation functions . . . . .	18
5.4	Parameters of the simulation . . . . .	18
5.5	Interpretation . . . . .	19
5.6	Comparison . . . . .	19
<b>6</b>	<b>Finite plates</b>	<b>22</b>
<b>7</b>	<b>Conclusion and Outlook</b>	<b>23</b>
<b>8</b>	<b>Annex</b>	<b>24</b>

# 1 Introduction

Proton- or generally ion dynamics in fluids directly imply fluctuations in the potential and conductivity of surrounded nano-scaled structures. The major processes which influence these measurable properties and which are considered here are **absorption** and **desorption** processes. The amount of mobile ions which are contained in the fluid are proportional to the fluids conductivity. An experimental way to detect this conductivity is by measuring the ionic current. And since ions carry charges, absorption of ions causes changes of the potential.

Since such fluctuations, could bring hints to extract physical, chemical or even geometrical information concerning the system, they are part of current investigations. Just now some of these fluctuations are the topic of current research<sup>1</sup>.

In this document the motion of one particle is considered as a brownian motion. Even though ions and especially protons are not necessarily smaller than the fluid molecules which is the reason why the diffusion constant does not fulfill EINSTEINS formula<sup>2</sup>, the general assumption that the motion is caused by random strokes is fulfilled for smaller particles as well. In this document a **Langevin-Simulation** of a particle is used to determine correlations of the absorption-desorption processes, which can be indirectly measured as conductivity and potential.

The simulations are based on previous analytic calculations by ROLAND NETZ which obtain these correlation functions for infinite cylindrical and planar geometries<sup>3</sup>. To determine the correctness of the **Langevin-Simulation** it is tested for these limiting cases as well as simple simulations for a free particle. Moreover the most important processes which take effect in the correlation functions and methods to describe and implement these in the simulation are briefly summarized and explained.

The title image shows one visualized particle on a random walk between two plates in 2-d for different times. The time increases from left to right.

---

<sup>1</sup>[5] Simon Gravelle and Bocquet “Pink noise of ionic current, theory and modelisation” 2014

<sup>2</sup>[4] “Notes on Brownian Motion” 2011

<sup>3</sup>[3] Netz and Bocquet “Conductivity Fluctuations in Cylindrical Pores” 2014

## 2 Analytic calculations

The analytic calculations are mostly inherited from [3] and [2]. To understand the used properties in the simulation later on, the key principles should be explained briefly.

Considered should be  $N$  charges carriers such as ions or protons. For every particle  $i \in \{1, 2, \dots, N\}$  the binary function  $n_j^i(t)$  should stand for the absorption state of the particle on the surface  $j$ .  $n_A^i(t) = 1$  if the particle is absorbed on Plate  $A$  and  $n_A^i(t) = 0$  if it is not.

Of course the number of particles absorbed on plate  $A$  is given by:

$$N_A(t) = \sum_{i=1}^N n_A^i(t) \quad (1)$$

Since experimental investigation measure fluctuations in the potential of nano-structures, especially the auto-correlation function of  $N_A(t)$ , is of essential importance.

$$\langle N_A(0)N_A(t) \rangle = \sum_{i,j=1}^N \langle n_A^i(0)n_A^j(t) \rangle \quad (2)$$

$$= \sum_{i \neq j}^N \langle n_A^i(0)n_A^j(t) \rangle + \sum_{i=1}^N \langle n_A^i(0)n_A^i(t) \rangle \quad (3)$$

The particles are usually uncorrelated among each other

$$\langle n_A^i(0)n_A^j(t) \rangle = \langle n_A(0) \rangle \langle n_A(t) \rangle \quad (4)$$

so let the expectation value of the absorption state for an single particle be:

$$\langle n_A^i(t) \rangle = \langle n_A^i(0) \rangle = p_A \quad (5)$$

$$\Rightarrow \langle N_A(0)N_A(t) \rangle = N(N-1)p_A^2 + N\langle n_A(0)n_A(t) \rangle \quad (6)$$

The only non-trivial quantity to calculate can be defined via the single-ion correlation function  $C_{AA}(t)$  which is the probability that a particle is absorbed at time  $t$  given it is absorbed at time  $t = 0$ .

$$C_{AA}(t) = \langle n_A(0)n_A(t) \rangle / \langle n_A(0) \rangle \quad (7)$$

$$\Rightarrow \langle N_A(0)N_A(t) \rangle = N^2 p_A^2 + N p_A C_{AA}(t) - N p_A^2 \quad (8)$$

## 2.1 Constructing the single-ion correlation function

### 2.1.1 One surface

For a single surface  $A$ , one particle that is absorbed at time  $t = 0$  can basically keep absorbed until time  $t$  or desorb  $n$ -times before it is absorbed again, if it should be absorbed at time  $t$ .




Figure 1: Illustrated construction of  $C_{AA}(t)$

As illustrated in the figure above  $C_{AA}(t)$  as the sum of different paths. The distribution  $P(t)$  is the Probability for an absorbed ion to desorb at time  $t$  for the first time.  $Q_A(t)$  is the survival distribution for an ion to be absorbed on the surface  $A$  over the time span  $t = 0$  to  $t$ . Note that this equals the probability that ion desorbes at some time  $> t$ . It is given by:

$$Q_A(t) = \int_t^\infty P(t') dt' \quad (9)$$

$J_{AA}(t)$  is the probability for an ion to desorb at time  $t = 0$  and to return to the surface  $A$  at time  $t$ .

The probability that a particle is absorbed at time  $t$  given it is absorbed at time  $t = 0$  is the sum over the probabilities of all ways on which a particle is absorbed at time  $t = 0$  and on time  $t$ . The probability of one way on the other hand is the product of the probabilities of its parts  $Q_A$ ,  $P_A$ , and  $J_{AA}$ .

Generally all that stages in  $C_{AA}(t)$  of particles could have every time  $< t$ . Therefore it is necessary to vary all these times.

The construct of  $C_{AA}$  below sums up all variations illustrated in *Figure 1* with general times  $t_e$  for  $Q(t)$ ,  $t_j$  for  $P_A(t)$  and  $t'_j$  for  $J_{AA}(t)$ . In order to obtain the probability at time  $t$  the **Dirac-Delta-Function** picks out variants with  $\sum_j (t_j + t'_j) + t_e = t$ .

$$C_{AA}(t) = \sum_{i=0}^{\infty} \left\{ \int_0^\infty dt_e Q_A(t_e) \prod_{j=1}^i \left[ \int_0^\infty dt_j P_A(t_j) \int_0^\infty dt'_j J_{AA}(t'_j) \right] \delta \left( \sum_{k=1}^i (t_k + t'_k) + t_e - t \right) \right\} \quad (10)$$

This construct itself is very though to analyze. But there is common trick to derive the **Laplace-transform**  $\tilde{C}_{AA}(\omega)$  of  $C_{AA}(t)$ .

$$\tilde{C}_{AA}(\omega) = \int_0^\infty e^{-\omega t'} \cdot C_{AA}(t') dt' \quad (11)$$

Remember that the integrand in *Equation 10* has only non-zero values for  $\sum_j (t_j + t'_j) + t_e = t$ . Therefore it is possible to split the  $e^{-\omega t'}$ -term.

$$e^{-\omega t'} = e^{-\omega t_e} \cdot \prod_{j=1}^i e^{-\omega t_j} \cdot e^{-\omega t'_j} \quad (12)$$

$$\Rightarrow \tilde{C}_{AA}(\omega) = \sum_{i=0}^{\infty} \left\{ \int_0^\infty dt_e e^{-\omega t_e} Q_A(t_e) \prod_{j=1}^i \left[ \int_0^\infty dt_j e^{-\omega t_j} P_A(t_j) \int_0^\infty dt'_j e^{-\omega t'_j} J_{AA}(t'_j) \right] \delta \left( \sum_{k=1}^i (t_k + t'_k) + t_e - t \right) \right\} \quad (13)$$

$$= \sum_{i=0}^{\infty} \left\{ \tilde{Q}_A(\omega) \prod_{j=1}^i [\tilde{P}_A(\omega) \cdot \tilde{J}_{AA}(\omega)] \right\} \quad (14)$$

$$= \sum_{i=0}^{\infty} \tilde{Q}_A(\omega) [\tilde{P}_A(\omega) \cdot \tilde{J}_{AA}(\omega)]^i \quad (15)$$

With the formula for geometric series

$$\sum_{i=0}^{n-1} q^i = \frac{1 - q^n}{1 - q} \quad (16)$$

this becomes the following expression:

$$\tilde{C}_{AA}(\omega) = \frac{\tilde{Q}_A(\omega)}{1 - \tilde{P}_A(\omega) \cdot \tilde{J}_{AA}(\omega)} \quad (17)$$

### 2.1.2 Two surfaces

The calculation of  $\tilde{C}_{AA}(\omega)$  for one surface should give a general idea how the laplace-transform of The correlation functions look like.

For instance  $\overline{C}_{AA}(\omega)$ ,  $\overline{C}_{BB}(\omega)$  which omit the  $\tilde{Q}_{A/B}(\omega)$  terms, represent ways where the particle is absorbed at time  $t = 0$  and exactly absorbs at time  $t$ .

$$\overline{C}_{AA}(\omega) = \frac{1}{1 - \tilde{P}_A(\omega) \cdot \tilde{J}_{AA}(\omega)} \quad (18)$$

$$\overline{C}_{BB}(\omega) = \frac{1}{1 - \tilde{P}_B(\omega) \cdot \tilde{J}_{BB}(\omega)} \quad (19)$$

Parts which are performed one time are in the nominator, parts which are performed  $n$ -times are in the denominator. Parts which are performed equal times can be just multiplied in the laplace-transform.

$\tilde{C}_{AA}(\omega)$  can be constructed as:

$$\tilde{C}_{AA}(\omega) = \frac{\overline{C}_{AA}(\omega) \tilde{Q}_A(\omega)}{1 - \overline{C}_{AA}(\omega) \tilde{P}_A(\omega) \tilde{J}_{AB}(\omega) \overline{C}_{BB}(\omega) \tilde{P}_B(\omega) \tilde{J}_{BA}(\omega)} \quad (20)$$

$$= \frac{\tilde{Q}_A(\omega) \left(1 - \tilde{P}_A(\omega) \tilde{J}_{AA}(\omega)\right)}{\left(1 - \tilde{P}_A(\omega) \tilde{J}_{AA}(\omega)\right) \left(1 - \tilde{P}_B(\omega) \tilde{J}_{BB}(\omega)\right) - \tilde{P}_A(\omega) \tilde{J}_{AB}(\omega) \tilde{P}_B(\omega) \tilde{J}_{BA}(\omega)} \quad (21)$$

## 2.2 Limiting case

## 3 Brownian Dynamics

The motion of a particle such as a proton in a fluid is primarily caused by collisions with other particles. This motion firstly was described by ROBERT BROWN who observed the motion of minute particles of pollen in water and is widely known as **brownian motion**.

### 3.1 Langevin Symulation

#### 3.1.1 Priciple

The principle of a Langevin simulation generally is to consider collisions with other particles as a random force. And propergate the position of a particle over small time steps  $\delta t$ .

#### 3.1.2 Physical background

An approach to describe situations with brownian motion was suggested by PAUL LANGEVIN who added the random force  $\mathbf{Z}(t)$  in newtons equation of motion. This stochastic term represents the collision driven force. His equation, the Langevin equation reads:

$$m\ddot{\mathbf{r}} = -\lambda\dot{\mathbf{r}} + \mathbf{Z}(t) \quad (22)$$

where  $m$  is the mass of the particle,  $\mathbf{r}$  the position of the particle and  $\lambda$  The friction constant.

Brownian dynamics can be represented with the so called **overdamped langevin equation** where the  $m\ddot{\mathbf{r}}$  term is neglected. The equation for the prevailing situation therefore is:

$$\lambda\dot{\mathbf{r}} = \mathbf{Z}(t) \quad (23)$$

In order to get an iterable expression this expression is discretised in time intervals  $\Delta t$ :



$$\int_t^{t+\delta t} \dot{\mathbf{r}}(t') dt' = \int_t^{t+\Delta t} \frac{\mathbf{Z}(t')}{\lambda} dt' \quad (24)$$

$$\mathbf{r}(t + \delta t) - \mathbf{r}(t) = \frac{1}{\lambda} \int_t^{t+\Delta t} \mathbf{Z}(t') dt' \quad (25)$$

$$\mathbf{r}(t + \delta t) - \mathbf{r}(t) \cong \boldsymbol{\zeta}(t, \varepsilon) \quad (26)$$

In the discussed 1-dimensional case  $\varepsilon$  is the mean step size and  $\boldsymbol{\zeta}(t, \varepsilon)$  is one random vector. For a simple random walk with unique step size this will be  $\varepsilon$  or  $-\varepsilon$  with a chance of 50% each. For a gaussian random walk this will be some value  $s$  with a probability:

$$P(s) = \frac{1}{\varepsilon \sqrt{2\pi}} e^{-s^2/2\varepsilon^2} \quad (27)$$

The position of a particle at time  $t + \Delta t$  can easily derived from its position at time  $t$ .

$$\mathbf{r}(t + \delta t) = \mathbf{r}(t) + \boldsymbol{\zeta}(t, \varepsilon) \quad (28)$$

### 3.2 Connection to diffusion

The probability density function for both mentioned types of **random walks** fulfill the diffusion equation (*Note:  $n = t/\delta t$* ):

$$\frac{\partial}{\partial t} \rho(x, t) = D \frac{\partial^2}{\partial x^2} \rho(x, t) \quad * \quad (29)$$

$$\text{with } D = \frac{\varepsilon^2}{2\delta t} \quad (30)$$

What means that a random walk can also be seen as a diffusion process. The exact form of the probability density functions is discussed below.

#### 3.2.1 Fixed step size

The motion which is performed in the langevin simulation is widely known as a **random walk**. The simplest variant is as mentioned before to choose  $\varepsilon$  or  $-\varepsilon$  with a chance of 50% each. The probability  $p(x, n)$  that a random walk with discrete steps of length  $\varepsilon$  comes to a position  $x$  after  $n$  steps. Is given by:

$$p(x, n) = \frac{\text{Number of ways to position } x}{\text{Total number of ways}} = \frac{N_x}{N} \quad (31)$$

Since there are 2 possible successors for every position the total number of ways doubles every step.

$$N = 2^n \quad (32)$$

The number of ways to the position  $x$  after  $n$  steps can be obtained by **Pascal's triangle**.




Figure 2: Number of ways to different distances

$$N_x = \binom{n}{k} \quad (33)$$

$$\text{with } k = \frac{1}{2} \left( \frac{x}{\varepsilon} + n \right) \quad (34)$$

Therefore  $p(x, n)$  follows as:

$$p(x, n) = \binom{n}{k} \cdot 2^{-n} \quad (35)$$

A consequence of the **de Moivre–Laplace theorem** is that for large numbers of steps ( $n \rightarrow \infty$ ) this expression can be approximated with the following gaussian curve:

$$p(x, n) \cong \sqrt{\frac{2}{n\pi}} e^{-x^2/2\varepsilon^2 n} \quad * \quad (36)$$

The probability function  $p(x, n)$  has valid values only for values

$$x \in \{x \mid x = z \cdot 2\varepsilon + \varepsilon (n \bmod 2), z \in \mathbb{Z} \wedge |x| \leq n\varepsilon\} \quad (37)$$

For other values  $x$  the probability is 0. This means that there is only one value per  $2\varepsilon$  interval, therefore the probability density function  $\rho(x, n)$  is:

$$\rho(x, n) = \frac{1}{2\varepsilon} p(x, n) = \frac{1}{\varepsilon\sqrt{2\pi n}} e^{-x^2/2\varepsilon^2 n} \quad (38)$$

Which is the normal distribution with  $\mu = 0$ ;  $\sigma = \varepsilon\sqrt{n}$ .

### 3.2.2 Gaussian random walk

The probability density function for a gaussian random walk with mean step size  $\varepsilon$  has exactly the same form as the for fixed step size  $\varepsilon$ .

$$\rho(x, n) = \frac{1}{\varepsilon\sqrt{2\pi n}} e^{-x^2/2\varepsilon^2 n} \quad (39)$$

This can easily be shown with a mathematical induction for  $p(x, n)$ . Note that because every position is possible the probability

$$p(x, n) = 0 \quad (40)$$

because that means that the number of possible positions is infinite.

**Base case** ( $n = 1$ ):

$$\rho(x, 1) = \frac{1}{\varepsilon\sqrt{2\pi}} e^{-x^2/2\varepsilon^2} \quad (41)$$

Which is exactly the expression of a gaussian distributed 1-dimensional random vector with  $\varepsilon$  variance. ✓

**Inductive step** ( $n + 1$ ):

$$\rho(x, n + 1) = \int_{-\infty}^{\infty} \frac{1}{\varepsilon\sqrt{2n\pi}} e^{-x'^2/2n\varepsilon^2} \cdot \frac{1}{\varepsilon\sqrt{2\pi}} e^{-(x - x')^2/2\varepsilon^2} dx' \quad (42)$$

$$= \frac{1}{\varepsilon\sqrt{2(n+1)\pi}} e^{-x^2/2(n+1)\varepsilon^2} \quad * \quad (43)$$

and since the calculated  $\rho(x, n + 1)$  is the same as  $\rho(x, n)$  in *Equation 38* the expression is proofed. ✓

### 3.3 Mean square distance

Obviously the expectation value for  $x$  after  $n$  steps  $\langle x_n \rangle = 0$ . What can be derived from:

$$\langle x_n \rangle = \int_{-\infty}^{\infty} x \cdot \rho(x, t) \, dx = 0 \quad (44)$$

But the **Mean Square Distance**  $\langle x_n^2 \rangle$  (MSD) fulfills the following expression:

$$\langle x_n^2 \rangle = \int_{-\infty}^{\infty} x^2 \cdot \rho(x, t) \, dx = 2Dt \quad * \quad (45)$$

## 4 Free particle - MSD

A very simple test is to check if the simulation of a particle without any barriers has a linear growing MSD in time steps. *Figure 3* shows the average squared distance  $x_n^2$  for different numbers of runs compared to the analytic expression. The parameters were:

Parameter	Value
$D$	0.1
$\delta t$	20
$x(n=0)$	0
$n_{\max}$	400

Note that  $\varepsilon$  is determined by *Equation 29*.

The expectation is that for large numbers of runs the mean square converges against the following analytic expression derived from *Equation 44*:

$$\langle x_n^2 \rangle(t) = 2Dt \quad (46)$$

$$\Rightarrow \langle x_n^2 \rangle(n) = 2D\delta t \cdot n \quad (47)$$




Figure 3: MSD over time for different numbers of runs

## 5 Infinite plates

The system which is simulated consists of two infinite plates A and B with distance  $H$ . Between the two plates is some fluid which surrounds the simulated ion. The ion absorbs at the surface of the two plates with the probability  $p$  and desorbs with an exponential decaying probability  $P(t)$  given by:

$$P(t) = \frac{1}{\tau} e^{-t/\tau} \quad (48)$$




Figure 4: Schematic structure of the system

### 5.1 Simple random walk vs. Gaussian random walk

For all relevant simulations a gaussian random walk was used. Even if the distribution becomes equivalent for large numbers of steps the gauss distribution represents a much more natural motion of the particle. Moreover much smaller step sizes what means more steps in the simulation are necessary to allow small motions of the particle. And since performance of the simulation is not an issue the gaussian random walk is just superior the simple random walk. Below figures show example random walks between two plates at  $x = \pm 1$ .




Figure 5: Position of a particle on a simple random walk




Figure 6: Position of a particle on a gaussian random walk

## 5.2 Probability density

The probability density for two infinite plates where the particle can absorb looks for  $t \rightarrow \infty$  is shown in the figure below. The plates are again at  $x = \pm 1$ .






Figure 7: Probability density for two plates

The exception for a reflecting plate would be a constant probability density function. The given probability density function shows some important differences which may due to the implementation of the surface reaction boundary condition.

It was implemented that the particle absorbs when it crosses one plate with the probability  $p$  which was chosen as 0.3 here. If the particle absorbs it keeps absorbed for an exponential distributed time as *Equation 47*. After the time the particle keeps absorbed the langevin simulation continues from the point on the plate. If the first step goes through the plate again the particle absorbs again in the next step.

This directly implies that there will be significant probability that the particle is at the position of a plate what causes the very high values at  $x = \pm 1$ . On the other hand the probability that the particle is next to a plate but not absorbed is significant smaller than that it is somewhere else between the plates. This may due to the implementation of the boundary condition as well because all movements that would crosses a plate end up on the position of the plate here. So ways that would include a reflection process at the boundaries drop out for absorbing plates.

### 5.3 Correlation functions

For a geometry with two plates there is an analytic expression for the Correlation functions  $C_{AA}(t)$  and  $C_{AB}(t)$  which represent the probability for an ion to be absorbed at time  $t$  on plate A/B given that it is absorbed on plate A at time  $t = 0$ .

These correlation functions can easily be obtained from the langevin simulation by noting if and where the particle is absorbed over the time.

Afterwards the values of the correlation function  $C_{AA}(t)$  can be calculated as the number of time intervals with length  $t$  where the ion is absorbed on plate A at the beginning and at the end over the total number of intervals.

Similarly the values of the correlation function  $C_{AB}(t)$  can be calculated as the number of time intervals with length  $t$  where the ion is absorbed on plate A at the beginning and on plate B at the end over the total number of intervals.

The correlation functions were obtained out of 1 simulation. For all correlation functions the values of the correlation function  $C(t)$  for different times  $t$  were calculated as:

$$C_{ij}(t) = \frac{\int_0^{t_{max}-t} S_i(t') \cdot S_j(t' + t) dt'}{\int_0^{t_{max}-t} S_i(t') dt'} \quad (49)$$

Where  $S_{i/j}(t) = 1$  if the particle has the state of interest at the beginning/end of an interval and  $S_{i/j}(t) = 0$  otherwise.

### 5.4 Parameters of the simulation

The run for the comparison was performed with the following parameters:

Parameter	Value
$D$	0.1
$\delta t$	0.1
$\tau$	1.0
$p$	1.0
$H$	1.0

where  $D$  the diffusion constant,  $\delta t$  the step size,  $\tau$  the exponential decay constant for the desorption probability,  $p$  the probability that a particle at the boundary absorbs and  $s$  is the plate distance. The start position for this langevin-simulation is not relevant for the correlation functions.

## 5.5 Interpretation

Also the behavior of the correlation functions in *Figure 8* is easily explainable. Obviously  $C_{AA}(t)$  starts from 1 because after its definition it is definitely absorbed at time  $t = 0$  on Plate *A*.  $C_{AB}(t)$  and  $C_{AD}(t)$  start from 0 because if a particle is absorbed on plate *A* it is neither possible that it is absorbed on plate *B* nor it is desorbed. For  $t \rightarrow \infty$  the absorbed-desorbition-state wont depend on the state at time  $t = 0$  therefore the correlation functions  $C_{AA}(t)$  and  $C_{AB}(t)$  become stationary on some finite probability. Also the relation  $C_{AD}(t) = 1 - C_{AA}(t) - C_{AB}(t)$  is fulfilled what implies that  $C_{AD}(t)$  becomes stationary for  $t \rightarrow \infty$  as well.

## 5.6 Comparison

In order to compare these correlation functions with its analytic equivalents they need to be Laplace transformed. To obtain easy expressions for the Laplace transformed functions they are fitted with the functions of the following type:

$$C(t) = \sum_{i=0}^n c_i \cdot e^{b_i \cdot t} \quad (50)$$

To determine the constants  $c_i$  and  $b_i$  the `scipy.optimize.curve_fit` in Python was used which uses non-linear least squares to fit a function<sup>4</sup>. For  $C_{AA}$  4 terms of the and for  $C_{AB}$  2 terms of *Equation 49* function were used.

---

<sup>4</sup>[1] *Doc: scipy.optimize.curve\_fit* 2014




Figure 8: Correlation Functions and its fits

For the simple exponential expression in *Equation 49* the Laplace transformed can be obtained via:

$$F(t) = c_i \cdot e^{b_i \cdot t} \quad (51)$$

$$\tilde{F}(\omega) = \int_0^\infty c_i \cdot e^{b_i \cdot x} \cdot e^{-\omega \cdot t} dt \quad (52)$$

$$= \frac{c_i}{\omega - b_i} \quad (\text{only for } b_i > \omega) \quad (53)$$

The Laplace transform correlation functions follow as:

$$\tilde{C}(\omega) = \sum_{i=0}^n \frac{c_i}{\omega - b_i} \quad (54)$$

To get an idea of the accuracy of the Laplace transform of the fitted correlation functions also the error of the determined parameters  $c_i$  and  $b_i$  should be considered here.

$$\Delta \tilde{C}(\omega) = \sqrt{\sum_{i=0}^n \left( \frac{\Delta c_i}{c_i} \right)^2 + \left( \frac{\Delta b_i}{(\omega - b_i)^2} \right)^2} \cdot \tilde{C}(\omega) \quad (55)$$

The comparison between that and the analytic result for  $\tilde{C}_{AA}(\omega)$  is shown on the log-log-plot below. The analytic laplace transform correlation function is given by:

$$\Delta\tilde{C}_{AA}(\omega) = 1/\omega \quad (56)$$




Figure 9: Comparison of the laplace transform correlation function  $\tilde{C}_{AA}(\omega)$

The errorbars in the figure above are obtained from the variance of the fitting parameters  $c_i$  and  $b_i$  which are defined for every time  $t$  so there generally has to be an error bar for every time  $t$ . The ones which are considered here, stand only exemplary for these other errorbars. The positions of the errorbars is just chosen to get a nice plot they are not the only data points.

The figure above makes clear that in the uncertainty of the fitting curve the results are identical. Note that for  $\omega \rightarrow b_i$  The fitting function is not properly defined (See *Equation 52*). That means that the divergence at these values may only be caused by the fitting method.

## 6 Finite plates




Figure 10: Schematic structure of the finite plate system

## 7 Conclusion and Outlook

## 8 Annex

Equation 28

$$\rho(x, t) = \frac{1}{\varepsilon \sqrt{2\pi t / \delta t}} e^{-\frac{x^2 \delta t}{2\varepsilon^2 t}} \quad (57)$$

$$\frac{\partial}{\partial t} \rho(x, t) = -\frac{1}{2t} \rho(x, t) + \frac{x^2 \delta t}{\varepsilon^2 t^2} \rho(x, t) \quad (58)$$

$$= \frac{1}{2} \left( \frac{x^2 \delta t}{\varepsilon^2 t^2} - \frac{1}{t} \right) \rho(x, t) \quad (59)$$

$$\frac{\partial^2}{\partial x^2} \rho(x, t) = \frac{\partial}{\partial x} \left( -\frac{x \delta t}{\varepsilon^2 t} \right) \rho(x, t) \quad (60)$$

$$= -\frac{\delta t}{\varepsilon^2 t} \rho(x, t) + \frac{x^2 \delta t^2}{\varepsilon^4 t^2} \rho(x, t) \quad (61)$$

$$= \frac{\delta t}{\varepsilon^2} \left( \frac{x^2 \delta t}{\varepsilon^2 t^2} - \frac{1}{t} \right) \rho(x, t) \quad (62)$$

$$\frac{\partial}{\partial t} \rho(x, t) = D \frac{\partial^2}{\partial x^2} \rho(x, t) \quad (63)$$

$$\Rightarrow \frac{1}{2} = D \cdot \frac{\delta t}{\varepsilon^2} \quad (64)$$

$$\Leftrightarrow D = \frac{\varepsilon^2}{2\delta t} \quad (65)$$



Equation 35

$$p(x, n) = \binom{n}{k} \cdot 2^{-n} \quad \text{with} \quad k = \frac{1}{2} \left( n + \frac{x}{\varepsilon} \right) \quad (66)$$

$$= \frac{n!}{k!(n-k)!} \cdot 2^{-n} \quad (67)$$

$$\text{with } n! \cong \sqrt{2\pi n} n^n e^{-n} \quad (\text{Stirling's formula})$$

$$\Rightarrow p(x, n) \cong \frac{\sqrt{2\pi n} n^n e^{-n}}{\sqrt{2\pi k} k^k e^{-k} \sqrt{2\pi(n-k)} (n-k)^{n-k} e^{k-n}} \cdot 2^{-n} \quad (68)$$

$$= \left( \frac{\sqrt{2\pi n}}{\sqrt{2\pi k} \sqrt{2\pi(n-k)}} \right) \underbrace{\left( \frac{e^{-n}}{e^{-k} e^{k-n}} \right)}_{=1} \left( \frac{n^n}{k^k (n-k)^{n-k}} \right) \cdot 2^{-n} \quad (69)$$

$$= \sqrt{\frac{2n}{\pi(n+x/\varepsilon)(n-x/\varepsilon)}} \cdot \left( \frac{n}{2} \right)^k \cdot \left( \frac{n}{2} \right)^{n-k} \cdot \left( \frac{1}{2} (n+x/\varepsilon) \right)^{-\frac{1}{2}(n+x/\varepsilon)} \cdot \left( \frac{1}{2} (n-x/\varepsilon) \right)^{-\frac{1}{2}(n-x/\varepsilon)} \quad (70)$$

$$= \sqrt{\frac{2n}{\pi(n^2 - (x/\varepsilon)^2)}} \cdot \left( \frac{n}{n+x/\varepsilon} \right)^{-\frac{1}{2}(n+x/\varepsilon)} \cdot \left( \frac{n}{n-x/\varepsilon} \right)^{-\frac{1}{2}(n-x/\varepsilon)} \quad (71)$$

$$= \sqrt{\frac{2n}{\pi(n^2 - (x/\varepsilon)^2)}} \cdot \left( 1 + \frac{x}{\varepsilon n} \right)^{-\frac{1}{2}(n+x/\varepsilon)} \cdot \left( 1 - \frac{x}{\varepsilon n} \right)^{-\frac{1}{2}(n-x/\varepsilon)} \quad (72)$$

$$\text{use } x = e^{\ln(x)} \quad (73)$$

$$\Rightarrow p(x, n) \cong \sqrt{\frac{2}{\pi n \left( 1 - \left( \frac{x/\varepsilon}{n} \right)^2 \right)}} \cdot \exp \left\{ -\frac{n}{2} \left( 1 + \frac{x/\varepsilon}{n} \right) \ln \left( 1 + \frac{x/\varepsilon}{n} \right) - \frac{1}{2} (n - x/\varepsilon) \ln \left( 1 - \frac{x/\varepsilon}{n} \right) \right\} \quad (74)$$

$$\text{for } n \rightarrow \infty \Rightarrow \alpha \rightarrow 0 \quad \alpha := \frac{x/\varepsilon}{n} \quad (75)$$

$$\Rightarrow 1 - \alpha^2 \rightarrow 1 + \dots \quad (76)$$

$$\text{and } \ln(1 + \alpha) \rightarrow \alpha - \frac{1}{2}\alpha^2 + \dots \quad (77)$$

$$\Rightarrow p(x, n) \cong \sqrt{\frac{2}{n\pi}} \cdot \exp \left\{ -\frac{n}{2} \left[ (1 + \alpha) \left( \alpha - \frac{1}{2}\alpha^2 \right) + (1 - \alpha) \left( -\alpha - \frac{1}{2}\alpha^2 \right) \right] \right\} \quad (78)$$

$$= \sqrt{\frac{2}{n\pi}} \cdot \exp \left\{ -\frac{n}{2} \left[ \alpha - \frac{1}{2}\alpha^2 + \alpha^2 - \alpha - \frac{1}{2}\alpha^2 + \alpha^2 + \mathcal{O}(\alpha^3) \right] \right\} \quad (79)$$

$$\cong \sqrt{\frac{2}{n\pi}} \cdot \exp \left\{ -\frac{n}{2} [\alpha^2] \right\} \quad (80)$$

$$= \sqrt{\frac{2}{n\pi}} \cdot e^{-x^2/2\varepsilon^2 n} \quad (81)$$

Equation 42

$$p(x, n+1) = \int_{-\infty}^{\infty} \frac{1}{\varepsilon\sqrt{2n\pi}} e^{-x'^2/2n\varepsilon^2} \cdot \frac{1}{\varepsilon\sqrt{2\pi}} e^{-(x-x')^2/2\varepsilon^2} dx' \quad (82)$$

$$= \frac{1}{\varepsilon^2\sqrt{4\pi^2n}} \int_{-\infty}^{\infty} e^{-\frac{x'^2+x^2n-2xx'n+x'^2n}{2n\varepsilon^2}} dx' \quad (83)$$

$$= \frac{1}{\varepsilon^2\sqrt{4\pi^2n}} \int_{-\infty}^{\infty} e^{-\frac{(n+1)x'^2-2xx'n+x^2n^2/(n+1)+x^2n-x^2n^2/(n+1)}{2n\varepsilon^2}} dx' \quad (84)$$

$$= \frac{1}{\varepsilon^2\sqrt{4\pi^2n}} \int_{-\infty}^{\infty} e^{-\frac{(\sqrt{n+1}x-xn/\sqrt{n+1})^2}{2n\varepsilon^2}} \cdot e^{-\frac{x^2-x^2n/(n+1)}{2\varepsilon^2}} dx' \quad (85)$$

$$\text{let } \alpha = \frac{\sqrt{n+1}x' - xn/\sqrt{n+1}}{\sqrt{2n\varepsilon}} \Rightarrow d\alpha = \sqrt{\frac{n+1}{2n\varepsilon^2}} dx'$$

$$\begin{aligned} p(x, n+1) &= \frac{1}{\varepsilon^2\sqrt{4\pi^2n}} \sqrt{\frac{2n\varepsilon^2}{n+1}} \cdot e^{-\frac{(n+1)x^2-x^2n}{2(n+1)\varepsilon^2}} \int_{-\infty}^{\infty} e^{-\alpha^2} d\alpha \\ &= \frac{1}{\varepsilon\sqrt{2(n+1)\pi}} \cdot e^{-\frac{x^2}{2(n+1)\varepsilon^2}} \end{aligned}$$

Equation 44

$$\langle x_n^2 \rangle = \int_{-\infty}^{\infty} x^2 \cdot \frac{1}{\varepsilon\sqrt{2\pi n}} e^{-x^2/2\varepsilon^2n} dx \quad (86)$$

$$= \frac{1}{\varepsilon\sqrt{2\pi n}} \int_{-\infty}^{\infty} x^2 \cdot e^{-x^2/2\varepsilon^2n} dx \quad (87)$$

$$\text{let } \alpha = \frac{x}{\sqrt{2\varepsilon^2n}} \Rightarrow dx = \sqrt{2\varepsilon^2n} d\alpha \quad (88)$$

$$\langle x_n^2 \rangle = \frac{\sqrt{2\varepsilon^2n}}{\varepsilon\sqrt{2\pi n}} \int_{-\infty}^{\infty} 2\varepsilon^2n\alpha^2 e^{-\alpha^2} d\alpha \quad (89)$$

$$= \frac{2\varepsilon^2n}{\sqrt{\pi}} \int_{-\infty}^{\infty} \alpha^2 e^{-\alpha^2} d\alpha \quad (90)$$

$$= \frac{\varepsilon^2n}{\sqrt{\pi}} \int_{-\infty}^{\infty} \alpha \cdot 2\alpha e^{-\alpha^2} d\alpha \quad (91)$$

$$= \frac{\varepsilon^2n}{\sqrt{\pi}} \left( -\alpha e^{-\alpha^2} \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} 1 \cdot (-e^{-\alpha^2}) d\alpha \right) \quad (92)$$

$$= \frac{\varepsilon^2n}{\sqrt{\pi}} (0 + \sqrt{\pi}) \quad (93)$$

$$= \varepsilon^2n \quad (94)$$

$$= 2D\delta tn \quad (95)$$

$$= 2Dt \quad (96)$$

## References

- [1] *Doc: scipy.optimize.curve\_fit*. 2014. URL: [http://docs.scipy.org/doc/scipy-0.14.0/reference/generated/scipy.optimize.curve\\_fit.html](http://docs.scipy.org/doc/scipy-0.14.0/reference/generated/scipy.optimize.curve_fit.html).
- [2] Roland Netz. “N/A”. In: (2014).
- [3] Roland Netz and Lydric Bocquet. “Conductivity Fluctuations in Cylindrical Pores”. In: (2014).
- [4] “Notes on Brownian Motion”. In: (2011).
- [5] Roland R. Netz Simon Gravelle Laurent Joly and Lydric Bocquet. “Pink noise of ionic current, theory and modelisation”. In: (2014).