

NUEN 301

Homework 1

Donald Doyle UIN: 128005953

November 1, 2020

Exercise 1 [20 pts]: [computing atom densities and macroscopic XS]

Compute the total, elastic, fission, and radiative capture macroscopic cross sections for UO_2 (4 types of macroscopic XS in total).

- Assume UO_2 is only made of U-235, U-238, and O-16. U-235 has been enriched at 25 weight%. Density of UO_2 is 10.4 [g/cc].

- We are interested in computing macroscopic cross sections for both the thermal range (Maximillian average) and the fission spectrum range. Therefore, you will have to procedure 8 macroscopic cross sections in total.

a. [7 pts] Fill the table below:

variable	value	unit
Molar mass of U-235	235.0439282	g/mol
Molar mass of U-238	238.050787	g/mol
Molar mass of O-16	15.99491462	g/mol
Avogadro number	$6.0221409 * 10^{23}$	atoms/mol
(n,total) microscopic XS for U-235 (Maxwellian average)	686.2	b
(n,elastic) microscopic XS for U-235 (Maxwellian average)	16.95	b
(n,fission) microscopic XS for U-235 (Maxwellian average)	571.4	b
(n, γ) microscopic XS for U-235 (Maxwellian average)	97.83	b
(n,total) microscopic XS for U-238 (Maxwellian average)	13.19	b
(n,elastic) microscopic XS for U-238 (Maxwellian average)	10.50	b
(n,fission) microscopic XS for U-238 (Maxwellian average)	$1.683 * 10^{-5}$	b
(n, γ) microscopic XS for U-238 (Maxwellian average)	2.690	b
(n,total) microscopic XS for O-16 (Maxwellian average)	4.474	b
(n,elastic) microscopic XS for O-16 (Maxwellian average)	4.474	b
(n,fission) microscopic XS for O-16 (Maxwellian average)	0	b
(n, γ) microscopic XS for O-16 (Maxwellian average)	$1.9 * 10^{-4}$	b
(n,total) microscopic XS for U-235 (fission spectrum average)	7.628	b
(n,elastic) microscopic XS for U-235 (fission spectrum average)	4.451	b
(n,fission) microscopic XS for U-235 (fission spectrum average)	1.218	b
(n, γ) microscopic XS for U-235 (fission spectrum average)	0.08741	b
(n,total) microscopic XS for U-238 (fission spectrum average)	7.759	b
(n,elastic) microscopic XS for U-238 (fission spectrum average)	4.827	b
(n,fission) microscopic XS for U-238 (fission spectrum average)	0.3064	b
(n, γ) microscopic XS for U-238 (fission spectrum average)	0.07016	b
(n,total) microscopic XS for O-16 (fission spectrum average)	2.764	b
(n,elastic) microscopic XS for O-16 (fission spectrum average)	2.751	b
(n,fission) microscopic XS for O-16 (fission spectrum average)	0	b
(n, γ) microscopic XS for O-16 (fission spectrum average)	$1.136 * 10^{-4}$	b

b. [4 pts] Compute the molar mass of UO_2 .

$$M_{H_2O} = \gamma M_{U-235} + (1 - \gamma) M_{U-238} + 2M_O = .25(235.0439282) + .75(238.050787) + 2(15.99491462) = 269.2889015$$

c. [5 pts] Compute the atom densities for O-16, U-235, U-238.

$$N_{U-235} = \frac{0.25(10.4)(6.0221409 * 10^{23})}{235.0439282} = 6.661549 * 10^{21} [atoms/cm^3]$$

$$N_{U-237} = \frac{0.75(10.4)(6.0221409 * 10^{23})}{238.050787} = 1.973222 * 10^{22} [atoms/cm^3]$$

$$N_{O-16} = \frac{2(10.4)(6.0221409 * 10^{23})}{15.99491462} = 7.831272 * 10^{23} [atoms/cm^3]$$

d. [4 pts] Compute the 8 requested macroscopic cross sections

$$\Sigma_{total} \text{ (Maxwellian average)} = (686.2)(10^{-24})(6.661549 \times 10^21) + (13.19)(10^{-24})(1.973222 \times 10^22) + (4.474)(10^{-24})(7.831272 \times 10^23) = 4.693436359[cm^{-1}]$$

$$\Sigma_{elastic} \text{ (Maxwellian average)} = (16.95)(10^{-24})(6.661549 \times 10^21) + (10.50)(10^{-24})(1.973222 \times 10^22) + (4.474)(10^{-24})(7.831272 \times 10^23) = 3.533098798[cm^{-1}]$$

$$\Sigma_{fission} \text{ (Maxwellian average)} = (571.4)(10^{-24})(6.661549 \times 10^21) + (1.683 \times 10^{-5})(10^{-24})(1.973222 \times 10^22) + (0)(10^{-24})(7.831272 \times 10^23) = 0.990686414[cm^{-1}]$$

$$\Sigma_{\gamma} \text{ (Maxwellian average)} = (97.83)(10^{-24})(6.661549 \times 10^21) + (2.690)(10^{-24})(1.973222 \times 10^22) + (1.9 \times 10^{-4})(10^{-24})(7.831272 \times 10^23) = 0.169765266[cm^{-1}]$$

$$\Sigma_{total} \text{ (fission spectrum average)} = (7.628)(10^{-24})(6.661549 \times 10^21) + (7.759)(10^{-24})(1.973222 \times 10^22) + (2.764)(10^{-24})(7.831272 \times 10^23) = 2.177788915[cm^{-1}]$$

$$\Sigma_{elastic} \text{ (fission spectrum average)} = (4.451)(10^{-24})(6.661549 \times 10^21) + (4.827)(10^{-24})(1.973222 \times 10^22) + (2.751)(10^{-24})(7.831272 \times 10^23) = 2.132100017[cm^{-1}]$$

$$\Sigma_{fission} \text{ (fission spectrum average)} = (1.218)(10^{-24})(6.661549 \times 10^21) + (0.3064)(10^{-24})(1.973222 \times 10^22) + (0)(10^{-24})(7.831272 \times 10^23) = 0.002111754[cm^{-1}]$$

$$\Sigma_{\gamma} \text{ (fission spectrum average)} = (0.08741)(10^{-24})(6.661549 \times 10^21) + (0.07016)(10^{-24})(1.973222 \times 10^22) + (1.136 \times 10^{-4})(10^{-24})(7.831272 \times 10^23) = 2.405136515 \times 10^{-4}[cm^{-1}]$$

Exercise 2 [10 pts]: [on the meaning of microscopic XS]

a. [3 pts] Assume a homogeneous material made of U-235. Use the data collected in Exercise 1. What is the probability that the outcome of collision for an incident neutron of thermal energy (Maxwell average) is a

1. Fission

$$\frac{16.95}{686.2} = 0.0247$$

2. Radiative capture

$$\frac{571.4}{686.2} = 0.8327$$

3. Elastic Scattering

$$\frac{97.83}{686.2} = 0.1426$$

b. [3 pts] Repeat Question a) but assume that the neutron is a fast neutron (fission spectrum average)

$$\frac{4.451}{7.628} = 0.5835$$

$$\frac{1.218}{7.628} = 0.1597$$

$$\frac{0.08741}{7.628} = 0.0115$$

c. [2 pts] Now assume that the homogeneous material is O-16. What is the most probable outcome of a collision on O-16 if the incident neutron is thermal (Maxwell average). What is the probability for that outcome?

The most likely event is elastic scattering $\frac{4.474}{4.474} = 1$

d. [2 pts] Still assuming that the homogeneous material is O-16. What are the two most probable outcome of a collision on O-16 if the incident neutron is fast (fission spectrum average). What are the probabilities for these two outcomes?

The most likely event is elastic scattering $\frac{2.751}{2.764} = 0.9953$

Exercise 3 [20 pts]: [computing atom densities and macroscopic XS]

A boiling water reactor operates at 1000 psi. At that pressure the density of water and of steam are, respectively, 0.74 [g/cm³] and 0.036 [g/cm³]. Use these entries in the database and pick the thermal (Maxwellian) average values.

a. [5 pts] What is the macroscopic total cross section of the water?

$$\Sigma_{water} = \frac{(0.74)(6.0221409 \times 10^{23})}{2(1.00794) + 15.9994} (2(33.15 \times 10^{-24}) + (4.463 \times 10^{-24})) = 1.750442512 [cm^{-1}]$$

b. [5 pts] What is the macroscopic total cross section of the steam?

$$\Sigma_{steam} = \frac{(0.036)(6.0221409 \times 10^{23})}{2(1.00794) + 15.9994} (2(33.15 \times 10^{-24}) + (4.463 \times 10^{-24})) = 0.0851566628 [cm^{-1}]$$

c. [5 pts] If, on average, $40\Sigma_{total} = .4(0.0851566628) + .6(1.750442512) = 1.084328172 [cm^{-1}]$

d. [5 pts] What is the macroscopic total cross section of water under atmospheric con-

ditions at room temperature?

$$\Sigma_{water} = \frac{(1)(6.0221409 \times 10^{23})}{2(1.00794) + 15.9994} (2(33.15 \times 10^{-24}) + (4.463 \times 10^{-24})) = 2.365462854 [cm^{-1}]$$

Exercise 4 [25 pts]: [reaction rates for uncollided beam, thin and thick target meanings]
A beam of neutrons of energy E_1 is normally incident on a homogeneous slab 3 [cm] thick. The area of the slab covered by the beam is 6[cm²]. The intensity of neutrons transmitted through the slab without interactions is found to be 25a. [2 pts] What is the total macroscopic cross section for the slab material at energy E_1 ?

$$I(x) = I_0 e^{-\Sigma_t x} \Rightarrow \Sigma_t = \frac{-\ln(.25)}{3} = 0.462098 [cm^{-1}]$$

b. [2 pts] What is the average distance a neutron travels in this material before undergoing an interaction?

$$\text{Mean free path} = \frac{1}{\Sigma_t} = \frac{1}{0.462098} = 2.16404 [cm]$$

c. [2 pts] At another neutron energy E_2 , the macroscopic cross section is found to be 50 times smaller than in (a). What is the fraction of neutrons of that energy that would be transmitted through the slab without interactions?

$$\Sigma_t = \frac{0.462098}{50} = 0.00924196 [cm^{-1}] \Rightarrow I = I_0 e^{-0.00924196(3)} = 0.9727$$

In what follows, assume the incident beam intensity to be $I_0(E_1) = I_0(E_2) = 10^6 [n/cm^2-s]$

d. [4 pts] Compute the reaction rate in the slab using the thin-target approximation at both energies, E_1 and E_2 .

$$RR_{E_1} = I \Sigma_t = \Sigma I A \Delta x = (10^6)(0.462098)(6)(3) = 8317764 [reactions/s]$$

$$RR_{E_2} = I \Sigma_t = \Sigma I A \Delta x = (10^6)(0.00924196)(6)(3) = 166355 [reactions/s]$$

e. [8 pts] Compute the reaction rate in the slab using the thick-target approximation at both energies, E_1 and E_2 .

$$RR_{E_1} = I A \frac{1 - e^{-\Sigma_t L}}{\Sigma_t} = (10^6)(6) \frac{1 - e^{-(0.462098)(3)}}{(0.462098)} = 9738163 [reactions/s]$$

$$RR_{E_1} = I A \frac{1 - e^{-\Sigma_t L}}{\Sigma_t} = (10^6)(6) \frac{1 - e^{-(0.00924196)(3)}}{(0.00924196)} = 17752757 [reactions/s]$$

f. [2 pts] In which case is the thin target a good approximation? Provide the relative errors, at both energies, E_1 and E_2 .

$$\text{Error}_{E_1} = \frac{|8317764 - 9738163|}{9738163} * 100 = 14.5859\%$$

$$\text{Error}_{E_2} = \frac{|166355 - 17752757|}{17752757} * 100 = 99.0629\%$$

g. [1 pts] To further understand what is going on, write the Taylor Series expansion for the function $f(x) = 1 - \exp(-x)$ around zero.


h. [4 pts] Show that the thick-target expression for the reaction rate becomes the thin-target expression when “something” is small. What is that “something”?

Exercise 5 [25 pts]: [on the thermal motion of the target nucleus and the averaging of XS]

Consider a hypothetical material whose nuclei are moving with a velocity \vec{V} as follows:


- all nuclei have speed $V = \|\vec{V}\| = 10,000[\frac{cm}{s}]$
- at any given time $1/5$ of nuclei are moving in direction $+\hat{i}$
- at any given time $1/5$ of nuclei are moving in direction $-\hat{i}$
- at any given time $3/5$ of nuclei are at rest.

Here $\hat{i}\hat{j}\hat{k}$ are unit vectors along the x, y, and z axes. Suppose the cold cross section for this material is as given in the following table and plot. Recall that the cold cross section is the cross section when there is no thermal motion.



v_{rel} = relative speed	$\sigma_{t,cold}(v_{rel})$
$v_{rel} < 1 \times 10^6$ cm/s	45 b
$1 \times 10^6 \leq v_{rel} \leq 1.01 \times 10^6$ cm/s	90 b
$v_{rel} > 1.01 \times 10^6$ cm/s	35 b

Consider a mono-directional beam of neutrons of intensity $I = n \times v_{tab}$ incident from the left on a thin slice of this material (neutrons are traveling along the x-axis, in direction $+\hat{i}$). This beam has a uniform lab-frame speed distribution, i.e., $0.95106 \leq v_{tab} \leq 1.1106[\frac{cm}{s}]$ Compute the appropriate averaged cross section for each neutron lab-frame speed and plot that average cross section on the same plot as the cold cross section.



Exercise 6 [10 EXTRA pts, MADATORY for Honors]: [Monte Carlo] Implement 2-stream scattering in your Monte Carlo code. Use the following values. $\Sigma_t = 10[cm^{-1}]$, $\Sigma_a = 4[cm^{-1}]$, slab thickness = 0.5 cm. Normally incident beam on the left with intensity = 1000.