

Each question is worth 5 points for a total of 30 points. The Extra Credit problem can get you up to an additional 5 points.

Problem 1

Two workers pull horizontally on a heavy box (this just means they're not lifting it), but one pulls twice as hard as the other. The larger pull is directed at 30.0° west of north, and the resultant of these two pulls is 480.0 N directly northward. Use vector components to find the magnitude of each of these pulls and the direction of the smaller pull.

Problem 2

A 1600-kg blue convertible is traveling south, and a 2100-kg red SUV is traveling west. If the total momentum of the system consisting of the two cars is 8200 kg m/s directed at 60.0° west of south, what is the speed of each vehicle?

Problem 3

A hunter and a 2.5 kg monkey are each 30 m above the ground in trees 80 m apart. Just as the hunter shoots horizontally at the monkey, the monkey drops from the tree in a vain attempt to escape being hit. What must the minimum muzzle velocity of the hunter's bullet be for the hunter to hit the monkey before it reaches the ground?

Problem 4

A 392-N wheel comes off a moving truck and rolls without slipping along a highway. At the bottom of a hill it is rotating at 25.0 rad/s. The radius of the wheel is 0.600 m, and its moment of inertia about its rotation axis is $0.800MR^2$. Friction does work on the wheel as it rolls up the hill to a stop, a height h above the bottom of the hill; this work has absolute value 4000 J. Calculate h .

Problem 5

A mysterious rocket-propelled object of mass 45.0 kg is initially at rest in the middle of the horizontal, frictionless surface of an ice-covered lake. Then a force directed east and with magnitude $F(t) = (120.0 \text{ N/s})t$ is applied. How far does the object travel in the first 5.00 s after the force is applied?

Problem 6

A pump is required to lift 800 kg of water (about 210 gallons) per minute from a well 17.0 m deep and eject it with a speed of 21.0 m/s. (a) How much work is done per minute in lifting the water? (b) How much work is done in giving the water the kinetic energy it has when ejected? (c) What must be the power output of the pump?

Extra Credit:

We usually ignore the kinetic energy of the moving coils of a spring, but let's try to get a reasonable approximation to this. Consider a spring of mass M , equilibrium length L_0 , and spring constant k . The work done to stretch or compress the spring by a distance L is $\frac{1}{2}kX^2$, where $X = L - L_0$. Consider a spring, as described above, that has one end fixed and the other end moving with speed v . Assume that the speed of points along the length of the spring varies linearly with distance l from the fixed end. Assume also that the mass M of the spring is distributed uniformly along the length of the spring. (a) Calculate the kinetic energy of the spring in terms of M and v . (Hint: Divide the spring into pieces of length dl ; find the speed of each piece in terms of l , v , and L ; find the mass of each piece in terms of dl , M , and L ; and integrate from 0 to L . The result is *not* $\frac{1}{2}Mv^2$, since not all of the spring moves with the same speed.) In a spring gun, a spring of mass 0.243 kg and force constant 3200 N/m is compressed 2.50 cm from its unstretched length. When the trigger is pulled, the spring pushes horizontally on a 0.053-kg ball. The work done by friction is negligible. Calculate the ball's speed when the spring reaches its uncompressed length (b) ignoring the mass of the spring and (c) including the mass of the spring (use the results of part (a) to solve this). (d) In part (c), what is the final kinetic energy of the ball and of the spring?