

WORKBOOK EXAMPLES

CHAPTER 4

MATH 1100

Don D. Kim

29 March 2020



Mizzou
University of Missouri

OUTLINE

1 §4.1: POLYNOMIAL FUNCTIONS AND MODELS

OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

Factor polynomial functions, and find the zeros and their multiplicities.

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where the coefficients $a_n, a_{n-1}, \dots, a_1, a_0$ are real numbers and the exponents are whole numbers.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

EXAMPLE


Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

EXAMPLE

Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

FIGURE: $f(x) = x^2 - 2x - 3$ 

EXAMPLE (CONT.)

find the zeros of $f(x)$:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

SOLUTION: $[-4, \infty)$.

EXAMPLE


Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

EXAMPLE

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

FIGURE: $g(x) = x^3 + 2x^2 - 11x - 12$ 

EXAMPLE (CONT.)

find the zeros of $g(x)$:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:





SOLUTION: $(-\infty, \infty)$;

the range:

SOLUTION: $(-\infty, \infty)$.





EXAMPLES OF POLYNOMIAL FUNCTIONS

FIGURE: Polynomial Functions



EXAMPLES OF NONPOLYNOMIAL FUNCTIONS

FIGURE: Nonpolynomial Functions



POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

Furthermore, the domain of a polynomial function is the set of all real numbers.

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function,

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function,
then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$





THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function, then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$ can be described in one of the four following ways.

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function, then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$ can be described in one of the four following ways.

FIGURE:

n	$a_n > 0$	$a_n < 0$
Even		
Odd		

EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

FIGURE:


A.




B.



C.



D.



EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d. $f(x) = -x^6 + x^5 - 4x^3$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d. $f(x) = -x^6 + x^5 - 4x^3$

SOLUTION: **C**

EXAMPLE

Describe the end behavior of the graph of the following functions.

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $f(x) = -x^6 + \frac{3}{4}x^4$

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $f(x) = -x^6 + \frac{3}{4}x^4$

SOLUTION: $\frac{3}{4}x^4$ is the leading term. So $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $f(x) = -x^6 + \frac{3}{4}x^4$

SOLUTION: $\frac{3}{4}x^4$ is the leading term. So $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

(2) $f(x) = 3 - \frac{1}{10}x + x^5$

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $f(x) = -x^6 + \frac{3}{4}x^4$

SOLUTION: $\frac{3}{4}x^4$ is the leading term. So $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

(2) $f(x) = 3 - \frac{1}{10}x + x^5$

SOLUTION: x^5 is the leading term. So $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$, and $f(x) \rightarrow \infty$ as $x \rightarrow \infty$.

EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION: $2x^4$ is the leading term. So $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ and as $x \rightarrow -\infty$.

EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION: $2x^4$ is the leading term. So $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ and as $x \rightarrow -\infty$.

$$(4) f(x) = x^2 - x^3 - 2x + 4.$$

EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION: $2x^4$ is the leading term. So $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ and as $x \rightarrow -\infty$.

$$(4) f(x) = x^2 - x^3 - 2x + 4.$$

SOLUTION: $-x^3$ is the leading term. So $f(x) \rightarrow \infty$ as $x \rightarrow -\infty$, and $f(x) \rightarrow -\infty$ as $x \rightarrow \infty$.

EXAMPLE

Choose the end behavior diagram that best describes the function $f(x) = -3.9x^4 + x^6 + 0.1x^7$.

EXAMPLE

Choose the end behavior diagram that best describes the function $f(x) = -3.9x^4 + x^6 + 0.1x^7$.

FIGURE:

Choose the correct diagram below.

☐ A.



☐ C.



☐ B.



☐ D.



EXAMPLE

Choose the end behavior diagram that best describes the function $f(x) = -3.9x^4 + x^6 + 0.1x^7$.

FIGURE:

Choose the correct diagram below.

☐ A.



☐ B.



☐ C.



☐ D.



SOLUTION: $f(x) \approx x^7 \Rightarrow \text{C.}$

EXAMPLE



Choose the end behavior diagram that best describes the function $f(x) = 5 + \frac{1}{5}x^4 - \frac{1}{2}x^3$.

EXAMPLE

Choose the end behavior diagram that best describes the function $f(x) = 5 + \frac{1}{5}x^4 - \frac{1}{2}x^3$.

FIGURE:

Choose the correct diagram below.





EXAMPLE

Choose the end behavior diagram that best describes the function $f(x) = 5 + \frac{1}{5}x^4 - \frac{1}{2}x^3$.

FIGURE:

Choose the correct diagram below.



SOLUTION: $f(x) \approx x^4 \Rightarrow$ D.

EXAMPLE

Use the leading term test to match the function
 $f(x) = x^5 + \frac{1}{14}x - 3$ with one of the given graphs.


EXAMPLE

Use the leading term test to match the function $f(x) = x^5 + \frac{1}{14}x - 3$ with one of the given graphs.


FIGURE:

Choose the correct graph below.


☐ A.




☐ B.



☐ C.



☐ D.




EXAMPLE

Use the leading term test to match the function $f(x) = x^5 + \frac{1}{14}x - 3$ with one of the given graphs.


FIGURE:

Choose the correct graph below.


☐ A.




☐ B.



☐ C.



☐ D.



SOLUTION: $f(x) \approx x^5 \Rightarrow \text{B.}$

FINDING ZEROS OF POLYNOMIAL FUNCTIONS

If c is a real zero of a function (that is, $f(c) = 0$),

FINDING ZEROS OF POLYNOMIAL FUNCTIONS

If c is a real zero of a function (that is, $f(c) = 0$), then $(c, 0)$ is an x -intercept of the graph of the function.

EVEN AND ODD MULTIPLICITY

If $(x - c)^k$, $k \geq 1$, is a factor of a polynomial function $P(x)$ and $(x - c)^{k+1}$ is not a factor and:

EVEN AND ODD MULTIPLICITY

If $(x - c)^k$, $k \geq 1$, is a factor of a polynomial function $P(x)$ and $(x - c)^{k+1}$ is not a factor and:

(i) k is odd, then the graph crosses the x -axis at $(c, 0)$;

EVEN AND ODD MULTIPLICITY

If $(x - c)^k$, $k \geq 1$, is a factor of a polynomial function $P(x)$ and $(x - c)^{k+1}$ is not a factor and:

- (i) k is odd, then the graph crosses the x -axis at $(c, 0)$;
- (ii) k is even, then the graph is tangent to the x -axis at $(c, 0)$.

EXAMPLE

Use substitution to determine whether 3 is a zero of

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

$$\Rightarrow f(3) = 27 - 90 + 51 + 12 = 0$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

$$\Rightarrow f(3) = 27 - 90 + 51 + 12 = 0$$

Thus 3 is a zero of $f(x)$.

EXAMPLE

Use substitution to determine whether 3 is a zero of

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^2) - 2(3) - 15$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^2) - 2(3) - 15$$

$$\Rightarrow f(3) = -75$$

EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^2) - 2(3) - 15$$

$$\Rightarrow f(3) = -75$$

Thus 3 is *not* a zero of $f(x)$.

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

Rewrite $f(x) = 5(x - 2)^3(x + 1)$.

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

Rewrite $f(x) = 5(x - 2)^3(x + 1)$.

$x = 2$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

Rewrite $f(x) = 5(x - 2)^3(x + 1)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 3.

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

Rewrite $f(x) = 5(x - 2)^3(x + 1)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 3.

$x = -1$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$ and determine the multiplicity of each.

SOLUTION:

Rewrite $f(x) = 5(x - 2)^3(x + 1)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 3.

$x = -1$ is a zero of $f(x)$

with a multiplicity of 1

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

SOLUTION:

$x = 1$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

SOLUTION:

$x = 1$ is a zero of $f(x)$
with a multiplicity of 2.

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

SOLUTION:

$x = 1$ is a zero of $f(x)$

with a multiplicity of 2.

$x = -2$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = -(x - 1)^2(x + 2)^2$ and determine the multiplicity of each.

SOLUTION:

$x = 1$ is a zero of $f(x)$

with a multiplicity of 2.

$x = -2$ is a zero of $f(x)$

with a multiplicity of 2

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 1.

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 3$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 3$ is a zero of $f(x)$

with a multiplicity of 1

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 3$ is a zero of $f(x)$

with a multiplicity of 1

$x = -3$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = x^3 - 2x^2 - 9x + 18$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$.

$x = 2$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 3$ is a zero of $f(x)$

with a multiplicity of 1

$x = -3$ is a zero of $f(x)$

with a multiplicity of 1.

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

$$\text{Factor } f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7).$$

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

with a multiplicity of 1.

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$ is a zero of $f(x)$

with a multiplicity of 1

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$ is a zero of $f(x)$

with a multiplicity of 1

$x = -\sqrt{7}$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = 2x^3 - x^2 - 14x + 7$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$.

$x = \frac{1}{2}$ is a zero of $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$ is a zero of $f(x)$

with a multiplicity of 1

$x = -\sqrt{7}$ is a zero of $f(x)$

with a multiplicity of 1.

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 1)$.

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 1)$.

$x = 1$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 1)$.

$x = 1$ is a zero of $f(x)$

with a multiplicity of 1.

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 1)$.

$x = 1$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 0$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = x^3 - x^2$ and determine the multiplicity of each.

SOLUTION:

Factor $f(x) = x^2(x - 1)$.

$x = 1$ is a zero of $f(x)$

with a multiplicity of 1.

$x = 0$ is a zero of $f(x)$

with a multiplicity of 2

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

SOLUTION:

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

SOLUTION:

$x = -1$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

SOLUTION:

$x = -1$ is a zero of $f(x)$
with a multiplicity of 4.

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

SOLUTION:

$x = -1$ is a zero of $f(x)$

with a multiplicity of 4.

$x = 3$ is a zero of $f(x)$

EXAMPLE

Find the zeros of $f(x) = (x + 1)^4(x - 3)$ and determine the multiplicity of each.

SOLUTION:

$x = -1$ is a zero of $f(x)$

with a multiplicity of 4.

$x = 3$ is a zero of $f(x)$

with a multiplicity of 1

EXAMPLE

Determine whether the following statement is true or false.

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

SOLUTION:

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

SOLUTION:

$P(x)$ has a zero at $x = -5$.

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

SOLUTION:

$P(x)$ has a zero at $x = -5$.

This zero has a multiplicity of 2, an even number.

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

SOLUTION:

$P(x)$ has a zero at $x = -5$.

This zero has a multiplicity of 2, an even number.

Thus $P(x)$ is tangent to the x -axis at $(-5, 0)$

EXAMPLE

Determine whether the following statement is true or false.

“If $P(x) = (x - 7)^7(x + 5)^2$, then the graph of the polynomial function $y = P(x)$ is tangent to the x -axis at $(-5, 0)$. ”

SOLUTION:

$P(x)$ has a zero at $x = -5$.

This zero has a multiplicity of 2, an even number.

Thus $P(x)$ is tangent to the x -axis at $(-5, 0)$

The statement is true (Statement B).

EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

$$f(x) \sim -x^4.$$

EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

$$f(x) \sim -x^4.$$

Choose D.