

# WORKBOOK EXAMPLES

## CHAPTER 4

### MATH 1100

Course Coordinator: Luiza DeSouza

22 March 2020



# OUTLINE

## 1 §4.1: POLYNOMIAL FUNCTIONS AND MODELS

# OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

# OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

Factor polynomial functions, and find the zeros and their multiplicities.

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where the coefficients  $a_n, a_{n-1}, \dots, a_1, a_0$  are real numbers and the exponents are whole numbers.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of  $g(x)$  is 3.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of  $g(x)$  is 3.

The polynomial is a cubic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of  $g(x)$  is 4.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of  $g(x)$  is 4.

The polynomial is a quartic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of  $g(x)$  is 3.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of  $g(x)$  is 3.

The polynomial is a cubic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of  $g(x)$  is 4.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of  $g(x)$  is 4.

The polynomial is a quartic polynomial.

## EXAMPLE


Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

## EXAMPLE

Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

FIGURE:  $f(x) = x^2 - 2x - 3$ 

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:



## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:



## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

SOLUTION:  $[-4, \infty)$ .

## EXAMPLE


Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

## EXAMPLE

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

FIGURE:  $g(x) = x^3 + 2x^2 - 11x - 12$ 

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:



## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:



## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:





SOLUTION:  $(-\infty, \infty)$ ;

the range:

SOLUTION:  $(-\infty, \infty)$ .





# EXAMPLES OF POLYNOMIAL FUNCTIONS

FIGURE: Polynomial Functions



# EXAMPLES OF NONPOLYNOMIAL FUNCTIONS

FIGURE: Nonpolynomial Functions



# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

Furthermore, the domain of a polynomial function is the set of all real numbers.



# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function,

# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function,  
then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$





# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function, then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$  can be described in one of the four following ways.

# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function, then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$  can be described in one of the four following ways.

FIGURE:

$n$	$a_n > 0$	$a_n < 0$
Even		
Odd		

# EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

## EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

FIGURE:


**A.**




**B.**



**C.**



**D.**



## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**



## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d.  $f(x) = -x^6 + x^5 - 4x^3$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d.  $f(x) = -x^6 + x^5 - 4x^3$

SOLUTION: **C**