

# WORKBOOK EXAMPLES

## CHAPTER 4

### MATH 1100

Don D. Kim

6 April 2020



# OUTLINE

## 1 §4.1: POLYNOMIAL FUNCTIONS AND MODELS

# OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

# OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

Factor polynomial functions, and find the zeros and their multiplicities.

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

# POLYNOMIAL FUNCTION

A *polynomial function*  $P$  is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where the coefficients  $a_n, a_{n-1}, \dots, a_1, a_0$  are real numbers and the exponents are whole numbers.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of  $g(x)$  is 3.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of  $g(x)$  is 3.

The polynomial is a cubic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of  $g(x)$  is 4.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of  $g(x)$  is 4.

The polynomial is a quartic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of  $g(x)$  is 3.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of  $g(x)$  is 3.

The polynomial is a cubic polynomial.

# EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is



## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of  $g(x)$  is 4.

## EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of  $g(x)$  is 4.

The polynomial is a quartic polynomial.

## EXAMPLE


Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

## EXAMPLE

Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

FIGURE:  $f(x) = x^2 - 2x - 3$ 

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:



## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:



## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

## EXAMPLE (CONT.)

find the zeros of  $f(x)$ :

SOLUTION:  $x = -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -3$ ;

the minimum:

SOLUTION:  $y = -4$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

SOLUTION:  $[-4, \infty)$ .

## EXAMPLE


Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

## EXAMPLE

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

FIGURE:  $g(x) = x^3 + 2x^2 - 11x - 12$ 

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:



## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:



## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:

SOLUTION:  $(-\infty, \infty)$ ;

the range:

## EXAMPLE (CONT.)

find the zeros of  $g(x)$ :

SOLUTION:  $x = -4, -1, 3$ ;

the  $x$ -intercepts:

SOLUTION:  $x = -4, -1, 3$ ;

the  $y$ -intercept:

SOLUTION:  $y = -12$ ;

the minimum:

SOLUTION:  $-\infty$ ;

the maximum:

SOLUTION:  $\infty$ ;

the domain:





SOLUTION:  $(-\infty, \infty)$ ;

the range:

SOLUTION:  $(-\infty, \infty)$ .





# EXAMPLES OF POLYNOMIAL FUNCTIONS

FIGURE: Polynomial Functions



# EXAMPLES OF NONPOLYNOMIAL FUNCTIONS

FIGURE: Nonpolynomial Functions



# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

# POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

Furthermore, the domain of a polynomial function is the set of all real numbers.



# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function,

# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function,  
then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$





# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function, then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$  can be described in one of the four following ways.

# THE LEADING-TERM TEST

If  $a_n x^n$  is the leading term of a polynomial function, then the behavior of the graph as  $x \rightarrow \infty$  or as  $x \rightarrow -\infty$  can be described in one of the four following ways.

FIGURE:

$n$	$a_n > 0$	$a_n < 0$
Even		
Odd		

# EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

## EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

FIGURE:


**A.**




**B.**



**C.**



**D.**



## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**



## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d.  $f(x) = -x^6 + x^5 - 4x^3$

## EXAMPLE (CONT.)

a.  $f(x) = 3x^4 - 2x^3 + 3$ .

SOLUTION: **D**

b.  $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c.  $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d.  $f(x) = -x^6 + x^5 - 4x^3$

SOLUTION: **C**

# EXAMPLE

Describe the end behavior of the graph of the following functions.

## EXAMPLE

Describe the end behavior of the graph of the following functions.

(1)  $f(x) = -x^6 + \frac{3}{4}x^4$



## EXAMPLE

Describe the end behavior of the graph of the following functions.

(1)  $f(x) = -x^6 + \frac{3}{4}x^4$

SOLUTION:  $-x^6$  is the leading term. So  $f(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$  and as  $x \rightarrow \infty$ .

## EXAMPLE

Describe the end behavior of the graph of the following functions.

$$(1) f(x) = -x^6 + \frac{3}{4}x^4$$

SOLUTION:  $-x^6$  is the leading term. So  $f(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$  and as  $x \rightarrow \infty$ .

$$(2) f(x) = 3 - \frac{1}{10}x + x^5$$

## EXAMPLE

Describe the end behavior of the graph of the following functions.

$$(1) f(x) = -x^6 + \frac{3}{4}x^4$$

SOLUTION:  $-x^6$  is the leading term. So  $f(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$  and as  $x \rightarrow \infty$ .

$$(2) f(x) = 3 - \frac{1}{10}x + x^5$$

SOLUTION:  $x^5$  is the leading term. So  $f(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$ , and  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$ .

## EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

## EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION:  $2x^4$  is the leading term. So  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$  and as  $x \rightarrow -\infty$ .

## EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION:  $2x^4$  is the leading term. So  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$  and as  $x \rightarrow -\infty$ .

$$(4) f(x) = x^2 - x^3 - 2x + 4.$$

## EXAMPLE (CONT.)

$$(3) f(x) = 2x^4 - x^2 + 1$$

SOLUTION:  $2x^4$  is the leading term. So  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$  and as  $x \rightarrow -\infty$ .

$$(4) f(x) = x^2 - x^3 - 2x + 4.$$

SOLUTION:  $-x^3$  is the leading term. So  $f(x) \rightarrow \infty$  as  $x \rightarrow -\infty$ , and  $f(x) \rightarrow -\infty$  as  $x \rightarrow \infty$ .

## EXAMPLE

Choose the end behavior diagram that best describes the function  $f(x) = -3.9x^4 + x^6 + 0.1x^7$ .



## EXAMPLE

Choose the end behavior diagram that best describes the function  $f(x) = -3.9x^4 + x^6 + 0.1x^7$ .

FIGURE:

Choose the correct diagram below.

☐ A.



☐ C.



☐ B.



☐ D.



## EXAMPLE

Choose the end behavior diagram that best describes the function  $f(x) = -3.9x^4 + x^6 + 0.1x^7$ .

FIGURE:

Choose the correct diagram below.

☐ A.



☐ B.



☐ C.



☐ D.



SOLUTION:  $f(x) \approx x^7 \Rightarrow \text{C.}$

## EXAMPLE



Choose the end behavior diagram that best describes the function  $f(x) = 5 + \frac{1}{5}x^4 - \frac{1}{2}x^3$ .

## EXAMPLE

Choose the end behavior diagram that best describes the function  $f(x) = 5 + \frac{1}{5}x^4 - \frac{1}{2}x^3$ .

FIGURE:

Choose the correct diagram below.





## EXAMPLE

Use the leading term test to match the function  
 $f(x) = x^5 + \frac{1}{14}x - 3$  with one of the given graphs.


## EXAMPLE

Use the leading term test to match the function  $f(x) = x^5 + \frac{1}{14}x - 3$  with one of the given graphs.


FIGURE:

Choose the correct graph below.


☐ A.




☐ B.



☐ C.



☐ D.




## EXAMPLE

Use the leading term test to match the function  $f(x) = x^5 + \frac{1}{14}x - 3$  with one of the given graphs.


FIGURE:

Choose the correct graph below.


☐ A.




☐ B.



☐ C.



☐ D.



SOLUTION:  $f(x) \approx x^5 \Rightarrow \text{B.}$



# FINDING ZEROS OF POLYNOMIAL FUNCTIONS

If  $c$  is a real zero of a function (that is,  $f(c) = 0$ ),

# FINDING ZEROS OF POLYNOMIAL FUNCTIONS

If  $c$  is a real zero of a function (that is,  $f(c) = 0$ ), then  $(c, 0)$  is an  $x$ -intercept of the graph of the function.

# EVEN AND ODD MULTIPLICITY

If  $(x - c)^k$ ,  $k \geq 1$ , is a factor of a polynomial function  $P(x)$  and  $(x - c)^{k+1}$  is not a factor and:

# EVEN AND ODD MULTIPLICITY

If  $(x - c)^k$ ,  $k \geq 1$ , is a factor of a polynomial function  $P(x)$  and  $(x - c)^{k+1}$  is not a factor and:

(i)  $k$  is odd, then the graph crosses the  $x$ -axis at  $(c, 0)$ ;

# EVEN AND ODD MULTIPLICITY

If  $(x - c)^k$ ,  $k \geq 1$ , is a factor of a polynomial function  $P(x)$  and  $(x - c)^{k+1}$  is not a factor and:

- (i)  $k$  is odd, then the graph crosses the  $x$ -axis at  $(c, 0)$ ;
- (ii)  $k$  is even, then the graph is tangent to the  $x$ -axis at  $(c, 0)$ .

# EXAMPLE

Use substitution to determine whether 3 is a zero of

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:



## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

$$\Rightarrow f(3) = 27 - 90 + 51 + 12 = 0$$

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(a) f(x) = x^3 - 10x^2 + 17x + 12.$$

SOLUTION:

$$f(3) = 3^3 - 10(3^2) + 17(3) + 12$$

$$\Rightarrow f(3) = 27 - 90 + 51 + 12 = 0$$

Thus 3 is a zero of  $f(x)$ .

# EXAMPLE

Use substitution to determine whether 3 is a zero of

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^3) - 2(3) - 15$$

## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^3) - 2(3) - 15$$

$$\Rightarrow f(3) = -75$$



## EXAMPLE

Use substitution to determine whether 3 is a zero of

$$(b) f(x) = x^4 - 5x^3 - 2x - 15.$$

SOLUTION:

$$f(3) = 3^4 - 5(3^3) - 2(3) - 15$$

$$\Rightarrow f(3) = -75$$

Thus 3 is *not* a zero of  $f(x)$ .

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

Rewrite  $f(x) = 5(x - 2)^3(x + 1)$ .

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

Rewrite  $f(x) = 5(x - 2)^3(x + 1)$ .

$x = 2$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

Rewrite  $f(x) = 5(x - 2)^3(x + 1)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 3.

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

Rewrite  $f(x) = 5(x - 2)^3(x + 1)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 3.

$x = -1$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = 5(x - 2)(x - 2)(x - 2)(x + 1)$  and determine the multiplicity of each.

SOLUTION:

Rewrite  $f(x) = 5(x - 2)^3(x + 1)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 3.

$x = -1$  is a zero of  $f(x)$

with a multiplicity of 1



## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

SOLUTION:

## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

SOLUTION:

$x = 1$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

SOLUTION:

$x = 1$  is a zero of  $f(x)$   
with a multiplicity of 2.

## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

SOLUTION:

$x = 1$  is a zero of  $f(x)$

with a multiplicity of 2.

$x = -2$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = -(x - 1)^2(x + 2)^2$  and determine the multiplicity of each.

SOLUTION:

$x = 1$  is a zero of  $f(x)$

with a multiplicity of 2.

$x = -2$  is a zero of  $f(x)$

with a multiplicity of 2

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:



## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 1.

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = 3$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = 3$  is a zero of  $f(x)$

with a multiplicity of 1

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = 3$  is a zero of  $f(x)$

with a multiplicity of 1

$x = -3$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = x^3 - 2x^2 - 9x + 18$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 2) - 9(x - 2) = (x - 2)(x - 3)(x + 3)$ .

$x = 2$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = 3$  is a zero of  $f(x)$

with a multiplicity of 1

$x = -3$  is a zero of  $f(x)$

with a multiplicity of 1.

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.



## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

$$\text{Factor } f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7).$$

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

with a multiplicity of 1.

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$  is a zero of  $f(x)$

with a multiplicity of 1

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$  is a zero of  $f(x)$

with a multiplicity of 1

$x = -\sqrt{7}$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = 2x^3 - x^2 - 14x + 7$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(2x - 1) - 7(2x - 1) = (2x - 1)(x^2 - 7)$ .

$x = \frac{1}{2}$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = \sqrt{7}$  is a zero of  $f(x)$

with a multiplicity of 1

$x = -\sqrt{7}$  is a zero of  $f(x)$

with a multiplicity of 1.



## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 1)$ .

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 1)$ .

$x = 1$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 1)$ .

$x = 1$  is a zero of  $f(x)$

with a multiplicity of 1.

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 1)$ .

$x = 1$  is a zero of  $f(x)$

with a multiplicity of 1.

$x = 0$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = x^3 - x^2$  and determine the multiplicity of each.

SOLUTION:

Factor  $f(x) = x^2(x - 1)$ .

$x = 1$  is a zero of  $f(x)$   
with a multiplicity of 1.

$x = 0$  is a zero of  $f(x)$   
with a multiplicity of 2

## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.



## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.

SOLUTION:

## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.

SOLUTION:

$x = -1$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.

SOLUTION:

$x = -1$  is a zero of  $f(x)$   
with a multiplicity of 4.

## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.

SOLUTION:

$x = -1$  is a zero of  $f(x)$

with a multiplicity of 4.

$x = 3$  is a zero of  $f(x)$

## EXAMPLE

Find the zeros of  $f(x) = (x + 1)^4(x - 3)$  and determine the multiplicity of each.

SOLUTION:

$x = -1$  is a zero of  $f(x)$

with a multiplicity of 4.

$x = 3$  is a zero of  $f(x)$

with a multiplicity of 1

## EXAMPLE

Determine whether the following statement is true or false.

## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

SOLUTION:



## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

SOLUTION:

$P(x)$  has a zero at  $x = -5$ .

## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

SOLUTION:

$P(x)$  has a zero at  $x = -5$ .

This zero has a multiplicity of 2, an even number.

## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

SOLUTION:

$P(x)$  has a zero at  $x = -5$ .

This zero has a multiplicity of 2, an even number.

Thus  $P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$

## EXAMPLE

Determine whether the following statement is true or false.

“If  $P(x) = (x - 7)^7(x + 5)^2$ , then the graph of the polynomial function  $y = P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$ . ”

SOLUTION:

$P(x)$  has a zero at  $x = -5$ .

This zero has a multiplicity of 2, an even number.

Thus  $P(x)$  is tangent to the  $x$ -axis at  $(-5, 0)$

The statement is true (Statement B).

# EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

## EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

## EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

## EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

$$f(x) \sim -x^4.$$



## EXAMPLE

Find the correct end behavior diagram for the given polynomial function.

$$f(x) = -2.76x^4 - x^3 + x^2 - 2x + 1.$$

SOLUTION:

$$f(x) \sim -x^4.$$

Choose D.