

WORKBOOK EXAMPLES

CHAPTER 4

MATH 1100

Course Coordinator: Luiza DeSouza

22 March 2020



OUTLINE

1 §4.1: POLYNOMIAL FUNCTIONS AND MODELS

OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

OBJECTIVES

Determine the behavior of the graph of a polynomial function using the leading term test.

Factor polynomial functions, and find the zeros and their multiplicities.

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

POLYNOMIAL FUNCTION

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where the coefficients $a_n, a_{n-1}, \dots, a_1, a_0$ are real numbers and the exponents are whole numbers.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

EXAMPLE

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

SOLUTION:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

EXAMPLE


Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

EXAMPLE

Given

$$f(x) = x^2 - 2x - 3 = (x + 1)(x - 3),$$

FIGURE: $f(x) = x^2 - 2x - 3$ 

EXAMPLE (CONT.)

find the zeros of $f(x)$:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the x -intercepts:

SOLUTION: $x = -1, 3$;

the y -intercept:

SOLUTION: $y = -3$;

the minimum:

SOLUTION: $y = -4$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

SOLUTION: $[-4, \infty)$.

EXAMPLE


Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

EXAMPLE

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

FIGURE: $g(x) = x^3 + 2x^2 - 11x - 12$ 

EXAMPLE (CONT.)

find the zeros of $g(x)$:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:

SOLUTION: $(-\infty, \infty)$;

the range:

EXAMPLE (CONT.)

find the zeros of $g(x)$:

SOLUTION: $x = -4, -1, 3$;

the x -intercepts:

SOLUTION: $x = -4, -1, 3$;

the y -intercept:

SOLUTION: $y = -12$;

the minimum:

SOLUTION: $-\infty$;

the maximum:

SOLUTION: ∞ ;

the domain:





SOLUTION: $(-\infty, \infty)$;

the range:

SOLUTION: $(-\infty, \infty)$.





EXAMPLES OF POLYNOMIAL FUNCTIONS

FIGURE: Polynomial Functions



EXAMPLES OF NONPOLYNOMIAL FUNCTIONS

FIGURE: Nonpolynomial Functions



POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

POLYNOMIAL FUNCTIONS

The graph of a polynomial function is *continuous*; that is, it has no holes or breaks.

It is also smooth; there are no "sharp" corners.

Furthermore, the domain of a polynomial function is the set of all real numbers.

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function,

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function,
then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$





THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function, then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$ can be described in one of the four following ways.

THE LEADING-TERM TEST

If $a_n x^n$ is the leading term of a polynomial function, then the behavior of the graph as $x \rightarrow \infty$ or as $x \rightarrow -\infty$ can be described in one of the four following ways.

FIGURE:

n	$a_n > 0$	$a_n < 0$
Even		
Odd		

EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

EXAMPLE


Using the leading term test, match each of the following functions with one of the graphs **A–D** that follow.

FIGURE:


A.




B.



C.



D.



EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3$.

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d. $f(x) = -x^6 + x^5 - 4x^3$

EXAMPLE (CONT.)

a. $f(x) = 3x^4 - 2x^3 + 3.$

SOLUTION: **D**

b. $f(x) = -5x^3 - x^2 + 4x + 2$

SOLUTION: **B**

c. $f(x) = x^5 + \frac{1}{4}x + 1$

SOLUTION: **A**

d. $f(x) = -x^6 + x^5 - 4x^3$

SOLUTION: **C**

EXAMPLE

Describe the end behavior of the graph of the following functions.

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $-x^6 + \frac{3}{4}x^4$

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $-x^6 + \frac{3}{4}x^4$

SOLUTION: The $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $-x^6 + \frac{3}{4}x^4$

SOLUTION: The $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

(2) $3 - \frac{1}{10}x + x^5$

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $-x^6 + \frac{3}{4}x^4$

SOLUTION: The $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

(2) $3 - \frac{1}{10}x + x^5$

SOLUTION: x^5 is the leading term.

EXAMPLE

Describe the end behavior of the graph of the following functions.

(1) $-x^6 + \frac{3}{4}x^4$

SOLUTION: The $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$ and as $x \rightarrow \infty$.

(2) $3 - \frac{1}{10}x + x^5$

SOLUTION: x^5 is the leading term.

So $f(x) \rightarrow -\infty$ as $x \rightarrow -\infty$, and $f(x) \rightarrow \infty$ as $x \rightarrow \infty$.