Workbook Examples
Chapter 4
Math 1100

Don D. Kim

21 March 2020

Outline

Objectives

Determine the behavior of the graph of a polynomial function using the leading term test.

Objectives

Determine the behavior of the graph of a polynomial function using the leading term test.

Factor polynomial functions, and find the zeros and their multiplicities.

Polynomial Function

A *polynomial function* P is given by

Polynomial Function

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

Polynomial Function

A *polynomial function* P is given by

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where the coefficients $a_n, a_{n-1}, \ldots, a_1, a_0$ are real numbers and the exponents are whole numbers.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Solution:

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Solution:

The leading term, i.e. the term with the highest degree, is

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -\frac{1}{6}x^3 - 4x + 8.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$\frac{1}{6}x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Solution:

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Solution:

The leading term, i.e. the term with the highest degree, is

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = 2.4x^4 + 5x^2 - x + \frac{5}{8}.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$2.4x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Solution:

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -3x + 7x^2 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$x^3.$$

The degree of $g(x)$ is 3.

The polynomial is a cubic polynomial.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Solution:

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

Example

Determine the leading term, the leading coefficient, and the degree of the polynomial. Then classify the polynomial as a constant, linear, quadratic, cubic or quartic.

$$g(x) = -4x^2 + 9x^4 + x^3.$$

Solution:

The leading term, i.e. the term with the highest degree, is

$$9x^4.$$

The degree of $g(x)$ is 4.

The polynomial is a quartic polynomial.

Example

Given

$$f(x) = x^2 - 2x - 3 = (x+1)(x-3),$$

Example

Given

$$f(x) = x^2 - 2x - 3 = (x+1)(x-3),$$

Figure: $f(x) = x^2 - 2x - 3$



Example (cont.)

find the zeros of $f(x)$:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

EXAMPLE (CONT.)

find the zeros of $f(x)$:

SOLUTION: $x = -1, 3$;

the $x-$intercepts:

SOLUTION: $x = -1, 3$;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Solution: ∞;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Solution: ∞;

the domain:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

the range:

Example (cont.)

find the zeros of $f(x)$:

Solution: $x = -1, 3$;

the $x-$intercepts:

Solution: $x = -1, 3$;

the $y-$intercept:

Solution: $y = -3$;

the minimum:

Solution: $y = -4$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

the range:

Solution: $[-4, \infty)$.

Example

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

Example

Given

$$g(x) = x^3 + 2x^2 - 11x - 12 = (x + 4)(x + 1)(x - 3),$$

Figure: $g(x) = x^3 + 2x^2 - 11x - 12$



Example (cont.)

find the zeros of $g(x)$:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3;$

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Solution: ∞;

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Solution: ∞;

the domain:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

the range:

Example (cont.)

find the zeros of $g(x)$:

Solution: $x = -4, -1, 3$;

the $x-$intercepts:

Solution: $x = -4, -1, 3$;

the $y-$intercept:

Solution: $y = -12$;

the minimum:

Solution: $-\infty$;

the maximum:

Solution: ∞;

the domain:

Solution: $(-\infty, \infty)$;

the range:

Solution: $(-\infty, \infty)$.

Examples of Polynomial Functions

Figure: Polynomial Functions



Examples of Nonpolynomial Functions

Figure: Nonpolynomial Functions