

Determinants

Brian D. Fitzpatrick

[1, §1.5]

DEF. The (i, j) -submatrix of a $n \times n$ matrix A is the $(n - 1) \times (n - 1)$ matrix obtained by deleting the i th row and j th column of A . We denote the (i, j) -submatrix of A by A_{ij} .

$$\text{Ex. } A = \begin{bmatrix} 1 & 2 & 3 \\ 7 & 0 & 10 \\ 0 & -1 & 6 \end{bmatrix} \implies A_{21} = \begin{bmatrix} 2 & 3 \\ -1 & 6 \end{bmatrix}$$

DEF. The determinant of a 1×1 matrix $A = [a_{11}]$ is $\det(A) = a_{11}$.

DEF. The determinant of a $n \times n$ matrix A is $\det(A) = \sum_{k=1}^n (-1)^{1+k} a_{1k} \det(A_{1k})$

NOTE. We often write $\det(A) = |A|$.

$$\text{Ex. Compute } \det \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

SOL.

$$\begin{aligned} \det(A) &= \sum_{k=1}^2 (-1)^{1+k} a_{1k} \det(A_{1k}) \\ &= (-1)^{1+1} a_{11} \det(A_{11}) + (-1)^{1+2} a_{12} \det(A_{12}) \\ &= a \det([d]) - b \det([c]) \\ &= ad - bc \end{aligned}$$

Ex. Compute $\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$.

SOL.

$$\begin{aligned} \det(A) &= \sum_{k=1}^3 (-1)^{1+k} a_{1k} \det(A_{1k}) \\ &= (-1)^{1+1} a_{11} \det(A_{11}) + (-1)^{1+2} a_{12} \det(A_{12}) + (-1)^{1+3} a_{13} \det(A_{13}) \\ &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \end{aligned}$$

DEF. The (i, j) -minor of A is $M_{ij} = \det(A_{ij})$. The (i, j) -cofactor of A is $C_{ij} = (-1)^{i+j}M_{ij}$.

NOTE. We this notation, our formula for determinants is

$$\det(A) = \sum_{k=1}^n a_{1k}C_{1k}$$

Ex. For $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ we have

$$\begin{aligned} C_{11} &= (-1)^{1+1} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} \\ C_{12} &= (-1)^{1+2} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} = - \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} \\ C_{32} &= (-1)^{3+2} \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} = - \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \end{aligned}$$

NOTE. The signs of the cofactors can be remembered by

$$\begin{array}{ccccc} + & - & + & - & \cdots \\ - & + & - & + & \cdots \\ + & - & + & - & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{array}$$

THM (Laplace Cofactor Expansion Theorem, Theorem 1.16 in [1]).

For any $1 \leq i \leq n$ we have

$$\det(A) = \sum_{k=1}^n a_{ik} C_{ik} \quad (\textit{ith row expansion})$$

$$\det(A) = \sum_{k=1}^n a_{ki} C_{ki} \quad (\textit{ith column expansion})$$

IDEA. The Laplace Cofactor Expansion Theorem allows us to compute determinants by “expanding” about any row or column. This is nice because some rows or columns may be easier to expand about than others.

Ex. Compute $\begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix}.$

SOL. Expansion about Row₂ gives

$$\begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} = (-4) \begin{vmatrix} 2 & 3 \\ 8 & 9 \end{vmatrix} + (5) \begin{vmatrix} 1 & 3 \\ 7 & 9 \end{vmatrix} + (-6) \begin{vmatrix} 1 & 2 \\ 7 & 8 \end{vmatrix} = 0$$

Expansion about Col₃ gives

$$\begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} = (3) \begin{vmatrix} 4 & 5 \\ 7 & 8 \end{vmatrix} + (-6) \begin{vmatrix} 1 & 2 \\ 7 & 8 \end{vmatrix} + (9) \begin{vmatrix} 1 & 2 \\ 4 & 5 \end{vmatrix} = 0$$

Ex. Compute
$$\begin{vmatrix} 7 & -3 & 0 & 4 \\ 0 & 1 & 0 & 3 \\ 2 & 1 & -2 & -5 \\ 0 & 4 & 0 & 6 \end{vmatrix}.$$

SOL. Expansion about Col_3 gives

$$\begin{vmatrix} 7 & -3 & 0 & 4 \\ 0 & 1 & 0 & 3 \\ 2 & 1 & -2 & -5 \\ 0 & 4 & 0 & 6 \end{vmatrix} = (-2) \begin{vmatrix} 7 & -3 & 4 \\ 0 & 1 & 3 \\ 0 & 4 & 6 \end{vmatrix}$$

Then expand about Col_1 to obtain

$$\begin{vmatrix} 7 & -3 & 4 \\ 0 & 1 & 3 \\ 0 & 4 & 6 \end{vmatrix} = (7) \begin{vmatrix} 1 & 3 \\ 4 & 6 \end{vmatrix} = 7(6 - 12) = -42$$

Hence

$$\begin{vmatrix} 7 & -3 & 0 & 4 \\ 0 & 1 & 0 & 3 \\ 2 & 1 & -2 & -5 \\ 0 & 4 & 0 & 6 \end{vmatrix} = (-2)(-42) = 84$$

THM (Corollary 1.17 in [1]). *If A has a row or column of zeros, then $\det(A) = 0$.*

THM (Corollary 1.18 in [1]). *The determinant of a triangular matrix is the product of its diagonal entries.*

THM (Theorem 1.19 in [1]). $\det(A^\top) = \det(A)$

THM (Theorem 1.20 in [1]).

1. $A \xrightarrow{R_i \leftrightarrow R_j} B \implies \det(B) = -\det(A)$
2. $A \xrightarrow{\lambda R_i \rightarrow R_i} B \implies \det(B) = \lambda \det(A)$
3. $A \xrightarrow{R_i + \lambda R_j \rightarrow R_i} B \implies \det(B) = \det(A)$

IDEA. Determinants play nicely with elementary row operations.

REFERENCES

- [1] G.L. Peterson and J.S. Sochacki. Linear Algebra and Differential Equations. Addison-Wesley, 2002.