

A
B
CD4T
CD8T
Mono
NK
Micro
Oligo
cis-pQTL → protein → AD
Protein-to-AD MR
Top 50%
Bellenguez
FinnGen
Gene / protein identity
Both outcomes
BIN1
CR1
PILRB
GRN
BLNK
Union only
CD55
ACP5
ST13
NESP55
BST2
B
Network top 50%
Bellenguez MR
FinnGen MR
Outcome direction
Platform direction
Bellenguez coloc
FinnGen coloc
DrugCLIP top 20
Docking pose QC
Met
Not met
Unresolved
All 9 met
PILRA
LRRC25
CTSH
CD33
SIGLEC9
LILRB5
DHRS4L2
SULT1A1
SH3BP1
TREM1
C
Entity correspondence
ACP5 | AGES
ACP5 | UKB
BIN1 | AGES
BLNK | UKB
BST2 | UKB
CD55 | AGES
CD55 | UKB
CR1 | UKB
GRN | AGES
GRN | UKB
NESP55 | UKB
PILRB | UKB
ST13 | UKB
O00499
O95467
P08174
P13686
P17927
P28799
P50502
Q10589
Q8WV28
Q9UKJ0
Model human members
Protein assay
Structure UniProt
D
Assay q
Bellenguez log(OR), 95% CI
E
FinnGen log(OR), 95% CI
F
rs6733839
−log10(P)
BIN1 region: chromosome 2 (Mb)
G
rs5848
GRN region: chromosome 17 (Mb)
H
BIN1
GRN
Colocalization PP(H4)
Shared−association prior p12
Bellenguez
FinnGen
I
Mean evidence percentile
53.7
q = 0.0075
46.0
Model
Blood AD difference
Top−decile targets (%)
11.5%
2.08−fold
5.6%
J
GRN | aged Grn−KO microglia
GPNMB
CTSD
CTSB
LAMP1
CST7
APOE
TYROBP
Grn−KO response (log expression)
BIN1 | microglial Bin1cKO
CX3CR1
APOE
TREM2
TYROBP
CST3
CTSD
IFITM3
Bin1cKO response (log expression)