



Microglia Sing the Prelude of Neuroinflammation-Associated Depression

Qingqing Li¹ · Ying Xie¹ · Jinyi Lin¹ · Miaomiao Li¹ · Ziyang Gu¹ · Tianli Xin¹ · Yang Zhang¹ · Qixia Lu¹ · Yihui Guo¹ · Yanhong Xing¹ · Wuyang Wang¹

Received: 14 June 2023 / Accepted: 21 October 2024

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2024

Abstract

Major depressive disorder (MDD) is a psychiatric condition characterized by sadness and anhedonia and is closely linked to chronic low-grade neuroinflammation, which is primarily induced by microglia. Nonetheless, the mechanisms by which microglia elicit depressive symptoms remain uncertain. This review focuses on the mechanism linking microglia and depression encompassing the breakdown of the blood–brain barrier, the hypothalamic–pituitary–adrenal axis, the gut–brain axis, the vagus and sympathetic nervous systems, and the susceptibility influenced by epigenetic modifications on microglia. These pathways may lead to the alterations of microglia in cytokine levels, as well as increased oxidative stress. Simultaneously, many antidepressant treatments can alter the immune phenotype of microglia, while anti-inflammatory treatments can also have antidepressant effects. This framework linking microglia, neuroinflammation, and depression could serve as a reference for targeting microglia to treat depression.

Keywords Major depressive disorder · Microglia · Neuroinflammation · Immunity

Abbreviations

5-HT	5-Hydroxy tryptamine	CCL9	C-C motif chemokine ligand 9
ACC	Anterior cingulate cortex	CCR10	C-C motif chemokine receptor 10
Ach	Acetylcholine	CD	Cluster of differentiation
ACTH	Adrenocorticotrophic hormone	CNS	Central nervous system
AG	Adrenal gland	COX2	Cyclooxygenase-2
Akt1/Akt2	AKT serine/threonine kinase 1/2	CPA	Conditioned place aversion
Amy	Amygdala	CRH	Corticotropin-releasing hormone
Arg1	Arginase-1	CRP	C-reactive protein
BBB	Blood-brain barrier	CSDS	Chronic social defeat stress
BDNF	Brain-derived neurotrophic factor	CSF	Cerebrospinal fluid
cAMP	Cyclic adenosine monophosphate	CUMS	Chronic unpredictable mild stress
CCL2	C-C motif chemokine ligand 2	CUS	Chronic unpredictable stress
CCL5	C-C motif chemokine ligand 5	CX1CR1	C-X-C chemokine receptor 1
		CX3CR1	C-X3-C chemokine receptor 1
		DBS	Deep brain stimulation
		DHEA	Dehydroepiandrosterone
		DNMT	DNA methyltransferase
		DNMT3B	DNA methyltransferase 3 beta
		IL-1R1	IL-1 receptor type I
		IL-6	Interleukin-6
		iNOS	Inducible nitric oxide synthase
		JMJD3	Jumonji domain protein 3
		JSE	Juvenile social exploration
		KYN	Kynurenine
		LAG3	Lymphocyte-activating gene-3

✉ Yanhong Xing
yanhong646@163.com

✉ Wuyang Wang
wuyangwang80@gmail.com

¹ Jiangsu Province Key Laboratory of Anesthesiology, Jiangsu Province Key Laboratory of Anesthesia and Analgesia Application Technology, NMPA Key Laboratory for Research and Evaluation of Narcotic and Psychotropic Drugs, Xuzhou Medical University, 209 Tongshan Rd, Xuzhou 221004, Jiangsu, China

LC	Locus coeruleus
LDT	Light/dark box test
Lgals3	Advanced glycation end-product receptor 3
LHb	Lateral habenula
LPS	Lipopolysaccharide
MAOI	Monoamine oxidase inhibitor
MBT	Marble burying test
MDD	Major depressive disorder
MHC	Major histocompatibility complex
mPFC	Medial prefrontal cortex
NE	Norepinephrine
NF- κ B	Nuclear factor-kappa B
NLRP3	NOD-, LRR-, and pyrin domain-containing protein 3
NO	Nitrogen monoxide
Nrf2	Nuclear factor erythroid 2-related factor 2
NSAIDs	Nonsteroidal anti-inflammatory drugs
OFT	Open field test
P2RY12	Purinergic receptor P2Y12
PET	Positron emission computed tomography
PFC	Prefrontal cortex
PGE2	Prostaglandin E2
PKA	Protein kinase A
PKB	Protein kinase B
PVN	Paraventricular nucleus of the hypothalamus

Introduction

Major depressive disorder (MDD) is a complicated condition that manifests as a combination of various signs and symptoms, mainly characterized by sadness and anhedonia. The impact of depression on individuals can lead to significant health loss [1, 2]. According to the World Health Organization, depression will be the primary cause of the worldwide disease burden by 2030 [3]. Due to the diverse symptoms and pathophysiology of the condition, various biological mechanisms are involved in its etiology. These mechanisms include dysregulation of the hypothalamic–pituitary–adrenal (HPA) axis, maladjustment of monoamine neurotransmitters, abnormal neuroplasticity and neurogenesis, and inflammatory imbalance [4]. Previous studies on depression have primarily focused on neuronal changes in the brains of both depressed patients and animal models. As a result, many antidepressants have been developed to modulate the level of 5-HT, dopamine, norepinephrine, and their receptors. However, despite following correct medical advice, approximately two-thirds of diagnosed patients still fail to see improvement or even experience relapse [4].

Microglia, which are the primary immune cells of the central nervous system (CNS), have a crucial role in monitoring and modulating the immune system as neurons undergo maturation, damage, aging, etc. [5, 6]. Accumulating evidence

suggests that chronic low-grade neuroinflammation significantly contributes to the pathogenesis of mental health disorders [7, 8]. Chronic low-grade neuroinflammation is also described as “para-inflammation”, which means the adaptive response that is intermediate between basal and inflammatory states. Para-inflammation represents an adaptive response intermediate between basal and inflammatory states, with graded levels ranging from near-basal to more inflammatory responses triggered by infection or tissue damage. This adaptive process helps tissues adapt to adverse conditions and promotes functional tissue restoration [7, 8]. Under normal conditions, short-term inflammation acts as a guardian to prevent the spread of pathogenic microorganisms, remove necrotic tissue, and promote tissue repair [9]. However, prolonged tissue dysfunction can perpetuate low-grade inflammation, potentially leading to a chronic state where inflammation can be the real culprit in triggering mood disorders [10]. As the major locus of neuroinflammation response, microglia are possibly involved in modulating neurobiochemistry, neuroendocrine, and behavioral alterations associated with depressive symptoms [11, 12]. Clinical evidence also demonstrates microglial dynamics in the brains of patients with depression, supporting the idea that depression may be a microglial disease [13]. Nevertheless, the underlying mechanism concerning the relationship between microglia-induced neuroinflammation and MDD remains uncertain, and the understanding of the intersection of immunology and neuroscience remains pending, which is the subject of a current review.

The Immunomodulation of Microglia with Depression

Immune Disorders in MDD

Factors such as environment, genetics, stress, and infection can affect individual immunity to varying degrees, and eventually trigger the remodeling of the inflammatory phenotype in MDD patients, manifested by the elevated inflammation scores in both pro-inflammatory and anti-inflammatory cytokines. The pro-inflammatory markers interleukin (IL)-6 (IL-6), tumor necrosis factor (TNF)- α (TNF- α), IL-1 β , and C-reaction protein (CRP, a biomarker of systemic inflammation) are reliable biomarkers of immune disorders occurring in MDD [14–16]. A clinical study of 89 MDD patients showed that cerebrospinal fluid (CSF) IL-6, TNF- α , and IL-1 β were associated with high plasma CRP (> 3 mg/L), as well as the severity of depressive symptoms [17]. It has been found that peripheral monocytes in MDD were less responsive to lipopolysaccharide (LPS) [18]. Similarly, MDD plasma also has an immunosuppressive effect on peripheral blood mononuclear cells from healthy donors, resulting in

reduced activation of monocyte/dendritic cells and B cells and reduced T cell memory [19]. The aforementioned evidence strongly indicates the potential existence of immune tolerance among individuals experiencing depression. Furthermore, high levels of cytokines alone can also lead to depressive symptoms. Striking examples are two reports of depression in 45% of patients with malignant melanoma and 35% of patients with hepatitis C virus after treatment with high doses of the pro-inflammatory cytokine interferon (IFN)- α (IFN- α) [20, 21]. These human data are consistent with the literature in many experimental animals, such as the LPS-induced depression model, suggesting that cytokines can induce a range of behavioral changes that overlap with those observed in depression, including anhedonia, cognitive dysfunction, and reduced activity [22–24].

In addition to elevated cytokines in peripheral blood and CSF, there is also direct evidence of inflammation in depressed patients' brains by imaging. Translocator protein (TSPO) is highly expressed in activated microglia during neuroinflammation, making it a biomarker of brain inflammation [25], which also supports the neuroinflammatory hypothesis of depression [26, 27]. However, using TSPO binding as an indicator of activated microglia is still controversial. Using the positron emission computed tomography (PET) ligand [^{18}F], FEPPA found that the TSPO distribution volume in MDD was increased in the anterior cingulate cortex (ACC), prefrontal cortex (PFC), and insula [28]. Using the ligand [^{11}C]PK-11195 exhibited that there is a raised TSPO distribution volume in the ACC of MDD and a substantial increase in depression with suicidal tendencies [29]. Although there is no direct evidence that TSPO represents pro-inflammatory or anti-inflammatory microglia activation, several animal and human studies have shown that elevated TSPO correlates with behavioral manifestations of upward pro-inflammatory cytokines in depression. Nonetheless, several PET analyses have examined the microglial status of postmortem whole-brain samples in individuals with depression, yielding negative results, which is likely because depression is predominantly linked to microglial activation or reduction in particular brain regions, rendering the overall status of microglia in the entire brain sample unreliable [13]. In conclusion, because of limited ethical and inadequate post-mortem brain samples, as well as many confounding variables including medication, age, and severity, the TSPO results should be taken with care. Thus, either central or peripheral immune system dysregulation can mediate depressive-like symptoms.

Microglia in CNS

Microglia are resident innate immune cells that serve as the first line of immune janitors, constituting 5 to 10% of the CNS population [30]. They are derived from immature yolk

sac progenitor cells during early embryogenesis and persist throughout life [31]. In adults, regeneration occurs solely on residual microglia without the presence of de novo progenitors [32]. As the neurons' inseparable partners, microglia are involved in adjusting neuronal activity through various mechanisms [33], covering neurogenesis, synaptic and neuronal plasticity [34], phagocytosis, antigen-presentation, cytokines release, metabolic support, neurotransmitter release and uptake, extracellular matrix reconstruction [35], and signal feedback [36]. According to the morphology, function, and specific position, microglia can be classified into the following forms: ramified microglia, amoeboid microglia, hypertrophic microglia, rod microglia, dystrophic microglia, and satellite microglia [37]. Under normal circumstances, microglia are in a "resting" state, with a ramified morphology. In response to tissue injury or pathogen invasion, ramified microglia proliferate and transform into activated "brain macrophages," also known as "reactive microglia," or hypertrophic microglia, eventually obtaining an amoeboid shape with enlarged somas and sprout, an excess of short and thick protrusions [37, 38].

Referencing the dichotomy of macrophage activation, microglia are classified into M1 and M2 types based on distinct stimulatory factors, although their transcriptional adaptations differ from those of peripheral macrophages [39]. M1 phenotype is induced by LPS and/or IFN- γ stimulation, which activates toll-like receptor 4 (TLR4) or IFN- γ receptors 1 and 2, respectively, followed by activation of transcription factors the nuclear factor-kappa B (NF- κ B) and signal transducer and activator of transcription (STAT) 1 and raised levels of CD86 and MHC II, at which point M1 phenotype markers include CD16, CD32, CD86, inducible nitric oxide synthase (iNOS), and MHCII. M1 secrete pro-inflammatory cytokines (IL-1 α , IL-1 β , TNF- α , IL-6, IL-12, IL-23), chemokines (CCL2, CCL9, CCR10), cyclooxygenase-2 (COX2), reactive oxygen species (ROS), etc. M2 phenotype is induced by IL-4/IL-13 stimulation, in which IL-4 binding to its receptor triggers activation of STAT6, which lets microglia shift toward anti-inflammatory phenotype with increased expression of arginase-1 (Arg1), CD206, mannose receptor, and release of anti-inflammatory factors (IL-4, IL-10, IL-13, IL-1RA) [40]. During the progression of the disease, activated microglia can switch from M2 to M1 phenotype. A transient neuroinflammatory response is protective during stress and tissue damage under normal conditions, as M1 microglia release chemokines to move rapidly and purposefully to the site of damage, while M2 microglia release trophic factors such as brain-derived neurotrophic factor (BDNF) to promote neuronal microenvironmental homeostasis. In abnormal contexts, microglia become dysfunctional (e.g., suppressed microglia function, reduced numbers, senescence) and are unable to "put out the fire" after inflammation, leading to a constant and chronic

CNS inflammatory state, causing neuronal excitotoxicity, excessive synapse pruning, demyelination, and death, which disrupts mood and cognition and causes various behaviors associated with depression [41].

Two-photon microscopy techniques have directly revealed the activity state of microglia that always performs to have highly motile protrusions, even in the absence of conventional stimulus signals [22]. Since there is no absolute concept of the “resting” state of microglia, the classification of these cells into M1 and M2 subtypes is no longer applicable. Theoretically, the distinction between M1 and M2 serves to facilitate the description of microglial inflammatory features and to investigate the mechanisms of microglial dual regulation, and microglia are always in kinetic equilibrium to accomplish their mission due to their high susceptibility to the fluctuating microenvironment [42]. Hence, microglia are morphologically and phenotypically distinct at various phases of life and in various brain regions [43]. For example, the aging brain contains a specific subset of dysfunctional lipid-droplet-accumulating microglia that exhibit pro-inflammatory effects and weak phagocytosis [44].

Immunological Alterations of Microglia in Different Brain Regions

The brain regions linked to depression can be broadly categorized as the PFC, which governs cognition, and the limbic system, which controls emotional responses [45–47]. The medial prefrontal cortex (mPFC) has been linked to affective experiences such as anxiety, guilt, and humiliation in individuals with MDD [48]. In the depression model, there is an elevation in pro-inflammatory markers on microglia, commonly observed cytokines comprise IL-1 α , TNF- α , IL-1 β , and IL-6, while anti-inflammatory markers such as Arg1, IL-4, and IL-10 experience a decrease [49–54]. Exposure to an inflammatory environment has been observed to lead to a decrease in BDNF, as well as a reduction in neuronal response, dendritic atrophy, synapse loss, and an imbalance in local excitatory and inhibitory transmission. These changes can ultimately contribute to the development of depressive symptoms [49–54]. The ACC is linked to the cognitive respects of emotions, while activated microglia in the ACC are known to have a significant impact on mood disorders [55]. The occurrence of an inflammatory response in the ACC during the early stages of childhood boosts the likelihood of an individual developing depression in adulthood. This is because early inflammation triggers excessive phagocytosis of the glutamatergic neuronal spines by microglia, subsequently enduring maladaptation of neurons to stress, thereby facilitating the emergence of symptoms resembling depression during adolescence [22].

The limbic system is accountable for the regulation of emotional responses and is linked to states of physiological

and psychological activity, such as excitement, fear, anxiety, and emotional memories [56, 57]. Specifically, the amygdala dominates anxiety, fear, and other negative emotions. In models of depression induced by acute or chronic stress, microglia located in the amygdala demonstrate an activated state characterized by an increase in numbers, enlarged cytosomes, and a reduction in total protrusion length, with inflammation-related markers including CD14, CD86, and TLR4 on their surface [58, 59]. The depressive symptoms may be attributed to the dysregulation of synaptic and non-synaptic transmission through inflammation in the amygdala [58, 59]. The hippocampus preserves emotional memory. The activation of microglia in the hippocampus shows a notable increase of several pro-inflammatory markers, including NLRP3 inflammasome, IL-1 β , IL-6, and IL-18, which have been observed to impact neurogenesis in the hippocampus within a depression model [60]. The striatum has been linked to impulsivity, fatigue, and negative affect in the context of depression [58, 61]. The induction of aversion in mice can be achieved through the targeted activation of striatal microglia. This process relies on the critical involvement of IL-6 signaling and COX-1-mediated prostaglandin synthesis within the microglia [62]. Studies have indicated a correlation between depression and activated insula, showing atypical somatic sensitivity. Furthermore, it has been observed that individuals suffering from depression exhibit a reduction in connectivity between the insula and ACC, which is concomitant with an increase in microglia activation [63]. The lateral habenula (LHb) has been connected to feelings of helplessness, anhedonia, and excessive concern with negative stimuli [64]. The activation of microglia in the LHb can be also found in repetitive social defeat stress and LPS-induced depression models [65] (Table 1).

Microglia Link Bidirectional Immune Communication in Depression

Activated microglia are considered the manifestation of neuroinflammation, the pivotal convergence of various stimulation-induced neuroinflammatory responses, and an important source of pro-inflammatory factors and oxidative stress [73]. There are multiple pathways to activate microglia.

BBB

As a highly selective and dynamic interface between blood and brain tissue with monitoring brain immune dynamics [74], the blood–brain barrier (BBB) is a territorial boundary between the brain and inflammatory signals in the circulating blood [75] and is mainly composed of vascular endothelial cells with tight junctions [76]. The dramatically increased microglia and macrophage recruitment observed in the dorsal ACC of depressed suicidal individuals may be linked

Table 1 Immunological alterations of microglia in different depressed brain regions

Brain region	Model of depression	Depressive-like behavior	State of microglia	Alteration relating to neuron	Alteration relating to immunity	Reference
ACC	Early-life inflammation induced by LPS	Anhedonia (SPT); Despair (FST, TST)	IF: Iba1↑ (number↑, soma size↓, total process length↓, branch points↓), CD68↑, MHCII↑; Protein: TLR4↑	LPS increases the extent of microglial engulfment around neuronal spines in adolescence	Protein: p-NF-κB p65↑, CX3CR1↑; mRNA and protein: IL-1β↑, IL-6↑, TNF-α↑	[22]
ACC	Chronic LPS stress	Anhedonia (SPT); Anxiety (OFT, EPM); Despair (FST, TST); Body weight loss	IF: Iba1↑ (number↑, soma size↓, total process length↓, branch points↓); mRNA: Iba1↑, CD68↑	Dendritic spines↓	mRNA: CX3CR1↑, caspase4↑, CCL5↑, TLR1↑, TLR9↑, TLR13↑	[66]
ACC	MDD with suicidal ideation	Suicidal ideation	TSPO: volume↑	-	-	[29]
Amy	LPS stress	Anhedonia (SPT); Anxiety (EPM, OFT); Despair (FST, TST)	IF: Iba-1 (number↑, soma size↓, total process length↓)	Presynaptic glutamate↑; Intrinsic neuronal excitability↑	ELISA: IL-1β↑, TNF-α↑; Protein: IL-1β↑, TNF-α↑, TNF-α receptor↑	[59]
Amy PFC	Social disruption stress	Anxiety (LDT)	IH/IF: Iba-1 (number↑); Flow cytometry: CD14↑, CD86↑, TLR4↑	-	mRNA: IL-1β↑, GC responsive genes↓; ELISA: IL-6↑	[58]
Hip	Chronic mild stress	Anxiety (EPM, OFT); Despair (FST)	IF: Iba-1 (number↑), TSPO↑	-	Protein: NLRP3 inflammatory↑; some↑; mRNA: IL-1β↑, IL-6↑, IL-18↑, IL-4↑, IL-10↑	[67]
Hip	Chronic social defeat	Aberrant socialization (SIT); Anxiety (LDT)	IF: Iba-1 (number↑)	-	ROS activity↑	[68]
Hip	Chronic social defeat stress	Social avoidance (SAT); Body weight loss	IH: Iba-1 (number↑)	Neurogenesis↓	IF: NLRP3↑, CX1CR1↓; ELISA: IL-6↑	[69]
Hip	Chronic jetlag with a chronic phase advance model	Anhedonia (SPT); Anxiety (OFT, EPM, JSE); Despair (FST); Sleep disturbance	mRNA: CD68↑	BDNF↓	ELISA: GC↑; mRNA: IL-1β↓, Arg1↑	[70]
Hip	An experimental model of shift-work	Anhedonia (SPT); Anxiety (OFT); Hypoactivity (24-h activity)	IH: Iba-1 (number↑)	-	-	[71]
Hip	LPS stress	Anxiety (OFT); Despair (FST)	IH: Iba-1 (number↑); mRNA: Iba1↑	-	mRNA: IL-1β↑, IL-6↑, TNF-α↑	[72]
LH	Repeated social defeat stress/LPS stress	Anhedonia (SPT); Despair (FST, TST); Social avoidance (SIT)	IF: Iba1↑ (soma size↑, total process length↓)	Neuronal excitability↑	mRNA: TNF-α↑, IL-1β↑, IL-6↑; Protein: iNOS↑	[65]
mPFC	IL-10 knockout mice	Anhedonia (SPT); Anxiety (MBT); Despair (FST, TST)	IF: Iba-1 (number↑)	Protein: GAD67↓, parvalbumin↓;	Protein: IL-1β↑, IL-6↑, IL-10↓	[51]
mPFC	Chronic mild stress	Anhedonia (SPT); Despair (FST); Anxiety (OFT)	IF: Iba-1 (number↑), CD68↑	Synapse loss↑; Dendritic spine density↓; Neuronal activity↓	ELISA: TNF-α↑, IL-1β↑	[49]
PFC	Chronic stress	Anhedonia (SPT); Anxiety (FST, OFT); Body weight loss	IH: Iba-1 (number↑)	5-HT _{2A} ↑	mRNA: TNF-α↑, COX-1↑	[53]

Table 1 (continued)

Brain region	Model of depression	Depressive-like behavior	State of microglia	Alteration relating to neuron	Alteration relating to immunity	Reference
mPFC	Chronic social defeat stress	Anhedonia (SPT); Despair (FST); Social avoidance (SIT)	IF: Iba-1 (number↑), Arg1↓, TREM2; Protein: Arg1↓, TREM2↓	Protein: BDNF↓, PSD95↓	mRNA: IL-4↓, IL-10↓; IF: Nr2f2↓; Protein: Nr2f2↓	[50]
mPFC	Repeated social defeat stress	Anxiety (EPM); Social avoidance (SIT)	IF: CD68↑, Iba-1 (number↑), CD11b↑; mRNA: CD11b↑, TLR2↑, TLR4↑	Neuronal response attenuation; Dendritic atrophy	mRNA: IL-1α↑, TNF-α↑, IL-1R1↑	[52]
mPFC	Repeated social defeat stress	Anxiety (EPM); Social avoidance (SIT, submissive posture)	IF: Iba-1 (number↑)	Attenuation of the mesocortical dopaminergic pathway	ELISA: corticosterone↑; ELISA: PGE2↑	[54]
PFC ACC Insula	MDD episodes	Anhedonia; Anorexia; Low mood; Body weight loss	TSPO: volume↑	-	-	[28]
Striatum	LPS stress/ Chemogenetic activation	Aversion (CPA); Anhedonia (SPT); Anxiety (OFT); Despair (FST)	IF: Iba-1 (number↑)	Excitability of striatal medium spiny neurons↓	mRNA: IL-1↑, IL-6↑, TNF-α↑, CD14↑	[62]

ACC, anterior cingulate cortex; Amy, amygdala; Arg1, arginase-1; BDNF, brain-derived neurotrophic factor; CPA, conditioned place aversion; CX1CR1, C-X-C chemokine receptor 1; ELISA, enzyme-linked immunosorbent assay; EPM, elevated plus maze; FST, forced swimming test; GAD67, glutamate decarboxylase 67; GC, glucocorticoid; Hip, hippocampus; IL-1R1, IL-1 receptor type 1; iNOS, inducible nitric oxide synthase; IF, immunofluorescence staining; IH, immunohistochemistry staining; JSE, juvenile social exploration; LDT, light/dark box test; LH, lateral habenula; MBT, marble burying test; mPFC, medial prefrontal cortex; NLRP3, NOD-, LRR- and pyrin domain-containing protein 3; Nr2f2, nuclear factor erythroid-2-related factor 2; OFT, open field test; PFC, prefrontal cortex; SAT, social avoidance test; SPT, sucrose preference test; SIT, social interaction test; TREM2, triggering receptor expressed on myeloid cells-2; TSPO, translocator protein; TST, tail suspension test

to BBB disruption [77]. When the BBB compromised, periphery immune cells and cytokines entered the brain parenchyma and modulated the inflammatory response from within [78], producing much more inflammatory cytokines through mutual action with microglia and astrocytes during the process [79], and elevated levels of inflammatory cytokines such as TNF- α and IL-1 β in the brain able to cause prolonged mouse depression-like behavior [80–82]. Changes in the BBB's tight junctions allow the entry of monocytes/macrophages and circulating inflammatory cytokines into the brain, while endothelial cell dysfunction increases the adherence of leukocytes and yields conditions that induce inflammation [83]. Meanwhile, endothelial cells can transmit peripheral inflammatory signals directly to astrocytes and microglia, while activated microglia can recruit more monocytes for transport to the brain by releasing CCL2 [84], without BBB destruction [85]. For instance, under conditions of chronic stress, microglia can selectively increase CCL2 expression, thereby attracting monocytes that subsequently express IL-1 β to bind to IL-1R1 in brain endothelial cells [86]. Systemic inflammation can also induce microglia migration to the vasculature via the chemokine CCL5, while early inflammation promotes microglia expression of the tight junction protein cln5 to preserve BBB integrity and physical contact with endothelial cells, which may refer to the P2RY12 receptor of microglia [87]. However, persistent inflammation leads to enhanced microglia phagocytosis of astrocyte end-foot and destructed BBB function, instead of limiting microglia activation [88].

Neuroimmune Interactions

Microglia are in a unique position as immune sensors of the stress response [89, 90], able to rapidly sense and respond to changes, which in turn adjust the CNS to respond to acute or chronic stress [91]. The stress involved in the mechanism of depression acts mainly through the HPA axis, which is consistent with the clinical finding of abnormalities in the HPA axis in depressed patients who manifested by elevated plasma cortisol levels. Two hypotheses could explain the paradox of both high GC levels and high levels of pro-inflammatory factors in depressed patients: GC resistance and GC synergy with pro-inflammatory cytokines. In the model of depression induced by LPS, GC combined with its receptor (GR) to inhibit the activity of NF- κ B in the early phase of the NF- κ B activity, but had no inhibitory effect in the late phase. The lack of inhibitory effect could be explained by the GC resistance which is probably mediated by selective accumulation of GR β protein [92]. Meanwhile, in the late phase of NF- κ B activity when depression-like behavior was evident, microglia in the infralimbic PFC significantly increased, emphasizing the key role of microglia immunity in the process of acute stress [93]. In addition, the

time effect of GC application modulates the cellular state to produce anti-inflammatory, pro-inflammatory, or combined effects. Short-term pre-exposure to GC inhibits pro-inflammatory responses, whereas long-term pre-exposure to GC induces increased NLRP3 expression, sensitizes microglia to extracellular ATP signaling molecules, and increases the release of ATP-mediated pro-inflammatory cytokines such as IL-1 β , TNF- α , and IL-6 [94, 95], or increases microglia proliferation through NMDA receptor activation [96]. Both the *in vivo* and *ex vivo* studies suggest that GC treatment sensitizes the TNF- α and IL-1 β response to LPS, and upregulates the expression of microglia activation markers [97] and receptors including MHCII, TLR2, and TLR4 in the hippocampal microglia [98]. The bidirectional regulation of inflammation by GC complicates its role in depression, and addressing the question of how to reconcile the time-dependency response of GC may help to normalize the immune system characteristics by modulating abnormally elevated GC in response to stress, thereby ameliorating MDD. The other way to regulate neuroimmune interactions is through sympathetic and parasympathetic systems. Chronic stress also affects peripheral immune cells by activating the sympathetic nervous system. Under stressful conditions, sympathetic nerves bias hematopoietic stem cells toward differentiation into GC-resistant myeloid immune cells, which can then be recruited to the CNS by chemokines secreted by microglia to participate in the inflammatory response [99]. Stress can also induce microglia activation via β -adrenergic receptors [100]. The vagus nerve suppresses inflammation by activating the HPA axis [101] and binding to the α 7 nicotinic acetylcholine receptor (α 7nAChR) of microglia through the cholinergic anti-inflammatory pathway [102]. The activation of α 7nAChR increases Ser3-phosphorylation of GSK9, which in turn activates PKB, and GSK3 β , inhibiting NF- κ B transcriptional activity and exerting anti-inflammatory effects [103].

Gut-Brain Axis

In the absence of a microbiome, microglia in the male embryonic stage showed substantial downregulation of gene expression associated with metabolic processes, cell and protein organization, and adaptive immune responses, while microglia in adult female mice exhibited dysregulation of genes associated with regulation of transcription, adaptive immune responses, cell migration, and chemotaxis, suggesting that microglia respond to the absence of a microbiome at the prenatal stage in a sex and time specific manner [104], and from this perspective, the increased baseline microglia activation status in adult females may partially explain why they show greater susceptibility to depression than adult males [105]. Microbiomes generate a variety of neuromodulators (e.g., catecholamine, aminobutyric acid,

5-HT), immune modulators (e.g., quinolinic acid (QUIN)), and short-chain fatty acids (SCFAs) (e.g., acetate, propionate, butyrate) that cross the BBB and exhibit immunomodulatory effects on microglia [106, 107]. Among them, SCFAs facilitate the maturation of microglia and possess properties that mitigate inflammation [108], resulting in a decrease in the levels of pro-inflammatory IL-1 β , CCL2, TNF- α , and an increase in the levels of anti-inflammatory TGF- β and IL-4 [109, 110]. Studies on adult mouse models of depression have found that depression-like behavior can be improved in mice by remodeling the microbiota [111], and its antidepressant effects are possibly related to promoting microglia to an anti-inflammatory phenotype [112]. However, it remains to be determined whether the effect of the microbiome on the immune system can ameliorate MDD. A meta-analysis of the use of probiotics in alleviating depressive symptoms showed that probiotic supplementation had an overall insignificant effect on mood, possibly limited by the discrepancies concerning probiotic dosing, bacterial strains, and strain combinations [113]. The antidepressant effect of minocycline, a microglia inhibitor, is associated with a reduction in the number of microglia in the PFC and changes in microbial composition and metabolites [114]. However, changes in the microbial composition may also be due to the direct inhibition of intestinal inflammation by minocycline as an antibiotic, rather than modulation by direct inhibition of microglia activation. Similarly, mice showed abnormal gut microbiota composition and SCFAs after repeated administration of colony-stimulating factor 1 receptor inhibitor PLX5622 to clear microglia [115]. These results mirror each other and suggest there is a tight immunomodulation between microglia and microbiota. In addition to interacting with microglia during development and adulthood, intestinal flora also triggers microglia mitochondrial dysfunction by increasing the microglial accumulation of N⁶-carboxymethyllysine in the aging brain's microglia [116]. In addition, there is an interaction between enteroendocrine cells and vagal afferent neuron function. Vagal afferent neurons expressing TLR4 can directly sense bacterial products such as LPS [117], and convey changes in intestinal inflammatory status to microglia, thereby inhibiting microglia proliferation and expression of pro-inflammatory cytokines [118], and restoring microglia to a resting state [119] (Fig. 1).

Microglia in Antidepressant Treatment

The Immunomodulatory Effects of Antidepressants

In the 1950s, it was discovered that iproniazid, an anti-tuberculosis drug, could increase 5-HT, dopamine, and norepinephrine levels in the brain by inhibiting monoamine oxidase, thereby improving mood, increasing appetite, making

sleep better, becoming energetic, and increasing social willingness, thus establishing the classic monoamine hypothesis of depression [120, 121]. Classical antidepressants modify neurotransmitters and directly eliminate the symptoms of “low monoamine neurotransmitter levels” in the brain, which seems to bypass the underlying cause of depression [122]. A major argument against the monoamine hypothesis is that although these antidepressants rapidly elevate inter-synaptic monoamine neurotransmitter concentrations, the antidepressants require 2–4 weeks to take effect, leading to the speculation that there may be downstream signaling molecules may be responsible for the therapeutic effects of antidepressants [123]. In light of pharmacogenomic approaches, most of the antidepressants' effects can be divided into two categories: reinforcing the process of neuroplasticity and neurotropy, and accommodating the inflammation, immunity, and redox reactions [124]. Numerous *in vivo* and *in vitro* studies on the effects of antidepressants on microglia have revealed that different classes of antidepressants have significant effects on alleviating depressive behaviors and the inflammatory environment of the brain, as well as on the immune and oxidative stress signaling of microglia [125]. Tricyclic antidepressants are thought to work through G_s-coupled microglial receptor activation, increase intracellular cAMP levels, and activate the cAMP-dependent protein kinase A (PKA), which blocks GSK3 enzyme function [103], while the blocked GSK3 β inhibits the subsequent transcriptional activity of NF- κ B, reducing the production of pro-inflammatory cytokines including IL-1 β , TNF- α , and IL-6 [126]. In particular, imipramine also prevents stress-related anxiety and depressive-like behaviors by preventing the accumulation of macrophages in the brain [127]. Selective serotonin reuptake inhibitors (SSRIs), similar to tricyclic antidepressants (TCA), demonstrate anti-inflammatory properties via PKA inhibition in microglia [128]. For instance, the administration of fluoxetine, paroxetine, sertraline, and citalopram has been observed to impede the production of nitrogen monoxide (NO) and TNF- α by microglia, as well as diminish the release of glutamate. These effects have been shown to facilitate the survival of neurons [129–131]. Tranylcypromine, a monoamine oxidase inhibitor, has been observed to hamper the synthesis of pro-inflammatory cytokines, including but not limited to IL-1 β , and IL-6 [132]. Additional classes of antidepressants can also regulate the inflammatory responses of microglia. Melatonin relieves LPS-induced depression by blocking the microglial NLRP3 inflammasome [133]. Ketamine, a blocker of NMDA receptors at glutamate receptors in the brain, can rapidly improve depressive symptoms and suicidal ideation within hours after a single dose, and its antidepressant action may be mediated by microglial production of QUIN [128, 134], the BDNF pathway [135], and TGF- β 1 [136].




Fig. 1 Microglia link bidirectional immune communication via multiple direct and indirect pathways in depression. (1) Impairment of the BBB leads to the infiltration of peripheral monocytes/macrophages and circulating inflammatory cytokines into the brain, resulting in the activation of microglia. The increased activity of IDO in microglia drives tryptophan metabolism toward the kynurenine pathway, which produces the neurotoxic QUIN and reduces the synthesis of 5-HT. Simultaneously, the dysfunction of endothelial cells results in heightened leukocyte adhesion, thereby creating an environment that triggers inflammation. Furthermore, the activation of microglia can attract a greater number of monocytes for transportation to the brain through the discharge of chemokines. (2) Microglia act as immune sensors of stress responses primarily through the HPA axis. The binding of GC to its receptors can result in anti-inflammatory, pro-inflammatory, or combined effects, which are dependent upon the duration of GC exposure. (3) An alternative approach to regulating neuroimmune interactions involves the sympathetic and parasympathetic nervous systems. The activation of microglia can be induced by sympathetic through the β -adrenergic receptors. (4) The inhibition of inflammation is facilitated by the binding of the vagus to the $\alpha 7$ nAChR of microglia through the cholinergic anti-inflammatory pathways. (5) The microbiota generates a diverse range of SCFAs, which traverse the BBB and demonstrate immunomodulatory properties on microglia. Additionally, they contribute to the maturation of microglia. (6) The activation of microglial PKA and PKB to block GSK3 β , which subsequently inhibits the transcriptional activity of

NF- κ B and ultimately reduces oxidation and inflammation, may account for the antidepressant effect of multiple drugs or pathways. (7) The oxidation and inflammation afterward stimulate additional pro-inflammatory cytokines, chemokines, ROS, and RNS. The cascade of events ultimately causes elevated levels of glutamate, QUIN, and GC, as well as decreased levels of 5-HT and BDNF, culminating in depressive symptoms. (8) Moreover, the susceptibility to depression is closely linked with epigenetic alterations in microglia. The aforementioned pathways represent three distinct modes of bidirectional immune communication between the peripheral and central systems. These include the humoral pathway, which is linked to the BBB, the neural pathway, which is associated with the vagus and sympathetic nerves, and the cellular pathway, which is connected to microglia. BDNF, brain-derived neurotrophic factor; TRP, tryptophan; IDO, indoleamine 2,3-dioxygenase; KYN, kynurenine; QUIN, quinolinic acid; LC, locus coeruleus; PVN, paraventricular nucleus of the hypothalamus; CRH, corticotropin-releasing hormone; ACTH, adrenocorticotrophic hormone; AG, adrenal gland; GC, glucocorticoid; GR, glucocorticoid receptor; β AR, β -adrenergic receptor; NE, norepinephrine; $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; ACh, acetylcholine; SCFAs, short-chain fatty acids; PKA, protein kinase A; PKB, protein kinase B; GSK3 β , glycogen synthase kinase 3 β ; NF- κ B, nuclear factor-kappa B; TNF- α , tumor necrosis factor; IL, interleukin; CCL2, C-C motif chemokine ligand 2; ROS, reactive oxygen species; RNS, reactive nitrogen species; 5-HT, 5-hydroxy tryptamine

The Antidepressant Effects of Anti-inflammatory Drugs

Numerous clinical studies find that inflammation-related diseases have a higher incidence of depression, such as chronic hepatitis [137] and autoimmune diseases [138]. An increase in inflammatory biomarkers may predict a lack of response

to antidepressants, one reason for which may be the presence of a defect in the anti-inflammatory response, which may also be a reason why traditional antidepressants do not work [19]. Therefore, anti-inflammatory treatments may be beneficial for depressed patients with increased immune baselines. Given the wide range of biological functions involved in inflammatory pathways, several clinical studies suggested

the possible clinical antidepressant benefit of targeting specific inflammatory cytokines [139]. A clinical trial targeting seven mechanisms of action (TNF- α , IL-12/23, IL-6, CD20, COX2, BL γ S, and P38/MAPK14) in depressed patients, who are diagnosed with one of nine inflammation-related diseases (ulcerative colitis, rheumatoid arthritis, psoriasis, asthma, ankylosing spondylitis, multicentric Castleman disease, osteoarthritis-knee, systemic lupus erythematosus, neuropathic pain), and the conclusion was that anti-inflammatory drugs significantly improved two core symptoms of depression: anhedonia and depressed mood, and these effects were not exclusively attributable to the therapeutic effect on physical health [140]. A systematic review and meta-analysis of clinical trials of chronic inflammatory conditions revealed that monoclonal antibodies against the IL-6 receptor and TNF- α ameliorated depressive symptoms in a statistically significant manner [139]. Despite there being controversial results [141], ineffective results obtained by targeting only a single cytokine and ignoring the complexity of inflammatory factor types and functions are not sufficient to negate the effectiveness of anti-inflammatory treatment, since cytokines do not act independently, but rather cause MDD synergistically. Although there is a wealth of clinical data suggesting that the treatment of depression with nonsteroidal anti-inflammatory drugs (NSAIDs) alone or in addition may have side effects such as gastrointestinal bleeding, studying the mechanisms of its antidepressant effects may be useful in elucidating the etiology of depression. The antibiotic minocycline is often tested as an antidepressant in animals [142, 143], and its antidepressant effects have been confirmed in clinical trials as an adjunctive therapy [144–146], and even in treatment-resistant depression [147–149]. Minocycline treats depression mainly through its immunomodulatory effects, including inhibition of microglia signaling in microglia such as MAPK, the release of cytokines such as TNF- α and IL-6 [150], and reduction of IDO activity, and impact on QUIN production. QUIN can interact with Fe²⁺ to increase ROS and RNS, limit SOD activity and GSH levels [151], impact 5-HT synthesis [152], and activate the HPA axis [153].

Microglia in Non-pharmacological Antidepressant Treatment

Electro-acupuncture treatment exhibited antidepressant effects associated with microglia, significantly reversed chronic unpredictable stress (CUS)-induced increases in IL-1 β , IL-18, TNF- α , and IL-6 expression, and alleviated neuroinflammation in the hippocampus [154, 155], while microglia-derived LAG3 may be involved in it [156]. In addition, repetitive transcranial magnetic stimulation functions as an antidepressant by switching the microglial phenotype from M1 to M2 without reducing the number of

microglia in the hippocampal and cortical regions [157]. Running exercise reduces the amount and activity of hippocampus microglia, which improves depressive-like behaviors caused by CUS [158]. Some articles suggest that the activation of microglia has antidepressant effects. For example, before starting the deep brain stimulation (DBS) treatment, the operation of electrode implantation already has an antidepressant effect by activated microglia, and the effect can be blocked by treatment with NSAIDs [159]. However, DBS is often used to treat treatment-resistant depression, where the microglia in the brain may have become senescent or even reduced in number due to the severity of the illness or the prolonged inflammation in the brain, so that external stimulation may alleviate depression by activating microglia or stimulating microglia proliferation [38]. Treating immune system dysregulation aids in the recovery of MDD, and both activating and inhibiting microglia have antidepressant effects, possibly because of differences in microglia status in MDD (Table 2).

Genetic and Epigenetic Regulations of Microglia in Depression

Neuroimmune and neuroendocrine responses elicited by prolonged stress are diverse due to individual differences, and early-life stress is associated with overt inflammation before the development of neuropsychiatric diseases, which presumably leads to behavioral vulnerability and resilience [164, 165]. Most people can adapt to the blow of stress by increasing their resilience, but some will collapse under similar circumstances and ultimately get depressed [166]. Whether the source of individual differences in depression susceptibility is related to gene and epigenetics alterations has been a hot issue in related research. In MDD, the most replicated and relevant genetic variants include polymorphisms in the genes for IL-1 β , IL-6, IL-10, CCL2, TNF- α , and CRP [167], and the best replication risk alleles for depression have pro-inflammatory or protective effects against pathogens [168]. Nevertheless, these genes do not show a substantial contribution to the occurrence of depression [169, 170].

In recent years, a growing body of research has demonstrated that distinct stress elicits unique epigenetic modifications in both brain neurons and the immune system. Notably, alterations in microglial epigenetics have been identified as a significant contributor to the preservation of brain health [171]. Under conditions of early-life stress, a significant reduction in DNA methylation occurs in microglia, which is linked to altered microglial activity [171]. The process of DNA demethylation increases the transcription of IL-1 β in microglia and regulates type I IFN responses induced by TLR-4 [172]. The *in vivo*

Table 2 Alterations of microglia in different types of antidepressant therapies

Treatment	Subject	Effect on microglia	Pro-inflammation	Anti-inflammation	Reference
Fluoxetine/norfluoxetine (SSRI)	Primary microglia with LPS	Inhibition	Gas chromatography: NO↓, TNF↓	-	[130]
Fluoxetine (SSRI)	BV2 with LPS	Inhibition	ELISA: TNF-α↓, IL-6↓; Griess reagent: NO↓; mRNA: TNF-α↓, IL-6↓, iNOS↓; Protein: iNOS↓; NF-κB↓	-	[160]
Vortioxetine (SSRI)	Rats with LPS	Inhibition	ELISA: IL-1β↓, IL-6↓, TNF-α↓	ELISA: IL-4↑	[161]
Escitalopram (SSRI)				-	
Amitriptyline (TCA)					
Tranylcypromine (MAOI)					
Paroxetine (SSRI)	BV2/primary microglia with LPS	Inhibition	ELISA: TNF-α↓, IL-1β↓; mRNA: TNF-α↓, IL-1β↓; Measurement of nitrite: NO↓; Protein: iNOS↓, TNF-α↓, IL-1β↓	-	[131]
Sertraline (SSRI)	Mice with CUMS	Inhibition	ELISA: NO↓, IL-1β↓, TNF-α↓; mRNA: Iba1↓; Protein: IL-1β↓, TNF-α↓, iNOS↓	-	[129]
Tranylcypromine (MAOI)	BV2 with LPS	Inhibition	mRNA: IL-1β↓, TNF-α↓, iNOS↓	-	[132]
Melatonin	Mice with LPS	Inhibition	IF: Iba-1↑; Protein: pro-IL-1β↓, IL-1β↓, NLRP3↓, cleaved caspase-1↓	-	[133]
Ketamine	Mice with LPS	Inhibition	Luminex assays: IL-1α↓, IL-6↓, Eotaxin↓, G-CSF↓, MIP1b↓, TNF-α↓; mRNA: Ptg2↓	mRNA: Lgals3↓, IL-1RA↓	[134]
(R)-ketamine	Mice with CSDS	Inhibition	mRNA: Tgfb1↑, Tgfb1↑, Tgfb2↑	-	[136]
Lithium	HM/primary microglia/hiPSC-derived microglia with IFN-γ/LPS	Inhibition	-	Multi-Spot Assay System: IL-10↑	[162]
taVNS	Rats with CUMS	Inhibition	Protein: IL-1β↓, NF-κB p65↓, p-NF-κB p65↓	Protein: α7nAChR↑	[163]
Running exercise	Rats with CUS	Inhibition	ELISA: IL-1β↓, iNOS↓, IL-6↓, TNF-α↓; mRNA: IL-1β↓, iNOS↓, IL-6↓, TNF-α↓	-	[158]
Imipramine (TCA)	Mice with RSD	Inhibition	ELISA: IL-6↓; mRNA: IL-1β↓, IL-6↓, TNF-α↓, TLR-4↓, CCL2↓, CX3CR1↓	-	[127]
EA	Rats with CUS	Inhibition	mRNA: IL-1β↓, IL-18↓, TNF-α↓; Protein: IL-1β, pro-IL-1β↓, NLRP3 inflammasome↓	-	[155]
ECS	Mice with CUS/BV2 with Poly I:C	Inhibition	Protein: TNF-α↓	-	[156]
rTMS	Mice with CUMS	Inhibition	ELISA: IL-6↓, IL-1β↓, TNF-α↓; mRNA: IL-6↓, IL-1β↓, TNF-α↓	-	[157]

α7nAChR, $\alpha 7$ nicotinic acetylcholine receptor; *CCL2*, C–C motif chemokine ligand 2; *CUMS*, chronic unpredictable mild stress; *CUS*, chronic unpredictable stress; *CSDS*, chronic social defeat stress; *CX3CR1*, C-X3-C chemokine receptor 1; *EA*, electro-acupuncture; *ECS*, electroconvulsive stimulation; *ELISA*, enzyme-linked immunosorbent assay; *hiPSC*, human-induced pluripotent stem cells; *HM*, immortalized human microglia; *IL*, interleukin; *iNOS*, inducible nitric oxide synthase; *Lgals3*, advanced glycation end-product receptor 3; *LPS*, lipopolysaccharide; *MAOI*, monoamine oxidase inhibitor; *NO*, nitrogen monoxide; *RSD*, repeated social defeat; *rTMS*, repetitive transcranial magnetic stimulation; *SSRIs*, selective serotonin reuptake inhibitors; *taVNS*, transcutaneous auricular vagus nerve stimulation; *TCA*, tricyclic antidepressants

experiments have also confirmed that the use of DNA demethylation drugs promotes LPS-induced pro-inflammatory responses in microglia [173]. Apart from DNA methylation, histone post-translational modifications also have a regulatory function in the immune function of microglia. The histone methyltransferase EZH2 has been implicated in susceptibility to neuroinflammation and depressive-like behaviors. Its upregulation of H3K27me3 exacerbates depression and neuroinflammation by inducing microglia M1-type polarization [174, 175]. While the utilization of the EZH2 inhibitor EPZ-6438 or methylation-reducing medications can impede the activation of M1 microglia, diminish the levels of pro-inflammatory cytokines, and augment the levels of anti-inflammatory cytokines in the brain. This, in turn, can alleviate depression-like behaviors and spatial memory impairment [176, 177]. The activation of microglia and expression of pro-inflammatory cytokines induced by LPS can be reduced by the SCFAs acetate and butyrate through increasing histone acetylation [110]. The deacetylase SIRT1 has been observed to induce a shift in microglial polarization toward the M2 phenotype through a decrease in the expression of IL-6, IL-1 β , and iNOS, as well as an increase in the expression of IL-10, TGF- β , and Arg1, and mitigate depressive-like behaviors [178]. Research has demonstrated that the administration of LPS *in vivo* can result in microglia retaining alterations in epigenetic modifications for an extended duration. Upon subsequent immune stimulation, microglia exhibit immune memory, which triggers the secretion of a greater quantity of pro-inflammatory cytokines [179]. Therefore, early life adversity may cause prolonged epigenetic modifications of microglia that consistently and profoundly alter individual stress susceptibility and stress resilience [180], which was consistent with the finding that childhood trauma increased immune activation in MDD [181]. However, it was also discovered that microglia, upon receiving secondary immune stimulation, exhibited a sluggish proinflammatory response. This could be attributed to the fact that the initial immune stimulation caused a decrease in active histone modifications in the promoters of the IL-1 β and TNF- α genes, while there was an increase in the amount of repressive histone modification in the IL-1 β promoter. This mechanism could be interpreted as a protective measure to prevent excessive damage to the CNS during recurrent peripheral inflammation [182]. In human studies, alterations in DNA methylation in the peripheral monocyte were associated with MDD, and the methylation status of peripheral macrophages may be a potential biomarker for detecting indicators of depression, and given the similarity of monocytes, macrophages, and microglia, brain microglia may also have similar methylation alterations [183] (Table 3).

Conclusion

Since microglia may be involved in many biological processes simultaneously, external determinants (stress), internal factors (gene, BBB, sympathetic and parasympathetic nervous systems, microbiota, and microglia), and the body's coping mechanisms (GC and cytokines) collectively influence the interaction between depression and immunity, playing different roles in the development of depression depending on a certain context, to varying degrees [185].

It is reasonable to speculate on the role of microglia-associated neuroinflammation in depression: Early-life stress induces long-term epigenetic changes in immune-related genes of microglia (e.g., DNA demethylation processes increase IL-1 β transcription in microglia [172]), profoundly altering an individual's stress susceptibility and resilience. Subsequently, when individuals face significant stress in adulthood, abnormal immune regulation fails to timely control immune responses, resulting in chronic low-grade neuroinflammation. In this state, the body's production of anti-inflammatory, pro-inflammatory, or combined effects through GC production regulated by the HPA axis may vary over time. Concurrently, pro-inflammatory cytokines (e.g., TNF- α , IL-6, IL-1 β), chemokines (e.g., CCL2, CCL5), ROS/RNS, etc., perpetuate a vicious cycle of oxidative stress and inflammation. This cascade leads to the production of excitotoxic substances such as QUIN, excessive glutamate causing excitotoxicity, and reduced release of 5-HT and BDNF, potentially accompanied by dysregulation of sympathetic and vagal function and imbalance of the microbiota. Ultimately, these mechanisms contribute to aberrant neuron-microglia interactions (e.g., maladjustment of neurotransmitters, abnormal neuroplasticity and neurogenesis, neuronal excitotoxicity), culminating in depressive symptoms in individuals. Therefore, the relationship between depression and inflammation likely involves multifaceted interplays among these factors rather than a simple one-way causative relationship.

Controlling neuroinflammation favors depression treatment for long-term benefits, although it is unclear whether inflammation in depression is only a consequence rather than a cause, the pro-inflammatory environment does have an exacerbating effect on the conditions, and chronic inflammation in the CNS drives disease progression, not just a secondary event [186, 187]. Interpreting depression from a microglial perspective, we are forced to face the thorny question of how abnormalities in microglia contribute to depression rather than other psychoneurological disorders. Indeed, as microglia are increasingly explored, there are many studies analyzing the role of microglia in

Table 3 Epigenetic regulations of microglia in depression

Epigenetic modification	Gene	Definition	Drug	Mechanism on microglia	Phenotype	Reference
DNA methylation	TET2	DNA demethylase	-	TET2 is upregulated in microglia and regulates the proinflammatory response induced by LPS in vitro and in vivo, and regulates TLR-4-induced type I IFN response and LPS-induced aerobic glycolysis	TET2 regulates the proinflammatory response in microglia of mice intraperitoneally injected with LPS	[172]
	-	-	5-azacytidine (a demethylating drug)	5-azacytidine decreases IL-1 β promoter DNA methylation and promotes the transcription of IL-1 β in microglia	Aged mice have decreased methylation of the IL-1 β gene promoter in primary microglia basally or following systemic LPS that is associated with increased IL-1 β mRNA, intracellular IL-1 β production, as well as prolonged sickness behavior	[173]
Histone modification	EZH2	Histone methyltransferase	EPZ-6438 (an inhibitor of EZH2)	EPZ-6438 inhibits M1 microglia activation, decreases levels of proinflammatory cytokines and P-p65 in the PFC and hippocampus in young and aged mice	EPZ-6438 relieves depression-like behaviors and spatial memory impairment in CUMS	[177]
	-	-	-	EZH2 inhibits miR-29b-3p expression by combining with miR-29b-3p promoter and H3K27me3 upregulation, and then elevates MMP2 transcription and triggers microglia M1-type polarization	EZH2 exacerbates depression-like behaviors and neuroinflammation of depression by modulating microglia polarization	[174]

Table 3 (continued)

Epigenetic modification	Gene	Definition	Drug	Mechanism on microglia	Phenotype	Reference
Histone modification	H3K9me3	Methylation	FA	FA promotes the expression of IL-13 by reducing the promoter binding of H3K9me3 in microglia	FA improves depressive- and anxiety-like behaviors in adults, accompanied by a decrease in the number of activated microglia, as well as a reduction in the inflammatory response in the cortex and hippocampus of neonatal mice	[176]
	H3K4me3	Methylation	-	LPS preconditioning is accompanied by a reduction in H3K4me3 in the promoters of the IL-1 β and TNF- α genes, and results in an increase in the amount of repressive histone modification H3K9me2 in the IL-1 β promoter	-	[182]
	JMJD3	Histone demethylase	DHEA (a steroid hormone that increases the expression of JMJD3)	DHEA regulates microglial inflammatory responses through phosphorylation of TrkA and subsequent activation of a pathway involving Akt1/Akt2 and cAMP response element-binding protein. The latter induces the expression of JMJD3, which thereby controls the expression of inflammation-related genes and microglial polarization	-	[175]
	H3K9ac	Acetylation	Acetate (a HDAC inhibitor)	Acetate increases histone acetylation and reduces LPS-induced microglial activation and IL-1 β expression	-	[110]

Table 3 (continued)

Epigenetic modification	Gene	Definition	Drug	Mechanism on microglia	Phenotype	Reference
Histone modification	SIRT1	Class III deacetylase	SRT2104 (a selective agonist of SIRT1)	SRT2104 abrogates the increased expression of M1 markers (IL-6, IL-1 β , and iNOS) and the decreased expression of M2 markers (IL-10, TGF- β , and Arginase1) induced by CUMS, shifting microglial polarization toward the M2 phenotype	SRT2104 ameliorates CUMS-induced depressive-like behaviors, as indicated by SPT, TST, and FST	[178]
	H3 or H4	Histone	SB (a histone deacetylase inhibitor)	SB suppresses the LPS-induced release of pro-inflammatory factors from microglia in vitro and increases histone acetylation in the hippocampus	SB abolishes LPS-induced depression-like behaviors and hippocampal microglial activation in mice	[72]
	H3S10phK14ac	Phosphorylation	-	H3S10phK14ac promotes inflammatory cytokines releases by driving the transcription of some of the principal genes mediating LPS response, such as c-Fos, IL-6, and iNOS	-	[184]

Akt1/Akt2, AKT serine/threonine kinase 1/2; *cAMP*, cyclic adenosine monophosphate; *CUMS*, chronic unpredictable mild stress; *DHEA*, dehydroepiandrosterone; *DNMT*, DNA methyltransferase; *DNMT3B*, DNA methyltransferase 3 beta; *EZH2*, enhancer of zeste 2 polycomb repressive complex 2 subunit; *FA*, folic acid; *HDAC*, histone deacetylase; *H3K4me1*, histone 3 lysine 4 monomethylation; *H3K4me3*, histone 3 lysine 4 trimethylation; *H3K9ac*, acetyl-histone H3 (Lys9); *H3K9me2*, histone 3 lysine 9 dimethylation; *H3K9me3*, histone 3 lysine 9 trimethylation; *H3K27me3*, trimethylation of histone H3 lysine 27 trimethylation; *H3S10phK14ac*, histone H3 phospho(Ser10)-acetylation(Lys14); *ICV*, intracerebroventricular; *IL*, interleukin; *JMJD3*, jumonji domain protein 3; *LPS*, lipopolysaccharide; *NF- κ B*, nuclear factor-kappa B; *SB*, sodium butyrate; *SIRT1*, sirtuin 1; *TET2*, tet methylcytosine dioxygenase 2; *TLR-4*, toll-like receptor 4; *TrkA*, tyrosine kinase receptor A

various CNS disorders, such as Alzheimer's disease, but whether there is a uniform paradigm for the microglia state in a particular brain region leading to depression remains unknown. At present, we are still facing some urgent questions. Firstly, are microglia initiators or passive receivers of signals in response to neuronal changes? Secondly, as epigenetic alterations in microglia can be triggered by stress or immune stimulation, whether there exist typical epigenetic modifications in MDD that are shared among the pathologies of this disorder, and what is the extent of microglia epigenetic involvement in each stage of depression? Finally, due to the dynamic and region-specific nature of microglia, how to track the dynamic changes of microglia in emotion-related brain regions in real-time, is the most important for further in-depth research. This framework linking microglia, neuroinflammation, and depression could serve as a reference for targeting microglia to treat depression.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12035-024-04575-w>.

Acknowledgements The authors appreciate the encouragement and helpful comments from other members of the Wang laboratory.

Author Contribution Qingqing Li: conceptualization and design, literature search and screening, writing and editing, figures and illustrations. Ying Xie, Jinyi Lin, Tianli Xin, Miaomiao Li, Ziyang Gu, Yang Zhang, Qixia Lu, and Yihui Guo: literature search and screening. Wuyang Wang: proofreading and reviewing, supervision and guidance, funding acquisition. Yanhong Xing: proofreading and reviewing, supervision and guidance. All authors commit to assume responsibility for the work's content.

Funding This work was supported by the Key University Science Research Project of Jiangsu Province (20KJA310001) and Major Projects of Jiangsu Provincial Health Commission (ZD2022051).

Availability of Data and Materials Not applicable.

Declarations

Ethics Approval Not applicable.

Consent to Participate Not applicable.

Consent for Publication Not applicable.

Competing Interests The authors declare no competing interests.

Disclaimer The funding organizations of this paper had no role in the design, article retrieval, writing, or the decision to submit the article for publication.

References

- Ganança L, Galfalvy HC, Cisneros-Trujillo S, Basseda Z, Cooper TB, Ren X, Figueira ML, Oquendo MA et al (2021) Relationships between inflammatory markers and suicide risk status in major depression. *J Psychiatr Res* 134:192–199. <https://doi.org/10.1016/j.jpsychires.2020.12.029>
- Jha MK, Cai L, Minhajuddin A, Fatt CC, Furman JL, Gadad BS, Mason BL, Greer TL et al (2020) Dysfunctional adaptive immune response in adolescents and young adults with suicide behavior. *Psychoneuroendocrinology* 111:104487. <https://doi.org/10.1016/j.psyneuen.2019.104487>
- Mathers CD, Loncar D (2006) Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 3(11):e442
- Malhi GS, Mann JJ (2018) Depression. *Lancet* 392(10161):2299–2312. [https://doi.org/10.1016/s0140-6736\(18\)31948-2](https://doi.org/10.1016/s0140-6736(18)31948-2)
- Wake H, Moorhouse AJ, Miyamoto A, Nabekura J (2013) Microglia: actively surveying and shaping neuronal circuit structure and function. *Trends Neurosci* 36(4):209–217
- Harry GJ (2013) Microglia during development and aging. *Pharmacol Ther* 139(3):313–326. <https://doi.org/10.1016/j.pharmthera.2013.04.013>
- Medzhitov R (2008) Origin and physiological roles of inflammation. *Nature* 454(7203):428–435. <https://doi.org/10.1038/nature07201>
- Nusslock R, Miller GE (2016) Early-life adversity and physical and emotional health across the lifespan: a neuroimmune network hypothesis. *Biol Psychiatr* 80(1):23–32. <https://doi.org/10.1016/j.biopsych.2015.05.017>
- Zindel J, Kubers P (2020) DAMPs, PAMPs, and LAMPs in immunity and sterile inflammation. *Annu Rev Pathol* 15:493–518. <https://doi.org/10.1146/annurev-pathmechdis-012419-032847>
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW (2008) From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* 9(1):46–56. <https://doi.org/10.1038/nrn2297>
- Miller AH, Maletic V, Raison CL (2009) Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol Psychiatr* 65(9):732–741
- Maes M (1995) Evidence for an immune response in major depression: a review and hypothesis. *Prog Neuropsychopharmacol Biol Psychiatry* 19(1):11–38
- Yirmiya R, Rimmerman N, Reshef R (2015) Depression as a microglial disease. *Trends Neurosci* 38(10):637–658. <https://doi.org/10.1016/j.tins.2015.08.001>
- Mac Giollabhui N, Ng TH, Ellman LM, Alloy LB (2020) The longitudinal associations of inflammatory biomarkers and depression revisited: systematic review, meta-analysis, and meta-regression. *Mol Psychiatr* 26. <https://doi.org/10.1038/s41380-020-00867-4>
- Colasanto M, Madigan S, Korczak DJ (2020) Depression and inflammation among children and adolescents: a meta-analysis. *J Affect Disord* 277:940–948. <https://doi.org/10.1016/j.jad.2020.09.025>
- Almeida IB, Gomes IA, Shanmugam S, de Moura TR, Magalhães LS, de Aquino LAG, de Souza Araújo AA, Oliveira PD et al (2020) Inflammatory modulation of fluoxetine use in patients with depression: a systematic review and meta-analysis. *Cytokine* 131:155100. <https://doi.org/10.1016/j.cyto.2020.155100>
- Felger JC, Haroon E, Patel TA, Goldsmith DR, Wommack EC, Woolwine BJ, Le NA, Feinberg R et al (2020) What does plasma CRP tell us about peripheral and central inflammation in depression? *Mol Psychiatry* 25(6):1301–1311. <https://doi.org/10.1038/s41380-018-0096-3>
- Zhang HX, Xu YQ, Li YY, Lu MF, Shi SX, Ji JL, Wang LW (2018) Difference in proinflammatory cytokines produced by monocytes between patients with major depressive disorder and healthy controls. *J Affect Disord* 234:305–310. <https://doi.org/10.1016/j.jad.2017.12.099>
- Syed SA, Beurel E, Loewenstein DA, Lowell JA, Craighead WE, Dunlop BW, Mayberg HS, Dhabhar F et al (2018) Defective

- inflammatory pathways in never-treated depressed patients are associated with poor treatment response. *Neuron* 5:914–924.e913
20. Alavi M, Grebely J, Matthews GV, Petoumenos K, Yeung B, Day C, Lloyd AR, Van Beek I et al (2012) Effect of pegylated interferon- α -2a treatment on mental health during recent hepatitis C virus infection. *J Gastroenterol Hepatol* 27(5):957–965. <https://doi.org/10.1111/j.1440-1746.2011.07035.x>
 21. Musselman DL, Lawson DH, Gumnick JF, Manatunga AK, Penna S, Goodkin RS, Greiner K, Nemeroff CB et al (2001) Paroxetine for the prevention of depression induced by high-dose interferon alfa. *N Engl J Med* 344(13):961–966. <https://doi.org/10.1056/nejm200103293441303>
 22. Cao P, Chen C, Liu A, Shan Q, Zhu X, Jia C, Peng X, Zhang M et al (2021) Early-life inflammation promotes depressive symptoms in adolescence via microglial engulfment of dendritic spines. *Neuron* 109. <https://doi.org/10.1016/j.neuron.2021.06.012>
 23. Walker AK, Wing EE, Banks WA, Dantzer R (2019) Leucine competes with kynurenine for blood-to-brain transport and prevents lipopolysaccharide-induced depression-like behavior in mice. *Mol Psychiatry* 24(10):1523–1532. <https://doi.org/10.1038/s41380-018-0076-7>
 24. Aguilar-Valles A, Kim J, Jung S, Woodside B, Luheshi GN (2014) Role of brain transmigration neutrophils in depression-like behavior during systemic infection. *Mol Psychiatry* 19(5):599–606. <https://doi.org/10.1038/mp.2013.137>
 25. Setiawan E, Attwells S, Wilson AA, Mizrahi R, Rusjan PM, Miler L, Xu C, Sharma S et al (2018) Association of translocator protein total distribution volume with duration of untreated major depressive disorder: a cross-sectional study. *Lancet Psychiatr* 5(4):339–347
 26. Zhang L, Hu K, Shao T, Hou L, Zhang S, Ye W, Josephson L, Meyer JH et al (2021) Recent developments on PET radiotracers for TSPO and their applications in neuroimaging. *Acta Pharm Sin B* 11(2):373–393
 27. Böttcher C, Fernández-Zapata C, Snijders GJL, Schlickeiser S, Sneebroeck MAM, Kunkel D, De Witte LD, Priller J (2020) Single-cell mass cytometry of microglia in major depressive disorder reveals a non-inflammatory phenotype with increased homeostatic marker expression. *Transl Psychiatry* 10(1):310. <https://doi.org/10.1038/s41398-020-00992-2>
 28. Setiawan E, Wilson AA, Mizrahi R, Rusjan PM, Miler L, Rajkowska G, Suridjan I, Kennedy JL et al (2015) Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. *JAMA Psychiatr* 72(3):268–275. <https://doi.org/10.1001/jamapsychiatry.2014.2427>
 29. Holmes SE, Hinz R, Conen S, Gregory CJ, Matthews JC, Anton-Rodriguez JM, Gerhard A, Talbot PS (2018) Elevated translocator protein in anterior cingulate in major depression and a role for inflammation in suicidal thinking: a positron emission tomography study. *Biol Psychiatr* 83(1):61–69. <https://doi.org/10.1016/j.biopsych.2017.08.005>
 30. Wang H, He Y, Sun Z, Ren S, Liu M, Wang G, Yang J (2022) Microglia in depression: an overview of microglia in the pathogenesis and treatment of depression. *J Neuroinflammation* 19(1):132. <https://doi.org/10.1186/s12974-022-02492-0>
 31. Priller J, Prinz M (2019) Targeting microglia in brain disorders. *Science* 365(6448):32–33
 32. Huang Y, Xu Z, Xiong S, Sun F, Qin G, Hu G, Wang J, Zhao L et al (2018) Repopulated microglia are solely derived from the proliferation of residual microglia after acute dep lesion. *Nat Neurosci* 21(4):530–540
 33. Li Q, Barres BA (2018) Microglia and macrophages in brain homeostasis and disease. *Nat Rev Immunol* 18(4):225–242
 34. Paolicelli R, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira T et al (2011) Synaptic pruning by microglia is necessary for normal brain development. *Science* 333(6048):1456–1458. <https://doi.org/10.1126/science.1202529>
 35. Nguyen PT, Dorman LC, Pan S, Vainchtein ID, Han RT, Nakao-Inoue H, Taloma SE, Barron JJ et al (2020) Microglial remodeling of the extracellular matrix promotes synapse plasticity. *Cell* 182(2):388–403.e315. <https://doi.org/10.1016/j.cell.2020.05.050>
 36. Badimon A, Strasburger HJ, Ayata P, Chen X, Nair A, Ikegami A, Hwang P, Chan AT et al (2020) Negative feedback control of neuronal activity by microglia. *Nature* 586(7829):417–423
 37. Savage JC, Carrier M, Tremblay M (2019) Morphology of microglia across contexts of health and disease. *Methods Mol Biol* 2034:13–26
 38. Kreisel T, Frank MG, Licht T, Reshef R, Ben-Menachem-Zidon O, Baratta MV, Maier SF, Yirmiya R (2014) Dynamic microglial alterations underlie stress-induced depressive-like behavior and suppressed neurogenesis. *Mol Psychiatry* 19(6):699–709. <https://doi.org/10.1038/mp.2013.155>
 39. Butovsky O, Jedrychowski MP, Moore CS, Cialic R, Lanser AJ, Gabriely G, Koeglsperger T, Dake B et al (2014) Identification of a unique TGF- β -dependent molecular and functional signature in microglia. *Nat Neurosci* 17(1):131–143. <https://doi.org/10.1038/nn.3599>
 40. Orihuela R, McPherson CA, Harry GJ (2016) Microglial M1/M2 polarization and metabolic states. *Br J Pharmacol* 173(4):649–665
 41. Leng F, Edison P (2021) Neuroinflammation and microglial activation in Alzheimer disease: where do we go from here? *Nat Rev Neurol* 17(3):157–172
 42. Cherry JD, Olschowka JA, O'Banion MK (2014) Are “resting” microglia more “m2”? *Front Immunol* 5:594. <https://doi.org/10.3389/fimmu.2014.00594>
 43. Mendes MS, Majewska AK (2021) An overview of microglia ontogeny and maturation in the homeostatic and pathological brain. *Eur J Neurosci* 53(11):3525–3547
 44. Marschallinger J, Iram T, Zardeneta M, Lee SE, Lehallier B, Haney MS, Pluvinaige JV, Mathur V et al (2020) Lipid-droplet-accumulating microglia represent a dysfunctional and proinflammatory state in the aging brain. *Nat Neurosci* 23(2):194–208
 45. Drevets WC (2000) Functional anatomical abnormalities in limbic and prefrontal cortical structures in major depression. *Prog Brain Res* 126:413–431. [https://doi.org/10.1016/s0079-6123\(00\)26027-5](https://doi.org/10.1016/s0079-6123(00)26027-5)
 46. Späti J, Hänggi J, Doerig N, Ernst J, Sambataro F, Brakowski J, Jäncke L, grosse Holtforth M et al (2015) Prefrontal thinning affects functional connectivity and regional homogeneity of the anterior cingulate cortex in depression. *Neuropsychopharmacology* 40(7):1640–1648. <https://doi.org/10.1038/npp.2015.8>
 47. Hultman R, Mague SD, Li Q, Katz BM, Michel N, Lin L, Wang J, David LK et al (2016) Dysregulation of prefrontal cortex-mediated slow-evolving limbic dynamics drives stress-induced emotional pathology. *Neuron* 91(2):439–452. <https://doi.org/10.1016/j.neuron.2016.05.038>
 48. Xu P, Chen A, Li Y, Xing X, Lu H (2019) Medial prefrontal cortex in neurological diseases. *Physiol Genom* 51(9):432–442. <https://doi.org/10.1152/physiolgenomics.00006.2019>
 49. Li C, Liu B, Xu J, Jing B, Guo L, Wang L, Wang M, Zhang H et al (2023) Phloretin decreases microglia-mediated synaptic engulfment to prevent chronic mild stress-induced depression-like behaviors in the mPFC. *Theranostics* 13(3):955–972. <https://doi.org/10.7150/thno.76553>
 50. He L, Zheng Y, Huang L, Ye J, Ye Y, Luo H, Chen X, Yao W et al (2022) Nrf2 regulates the arginase 1(+) microglia phenotype through the initiation of TREM2 transcription, ameliorating

- depression-like behavior in mice. *Transl Psychiatry* 12(1):459. <https://doi.org/10.1038/s41398-022-02227-y>
51. Yang L, Liu C, Li W, Ma Y, Huo S, Ozathaley A, Ren J, Yuan W et al (2021) Depression-like behavior associated with E/I imbalance of mPFC and amygdala without TRPC channels in mice of knockout IL-10 from microglia. *Brain Behav Immun* 97:68–78. <https://doi.org/10.1016/j.bbi.2021.06.015>
 52. Nie X, Kitaoka S, Tanaka K, Segi-Nishida E, Imoto Y, Ogawa A, Nakano F, Tomohiro A et al (2018) The innate immune receptors TLR2/4 mediate repeated social defeat stress-induced social avoidance through prefrontal microglial activation. *Neuron* 99(3):464–479.e467. <https://doi.org/10.1016/j.neuron.2018.06.035>
 53. Couch Y, Anthony DC, Dolgov O, Revischin A, Festoff B, Santos AI, Steinbusch HW, Strekalova T (2013) Microglial activation, increased TNF and SERT expression in the prefrontal cortex define stress-altered behaviour in mice susceptible to anhedonia. *Brain Behav Immun* 29:136–146. <https://doi.org/10.1016/j.bbi.2012.12.017>
 54. Tanaka K, Furuyashiki T, Kitaoka S, Senzai Y, Imoto Y, Segi-Nishida E, Deguchi Y, Breyer RM et al (2012) Prostaglandin E2-mediated attenuation of mesocortical dopaminergic pathway is critical for susceptibility to repeated social defeat stress in mice. *J Neurosci* 32(12):4319–4329. <https://doi.org/10.1523/jneurosci.5952-11.2012>
 55. Matisz CE, Gruber AJ (2022) Neuroinflammatory remodeling of the anterior cingulate cortex as a key driver of mood disorders in gastrointestinal disease and disorders. *Neurosci Biobehav Rev* 133:104497. <https://doi.org/10.1016/j.neubiorev.2021.12.020>
 56. Comte M, Schön D, Coull JT, Reynaud E, Khalfa S, Belzeaux R, Ibrahim EC, Guedj E et al (2016) Dissociating bottom-up and top-down mechanisms in the cortico-limbic system during emotion processing. *Cereb Cortex* 26(1):144–155. <https://doi.org/10.1093/cercor/bhu185>
 57. Catani M, Dell'acqua F, Thiebaut de Schotten M (2013) A revised limbic system model for memory, emotion and behaviour. *Neuroscience and biobehavioral reviews* 37(8):1724–1737. <https://doi.org/10.1016/j.neubiorev.2013.07.001>
 58. Wohleb ES, Hanke ML, Corona AW, Powell ND, Stiner LM, Bailey MT, Nelson RJ, Godbout JP et al (2011) β -Adrenergic receptor antagonism prevents anxiety-like behavior and microglial reactivity induced by repeated social defeat. *J Neurosci* 31(17):6277–6288. <https://doi.org/10.1523/jneurosci.0450-11.2011>
 59. Zheng ZH, Tu JL, Li XH, Hua Q, Liu WZ, Liu Y, Pan BX, Hu P et al (2021) Neuroinflammation induces anxiety- and depressive-like behavior by modulating neuronal plasticity in the basolateral amygdala. *Brain Behav Immun* 91:505–518. <https://doi.org/10.1016/j.bbi.2020.11.007>
 60. Zhang J, Rong P, Zhang L, He H, Zhou T, Fan Y, Mo L, Zhao Q et al (2021) IL4-driven microglia modulate stress resilience through BDNF-dependent neurogenesis. *Sci Adv* 7(12). <https://doi.org/10.1126/sciadv.abb9888>
 61. Castro DC, Bruchas MR (2019) A motivational and neuropeptidergic hub: anatomical and functional diversity within the nucleus accumbens shell. *Neuron* 102(3):529–552. <https://doi.org/10.1016/j.neuron.2019.03.003>
 62. Klawonn AM, Fritz M, Castany S, Pignatelli M, Canal C, Simil F, Tejeda HA, Levinsson J et al (2021) Microglial activation elicits a negative affective state through prostaglandin-mediated modulation of striatal neurons. *Immunity* 54(2):225–234.e226
 63. Cakmak JD, Liu L, Poirier SE, Schaefer B, Poolacherla R, Burhan AM, Sabesan P, St Lawrence K et al (2022) The functional and structural associations of aberrant microglial activity in major depressive disorder. *J Psychiatr Neurosci* 47(3):E197–e208. <https://doi.org/10.1503/jpn.210124>
 64. Yang Y, Wang H, Hu J, Hu H (2018) Lateral habenula in the pathophysiology of depression. *Curr Opin Neurobiol* 48:90–96. <https://doi.org/10.1016/j.conb.2017.10.024>
 65. Guan YF, Huang GB, Xu MD, Gao F, Lin S, Huang J, Wang J, Li YQ et al (2020) Anti-depression effects of ketogenic diet are mediated via the restoration of microglial activation and neuronal excitability in the lateral habenula. *Brain Behav Immun* 88:748–762. <https://doi.org/10.1016/j.bbi.2020.05.032>
 66. He Y, Wang Y, Yu H, Tian Y, Chen X, Chen C, Ren Y, Chen Z et al (2023) Protective effect of Nr4a2 (Nurr1) against LPS-induced depressive-like behaviors via regulating activity of microglia and CamkII neurons in anterior cingulate cortex. *Pharmacol Res* 191:106717. <https://doi.org/10.1016/j.phrs.2023.106717>
 67. Wang YL, Han QQ, Gong WQ, Pan DH, Wang LZ, Hu W, Yang M, Li B et al (2018) Microglial activation mediates chronic mild stress-induced depressive- and anxiety-like behavior in adult rats. *J Neuroinflammation* 15(1):21. <https://doi.org/10.1186/s12974-018-1054-3>
 68. Lehmann ML, Weigel TK, Poffenberger CN, Herkenham M (2019) The behavioral sequelae of social defeat require microglia and are driven by oxidative stress in mice. *J Neurosci* 39(28):5594–5605. <https://doi.org/10.1523/jneurosci.0184-19.2019>
 69. Ito N, Hirose E, Ishida T, Hori A, Nagai T, Kobayashi Y, Kiyohara H, Oikawa T et al (2017) Kososan, a Kampo medicine, prevents a social avoidance behavior and attenuates neuroinflammation in socially defeated mice. *J Neuroinflammation* 14(1):98. <https://doi.org/10.1186/s12974-017-0876-8>
 70. Chen R, Weitzner AS, McKennon LA, Fonken LK (2021) Chronic circadian phase advance in male mice induces depressive-like responses and suppresses neuroimmune activation. *Brain Behav Immun - Health* 17:100337. <https://doi.org/10.1016/j.bbih.2021.100337>
 71. Guerrero-Vargas NN, Zárate-Mozo C, Guzmán-Ruiz MA, Cárdenas-Rivera A, Escobar C (2021) Time-restricted feeding prevents depressive-like and anxiety-like behaviors in male rats exposed to an experimental model of shift-work. *J Neurosci Res* 99(2):604–620. <https://doi.org/10.1002/jnr.24741>
 72. Yamawaki Y, Yoshioka N, Nozaki K, Ito H, Oda K, Harada K, Shirawachi S, Asano S et al (2018) Sodium butyrate abolishes lipopolysaccharide-induced depression-like behaviors and hippocampal microglial activation in mice. *Brain Res* 1680:13–38. <https://doi.org/10.1016/j.brainres.2017.12.004>
 73. Chan KL, Cathomas F, Russo SJ (2019) Central and peripheral inflammation link metabolic syndrome and major depressive disorder. *Physiology (Bethesda)* 34(2):123–133. <https://doi.org/10.1152/physiol.00047.2018>
 74. Rustenhoven J, Drieu A, Mamuladze T, de Lima KA, Dykstra T, Wall M, Papadopoulos Z, Kanamori M et al (2021) Functional characterization of the dural sinuses as a neuroimmune interface. *Cell* 184(4):1000–1016.e1027
 75. Gu M, Mei XL, Zhao YN (2021) Sepsis and cerebral dysfunction: BBB damage, neuroinflammation, oxidative stress, apoptosis and autophagy as key mediators and the potential therapeutic approaches. *Neurotox Res* 39(2):489–503. <https://doi.org/10.1007/s12640-020-00270-5>
 76. Iadecola C (2013) The pathobiology of vascular dementia. *Neuron* 80(4):844–866. <https://doi.org/10.1016/j.neuron.2013.10.008>
 77. Torres-Platas SG, Cruceanu C, Chen GG, Turecki G, Mechawar N (2014) Evidence for increased microglial priming and macrophage recruitment in the dorsal anterior cingulate white matter of depressed suicides. *Brain Behav Immun* 42:50–59. <https://doi.org/10.1016/j.bbi.2014.05.007>
 78. Huang Y, Chen S, Luo Y, Han Z (2020) Crosstalk between Inflammation and the BBB in stroke. *Curr Neuropharmacol*

- 18(12):1227–1236. <https://doi.org/10.2174/1570159x18666200620230321>
79. Leonard BE (2018) Inflammation and depression: a causal or coincidental link to the pathophysiology? *Acta Neuropsychiatr* 30(1):1–16. <https://doi.org/10.1017/neu.2016.69>
80. Erickson MA, Banks WA (2018) Neuroimmune axes of the blood-brain barriers and blood-brain interfaces: bases for physiological regulation, disease states, and pharmacological interventions. *Pharmacol Rev* 70(2):278–314
81. Cheng Y, Dese S, Martinez A, Worthen RJ, Jope RS, Beurel E (2018) TNF α disrupts blood brain barrier integrity to maintain prolonged depressive-like behavior in mice. *Brain Behav Immun* 69:556–567
82. Sochocka M, Diniz BS, Leszek J (2017) Inflammatory response in the CNS: friend or foe? *Mol Neurobiol* 54(10):8071–8089
83. Kalkman HO (2020) The association between vascular inflammation and depressive disorder. Causality, biomarkers and targeted treatment. *Pharmaceuticals* 13(5):92. <https://doi.org/10.3390/ph13050092>
84. D'Mello C, Le T, Swain MG (2009) Cerebral microglia recruit monocytes into the brain in response to tumor necrosis factor- α signaling during peripheral organ inflammation. *J Neurosci* 29(7):2089–2102. <https://doi.org/10.1523/jneurosci.3567-08.2009>
85. Hu H, Yang X, He Y, Duan C, Sun N (2022) Psychological stress induces depressive-like behavior associated with bone marrow-derived monocyte infiltration into the hippocampus independent of blood-brain barrier disruption. *J Neuroinflammation* 19(1):208. <https://doi.org/10.1186/s12974-022-02569-w>
86. McKim D, Weber M, Niraula A, Sawicki C, Liu X, Jarrett B, Ramirez-Chan K, Wang Y et al (2018) Microglial recruitment of IL-1 β -producing monocytes to brain endothelium causes stress-induced anxiety. *Mol Psychiatry* 23(6):1421–1431. <https://doi.org/10.1038/mp.2017.64>
87. Bisht K, Okojie KA, Sharma K, Lentferink DH, Sun YY, Chen HR, Uweru JO, Amancherla S et al (2021) Capillary-associated microglia regulate vascular structure and function through PANX1-P2RY12 coupling in mice. *Nat Commun* 12(1):5289
88. Haruwaka K, Ikegami A, Tachibana Y, Ohno N, Konishi H, Hashimoto A, Matsumoto M, Kato D et al (2019) Dual microglia effects on blood brain barrier permeability induced by systemic inflammation. *Nat Commun* 10(1):5816
89. Weber MD, McKim DB, Niraula A, Witcher KG, Yin W, Sobol CG, Wang Y, Sawicki CM et al (2019) The influence of microglial elimination and repopulation on stress sensitization induced by repeated social defeat. *Biol Psychiatry* 85(8):667–678
90. Frank MG, Fonken LK, Watkins LR, Maier SF (2019) Microglia: neuroimmune-sensors of stress. *Semin Cell Dev Biol* 94:176–185. <https://doi.org/10.1016/j.semcdb.2019.01.001>
91. Nieto-Quero A, Chaves-Peña P, Santín LJ, Pérez-Martín M, Pedraza C (2021) Do changes in microglial status underlie neurogenesis impairments and depressive-like behaviours induced by psychological stress? A systematic review in animal models. *Neurobiol Stress* 15:100356. <https://doi.org/10.1016/j.ynstr.2021.100356>
92. Webster JC, Oakley RH, Jewell CM, Cidlowski JA (2001) Pro-inflammatory cytokines regulate human glucocorticoid receptor gene expression and lead to the accumulation of the dominant negative beta isoform: a mechanism for the generation of glucocorticoid resistance. *Proc Natl Acad Sci USA* 98(12):6865–6870. <https://doi.org/10.1073/pnas.121455098>
93. Han YM, Kim MS, Jo J, Shin D, Kwon SH, Seo JB, Kang D, Lee BD et al (2021) Decoding the temporal nature of brain GR activity in the NF κ B signal transition leading to depressive-like behavior. *Mol Psychiatry* 26(9):5087–5096. <https://doi.org/10.1038/s41380-021-01016-1>
94. Feng X, Zhao Y, Yang T, Song M, Wang C, Yao Y, Fan H (2019) Glucocorticoid-driven NLRP3 inflammasome activation in hippocampal microglia mediates chronic stress-induced depressive-like behaviors. *Front Mol Neurosci* 12:210. <https://doi.org/10.3389/fnmol.2019.00210>
95. Busillo JM, Azzam KM, Cidlowski JA (2011) Glucocorticoids sensitize the innate immune system through regulation of the NLRP3 inflammasome. *J Biol Chem* 286(44):38703–38713. <https://doi.org/10.1074/jbc.M111.275370>
96. Nair A, Bonneau RH (2006) Stress-induced elevation of glucocorticoids increases microglia proliferation through NMDA receptor activation. *J Neuroimmunol* 171(1–2):72–85. <https://doi.org/10.1016/j.jneuroim.2005.09.012>
97. Frank MG, Hershman SA, Weber MD, Watkins LR, Maier SF (2014) Chronic exposure to exogenous glucocorticoids primes microglia to pro-inflammatory stimuli and induces NLRP3 mRNA in the hippocampus. *Psychoneuroendocrinology* 40:191–200. <https://doi.org/10.1016/j.psyneuen.2013.11.006>
98. Frank MG, Miguel ZD, Watkins LR, Maier SF (2010) Prior exposure to glucocorticoids sensitizes the neuroinflammatory and peripheral inflammatory responses to E. coli lipopolysaccharide. *Brain Behav Immun* 24(1):19–30. <https://doi.org/10.1016/j.bbi.2009.07.008>
99. Weber MD, Godbout JP, Sheridan JF (2017) Repeated social defeat, neuroinflammation, and behavior: monocytes carry the signal. *Neuropsychopharmacology* 42(1):46–61
100. Sugama S, Takenouchi T, Hashimoto M, Ohata H, Takenaka Y, Kakinuma Y (2019) Stress-induced microglial activation occurs through β -adrenergic receptor: noradrenaline as a key neurotransmitter in microglial activation. *J Neuroinflammation* 16(1):266. <https://doi.org/10.1186/s12974-019-1632-z>
101. Hosoi T, Okuma Y, Nomura Y (2000) Electrical stimulation of afferent vagus nerve induces IL-1 β expression in the brain and activates HPA axis. *Am J Physiol Regul Integr Comp Physiol* 279(1):R141–147. <https://doi.org/10.1152/ajpregu.2000.279.1.R141>
102. Zhang Q, Lu Y, Bian H, Guo L, Zhu H (2017) Activation of the α 7 nicotinic receptor promotes lipopolysaccharide-induced conversion of M1 microglia to M2. *Am J Transl Res* 9(3):971–985
103. Kalkman HO (2023) Inhibition of microglial GSK3 β activity is common to different kinds of antidepressants: a proposal for an in vitro screen to detect novel antidepressant principles. *Biomedicines* 11(3). <https://doi.org/10.3390/biomedicines11030806>
104. Thion MS, Low D, Silvén A, Chen J, Grisel P, Schulte-Schrepping J, Blecher R, Ulas T et al (2018) Microbiome influences prenatal and adult microglia in a sex-specific manner. *Cell* 172(3):500–516.e516
105. Shea AK, Frey BN, Gervais N, Lopez A, Minuzzi L (2022) Depression in midlife women attending a menopause clinic is associated with a history of childhood maltreatment. *Climacteric* 25(2):203–207. <https://doi.org/10.1080/13697137.2021.1915270>
106. Gonzalez-Santana A, Diaz Heijtz R (2020) Bacterial peptidoglycans from microbiota in neurodevelopment and behavior. *Trends Mol Med* 26(8):729–743. <https://doi.org/10.1016/j.molmed.2020.05.003>
107. Valles-Colomer M, Falony G, Darzi Y, Tigchelaar EF, Wang J, Tito RY, Shiweck C, Kurilshikov A et al (2019) The neuroactive potential of the human gut microbiota in quality of life and depression. *Nat Microbiol* 4(4):623–632
108. Erny D, Dokalis N, Mezö C, Castoldi A, Mossad O, Staszewski O, Frosch M, Villa M et al (2021) Microbiota-derived acetate enables the metabolic fitness of the brain innate immune system during health and disease. *Cell Metab* 33(11):2260–2276.e2267. <https://doi.org/10.1016/j.cmet.2021.10.010>
109. Wenzel TJ, Gates EJ, Ranger AL, Klegeris A (2020) Short-chain fatty acids (SCFAs) alone or in combination regulate select

- immune functions of microglia-like cells. *Mol Cell Neurosci* 105:103493. <https://doi.org/10.1016/j.mcn.2020.103493>
110. Soliman ML, Puig KL, Combs CK, Rosenberger TA (2012) Acetate reduces microglia inflammatory signaling in vitro. *J Neurochem* 123(4):555–567. <https://doi.org/10.1111/j.1471-4159.2012.07955.x>
 111. D'Mello C, Ronaghan N, Zaheer R, Dicay M, Le T, MacNaughton WK, Surette MG, Swain MG (2015) Probiotics improve inflammation-associated sickness behavior by altering communication between the peripheral immune system and the brain. *J Neurosci* 35(30):10821–10830. <https://doi.org/10.1523/jneurosci.0575-15.2015>
 112. Li H, Xiang Y, Zhu Z, Wang W, Jiang Z, Zhao M, Cheng S, Pan F et al (2021) Rifaximin-mediated gut microbiota regulation modulates the function of microglia and protects against CUMS-induced depression-like behaviors in adolescent rat. *J Neuroinflammation* 18(1):254. <https://doi.org/10.1186/s12974-021-02303-y>
 113. Ng QX, Peters C, Ho CYX, Lim DY, Yeo WS (2018) A meta-analysis of the use of probiotics to alleviate depressive symptoms. *J Affect Disord* 228:13–19. <https://doi.org/10.1016/j.jad.2017.11.063>
 114. Schmidtner AK, Slattery DA, Gläsner J, Hiergeist A, Gryksa K, Malik VA, Hellmann-Regen J, Heuser I et al (2019) Minocycline alters behavior, microglia and the gut microbiome in a trait-anxiety-dependent manner. *Transl Psychiatry* 9(1):223. <https://doi.org/10.1038/s41398-019-0556-9>
 115. Yang Y, Ishima T, Wan X, Wei Y, Chang L, Zhang J, Qu Y, Hashimoto K (2021) Microglial depletion and abnormalities in gut microbiota composition and short-chain fatty acids in mice after repeated administration of colony stimulating factor 1 receptor inhibitor PLX5622. *Eur Arch Psychiatry Clin Neurosci* 272(3):483–495
 116. Mossad O, Batut B, Yilmaz B, Dokalis N, Mezö C, Nent E, Nabavi LS, Mayer M et al (2022) Gut microbiota drives age-related oxidative stress and mitochondrial damage in microglia via the metabolite N(6)-carboxymethyllysine. *Nat Neurosci* 25(3):295–305. <https://doi.org/10.1038/s41593-022-01027-3>
 117. Hosoi T, Okuma Y, Matsuda T, Nomura Y (2005) Novel pathway for LPS-induced afferent vagus nerve activation: possible role of nodose ganglion. *Auton Neurosci: Basic Clin* 120(1–2):104–107. <https://doi.org/10.1016/j.autneu.2004.11.012>
 118. Meneses G, Bautista M, Florentino A, Díaz G, Acero G, Besedovsky H, Meneses D, Fleury A et al (2016) Electric stimulation of the vagus nerve reduced mouse neuroinflammation induced by lipopolysaccharide. *J Inflamm* 13:33. <https://doi.org/10.1186/s12950-016-0140-5>
 119. Kaczmarczyk R, Tejera D, Simon BJ, Heneka MT (2017) Microglia modulation through external vagus nerve stimulation in a murine model of Alzheimer's disease. *J Neurochem*. <https://doi.org/10.1111/jnc.14284>
 120. Fantl K (1950) Psychiatry and tuberculosis. *Calif Med* 73(6):538–540
 121. Rybakowski J (2022) Infections and mental diseases: from tuberculosis to COVID-19. *Psychiatr Pol* 56(5):931–944. <https://doi.org/10.12740/pp/152125>
 122. Kroeze Y, Zhou H, Homberg JR (2012) The genetics of selective serotonin reuptake inhibitors. *Pharmacol Ther* 136(3):375–400. <https://doi.org/10.1016/j.pharmthera.2012.08.015>
 123. McCarron RM, Shapiro B, Rawles J, Luo J (2021) Depression. *Ann Intern Med* 174(5):Itc65–itc80
 124. Fabbri C, Kasper S, Zohar J, Souery D, Montgomery S, Albani D, Forloni G, Ferentinos P et al (2021) Drug repositioning for treatment-resistant depression: hypotheses from a pharmacogenomic study. *Prog Neuropsychopharmacol Biol Psychiatry* 104:110050. <https://doi.org/10.1016/j.pnpbp.2020.110050>
 125. Mariani N, Everson J, Pariante CM, Borsini A (2022) Modulation of microglial activation by antidepressants. *J Psychopharmacol* 36(2):131–150. <https://doi.org/10.1177/02698811211069110>
 126. Hoesel B, Schmid JA (2013) The complexity of NF- κ B signaling in inflammation and cancer. *Mol Cancer* 12:86. <https://doi.org/10.1186/1476-4598-12-86>
 127. Ramirez K, Sheridan JF (2016) Antidepressant imipramine diminishes stress-induced inflammation in the periphery and central nervous system and related anxiety- and depressive-like behaviors. *Brain Behav Immun* 57:293–303. <https://doi.org/10.1016/j.bbi.2016.05.008>
 128. Tynan RJ, Weidenhofer J, Hinwood M, Cairns MJ, Day TA, Walker FR (2012) A comparative examination of the anti-inflammatory effects of SSRI and SNRI antidepressants on LPS stimulated microglia. *Brain Behav Immun* 26(3):469–479. <https://doi.org/10.1016/j.bbi.2011.12.011>
 129. Lu Y, Xu X, Jiang T, Jin L, Zhao XD, Cheng JH, Jin XJ, Ma J et al (2019) Sertraline ameliorates inflammation in CUMS mice and inhibits TNF- α -induced inflammation in microglia cells. *Int Immunopharmacol* 67:119–128. <https://doi.org/10.1016/j.intimp.2018.12.011>
 130. Dhami KS, Churchward MA, Baker GB, Todd KG (2019) Fluoxetine and its metabolite norfluoxetine induce microglial apoptosis. *J Neurochem* 148(6):761–778. <https://doi.org/10.1111/jnc.14661>
 131. Liu RP, Zou M, Wang JY, Zhu JJ, Lai JM, Zhou LL, Chen SF, Zhang X et al (2014) Paroxetine ameliorates lipopolysaccharide-induced microglia activation via differential regulation of MAPK signaling. *J Neuroinflammation* 11:47. <https://doi.org/10.1186/1742-2094-11-47>
 132. Park H, Han KM, Jeon H, Lee JS, Lee H, Jeon SG, Park JH, Kim YG et al (2020) The MAO inhibitor tranylcypromine alters LPS- and A β -mediated neuroinflammatory responses in wild-type mice and a mouse model of AD. *Cells* 9(9). <https://doi.org/10.3390/cells9091982>
 133. Arioz BI, Tastan B, Tarakcioglu E, Tufekci KU, Olcum M, Ersoy N, Bagriyanik A, Genc K et al (2019) Melatonin attenuates LPS-induced acute depressive-like behaviors and microglial NLRP3 inflammasome activation through the SIRT1/Nrf2 pathway. *Front Immunol* 10:1511. <https://doi.org/10.3389/fimmu.2019.01511>
 134. Verdonk F, Petit AC, Abdel-Ahad P, Vinckier F, Jouvion G, de Maricourt P, De Medeiros GF, Danckaert A et al (2019) Microglial production of quinolinic acid as a target and a biomarker of the antidepressant effect of ketamine. *Brain Behav Immun* 81:361–373
 135. Yao W, Cao Q, Luo S, He L, Yang C, Chen J, Qi Q, Hashimoto K et al (2022) Microglial ERK-NRBP1-CREB-BDNF signaling in sustained antidepressant actions of (R)-ketamine. *Mol Psychiatry* 27(3):1618–1629. <https://doi.org/10.1038/s41380-021-01377-7>
 136. Zhang K, Yang C, Chang L, Sakamoto A, Suzuki T, Fujita Y, Qu Y, Wang S et al (2020) Essential role of microglial transforming growth factor- β 1 in antidepressant actions of (R)-ketamine and the novel antidepressant TGF- β 1. *Transl Psychiatry* 10(1):32. <https://doi.org/10.1038/s41398-020-0733-x>
 137. Su KP, Lai HC, Peng CY, Su WP, Chang JP, Pariante CM (2019) Interferon- α -induced depression: comparisons between early- and late-onset subgroups and with patients with major depressive disorder. *Brain Behav Immun* 80:512–518. <https://doi.org/10.1016/j.bbi.2019.04.032>
 138. Zhang C (2021) Flare-up of cytokines in rheumatoid arthritis and their role in triggering depression: shared common function and their possible applications in treatment (Review). *Biomed Rep* 14(1):16. <https://doi.org/10.3892/br.2020.1392>
 139. Kappellmann N, Lewis G, Dantzer R, Jones PB, Khandaker GM (2018) Antidepressant activity of anti-cytokine treatment: a systematic review and meta-analysis of clinical trials of chronic

- inflammatory conditions. *Mol Psychiatry* 23(2):335–343. <https://doi.org/10.1038/mp.2016.167>
140. Wittenberg GM, Stylianou A, Zhang Y, Sun Y, Gupta A, Jagannatha PS, Wang D, Hsu B et al (2020) Effects of immunomodulatory drugs on depressive symptoms: a mega-analysis of randomized, placebo-controlled clinical trials in inflammatory disorders. *Mol Psychiatry* 25(6):1275–1285. <https://doi.org/10.1038/s41380-019-0471-8>
 141. Knight JM, Costanzo ES, Singh S, Yin Z, Szabo A, Pawar DS, Hillard CJ, Rizzo JD et al (2021) The IL-6 antagonist tocilizumab is associated with worse depression and related symptoms in the medically ill. *Transl Psychiatry* 11(1):58. <https://doi.org/10.1038/s41398-020-01164-y>
 142. Bassett B, Subramaniam S, Fan Y, Varney S, Pan H, Carneiro AMD, Chung CY (2021) Minocycline alleviates depression-like symptoms by rescuing decrease in neurogenesis in dorsal hippocampus via blocking microglia activation/phagocytosis. *Brain Behav Immun* 91:519–530. <https://doi.org/10.1016/j.bbi.2020.11.009>
 143. Han Y, Zhang L, Wang Q, Zhang D, Zhao Q, Zhang J, Xie L, Liu G et al (2019) Minocycline inhibits microglial activation and alleviates depressive-like behaviors in male adolescent mice subjected to maternal separation. *Psychoneuroendocrinology* 107:37–45
 144. Dean OM, Kanchanatawan B, Ashton M, Mohebbi M, Ng CH, Maes M, Berk L, Sughondhabirum A et al (2017) Adjunctive minocycline treatment for major depressive disorder: a proof of concept trial. *Aust N Z J Psychiatry* 51(8):829–840. <https://doi.org/10.1177/0004867417709357>
 145. Emadi-Kouchak H, Mohammadinejad P, Asadollahi-Amin A, Rasoulinejad M, Zeinoddini A, Yalda A, Akhondzadeh S (2016) Therapeutic effects of minocycline on mild-to-moderate depression in HIV patients: a double-blind, placebo-controlled, randomized trial. *Int Clin Psychopharmacol* 31(1):20–26. <https://doi.org/10.1097/yic.0000000000000098>
 146. Miyaoka T, Wake R, Furuya M, Liaury K, Ieda M, Kawakami K, Tsuchie K, Taki M et al (2012) Minocycline as adjunctive therapy for patients with unipolar psychotic depression: an open-label study. *Prog Neuropsychopharmacol Biol Psychiatry* 37(2):222–226. <https://doi.org/10.1016/j.pnpbp.2012.02.002>
 147. Nettis MA, Lombardo G, Hastings C, Zajkowska Z, Mariani N, Nikkheslat N, Worrell C, Enache D et al (2021) Augmentation therapy with minocycline in treatment-resistant depression patients with low-grade peripheral inflammation: results from a double-blind randomised clinical trial. *Neuropsychopharmacology* 46(5):939–948. <https://doi.org/10.1038/s41386-020-00948-6>
 148. Husain MI, Chaudhry IB, Husain N, Khoso AB, Rahman RR, Hamirani MM, Hodsoll J, Qurashi I et al (2017) Minocycline as an adjunct for treatment-resistant depressive symptoms: a pilot randomised placebo-controlled trial. *J Psychopharmacol* 31(9):1166–1175. <https://doi.org/10.1177/0269881117724352>
 149. Husain MI, Chaudhry IB, Rahman RR, Hamirani MM, Qurashi I, Khoso AB, Deakin JF, Husain N et al (2015) Minocycline as an adjunct for treatment-resistant depressive symptoms: study protocol for a pilot randomised controlled trial. *Trials* 16:410. <https://doi.org/10.1186/s13063-015-0933-5>
 150. Dean OM, Data-Franco J, Giorlando F, Berk M (2012) Minocycline: therapeutic potential in psychiatry. *CNS Drugs* 26(5):391–401
 151. Kubicova L, Hadacek F, Chobot V (2013) Quinolinic acid: neurotoxin or oxidative stress modulator? *Int J Mol Sci* 14(11):21328–21338. <https://doi.org/10.3390/ijms141121328>
 152. Capuron L, Miller AH (2011) Immune system to brain signaling: neuropsychopharmacological implications. *Pharmacol Ther* 130(2):226–238
 153. Majidi J, Kosari-Nasab M, Salari AA (2016) Developmental minocycline treatment reverses the effects of neonatal immune activation on anxiety- and depression-like behaviors, hippocampal inflammation, and HPA axis activity in adult mice. *Brain Res Bull* 120:1–13. <https://doi.org/10.1016/j.brainresbull.2015.10.009>
 154. Stippl A, Kirkgöze FN, Bajbouj M, Grimm S (2020) Differential effects of electroconvulsive therapy in the treatment of major depressive disorder. *Neuropsychobiology* 79(6):408–416. <https://doi.org/10.1159/000505553>
 155. Yue N, Li B, Yang L, Han QQ, Huang HJ, Wang YL, Wang J, Yu R et al (2018) Electro-acupuncture alleviates chronic unpredictable stress-induced depressive- and anxiety-like behavior and hippocampal neuroinflammation in rat model of depression. *Front Mol Neurosci* 11:149. <https://doi.org/10.3389/fnmol.2018.00149>
 156. Rimmerman N, Verdiger H, Goldenberg H, Naggan L, Robinson E, Kozela E, Gelb S, Reshef R et al (2021) Microglia and their LAG3 checkpoint underlie the antidepressant and neurogenesis-enhancing effects of electroconvulsive stimulation. *Mol Psychiatry* 27(2):1120–1135
 157. Zuo C, Cao H, Feng F, Li G, Huang Y, Zhu L, Gu Z, Yang Y et al (2022) Repetitive transcranial magnetic stimulation exerts anti-inflammatory effects via modulating glial activation in mice with chronic unpredictable mild stress-induced depression. *Int Immunopharmacol* 109:108788. <https://doi.org/10.1016/j.intimp.2022.108788>
 158. Xiao K, Luo Y, Liang X, Tang J, Wang J, Xiao Q, Qi Y, Li Y et al (2021) Beneficial effects of running exercise on hippocampal microglia and neuroinflammation in chronic unpredictable stress-induced depression model rats. *Transl Psychiatry* 11(1):461. <https://doi.org/10.1038/s41398-021-01571-9>
 159. Perez-Caballero L, Pérez-Egea R, Romero-Grimaldi C, Puigdemont D, Molet J, Caso JR, Mico JA, Pérez V et al (2014) Early responses to deep brain stimulation in depression are modulated by anti-inflammatory drugs. *Mol psychiatr* 19(5):607–614
 160. Liu D, Wang Z, Liu S, Wang F, Zhao S, Hao A (2011) Anti-inflammatory effects of fluoxetine in lipopolysaccharide(LPS)-stimulated microglial cells. *Neuropharmacology* 61(4):592–599. <https://doi.org/10.1016/j.neuropharm.2011.04.033>
 161. Tomaz VS, Chaves Filho AJM, Cordeiro RC, Jucá PM, Soares MVR, Barroso PN et al (2020) Antidepressants of different classes cause distinct behavioral and brain pro- and anti-inflammatory changes in mice submitted to an inflammatory model of depression. *J Affect Disord* 268:188–200. <https://doi.org/10.1016/j.jad.2020.03.022>
 162. Göttert R, Fidzinski P, Kraus L, Schneider UC, Holtkamp M, Endres M et al (2022) Lithium inhibits tryptophan catabolism via the inflammation-induced kynurenine pathway in human microglia. *Glia* 70(3):558–571. <https://doi.org/10.1002/glia.24123>
 163. Chen Y, Zhang Y, Wang J, Li S, Wang Y, Zhang Z et al (2023) Anti-neuroinflammation effects of transcutaneous auricular vagus nerve stimulation against depression-like behaviors via hypothalamic $\alpha 7nAChR/JAK2/STAT3/NF-\kappa B$ pathway in rats exposed to chronic unpredictable mild stress. *CNS Neurosci Ther*. <https://doi.org/10.1111/cns.14207>
 164. Bauer ME, Teixeira AL (2019) Inflammation in psychiatric disorders: what comes first? *Ann N Y Acad Sci* 1437(1):57–67. <https://doi.org/10.1111/nyas.13712>
 165. Dantzer RCS, Russo SJ, Dinan TG (2018) Resilience and immunity. *Brain Behav Immun* 74:28–42
 166. Ménard C, Pfau ML, Hodes GE, Russo SJ (2017) Immune and neuroendocrine mechanisms of stress vulnerability and resilience. *Neuropsychopharmacology* 42(1):62–80. <https://doi.org/10.1038/npp.2016.90>

167. Barnes J, Mondelli V, Pariante CM (2017) Genetic contributions of inflammation to depression. *Neuropsychopharmacology* 42(1):81–98. <https://doi.org/10.1038/npp.2016.169>
168. Miller AH, Raison CL (2016) The role of inflammation in depression: from evolutionary imperative to modern treatment target. *Nat Rev Immunol* 16(1):22–34. <https://doi.org/10.1038/nri.2015.5>
169. Border R, Johnson EC, Evans LM, Smolen A, Berley N, Sullivan PF, Keller MC (2019) No support for historical candidate gene or candidate gene-by-interaction hypotheses for major depression across multiple large samples. *Am J Psychiatry* 176(5):376–387
170. Culverhouse RC, Saccone NL, Horton AC, Ma Y, Anstey KJ, Banaschewski T, Burmeister M, Cohen-Woods S et al (2018) Collaborative meta-analysis finds no evidence of a strong interaction between stress and 5-HTTLPR genotype contributing to the development of depression. *Mol Psychiatry* 23(1):133–142. <https://doi.org/10.1038/mp.2017.44>
171. Catale C, Bussone S, Lo Iacono L, Viscomi MT, Palacios D, Troisi A, Carola V (2020) Exposure to different early-life stress experiences results in differentially altered DNA methylation in the brain and immune system. *Neurobiol Stress* 13:100249. <https://doi.org/10.1016/j.ynstr.2020.100249>
172. Carrillo-Jimenez A, Deniz Ö, Niklison-Chirou MV, Ruiz R, Bezerra-Salomão K, Stratoulas V, Amouroux R, Yip PK et al (2019) TET2 regulates the neuroinflammatory response in microglia. *Cell Rep* 29(3):697–713.e698. <https://doi.org/10.1016/j.celrep.2019.09.013>
173. Matt SM, Lawson MA, Johnson RW (2016) Aging and peripheral lipopolysaccharide can modulate epigenetic regulators and decrease IL-1 β promoter DNA methylation in microglia. *Neurobiol Aging* 47:1–9. <https://doi.org/10.1016/j.neurobiolaging.2016.07.006>
174. Huang X, Yang Q, Xie L, Lei S (2022) Histone methyltransferase enhancer of zeste 2 polycomb repressive complex 2 subunit exacerbates inflammation in depression rats by modulating microglia polarization. *Bioengineered* 13(3):5509–5524. <https://doi.org/10.1080/21655979.2022.2036892>
175. Alexaki V, Fodelianaki G, Neuwirth A, Mund C, Kourgiantaki A, Ieronimaki E, Lyroni K, Troullinaki M et al (2018) DHEA inhibits acute microglia-mediated inflammation through activation of the TrkA-Akt1/2-CREB-Jmjd3 pathway. *Mol Psychiatry* 23(6):1410–1420. <https://doi.org/10.1038/mp.2017.167>
176. Zhao T, Wu D, Du J, Liu G, Ji G, Wang Z, Peng F, Man L et al (2022) Folic acid attenuates glial activation in neonatal mice and improves adult mood disorders through epigenetic regulation. *Front Pharmacol* 13:818423. <https://doi.org/10.3389/fphar.2022.818423>
177. Wang W, Qin X, Wang R, Xu J, Wu H, Khalid A, Jiang H, Liu D et al (2020) EZH2 is involved in vulnerability to neuroinflammation and depression-like behaviors induced by chronic stress in different aged mice. *J Affect Disord* 272:452–464. <https://doi.org/10.1016/j.jad.2020.03.154>
178. Duan CM, Zhang JR, Wan TF, Wang Y, Chen HS, Liu L (2020) SRT2104 attenuates chronic unpredictable mild stress-induced depressive-like behaviors and imbalance between microglial M1 and M2 phenotypes in the mice. *Behav Brain Res* 378:112296. <https://doi.org/10.1016/j.bbr.2019.112296>
179. Wendeln AC, Degenhardt K, Kaurani L, Gertig M, Ulas T, Jain G, Wagner J, Häslér LM et al (2018) Innate immune memory in the brain shapes neurological disease hallmarks. *Nature* 556(7701):332–338
180. Halaris A (2019) Inflammation and depression but where does the inflammation come from? *Curr Opin Psychiatry* 32(5):422–428. <https://doi.org/10.1097/ycp.0000000000000531>
181. Müller N, Krause D, Barth R, Myint AM, Weidinger E, Stettinger W, Zill P, Drexhage H et al (2019) Childhood adversity and current stress are related to pro- and anti-inflammatory cytokines in major depression. *J Affect Disord* 253:270–276. <https://doi.org/10.1016/j.jad.2019.04.088>
182. Schaafsma W, Zhang X, van Zomeren KC, Jacobs S, Georgieva PB, Wolf SA, Kettenmann H, Janova H et al (2015) Long-lasting pro-inflammatory suppression of microglia by LPS-preconditioning is mediated by RelB-dependent epigenetic silencing. *Brain Behav Immun* 48:205–221. <https://doi.org/10.1016/j.bbi.2015.03.013>
183. Zhu Y, Strachan E, Fowler E, Bacus T, Roy-Byrne P, Zhao J (2019) Genome-wide profiling of DNA methylome and transcriptome in peripheral blood monocytes for major depression: a monozygotic discordant twin study. *Transl Psychiatry* 9(1):215. <https://doi.org/10.1038/s41398-019-0550-2>
184. Riggio G, Vilella A, Benatti C, Schaeffer L, Brunello N, Blom JMC, Zoli M, Tascedda F (2018) LPS-induced histone H3 phospho(Ser10)-acetylation(Lys14) regulates neuronal and microglial neuroinflammatory response. *Brain Behav Immun* 74:277–290. <https://doi.org/10.1016/j.bbi.2018.09.019>
185. Zdanowicz N, Reynaert C, Jacques D, Dubois T (2017) Depression and immunity: a psychosomatic unit. *Psychiatr Danub* 29(Suppl 3):274–278
186. Chen WW, Zhang X, Huang WJ (2016) Role of neuroinflammation in neurodegenerative diseases (Review). *Mol Med Rep* 13(4):3391–3396
187. Katon WJ (2011) Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues Clin Neurosci* 13(1):7–23. <https://doi.org/10.31887/DCNS.2011.13.1/wkaton>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.