

Representation Theory of $GL_2(\mathbb{F}_q)$

Dongho Tommy Kim

December 2024

Representation Theory of Finite Groups

① $G = GL_2(\mathbb{F}_q)$

Representation Theory of Finite Groups

- 1 $G = GL_2(\mathbb{F}_q)$
- 2 $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$

Representation Theory of Finite Groups

- ① $G = GL_2(\mathbb{F}_q)$
- ② $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$
- ③ $\varphi_i : G \rightarrow GL(V_i)$,

$$\sum_i (\dim \varphi_i)^2 = |G|$$

Representation Theory of Finite Groups

- ① $G = GL_2(\mathbb{F}_q)$
- ② $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$
- ③ $\varphi_i : G \rightarrow GL(V_i)$,

$$\sum_i (\dim \varphi_i)^2 = |G|$$

- ④ # of irreducible representations = # of conjugacy classes

Representation Theory of Finite Groups

- ① $G = GL_2(\mathbb{F}_q)$
- ② $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$
- ③ $\varphi_i : G \rightarrow GL(V_i)$,

$$\sum_i (\dim \varphi_i)^2 = |G|$$

- ④ # of irreducible representations = # of conjugacy classes
- ⑤ $\|\chi_\varphi\|^2 = \langle \chi_\varphi, \chi_\varphi \rangle = \frac{1}{|G|} \sum_{c.c} (\chi_\varphi(c.c))^2 |c.c|$

Representation Theory of Finite Groups

- ① $G = GL_2(\mathbb{F}_q)$
- ② $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$
- ③ $\varphi_i : G \rightarrow GL(V_i)$,

$$\sum_i (\dim \varphi_i)^2 = |G|$$

- ④ # of irreducible representations = # of conjugacy classes
- ⑤ $\|\chi_\varphi\|^2 = \langle \chi_\varphi, \chi_\varphi \rangle = \frac{1}{|G|} \sum_{c.c} (\chi_\varphi(c.c))^2 |c.c|$
- ⑥ φ is irreducible $\Leftrightarrow \|\chi_\varphi\|^2 = 1$

Representation Theory of Finite Groups

- ① $G = GL_2(\mathbb{F}_q)$
- ② $\psi = \varphi \oplus \rho$, then $\chi_\psi = \chi_\varphi + \chi_\rho$
- ③ $\varphi_i : G \rightarrow GL(V_i)$,

$$\sum_i (\dim \varphi_i)^2 = |G|$$

- ④ # of irreducible representations = # of conjugacy classes
- ⑤ $\|\chi_\varphi\|^2 = \langle \chi_\varphi, \chi_\varphi \rangle = \frac{1}{|G|} \sum_{c.c} (\chi_\varphi(c.c))^2 |c.c|$
- ⑥ φ is irreducible $\Leftrightarrow \|\chi_\varphi\|^2 = 1$
- ⑦ Tensor product of representations give a representation

Size of $GL_2(\mathbb{F}_q)$

$$GL_2(\mathbb{F}_q) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0 \right\}$$

Size of $GL_2(\mathbb{F}_q)$

$$GL_2(\mathbb{F}_q) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0 \right\}$$

- $(q^2 - 1)$ choices for (a, b)

Size of $GL_2(\mathbb{F}_q)$

$$GL_2(\mathbb{F}_q) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0 \right\}$$

- $(q^2 - 1)$ choices for (a, b)
- $(q^2 - q)$ choices for (c, d)

Size of $GL_2(\mathbb{F}_q)$

$$GL_2(\mathbb{F}_q) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0 \right\}$$

- $(q^2 - 1)$ choices for (a, b)
- $(q^2 - q)$ choices for (c, d)

$$|GL_2(\mathbb{F}_q)| = (q^2 - 1)(q^2 - q).$$

Characteristic polynomial

Characteristic polynomial of $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{F}_q)$:

$$\lambda^2 - (a + d)\lambda + (ad - bc)$$

Roots: (λ_1, λ_2)

① Diagonalizable matrices

① Diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$

① Diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$
- Distinct eigenvalues: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$

① Diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$
- Distinct eigenvalues: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$

② Non-diagonalizable matrices

① Diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$
- Distinct eigenvalues: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$

② Non-diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$

① Diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$
- Distinct eigenvalues: Matrices of form $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$

② Non-diagonalizable matrices

- Repeated eigenvalue: Matrices of form $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$
- Distinct eigenvalues: Matrices of form $\begin{pmatrix} a & db \\ b & a \end{pmatrix}$

Conjugacy Classes

- Type A: $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$
- Type B: $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$
- Type C: $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$
- Type D: $\begin{pmatrix} a & db \\ b & a \end{pmatrix}$

Conjugacy Classes

- Type A: $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \longrightarrow (q-1) \text{ classes}$
- Type B: $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \longrightarrow \frac{(q-1)(q-2)}{2} \text{ classes}$
- Type C: $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix} \longrightarrow (q-1) \text{ classes}$
- Type D: $\begin{pmatrix} a & db \\ b & a \end{pmatrix} \longrightarrow \frac{q(q-1)}{2} \text{ classes}$

Partial Result:

$$\sum_{\text{matrix type}} (\# \text{ of conjugacy classes}) = q^2 - 1$$

Conjugacy Classes

Partial Result:

$$\sum_{\text{matrix type}} (\# \text{ of conjugacy classes}) = q^2 - 1$$

Number of irreducible representations: $q^2 - 1$

Size of Conjugacy Class

$$|\text{Orb}(x)| = \frac{|G|}{|\text{Stab}(x)|}$$

Size of Conjugacy Class

$$|\text{Orb}(x)| = \frac{|G|}{|\text{Stab}(x)|}$$

For our purposes:

$$|\text{Size of conjugacy class of } M| = \frac{|G|}{|\text{Matrices commuting with } M|}$$

Conjugacy Classes

Matrix Type	Element	No. of classes	Size of class
A	$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$	$q - 1$	1
B	$\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$	$\frac{(q-1)(q-2)}{2}$	$q(q+1)$
C	$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$	$(q - 1)$	$q(q+1)$
D	$\begin{pmatrix} a & db \\ b & a \end{pmatrix}$	$\frac{q^2-q}{2}$	$q(q-1)$

Character

Given the character table:

	A	B	C	D
χ_φ	x_A	x_B	x_C	x_D

We have:

$$\begin{aligned}\|\chi_\varphi\|^2 &= \langle \chi_\varphi, \chi_\varphi \rangle \\ &= \frac{1}{|G|} \sum (x_{\text{matrix type}})^2 \cdot (\text{Number of classes}) \cdot (\text{Size})\end{aligned}$$

Induced Representation

Given G , a finite group, $H \subseteq G$, and $\pi : H \rightarrow GL(\mathbb{C})$, a representation:

Definition

$$\text{Ind}_H^G(\pi) = \{f : G \rightarrow \mathbb{C} \mid \pi(h)f(g) = f(gh^{-1}) \ \forall h^{-1} \in H\}.$$

Induced Representation

Given G , a finite group, $H \subseteq G$, and $\pi : H \rightarrow GL(\mathbb{C})$, a representation:

Definition

$$\text{Ind}_H^G(\pi) = \{f : G \rightarrow \mathbb{C} \mid \pi(h)f(g) = f(gh^{-1}) \ \forall h^{-1} \in H\}.$$

Consider $\pi = \text{triv} : G \rightarrow GL(\mathbb{C})$.

Induced Representation

Given G , a finite group, $H \subseteq G$, and $\pi : H \rightarrow GL(\mathbb{C})$, a representation:

Definition

$$\text{Ind}_H^G(\pi) = \{f : G \rightarrow \mathbb{C} \mid \pi(h)f(g) = f(gh^{-1}) \ \forall h^{-1} \in H\}.$$

Consider $\pi = \text{triv} : G \rightarrow GL(\mathbb{C})$.

Definition

$$\begin{aligned}\text{Ind}_H^G(\text{triv}) &= \{f : G \rightarrow \mathbb{C} \mid f(g) = f(gh^{-1}) \ \forall h^{-1} \in H\} \\ &= \{f : G \rightarrow \mathbb{C} \mid f \text{ constant on } gH\} \\ &= \{f : G/H \rightarrow \mathbb{C}\}\end{aligned}$$

Consider family of matrices

$$T = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

We want $\text{Ind}_T^G(\text{triv}) = \{f : G/T \rightarrow \mathbb{C}\}.$

Consider family of matrices

$$T = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

We want $\text{Ind}_T^G(\text{triv}) = \{f : G/T \rightarrow \mathbb{C}\}$.

Problem: G/T is too large

Consider family of matrices (“Borel subgroup”):

$$T \subseteq B = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

“Inflating”

Consider family of matrices (“Borel subgroup”):

$$T \subseteq B = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

$$\mathrm{Ind}_B^G(\mathrm{triv}) = \{f : G/B \rightarrow \mathbb{C}\}$$

“Inflating”

Consider family of matrices (“Borel subgroup”):

$$T \subseteq B = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

$$\mathrm{Ind}_B^G(\mathrm{triv}) = \{f : G/B \rightarrow \mathbb{C}\}$$

$$\{gB\} \longleftrightarrow \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} / (\text{up to scaling})$$

“Inflating”

Consider family of matrices (“Borel subgroup”):

$$T \subseteq B = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \subseteq GL_2(\mathbb{F}_q)$$

$$\mathrm{Ind}_B^G(\mathrm{triv}) = \{f : G/B \rightarrow \mathbb{C}\}$$

$$\{gB\} \longleftrightarrow \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} / (\text{up to scaling})$$

$$\{gB\} \longleftrightarrow \mathbb{P}^1 \text{ in } \mathbb{F}_q^2$$

Permutation Representation

$$\text{Vector space } M \text{ over } \mathbb{P}^1(\mathbb{F}_q) = \text{span} \left\{ \begin{pmatrix} 1 \\ r \end{pmatrix} : r \in \mathbb{F}_q \cup \{\infty\} \right\}$$

Permutation Representation

$$\text{Vector space } M \text{ over } \mathbb{P}^1(\mathbb{F}_q) = \text{span} \left\{ \begin{pmatrix} 1 \\ r \end{pmatrix} : r \in \mathbb{F}_q \cup \{\infty\} \right\}$$

$$\left(\sum_{v \in \mathbb{P}^1} e_v \right) \text{ is a trivial subrepresentation of perm. rep}$$

Permutation Representation

$$\text{Vector space } M \text{ over } \mathbb{P}^1(\mathbb{F}_q) = \text{span} \left\{ \begin{pmatrix} 1 \\ r \end{pmatrix} : r \in \mathbb{F}_q \cup \{\infty\} \right\}$$

$$\left(\sum_{v \in \mathbb{P}^1} e_v \right) \text{ is a trivial subrepresentation of perm. rep}$$

$$\text{perm. rep} = \text{triv. subrep} \oplus \varphi$$

Character table

	A	B	C	D
χ_{perm}	$q + 1$	2	1	0
χ_{triv}	1	1	1	1
χ_{φ}	q	1	0	-1

Character table

	A	B	C	D
χ_{perm}	$q + 1$	2	1	0
χ_{triv}	1	1	1	1
χ_{φ}	q	1	0	-1

$$\begin{aligned}\|\chi_{\varphi}\|^2 &= \frac{1}{|G|} \sum (\text{char. value})^2 \cdot (\# \text{ of classes}) \cdot (\text{size of class}) \\ &= 1\end{aligned}$$

Other irreducible representations

① Type A matrices

Other irreducible representations

- ① Type A matrices
- ② Type B, C matrices: Principal series representations
- ③ Type D matrices: Cuspidal representations

- <https://arxiv.org/pdf/0712.4051>
- <https://math.ntnu.edu.tw/~li/note/REPGL2K.pdf>