Programming Report

Construct Bayes classifier using Iris Data set. The classifier treats the first twenty five vectors as traning set and the remaining vectors as test set for each class. Parameters of the features are estimated based on Maximum likelihood. The assumptions are that: 1) the conditional densities of features in the classes are multivariate normal; 2) the a priori probabilities of the classes are

Show the results such as correct classification rate for training and test data, respectively and confusion table.

List of Iris Data Set

List of the Data Set								Iris virginica			
lris setosa			Iris versicolor								
Sepal length	Sepal width	Petal length	Petal width	Sepal length	Sepal width	Petal length	Petal width	Sepal length	Sepal width	Petal length	Petal width
5.1	3.5	1.4	0.2	7.0	3.2	4.7	1.4	6.3	3.3	6.0	2.5
4.9	3.0	1.4	0.2	6.4	3.2	4.5	1.5	5.8		5.1	1.9
4.7	3.2	1.3	0.2	6.9	3.1	4.9	1.5	7.1	3.0	5.9	2.1
4.6	3.1	1.5	0.2	5.5	2.3	4.0	1.3	6.3	2.9	5.6	1.8
	3.6	1.4	0.2	6.5	2.8	4.6	1.5	6.5	3.0	5.8	2.2
5.0	3.9	1.7	0.4	5.7	2.8	4.5	1.3	7.6	3.0	6.6	2.1
	3.4	1.4	0.3	6.3	3.3	4.7	1.6	4.9	2.5	4.5	1.7
4.6 5.0	3.4	1.5	0.2	4.9	24	3.3	1.0	7.3	29	6.3	1.8
4.4	2.9	1.4	0.2	6.6	29	4.6	1.3	6.7	2.5	5.8	1.8
4.9	3.1	1.5	0.1	5.2	27	3.9	1.4	7.2	3.6	6.1	2.5
5.4	3.7	1.5	0.2	5.0	20	3.5	1.0	6.5	3.2	5.1	20
	3.4	1.6	0.2	5.9	3.0	4.2	1.5	6.4 .	27	5.3	1.9
4.8	3.0	1.4	0.1	6.0	2.2	4.0	1.0	6.8	3.0	5.5	2.1
4.8	3.0	1.1	0.1	6.1	2.9	4.7	1.4	5.7	2.5	5.0	2.0
4.3		1.2	0.2	5.6	29	3.6	1.3	5.8	2.8	5.1	2.4
5.8	4.0	1.5	0.4	6.7	3.1	4.4	1.4	6.4	3.2	5.3	2.3
5.7	4.4		0.4	5.6	3.0	4.5	1.5	6.5	3.0	5.5	1.8
5.4	3.9	1.3		5.8	2.7	4.1	1.0	7.7	3.8	6.7	2.2
5.1	3.5	1.4	0.3		2.2	4.5	1.5	7.7	2.6	6.9	2.3
5.7	3.8	1.7	0.3	6.2	2.5	3.9	1.1	6.0	2.2	5.0	1.5
5.1	3.8	1.5	0.3	5.6		4.8	1.8	6.9	3.2	5.7	2.3
3.4	3.4	1.7	0.2	5.9	3.2	4.0	1.3	5.6	2.8	4.9	2.0
5.1	3.7	1.5	0.4	6.1	2.8	4.9	1.5	7.7	2.8	6.7	2.0
4.6	3.6	1.0	0.2	6.3		4.7	1.2	6.3	2.7	4.9	1.8
5.1	3.3	1.7	0.5	6.1	2.8	4.3	1.3	6.7	3.3	5.7	2.1
4.8	3.4	1.9	0.2	6.4	3.0	4.4	1.4	7.2	3.2	6.0	1.8
5.0	3.0	1.6	0.2	6.6		4.8	1.4	6.2	2.8	4.8	1.8
5.0	3.4	1.6	0.4	6.8	2.8	5.0	1.7	6.1	3.0		1.8
5.2	3.5	1.5	0.2	6.7	3.0	4.5	1.5	5.4	2.8	5.5	2.1
5.2	3.4	1.4	0.2	6.0	20		1.0	7.2	3.0	5.8	1.6
4.7	3.2	1.6	0.2	5.7	2.6	3.5		7.4	2.8	6.1	1.5
4.8	3.1	1.6	0.2	5.5	2.4	3.8	1.1	7.9	3.8	6.4	2.0
5.4	3.4	1.5	0.4	5.5	2.4	3.7	1.0	6.4	2.8	5.6	2
5.2	4.1	1.5	0.1	5.8	2.7	3.9	1.2		2.8	5.1	1.
5.5	4.2	1.4	0.2	6.0	2.7	5.1	1.6	6.3		5.6	1.4
4.9	3.1	1.5	0.2	5.4	3.0	4.5	1.5	5.1	2.6		2.
5.0	3.2	1.2	0.2	6.0	3.4	4.5	1.6	7.7	3.0		2.
5.5	3.5	1.3	0.2	6.7	3.1	4.7	1.5	6.3	3.4		1.
4.9	3.6	1.4	0.1	6.3	23	4.4	1.3	6.4			
4.4	3.0	1.3	0.2	5.6	3.0	4.1	1.3	6.0			1.
5.1	3.4	1.5	0.2	5.5	2.5	4.0	1.3	6.9			2.
5.0	3.5		0.3	5.5	2.6	4.4	1.2	6.7	3.1		2
4.5	2.3	1.3	0.3	6.1	3.0		1.4	6.9			2
4.4	3.2	1.3	0.2	5.8	2.6			5.8			1.
5.0	3.5	1.6	0.6	5.0	23			6.8			
5.1	3.8	1.9	0.4	5.6	2.7	4.2	1.3	6.7			
4.8	3.0	1.4	0.3	5.7	3.0			6.7			
5.1	3.8	1.6	0.2	5.7	2.9			6.3			
4.6	3.2	1.4	0.2	6.2				6.5			
	3.6									5.4	. 2
5.3	3.7	1.5	0.2	5.1	2.5	3.0	1.1	1 0.2		5.1	

SOURCE: M. G. Kandall and A. Stuart. Advanced Theory of Statistics, vol. 3. Hafner, New York, 1966. P. 318.

Iris : 붓꽃속(屬)의 식물, 붓꽃 Sepal : 꽃받침 조각

Petal : 꽃잎