

1 Full publication list

References

- [1] **ALICE** Collaboration, S. Acharya *et al.*, “Z-boson production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV and Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **09** (2020) 076, [arXiv:2005.11126 \[nucl-ex\]](#).
- [2] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of J/ ψ production at midrapidity in pp collisions at $\sqrt{s} = 13$ TeV,” *Phys. Lett. B* (2020) 135758, [arXiv:2005.11123 \[nucl-ex\]](#).
- [3] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of nuclear effects on $\psi(2S)$ production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV,” *JHEP* **07** (2020) 237, [arXiv:2003.06053 \[nucl-ex\]](#).
- [4] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of π , K, and p production in pp collisions at $\sqrt{s} = 13$ TeV,” *Eur. Phys. J. C* **80** no. 8, (2020) 693, [arXiv:2003.02394 \[nucl-ex\]](#).
- [5] **ALICE** Collaboration, S. Acharya *et al.*, “Coherent photoproduction of ρ^0 vector mesons in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **06** (2020) 035, [arXiv:2002.10897 \[nucl-ex\]](#).
- [6] **ALICE** Collaboration, S. Acharya *et al.*, “Linear and non-linear flow modes of charged hadrons in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **05** (2020) 085, [arXiv:2002.00633 \[nucl-ex\]](#).
- [7] **ALICE** Collaboration, S. Acharya *et al.*, “Non-linear flow modes of identified particles in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **06** (2020) 147, [arXiv:1912.00740 \[nucl-ex\]](#).
- [8] **ALICE** Collaboration, S. Acharya *et al.*, “Investigation of the p- Σ^0 interaction via femtoscopy in pp collisions,” *Phys. Lett. B* **805** (2020) 135419, [arXiv:1910.14407 \[nucl-ex\]](#).
- [9] **ALICE** Collaboration, S. Acharya *et al.*, “Longitudinal and azimuthal evolution of two-particle transverse momentum correlations in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **804** (2020) 135375, [arXiv:1910.14393 \[nucl-ex\]](#).
- [10] **ALICE** Collaboration, S. Acharya *et al.*, “Evidence of rescattering effect in Pb-Pb collisions at the LHC through production of $K^*(892)^0$ and $\phi(1020)$ mesons,” *Phys. Lett. B* **802** (2020) 135225, [arXiv:1910.14419 \[nucl-ex\]](#).
- [11] **ALICE** Collaboration, S. Acharya *et al.*, “Centrality and transverse momentum dependence of inclusive J/ ψ production at midrapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **805** (2020) 135434, [arXiv:1910.14404 \[nucl-ex\]](#).

- [12] **ALICE** Collaboration, S. Acharya *et al.*, “Production of (anti-) ^3He and (anti-) ^3H in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. C* **101** no. 4, (2020) 044906, [arXiv:1910.14401](#) [[nucl-ex](#)].
- [13] **ALICE** Collaboration, S. Acharya *et al.*, “Underlying Event properties in pp collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **04** (2020) 192, [arXiv:1910.14400](#) [[nucl-ex](#)].
- [14] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of electrons from heavy-flavour hadron decays as a function of multiplicity in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **02** (2020) 077, [arXiv:1910.14399](#) [[nucl-ex](#)].
- [15] **ALICE** Collaboration, S. Acharya *et al.*, “Probing the effects of strong electromagnetic fields with charge-dependent directed flow in Pb-Pb collisions at the LHC,” *Phys. Rev. Lett.* **125** no. 2, (2020) 022301, [arXiv:1910.14406](#) [[nucl-ex](#)].
- [16] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of spin-orbital angular momentum interactions in relativistic heavy-ion collisions,” *Phys. Rev. Lett.* **125** no. 1, (2020) 012301, [arXiv:1910.14408](#) [[nucl-ex](#)].
- [17] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of $K^*(892)^0$ and $\phi(1020)$ production in pp collisions at $\sqrt{s} = 13$ TeV,” *Phys. Lett. B* **807** (2020) 135501, [arXiv:1910.14397](#) [[nucl-ex](#)].
- [18] **ALICE** Collaboration, S. Acharya *et al.*, “Jet-hadron correlations measured relative to the second order event plane in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **101** no. 6, (2020) 064901, [arXiv:1910.14398](#) [[nucl-ex](#)].
- [19] **ALICE** Collaboration, S. Acharya *et al.*, “ Υ production in pcollisions at $\sqrt{s_{\text{NN}}} = 8.16$ TeV,” *Phys. Lett. B* **806** (2020) 135486, [arXiv:1910.14405](#) [[nucl-ex](#)].
- [20] **ALICE** Collaboration, S. Acharya *et al.*, “Global baryon number conservation encoded in net-proton fluctuations measured in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **807** (2020) 135564, [arXiv:1910.14396](#) [[nucl-ex](#)].
- [21] **ALICE** Collaboration, S. Acharya *et al.*, “ $K^*(892)^0$ and $\phi(1020)$ production at midrapidity in pp collisions at $\sqrt{s} = 8$ TeV,” *Phys. Rev. C* **102** no. 2, (2020) 024912, [arXiv:1910.14410](#) [[nucl-ex](#)].
- [22] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of electrons from semileptonic heavy-flavour hadron decays at midrapidity in pp and Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **804** (2020) 135377, [arXiv:1910.09110](#) [[nucl-ex](#)].

- [23] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of the (anti-) ^3He elliptic flow in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **805** (2020) 135414, [arXiv:1910.09718](#) [nucl-ex].
- [24] **ALICE** Collaboration, S. Acharya *et al.*, “Production of charged pions, kaons, and (anti-)protons in Pb-Pb and inelastic pp collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. C* **101** no. 4, (2020) 044907, [arXiv:1910.07678](#) [nucl-ex].
- [25] **ALICE** Collaboration, S. Acharya *et al.*, “Measurements of inclusive jet spectra in pp and central Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. C* **101** no. 3, (2020) 034911, [arXiv:1909.09718](#) [nucl-ex].
- [26] **ALICE** Collaboration, S. Acharya *et al.*, “Studies of J/ψ production at forward rapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **02** (2020) 041, [arXiv:1909.03158](#) [nucl-ex].
- [27] **ALICE** Collaboration, S. Acharya *et al.*, “Global polarization of $\Lambda\bar{\Lambda}$ hyperons in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ and 5.02 TeV,” *Phys. Rev. C* **101** no. 4, (2020) 044611, [arXiv:1909.01281](#) [nucl-ex].
- [28] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of (multi-)strange hadron production in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *Eur. Phys. J. C* **80** no. 2, (2020) 167, [arXiv:1908.01861](#) [nucl-ex].
- [29] **ALICE** Collaboration, S. Acharya *et al.*, “ $^3_\Lambda\text{H}$ and $^3_\Lambda\bar{\text{H}}$ lifetime measurement in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV via two-body decay,” *Phys. Lett. B* **797** (2019) 134905, [arXiv:1907.06906](#) [nucl-ex].
- [30] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of $\Upsilon(1S)$ elliptic flow at forward rapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. Lett.* **123** no. 19, (2019) 192301, [arXiv:1907.03169](#) [nucl-ex].
- [31] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of prompt D^0 , D^+ , D^{*+} , and D_S^+ production in p collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **12** (2019) 092, [arXiv:1906.03425](#) [nucl-ex].
- [32] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of light (anti-)nuclei production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **800** (2020) 135043, [arXiv:1906.03136](#) [nucl-ex].
- [33] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of the inclusive isolated photon production cross section in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **79** no. 11, (2019) 896, [arXiv:1906.01371](#) [nucl-ex].
- [34] **ALICE** Collaboration, S. Acharya *et al.*, “Scattering studies with low-energy kaon-proton femtoscopy in proton-proton collisions at the LHC,” *Phys. Rev. Lett.* **124** no. 9, (2020) 092301, [arXiv:1905.13470](#) [nucl-ex].

- [35] **ALICE** Collaboration, S. Acharya *et al.*, “Charged-particle production as a function of multiplicity and transverse sphericity in pp collisions at $\sqrt{s} = 5.02$ and 13 TeV,” *Eur. Phys. J. C* **79** no. 10, (2019) 857, [arXiv:1905.07208 \[nucl-ex\]](#).
- [36] **ALICE** Collaboration, S. Acharya *et al.*, “Study of the Λ - Λ interaction with femtoscopy correlations in pp and p-Pb collisions at the LHC,” *Phys. Lett. B* **797** (2019) 134822, [arXiv:1905.07209 \[nucl-ex\]](#).
- [37] **ALICE** Collaboration, S. Acharya *et al.*, “Inclusive J/ψ production at mid-rapidity in pp collisions at $\sqrt{s} = 5.02$ TeV,” *JHEP* **10** (2019) 084, [arXiv:1905.07211 \[nucl-ex\]](#).
- [38] **ALICE** Collaboration, S. Acharya *et al.*, “Production of muons from heavy-flavour hadron decays in pp collisions at $\sqrt{s} = 5.02$ TeV,” *JHEP* **09** (2019) 008, [arXiv:1905.07207 \[nucl-ex\]](#).
- [39] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of the production of charm jets tagged with D^0 mesons in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **08** (2019) 133, [arXiv:1905.02510 \[nucl-ex\]](#).
- [40] **ALICE** Collaboration, S. Acharya *et al.*, “Exploration of jet substructure using iterative declustering in pp and Pb collisions at LHC energies,” *Phys. Lett. B* **802** (2020) 135227, [arXiv:1905.02512 \[nucl-ex\]](#).
- [41] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of charged jet cross section in pp collisions at $\sqrt{s} = 5.02$ TeV,” *Phys. Rev. D* **100** no. 9, (2019) 092004, [arXiv:1905.02536 \[nucl-ex\]](#).
- [42] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of jet radial profiles in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **796** (2019) 204–219, [arXiv:1904.13118 \[nucl-ex\]](#).
- [43] **ALICE** Collaboration, S. Acharya *et al.*, “First Observation of an Attractive Interaction between a Proton and a Cascade Baryon,” *Phys. Rev. Lett.* **123** no. 11, (2019) 112002, [arXiv:1904.12198 \[nucl-ex\]](#).
- [44] **ALICE** Collaboration, S. Acharya *et al.*, “Coherent J/ψ photoproduction at forward rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **798** (2019) 134926, [arXiv:1904.06272 \[nucl-ex\]](#).
- [45] **ALICE** Collaboration, S. Acharya *et al.*, “One-dimensional charged kaon femtoscopy in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. C* **100** no. 2, (2019) 024002, [arXiv:1903.12310 \[nucl-ex\]](#).
- [46] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of strange baryon interactions with femtoscopic correlations,” *Phys. Lett. B* **802** (2020) 135223, [arXiv:1903.06149 \[nucl-ex\]](#).

- [47] **ALICE** Collaboration, S. Acharya *et al.*, “Investigations of Anisotropic Flow Using Multiparticle Azimuthal Correlations in pp, p-Pb, Xe-Xe, and Pb-Pb Collisions at the LHC,” *Phys. Rev. Lett.* **123** no. 14, (2019) 142301, [arXiv:1903.01790](#) [[nucl-ex](#)].
- [48] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of (anti-)deuteron production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **794** (2019) 50–63, [arXiv:1902.09290](#) [[nucl-ex](#)].
- [49] **ALICE** Collaboration, S. Acharya *et al.*, “Calibration of the photon spectrometer PHOS of the ALICE experiment,” *JINST* **14** no. 05, (2019) P05025, [arXiv:1902.06145](#) [[physics.ins-det](#)].
- [50] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of D^0 , D^+ , D^{*+} and D_s^+ production in pp collisions at $\sqrt{s} = 5.02$ TeV with ALICE,” *Eur. Phys. J. C* **79** no. 5, (2019) 388, [arXiv:1901.07979](#) [[nucl-ex](#)].
- [51] **ALICE** Collaboration, S. Acharya *et al.*, “Event-shape and multiplicity dependence of freeze-out radii in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **09** (2019) 108, [arXiv:1901.05518](#) [[nucl-ex](#)].
- [52] **ALICE** Collaboration, S. Acharya *et al.*, “Real-time data processing in the ALICE High Level Trigger at the LHC,” *Comput. Phys. Commun.* **242** (2019) 25–48, [arXiv:1812.08036](#) [[physics.ins-det](#)].
- [53] **ALICE** Collaboration, S. Acharya *et al.*, “Charged-particle pseudorapidity density at mid-rapidity in p-Pb collisions at $\sqrt{s_{NN}} = 8.16$ TeV,” *Eur. Phys. J. C* **79** no. 4, (2019) 307, [arXiv:1812.01312](#) [[nucl-ex](#)].
- [54] **ALICE** Collaboration, S. Acharya *et al.*, “Study of J/ψ azimuthal anisotropy at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *JHEP* **02** (2019) 012, [arXiv:1811.12727](#) [[nucl-ex](#)].
- [55] **ALICE** Collaboration, S. Acharya *et al.*, “Jet fragmentation transverse momentum measurements from di-hadron correlations in $\sqrt{s} = 7$ TeV pp and $\sqrt{s_{NN}} = 5.02$ TeV p-Pb collisions,” *JHEP* **03** (2019) 169, [arXiv:1811.09742](#) [[nucl-ex](#)].
- [56] **ALICE** Collaboration, S. Acharya *et al.*, “ Λ_c^+ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **793** (2019) 212–223, [arXiv:1809.10922](#) [[nucl-ex](#)].
- [57] **ALICE** Collaboration, S. Acharya *et al.*, “Event-shape engineering for the D-meson elliptic flow in mid-central Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *JHEP* **02** (2019) 150, [arXiv:1809.09371](#) [[nucl-ex](#)].

- [58] **ALICE** Collaboration, S. Acharya *et al.*, “Measuring $K_S^0 K^\pm$ interactions using pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **790** (2019) 22–34, [arXiv:1809.07899 \[nucl-ex\]](#).
- [59] **ALICE** Collaboration, S. Acharya *et al.*, “Energy dependence of exclusive J/ψ photoproduction off protons in ultra-peripheral pcollisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J. C* **79** no. 5, (2019) 402, [arXiv:1809.03235 \[nucl-ex\]](#).
- [60] **ALICE** Collaboration, S. Acharya *et al.*, “Charged jet cross section and fragmentation in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. D* **99** no. 1, (2019) 012016, [arXiv:1809.03232 \[nucl-ex\]](#).
- [61] **ALICE** Collaboration, S. Acharya *et al.*, “Multiplicity dependence of light-flavor hadron production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. C* **99** no. 2, (2019) 024906, [arXiv:1807.11321 \[nucl-ex\]](#).
- [62] **ALICE** Collaboration, S. Acharya *et al.*, “Medium modification of the shape of small-radius jets in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **10** (2018) 139, [arXiv:1807.06854 \[nucl-ex\]](#).
- [63] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of dielectron production in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **99** no. 2, (2019) 024002, [arXiv:1807.00923 \[nucl-ex\]](#).
- [64] **ALICE** Collaboration, S. Acharya *et al.*, “p-p, p- Λ and Λ - Λ correlations studied via femtoscopy in pp reactions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. C* **99** no. 2, (2019) 024001, [arXiv:1805.12455 \[nucl-ex\]](#).
- [65] **ALICE** Collaboration, S. Acharya *et al.*, “Analysis of the apparent nuclear modification in peripheral Pbcollisions at 5.02 TeV,” *Phys. Lett. B* **793** (2019) 420–432, [arXiv:1805.05212 \[nucl-ex\]](#).
- [66] **ALICE** Collaboration, S. Acharya *et al.*, “ Υ suppression at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **790** (2019) 89–101, [arXiv:1805.04387 \[nucl-ex\]](#).
- [67] **ALICE** Collaboration, S. Acharya *et al.*, “Azimuthal Anisotropy of Heavy-Flavor Decay Electrons in p-Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **122** no. 7, (2019) 072301, [arXiv:1805.04367 \[nucl-ex\]](#).
- [68] **ALICE** Collaboration, S. Acharya *et al.*, “Dielectron and heavy-quark production in inelastic and high-multiplicity protoncollisions at $\sqrt{s_{NN}} = 13$ TeV,” *Phys. Lett. B* **788** (2019) 505–518, [arXiv:1805.04407 \[hep-ex\]](#).

- [69] **ALICE** Collaboration, S. Acharya *et al.*, “Centrality and pseudorapidity dependence of the charged-particle multiplicity density in Xecollisions at $\sqrt{s_{\text{NN}}} = 5.44 \text{ TeV}$,” *Phys. Lett. B* **790** (2019) 35–48, [arXiv:1805.04432 \[nucl-ex\]](#).
- [70] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of the inclusive J/ψ polarization at forward rapidity in pp collisions at $\sqrt{s} = 8 \text{ TeV}$,” *Eur. Phys. J. C* **78** no. 7, (2018) 562, [arXiv:1805.04374 \[hep-ex\]](#).
- [71] **ALICE** Collaboration, S. Acharya *et al.*, “Inclusive J/ψ production at forward and backward rapidity in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 8.16 \text{ TeV}$,” *JHEP* **07** (2018) 160, [arXiv:1805.04381 \[nucl-ex\]](#).
- [72] **ALICE** Collaboration, S. Acharya *et al.*, “Anisotropic flow of identified particles in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$,” *JHEP* **09** (2018) 006, [arXiv:1805.04390 \[nucl-ex\]](#).
- [73] **ALICE** Collaboration, S. Acharya *et al.*, “Direct photon elliptic flow in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$,” *Phys. Lett. B* **789** (2019) 308–322, [arXiv:1805.04403 \[nucl-ex\]](#).
- [74] **ALICE** Collaboration, S. Acharya *et al.*, “Dielectron production in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$,” *JHEP* **09** (2018) 064, [arXiv:1805.04391 \[hep-ex\]](#).
- [75] **ALICE** Collaboration, S. Acharya *et al.*, “Inclusive J/ψ production in Xecollisions at $\sqrt{s_{\text{NN}}} = 5.44 \text{ TeV}$,” *Phys. Lett. B* **785** (2018) 419–428, [arXiv:1805.04383 \[nucl-ex\]](#).
- [76] **ALICE** Collaboration, S. Acharya *et al.*, “Suppression of $\Lambda(1520)$ resonance production in central Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$,” *Phys. Rev. C* **99** (2019) 024905, [arXiv:1805.04361 \[nucl-ex\]](#).
- [77] **ALICE** Collaboration, S. Acharya *et al.*, “Production of the $\rho(770)^0$ meson in pp and Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$,” *Phys. Rev. C* **99** no. 6, (2019) 064901, [arXiv:1805.04365 \[nucl-ex\]](#).
- [78] **ALICE** Collaboration, S. Acharya *et al.*, “Two particle differential transverse momentum and number density correlations in p-Pb and Pb-Pb at the LHC,” *Phys. Rev. C* **100** no. 4, (2019) 044903, [arXiv:1805.04422 \[nucl-ex\]](#).
- [79] **ALICE** Collaboration, S. Acharya *et al.*, “Transverse momentum spectra and nuclear modification factors of charged particles in Xe-Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44 \text{ TeV}$,” *Phys. Lett. B* **788** (2019) 166–179, [arXiv:1805.04399 \[nucl-ex\]](#).
- [80] **ALICE** Collaboration, S. Acharya *et al.*, “Measurements of low- p_T electrons from semileptonic heavy-flavour hadron decays at mid-rapidity in pp and Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$,” *JHEP* **10** (2018) 061, [arXiv:1805.04379 \[nucl-ex\]](#).

- [81] **ALICE** Collaboration, S. Acharya *et al.*, “Anisotropic flow in Xe-Xe collisions at $\sqrt{s_{\text{NN}}} = 5.44$ TeV,” *Phys. Lett. B* **784** (2018) 82–95, [arXiv:1805.01832 \[nucl-ex\]](#).
- [82] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of D^0 , D^+ , D^{*+} and D_s^+ production in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **10** (2018) 174, [arXiv:1804.09083 \[nucl-ex\]](#).
- [83] **ALICE** Collaboration, S. Acharya *et al.*, “ ϕ meson production at forward rapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Eur. Phys. J. C* **78** no. 7, (2018) 559, [arXiv:1804.08906 \[nucl-ex\]](#).
- [84] **ALICE** Collaboration, S. Acharya *et al.*, “Energy dependence and fluctuations of anisotropic flow in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ and 2.76 TeV,” *JHEP* **07** (2018) 103, [arXiv:1804.02944 \[nucl-ex\]](#).
- [85] **ALICE** Collaboration, S. Acharya *et al.*, “Azimuthally-differential pion femtoscopy relative to the third harmonic event plane in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **785** (2018) 320–331, [arXiv:1803.10594 \[nucl-ex\]](#).
- [86] **ALICE** Collaboration, S. Acharya *et al.*, “Direct photon production at low transverse momentum in proton-proton collisions at $\sqrt{s} = 2.76$ and 8 TeV,” *Phys. Rev. C* **99** no. 2, (2019) 024912, [arXiv:1803.09857 \[nucl-ex\]](#).
- [87] **ALICE** Collaboration, S. Acharya *et al.*, “Neutral pion and η meson production at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **98** no. 4, (2018) 044901, [arXiv:1803.05490 \[nucl-ex\]](#).
- [88] **ALICE** Collaboration, S. Acharya *et al.*, “Transverse momentum spectra and nuclear modification factors of charged particles in pp, p-Pb and Pb-Pb collisions at the LHC,” *JHEP* **11** (2018) 013, [arXiv:1802.09145 \[nucl-ex\]](#).
- [89] **ALICE** Collaboration, S. Acharya *et al.*, “Prompt and non-prompt J/ψ production and nuclear modification at mid-rapidity in pcollisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Eur. Phys. J. C* **78** no. 6, (2018) 466, [arXiv:1802.00765 \[nucl-ex\]](#).
- [90] **ALICE** Collaboration, S. Acharya *et al.*, “Neutral pion and η meson production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Eur. Phys. J. C* **78** no. 8, (2018) 624, [arXiv:1801.07051 \[nucl-ex\]](#).
- [91] **ALICE** Collaboration, S. Acharya *et al.*, “ Λ_c^+ production in pp collisions at $\sqrt{s} = 7$ TeV and in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **04** (2018) 108, [arXiv:1712.09581 \[nucl-ex\]](#).

- [92] **ALICE** Collaboration, S. Acharya *et al.*, “Relative particle yield fluctuations in $Pb - Pb$ collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Eur. Phys. J. C* **79** no. 3, (2019) 236, [arXiv:1712.07929](#) [[nucl-ex](#)].
- [93] **ALICE** Collaboration, S. Acharya *et al.*, “Constraints on jet quenching in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured by the event-activity dependence of semi-inclusive hadron-jet distributions,” *Phys. Lett. B* **783** (2018) 95–113, [arXiv:1712.05603](#) [[nucl-ex](#)].
- [94] **ALICE** Collaboration, S. Acharya *et al.*, “First measurement of Ξ_c^0 production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **781** (2018) 8–19, [arXiv:1712.04242](#) [[hep-ex](#)].
- [95] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of Z^0 -boson production at large rapidities in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **780** (2018) 372–383, [arXiv:1711.10753](#) [[nucl-ex](#)].
- [96] **ALICE** Collaboration, S. Acharya *et al.*, “Longitudinal asymmetry and its effect on pseudorapidity distributions in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **781** (2018) 20–32, [arXiv:1710.07975](#) [[nucl-ex](#)].
- [97] **ALICE** Collaboration, S. Acharya *et al.*, “Production of ^4He and $^4\overline{\text{He}}$ in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV at the LHC,” *Nucl. Phys. A* **971** (2018) 1–20, [arXiv:1710.07531](#) [[nucl-ex](#)].
- [98] **ALICE** Collaboration, S. Acharya *et al.*, “Production of deuterons, tritons, ^3He nuclei and their antinuclei in pp collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV,” *Phys. Rev. C* **97** no. 2, (2018) 024615, [arXiv:1709.08522](#) [[nucl-ex](#)].
- [99] **ALICE** Collaboration, S. Acharya *et al.*, “Search for collectivity with azimuthal J/ψ -hadron correlations in high multiplicity p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ and 8.16 TeV,” *Phys. Lett. B* **780** (2018) 7–20, [arXiv:1709.06807](#) [[nucl-ex](#)].
- [100] **ALICE** Collaboration, S. Acharya *et al.*, “ J/ψ elliptic flow in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **119** no. 24, (2017) 242301, [arXiv:1709.05260](#) [[nucl-ex](#)].
- [101] **ALICE** Collaboration, S. Acharya *et al.*, “Constraining the magnitude of the Chiral Magnetic Effect with Event Shape Engineering in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **777** (2018) 151–162, [arXiv:1709.04723](#) [[nucl-ex](#)].
- [102] **ALICE** Collaboration, S. Acharya *et al.*, “The ALICE Transition Radiation Detector: construction, operation, and performance,” *Nucl. Instrum. Meth. A* **881** (2018) 88–127, [arXiv:1709.02743](#) [[physics.ins-det](#)].

- [103] **ALICE** Collaboration, S. Acharya *et al.*, “Kaon femtoscopy in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **96** no. 6, (2017) 064613, [arXiv:1709.01731 \[nucl-ex\]](#).
- [104] **ALICE** Collaboration, S. Acharya *et al.*, “Systematic studies of correlations between different order flow harmonics in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **97** no. 2, (2018) 024906, [arXiv:1709.01127 \[nucl-ex\]](#).
- [105] **ALICE** Collaboration, S. Acharya *et al.*, “ π^0 and η meson production in proton-proton collisions at $\sqrt{s} = 8$ TeV,” *Eur. Phys. J. C* **78** no. 3, (2018) 263, [arXiv:1708.08745 \[hep-ex\]](#).
- [106] **ALICE** Collaboration, S. Acharya *et al.*, “Charged-particle multiplicity distributions over a wide pseudorapidity range in proton-proton collisions at $\sqrt{s} = 0.9, 7$, and 8 TeV,” *Eur. Phys. J. C* **77** no. 12, (2017) 852, [arXiv:1708.01435 \[hep-ex\]](#).
- [107] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of deuteron spectra and elliptic flow in Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV at the LHC,” *Eur. Phys. J. C* **77** no. 10, (2017) 658, [arXiv:1707.07304 \[nucl-ex\]](#).
- [108] **ALICE** Collaboration, S. Acharya *et al.*, “Searches for transverse momentum dependent flow vector fluctuations in Pb-Pb and p-Pb collisions at the LHC,” *JHEP* **09** (2017) 032, [arXiv:1707.05690 \[nucl-ex\]](#).
- [109] **ALICE** Collaboration, S. Acharya *et al.*, “ D -meson azimuthal anisotropy in midcentral Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. Lett.* **120** no. 10, (2018) 102301, [arXiv:1707.01005 \[nucl-ex\]](#).
- [110] **ALICE** Collaboration, S. Acharya *et al.*, “Measuring $K_S^0 K^\pm$ interactions using Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **774** (2017) 64–77, [arXiv:1705.04929 \[nucl-ex\]](#).
- [111] **ALICE** Collaboration, S. Acharya *et al.*, “Linear and non-linear flow modes in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **773** (2017) 68–80, [arXiv:1705.04377 \[nucl-ex\]](#).
- [112] **ALICE** Collaboration, D. Adamová *et al.*, “ J/ψ production as a function of charged-particle pseudorapidity density in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **776** (2018) 91–104, [arXiv:1704.00274 \[nucl-ex\]](#).
- [113] **ALICE** Collaboration, J. Adam *et al.*, “Flow dominance and factorization of transverse momentum correlations in Pb-Pb collisions at the LHC,” *Phys. Rev. Lett.* **118** no. 16, (2017) 162302, [arXiv:1702.02665 \[nucl-ex\]](#).

- [114] **ALICE** Collaboration, D. Adamova *et al.*, “Azimuthally differential pion femtoscopy in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. Lett.* **118** no. 22, (2017) 222301, [arXiv:1702.01612](#) [[nucl-ex](#)].
- [115] **ALICE** Collaboration, S. Acharya *et al.*, “Production of muons from heavy-flavour hadron decays in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **770** (2017) 459–472, [arXiv:1702.01479](#) [[nucl-ex](#)].
- [116] **ALICE** Collaboration, S. Acharya *et al.*, “Production of π^0 and η mesons up to high transverse momentum in pp collisions at 2.76 TeV,” *Eur. Phys. J. C* **77** no. 5, (2017) 339, [arXiv:1702.00917](#) [[hep-ex](#)].
- [117] **ALICE** Collaboration, S. Acharya *et al.*, “First measurement of jet mass in Pb-Pb collisions at the LHC,” *Phys. Lett. B* **776** (2018) 249–264, [arXiv:1702.00804](#) [[nucl-ex](#)].
- [118] **ALICE** Collaboration, S. Acharya *et al.*, “Energy dependence of forward-rapidity J/ψ and $\psi(2S)$ production in pp collisions at the LHC,” *Eur. Phys. J. C* **77** no. 6, (2017) 392, [arXiv:1702.00557](#) [[hep-ex](#)].
- [119] **ALICE** Collaboration, S. Acharya *et al.*, “Measurement of D-meson production at mid-rapidity in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **77** no. 8, (2017) 550, [arXiv:1702.00766](#) [[hep-ex](#)].
- [120] **ALICE** Collaboration, J. Adam *et al.*, “ $K^*(892)^0$ and $\phi(1020)$ meson production at high transverse momentum in pp and Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **95** no. 6, (2017) 064606, [arXiv:1702.00555](#) [[nucl-ex](#)].
- [121] **ALICE** Collaboration, D. Adamova *et al.*, “Production of $\Sigma(1385)^\pm$ and $\Xi(1530)^0$ in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Eur. Phys. J. C* **77** no. 6, (2017) 389, [arXiv:1701.07797](#) [[nucl-ex](#)].
- [122] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of the pseudorapidity density distribution for charged particles in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **772** (2017) 567–577, [arXiv:1612.08966](#) [[nucl-ex](#)].
- [123] **ALICE** Collaboration, J. Adam *et al.*, “Insight into particle production mechanisms via angular correlations of identified particles in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **77** no. 8, (2017) 569, [arXiv:1612.08975](#) [[nucl-ex](#)]. [Erratum: *Eur.Phys.J.C* 79, 998 (2019)].
- [124] **ALICE** Collaboration, J. Adam *et al.*, “W and Z boson production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **02** (2017) 077, [arXiv:1611.03002](#) [[nucl-ex](#)].
- [125] **ALICE** Collaboration, J. Adam *et al.*, “Determination of the event collision time with the ALICE detector at the LHC,” *Eur. Phys. J. Plus* **132** no. 2, (2017) 99, [arXiv:1610.03055](#) [[physics.ins-det](#)].

- [126] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of the production of high- p_T electrons from heavy-flavour hadron decays in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **771** (2017) 467–481, [arXiv:1609.07104](#) [[nucl-ex](#)].
- [127] **ALICE** Collaboration, J. Adam *et al.*, “Anomalous evolution of the near-side jet peak shape in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **119** no. 10, (2017) 102301, [arXiv:1609.06643](#) [[nucl-ex](#)].
- [128] **ALICE** Collaboration, J. Adam *et al.*, “Evolution of the longitudinal and azimuthal structure of the near-side jet peak in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **96** no. 3, (2017) 034904, [arXiv:1609.06667](#) [[nucl-ex](#)].
- [129] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of electrons from beauty-hadron decays in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **07** (2017) 052, [arXiv:1609.03898](#) [[nucl-ex](#)].
- [130] **ALICE** Collaboration, J. Adam *et al.*, “Jet-like correlations with neutral pion triggers in pp and central Pb-Pb collisions at 2.76 TeV,” *Phys. Lett. B* **763** (2016) 238–250, [arXiv:1608.07201](#) [[nucl-ex](#)].
- [131] **ALICE** Collaboration, J. Adam *et al.*, “ J/ψ suppression at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **766** (2017) 212–224, [arXiv:1606.08197](#) [[nucl-ex](#)].
- [132] **ALICE** Collaboration, J. Adam *et al.*, “Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions,” *Nature Phys.* **13** (2017) 535–539, [arXiv:1606.07424](#) [[nucl-ex](#)].
- [133] **ALICE** Collaboration, J. Adam *et al.*, “Higher harmonic flow coefficients of identified hadrons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2016) 164, [arXiv:1606.06057](#) [[nucl-ex](#)].
- [134] **ALICE** Collaboration, J. Adam *et al.*, “Elliptic flow of electrons from heavy-flavour hadron decays at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2016) 028, [arXiv:1606.00321](#) [[nucl-ex](#)].
- [135] **ALICE** Collaboration, J. Adam *et al.*, “ D -meson production in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. C* **94** no. 5, (2016) 054908, [arXiv:1605.07569](#) [[nucl-ex](#)].
- [136] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of azimuthal correlations of D mesons and charged particles in pp collisions at $\sqrt{s} = 7$ TeV and p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J. C* **77** no. 4, (2017) 245, [arXiv:1605.06963](#) [[nucl-ex](#)].

- [137] **ALICE** Collaboration, J. Adam *et al.*, “Pseudorapidity dependence of the anisotropic flow of charged particles in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **762** (2016) 376–388, [arXiv:1605.02035 \[nucl-ex\]](#).
- [138] **ALICE** Collaboration, J. Adam *et al.*, “Correlated event-by-event fluctuations of flow harmonics in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. Lett.* **117** (2016) 182301, [arXiv:1604.07663 \[nucl-ex\]](#).
- [139] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of transverse energy at midrapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **94** no. 3, (2016) 034903, [arXiv:1603.04775 \[nucl-ex\]](#).
- [140] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of charged jet production in pcollisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Eur. Phys. J. C* **76** no. 5, (2016) 271, [arXiv:1603.03402 \[nucl-ex\]](#).
- [141] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of $\psi(2S)$ suppression in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **06** (2016) 050, [arXiv:1603.02816 \[nucl-ex\]](#).
- [142] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of D-meson production versus multiplicity in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **08** (2016) 078, [arXiv:1602.07240 \[nucl-ex\]](#).
- [143] **ALICE** Collaboration, J. Adam *et al.*, “Particle identification in ALICE: a Bayesian approach,” *Eur. Phys. J. Plus* **131** no. 5, (2016) 168, [arXiv:1602.01392 \[physics.data-an\]](#).
- [144] **ALICE** Collaboration, J. Adam *et al.*, “Anisotropic flow of charged particles in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. Lett.* **116** no. 13, (2016) 132302, [arXiv:1602.01119 \[nucl-ex\]](#).
- [145] **ALICE** Collaboration, J. Adam *et al.*, “Production of $K^* (892)^0$ and $\phi (1020)$ in pcollisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Eur. Phys. J. C* **76** no. 5, (2016) 245, [arXiv:1601.07868 \[nucl-ex\]](#).
- [146] **ALICE** Collaboration, J. Adam *et al.*, “Multiplicity dependence of charged pion, kaon, and (anti)proton production at large transverse momentum in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **760** (2016) 720–735, [arXiv:1601.03658 \[nucl-ex\]](#).
- [147] **ALICE** Collaboration, J. Adam *et al.*, “Multipion Bose-Einstein correlations in pp,p -Pb, and Pb-Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys. Rev. C* **93** no. 5, (2016) 054908, [arXiv:1512.08902 \[nucl-ex\]](#).
- [148] **ALICE** Collaboration, J. Adam *et al.*, “Multi-strange baryon production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **758** (2016) 389–401, [arXiv:1512.07227 \[nucl-ex\]](#).

- [149] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of the charged-particle multiplicity density at midrapidity in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Rev. Lett.* **116** no. 22, (2016) 222302, [arXiv:1512.06104](#) [[nucl-ex](#)].
- [150] **ALICE** Collaboration, J. Adam *et al.*, “Charge-dependent flow and the search for the chiral magnetic wave in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **93** no. 4, (2016) 044903, [arXiv:1512.05739](#) [[nucl-ex](#)].
- [151] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of an excess in the yield of J/ψ at very low p_{T} in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. Lett.* **116** no. 22, (2016) 222301, [arXiv:1509.08802](#) [[nucl-ex](#)].
- [152] **ALICE** Collaboration, J. Adam *et al.*, “Pseudorapidity and transverse-momentum distributions of charged particles in proton collisions at $\sqrt{s} = 13$ TeV,” *Phys. Lett. B* **753** (2016) 319–329, [arXiv:1509.08734](#) [[nucl-ex](#)].
- [153] **ALICE** Collaboration, J. Adam *et al.*, “Inclusive quarkonium production at forward rapidity in pp collisions at $\sqrt{s} = 8$ TeV,” *Eur. Phys. J. C* **76** no. 4, (2016) 184, [arXiv:1509.08258](#) [[hep-ex](#)].
- [154] **ALICE** Collaboration, J. Adam *et al.*, “Azimuthal anisotropy of charged jet production in $\sqrt{s_{\text{NN}}} = 2.76$ TeV Pb-Pb collisions,” *Phys. Lett. B* **753** (2016) 511–525, [arXiv:1509.07334](#) [[nucl-ex](#)].
- [155] **ALICE** Collaboration, J. Adam *et al.*, “Centrality evolution of the charged-particle pseudorapidity density over a broad pseudorapidity range in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **754** (2016) 373–385, [arXiv:1509.07299](#) [[nucl-ex](#)].
- [156] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of D_s^+ production and nuclear modification factor in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *JHEP* **03** (2016) 082, [arXiv:1509.07287](#) [[nucl-ex](#)].
- [157] **ALICE** Collaboration, J. Adam *et al.*, “Multiplicity and transverse momentum evolution of charge-dependent correlations in pp, p, and Pb collisions at the LHC,” *Eur. Phys. J. C* **76** no. 2, (2016) 86, [arXiv:1509.07255](#) [[nucl-ex](#)].
- [158] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of electrons from heavy-flavour hadron decays in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *Phys. Lett. B* **754** (2016) 81–93, [arXiv:1509.07491](#) [[nucl-ex](#)].
- [159] **ALICE** Collaboration, J. Adam *et al.*, “Charged-particle multiplicities in proton collisions at $\sqrt{s} = 0.9$ to 8 TeV,” *Eur. Phys. J. C* **77** no. 1, (2017) 33, [arXiv:1509.07541](#) [[nucl-ex](#)].

- [160] **ALICE** Collaboration, J. Adam *et al.*, “Direct photon production in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **754** (2016) 235–248, [arXiv:1509.07324](#) [nucl-ex].
- [161] **ALICE** Collaboration, J. Adam *et al.*, “Transverse momentum dependence of D-meson production in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *JHEP* **03** (2016) 081, [arXiv:1509.06888](#) [nucl-ex].
- [162] **ALICE** Collaboration, J. Adam *et al.*, “Coherent $\psi(2S)$ photo-production in ultra-peripheral Pb Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **751** (2015) 358–370, [arXiv:1508.05076](#) [nucl-ex].
- [163] **ALICE** Collaboration, J. Adam *et al.*, “Precision measurement of the mass difference between light nuclei and anti-nuclei,” *Nature Phys.* **11** no. 10, (2015) 811–814, [arXiv:1508.03986](#) [nucl-ex].
- [164] **ALICE** Collaboration, J. Adam *et al.*, “Study of cosmic ray events with high muon multiplicity using the ALICE detector at the CERN Large Hadron Collider,” *JCAP* **01** (2016) 032, [arXiv:1507.07577](#) [astro-ph.HE].
- [165] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of pion freeze-out radii in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **93** no. 2, (2016) 024905, [arXiv:1507.06842](#) [nucl-ex].
- [166] **ALICE** Collaboration, J. Adam *et al.*, “Event shape engineering for inclusive spectra and elliptic flow in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Rev. C* **93** no. 3, (2016) 034916, [arXiv:1507.06194](#) [nucl-ex].
- [167] **ALICE** Collaboration, J. Adam *et al.*, “Elliptic flow of muons from heavy-flavour hadron decays at forward rapidity in Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV,” *Phys. Lett. B* **753** (2016) 41–56, [arXiv:1507.03134](#) [nucl-ex].
- [168] **ALICE** Collaboration, J. Adam *et al.*, “ ϕ -meson production at forward rapidity in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV and in pp collisions at $\sqrt{s} = 2.76$ TeV,” *Phys. Lett. B* **768** (2017) 203–217, [arXiv:1506.09206](#) [nucl-ex].
- [169] **ALICE** Collaboration, J. Adam *et al.*, “Production of light nuclei and anti-nuclei in pp and Pb-Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys. Rev. C* **93** no. 2, (2016) 024917, [arXiv:1506.08951](#) [nucl-ex].
- [170] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of inclusive J/ψ production in p-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV,” *JHEP* **11** (2015) 127, [arXiv:1506.08808](#) [nucl-ex].

- [171] **ALICE** Collaboration, J. Adam *et al.*, “Differential studies of inclusive J/ψ and $\psi(2S)$ production at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **05** (2016) 179, [arXiv:1506.08804](#) [nucl-ex].
- [172] **ALICE** Collaboration, J. Adam *et al.*, “ ${}^3_\Lambda\text{H}$ and ${}^3_{\bar{\Lambda}}\bar{\text{H}}$ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **754** (2016) 360–372, [arXiv:1506.08453](#) [nucl-ex].
- [173] **ALICE** Collaboration, J. Adam *et al.*, “Forward-central two-particle correlations in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **753** (2016) 126–139, [arXiv:1506.08032](#) [nucl-ex].
- [174] **ALICE** Collaboration, J. Adam *et al.*, “One-dimensional pion, kaon, and proton femtoscopy in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **92** no. 5, (2015) 054908, [arXiv:1506.07884](#) [nucl-ex].
- [175] **ALICE** Collaboration, J. Adam *et al.*, “Search for weakly decaying $\bar{\Lambda}\text{n}$ and $\Lambda\Lambda$ exotic bound states in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **752** (2016) 267–277, [arXiv:1506.07499](#) [nucl-ex].
- [176] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of the nuclear modification factor of charged pions, kaons, and protons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **93** no. 3, (2016) 034913, [arXiv:1506.07287](#) [nucl-ex].
- [177] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of high- p_T D meson suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **11** (2015) 205, [arXiv:1506.06604](#) [nucl-ex]. [Addendum: *JHEP* **06**, 032 (2017)].
- [178] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of jet quenching with semi-inclusive hadron-jet distributions in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2015) 170, [arXiv:1506.03984](#) [nucl-ex].
- [179] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of charm and beauty production at central rapidity versus charged-particle multiplicity in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **09** (2015) 148, [arXiv:1505.00664](#) [nucl-ex].
- [180] **ALICE** Collaboration, J. Adam *et al.*, “Inclusive, prompt and non-prompt J/ψ production at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **07** (2015) 051, [arXiv:1504.07151](#) [nucl-ex].
- [181] **ALICE** Collaboration, J. Adam *et al.*, “Coherent ρ^0 photoproduction in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2015) 095, [arXiv:1503.09177](#) [nucl-ex].

- [182] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of pion, kaon and proton production in proton collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **75** no. 5, (2015) 226, [arXiv:1504.00024](#) [[nucl-ex](#)].
- [183] **ALICE** Collaboration, J. Adam *et al.*, “Rapidity and transverse-momentum dependence of the inclusive J/ψ nuclear modification factor in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *JHEP* **06** (2015) 055, [arXiv:1503.07179](#) [[nucl-ex](#)].
- [184] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of dijet k_T in p collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **746** (2015) 385–395, [arXiv:1503.03050](#) [[nucl-ex](#)].
- [185] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of charged jet production cross sections and nuclear modification in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **749** (2015) 68–81, [arXiv:1503.00681](#) [[nucl-ex](#)].
- [186] **ALICE** Collaboration, J. Adam *et al.*, “Measurement of jet suppression in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **746** (2015) 1–14, [arXiv:1502.01689](#) [[nucl-ex](#)].
- [187] **ALICE** Collaboration, J. Adam *et al.*, “Two-pion femtoscopy in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. C* **91** (2015) 034906, [arXiv:1502.00559](#) [[nucl-ex](#)].
- [188] **ALICE** Collaboration, J. Adam *et al.*, “Forward-backward multiplicity correlations in pp collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV,” *JHEP* **05** (2015) 097, [arXiv:1502.00230](#) [[nucl-ex](#)].
- [189] **ALICE** Collaboration, J. Adam *et al.*, “Centrality dependence of particle production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. C* **91** no. 6, (2015) 064905, [arXiv:1412.6828](#) [[nucl-ex](#)].
- [190] **ALICE** Collaboration, B. B. Abelev *et al.*, “Inclusive photon production at forward rapidities in proton-proton collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV,” *Eur. Phys. J. C* **75** no. 4, (2015) 146, [arXiv:1411.4981](#) [[nucl-ex](#)].
- [191] **ALICE** Collaboration, B. B. Abelev *et al.*, “Charged jet cross sections and properties in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. D* **91** no. 11, (2015) 112012, [arXiv:1411.4969](#) [[nucl-ex](#)].
- [192] **ALICE** Collaboration, B. B. Abelev *et al.*, “Production of inclusive $\Upsilon(1S)$ and $\Upsilon(2S)$ in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **740** (2015) 105–117, [arXiv:1410.2234](#) [[nucl-ex](#)].
- [193] **ALICE** Collaboration, B. B. Abelev *et al.*, “Event-by-event mean p_T fluctuations in pp and Pb-Pb collisions at the LHC,” *Eur. Phys. J. C* **74** no. 10, (2014) 3077, [arXiv:1407.5530](#) [[nucl-ex](#)].

- [194] **ALICE** Collaboration, B. B. Abelev *et al.*, “Exclusive J/ψ photoproduction off protons in ultra-peripheral p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **113** no. 23, (2014) 232504, [arXiv:1406.7819 \[nucl-ex\]](#).
- [195] **ALICE** Collaboration, B. B. Abelev *et al.*, “Multiplicity dependence of jet-like two-particle correlation structures in p collisions at $\sqrt{s_{NN}}=5.02$ TeV,” *Phys. Lett. B* **741** (2015) 38–50, [arXiv:1406.5463 \[nucl-ex\]](#).
- [196] **ALICE** Collaboration, B. B. Abelev *et al.*, “Production of $\Sigma(1385)^\pm$ and $\Xi(1530)^0$ in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **75** no. 1, (2015) 1, [arXiv:1406.3206 \[nucl-ex\]](#).
- [197] **ALICE** Collaboration, B. B. Abelev *et al.*, “Multiparticle azimuthal correlations in p -Pb and Pb-Pb collisions at the CERN Large Hadron Collider,” *Phys. Rev. C* **90** no. 5, (2014) 054901, [arXiv:1406.2474 \[nucl-ex\]](#).
- [198] **ALICE** Collaboration, B. B. Abelev *et al.*, “Elliptic flow of identified hadrons in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **06** (2015) 190, [arXiv:1405.4632 \[nucl-ex\]](#).
- [199] **ALICE** Collaboration, B. B. Abelev *et al.*, “Suppression of $\Upsilon(1S)$ at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **738** (2014) 361–372, [arXiv:1405.4493 \[nucl-ex\]](#).
- [200] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of electrons from semileptonic heavy-flavor hadron decays in pp collisions at $\sqrt{s} = 2.76$ TeV,” *Phys. Rev. D* **91** no. 1, (2015) 012001, [arXiv:1405.4117 \[nucl-ex\]](#).
- [201] **ALICE** Collaboration, B. B. Abelev *et al.*, “Beauty production in pp collisions at $\sqrt{s} = 2.76$ TeV measured via semi-electronic decays,” *Phys. Lett. B* **738** (2014) 97–108, [arXiv:1405.4144 \[nucl-ex\]](#).
- [202] **ALICE** Collaboration, B. B. Abelev *et al.*, “Neutral pion production at midrapidity in pp and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Eur. Phys. J. C* **74** no. 10, (2014) 3108, [arXiv:1405.3794 \[nucl-ex\]](#).
- [203] **ALICE** Collaboration, B. B. Abelev *et al.*, “Suppression of $\psi(2S)$ production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *JHEP* **12** (2014) 073, [arXiv:1405.3796 \[nucl-ex\]](#).
- [204] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of prompt D -meson production in $p - Pb$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **113** no. 23, (2014) 232301, [arXiv:1405.3452 \[nucl-ex\]](#).
- [205] **ALICE** Collaboration, B. B. Abelev *et al.*, “Transverse momentum dependence of inclusive primary charged-particle production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Eur. Phys. J. C* **74** no. 9, (2014) 3054, [arXiv:1405.2737 \[nucl-ex\]](#).

- [206] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of visible cross sections in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV in van der Meer scans with the ALICE detector,” *JINST* **9** no. 11, (2014) P11003, [arXiv:1405.1849 \[nucl-ex\]](#).
- [207] **ALICE** Collaboration, B. B. Abelev *et al.*, “Azimuthal anisotropy of D meson production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **90** no. 3, (2014) 034904, [arXiv:1405.2001 \[nucl-ex\]](#).
- [208] **ALICE** Collaboration, B. B. Abelev *et al.*, “Freeze-out radii extracted from three-pion cumulants in pp, p and Pb collisions at the LHC,” *Phys. Lett. B* **739** (2014) 139–151, [arXiv:1404.1194 \[nucl-ex\]](#).
- [209] **ALICE** Collaboration, B. B. Abelev *et al.*, “ $K^*(892)^0$ and $\phi(1020)$ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **91** (2015) 024609, [arXiv:1404.0495 \[nucl-ex\]](#).
- [210] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of quarkonium production at forward rapidity in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **74** no. 8, (2014) 2974, [arXiv:1403.3648 \[nucl-ex\]](#).
- [211] **ALICE** Collaboration, B. B. Abelev *et al.*, “Performance of the ALICE Experiment at the CERN LHC,” *Int. J. Mod. Phys. A* **29** (2014) 1430044, [arXiv:1402.4476 \[nucl-ex\]](#).
- [212] **ALICE** Collaboration, B. B. Abelev *et al.*, “Production of charged pions, kaons and protons at large transverse momenta in pp and Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **736** (2014) 196–207, [arXiv:1401.1250 \[nucl-ex\]](#).
- [213] **ALICE** Collaboration, B. B. Abelev *et al.*, “Measurement of charged jet suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **03** (2014) 013, [arXiv:1311.0633 \[nucl-ex\]](#).
- [214] **ALICE** Collaboration, B. B. Abelev *et al.*, “Centrality, rapidity and transverse momentum dependence of J/ψ suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **734** (2014) 314–327, [arXiv:1311.0214 \[nucl-ex\]](#).
- [215] **ALICE** Collaboration, B. B. Abelev *et al.*, “Two- and three-pion quantum statistics correlations in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV at the CERN Large Hadron Collider,” *Phys. Rev. C* **89** no. 2, (2014) 024911, [arXiv:1310.7808 \[nucl-ex\]](#).
- [216] **ALICE** Collaboration, B. B. Abelev *et al.*, “ J/ψ production and nuclear effects in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *JHEP* **02** (2014) 073, [arXiv:1308.6726 \[nucl-ex\]](#).

- [217] **ALICE** Collaboration, B. B. Abelev *et al.*, “Multiplicity Dependence of Pion, Kaon, Proton and Lambda Production in p-Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **728** (2014) 25–38, [arXiv:1307.6796](#) [nucl-ex].
- [218] **ALICE** Collaboration, B. B. Abelev *et al.*, “ K_S^0 and Λ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **111** (2013) 222301, [arXiv:1307.5530](#) [nucl-ex].
- [219] **ALICE** Collaboration, B. B. Abelev *et al.*, “Multi-strange baryon production at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **728** (2014) 216–227, [arXiv:1307.5543](#) [nucl-ex].
[Erratum: *Phys.Lett.B* 734, 409–410 (2014)].
- [220] **ALICE** Collaboration, B. B. Abelev *et al.*, “Long-range angular correlations of π , K and p in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **726** (2013) 164–177, [arXiv:1307.3237](#) [nucl-ex].
- [221] **ALICE** Collaboration, B. Abelev *et al.*, “Multiplicity dependence of two-particle azimuthal correlations in pp collisions at the LHC,” *JHEP* **09** (2013) 049, [arXiv:1307.1249](#) [nucl-ex].
- [222] **ALICE** Collaboration, B. B. Abelev *et al.*, “Energy Dependence of the Transverse Momentum Distributions of Charged Particles in pp Collisions Measured by ALICE,” *Eur. Phys. J. C* **73** no. 12, (2013) 2662, [arXiv:1307.1093](#) [nucl-ex].
- [223] **ALICE** Collaboration, B. B. Abelev *et al.*, “Multiplicity dependence of the average transverse momentum in pp, p-Pb, and Pb-Pb collisions at the LHC,” *Phys. Lett. B* **727** (2013) 371–380, [arXiv:1307.1094](#) [nucl-ex].
- [224] **ALICE** Collaboration, B. Abelev *et al.*, “Directed Flow of Charged Particles at Midrapidity Relative to the Spectator Plane in Pb-Pb Collisions at $\sqrt{s_{NN}}=2.76$ TeV,” *Phys. Rev. Lett.* **111** no. 23, (2013) 232302, [arXiv:1306.4145](#) [nucl-ex].
- [225] **ALICE** Collaboration, E. Abbas *et al.*, “Performance of the ALICE VZERO system,” *JINST* **8** (2013) P10016, [arXiv:1306.3130](#) [nucl-ex].
- [226] **ALICE** Collaboration, B. Abelev *et al.*, “D meson elliptic flow in non-central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **111** (2013) 102301, [arXiv:1305.2707](#) [nucl-ex].
- [227] **ALICE** Collaboration, E. Abbas *et al.*, “Charmonium and e^+e^- pair photoproduction at mid-rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV,” *Eur. Phys. J. C* **73** no. 11, (2013) 2617, [arXiv:1305.1467](#) [nucl-ex].

- [228] **ALICE** Collaboration, E. Abbas *et al.*, “Mid-rapidity anti-baryon to baryon ratios in pp collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV measured by ALICE,” *Eur. Phys. J. C* **73** (2013) 2496, [arXiv:1305.1562](#) [nucl-ex].
- [229] **ALICE** Collaboration, E. Abbas *et al.*, “Centrality dependence of the pseudorapidity density distribution for charged particles in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **726** (2013) 610–622, [arXiv:1304.0347](#) [nucl-ex].
- [230] **ALICE** Collaboration, E. Abbas *et al.*, “J/Psi Elliptic Flow in Pb-Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **111** (2013) 162301, [arXiv:1303.5880](#) [nucl-ex].
- [231] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality dependence of π , K, p production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **88** (2013) 044910, [arXiv:1303.0737](#) [hep-ex].
- [232] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality determination of Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE,” *Phys. Rev. C* **88** no. 4, (2013) 044909, [arXiv:1301.4361](#) [nucl-ex].
- [233] **ALICE** Collaboration, B. Abelev *et al.*, “Charge correlations using the balance function in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **723** (2013) 267–279, [arXiv:1301.3756](#) [nucl-ex].
- [234] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of the inclusive differential jet cross section in pp collisions at $\sqrt{s} = 2.76$ TeV,” *Phys. Lett. B* **722** (2013) 262–272, [arXiv:1301.3475](#) [nucl-ex].
- [235] **ALICE** Collaboration, B. Abelev *et al.*, “Charged kaon femtoscopic correlations in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. D* **87** no. 5, (2013) 052016, [arXiv:1212.5958](#) [hep-ex].
- [236] **ALICE** Collaboration, B. Abelev *et al.*, “Long-range angular correlations on the near and away side in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **719** (2013) 29–41, [arXiv:1212.2001](#) [nucl-ex].
- [237] **ALICE** Collaboration, B. Abelev *et al.*, “Transverse momentum distribution and nuclear modification factor of charged particles in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **110** no. 8, (2013) 082302, [arXiv:1210.4520](#) [nucl-ex].
- [238] **ALICE** Collaboration, B. Abelev *et al.*, “Pseudorapidity density of charged particles in p + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. Lett.* **110** no. 3, (2013) 032301, [arXiv:1210.3615](#) [nucl-ex].
- [239] **ALICE** Collaboration, B. Abelev *et al.*, “Coherent J/ψ photoproduction in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **718** (2013) 1273–1283, [arXiv:1209.3715](#) [nucl-ex].

- [240] **ALICE** Collaboration, B. Abelev *et al.*, “Production of $K^*(892)^0$ and $\phi(1020)$ in pp collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **72** (2012) 2183, [arXiv:1208.5717 \[hep-ex\]](#).
- [241] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of inelastic, single- and double-diffraction cross sections in proton–proton collisions at the LHC with ALICE,” *Eur. Phys. J. C* **73** no. 6, (2013) 2456, [arXiv:1208.4968 \[hep-ex\]](#).
- [242] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality Dependence of Charged Particle Production at Large Transverse Momentum in Pb–Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **720** (2013) 52–62, [arXiv:1208.2711 \[hep-ex\]](#).
- [243] **ALICE** Collaboration, B. Abelev *et al.*, “Pion, Kaon, and Proton Production in Central Pb–Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **109** (2012) 252301, [arXiv:1208.1974 \[hep-ex\]](#).
- [244] **ALICE** Collaboration, B. Abelev *et al.*, “ D_s^+ meson production at central rapidity in proton–proton collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **718** (2012) 279–294, [arXiv:1208.1948 \[hep-ex\]](#).
- [245] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of electrons from beauty hadron decays in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **721** (2013) 13–23, [arXiv:1208.1902 \[hep-ex\]](#). [Erratum: *Phys.Lett.B* 763, 507–509 (2016)].
- [246] **ALICE** Collaboration, B. Abelev *et al.*, “Net-Charge Fluctuations in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **110** no. 15, (2013) 152301, [arXiv:1207.6068 \[nucl-ex\]](#).
- [247] **ALICE** Collaboration, B. Abelev *et al.*, “Charge separation relative to the reaction plane in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **110** no. 1, (2013) 012301, [arXiv:1207.0900 \[nucl-ex\]](#).
- [248] **ALICE** Collaboration, B. Abelev *et al.*, “ $K_s^0 - K_s^0$ correlations in pp collisions at $\sqrt{s} = 7$ TeV from the LHC ALICE experiment,” *Phys. Lett. B* **717** (2012) 151–161, [arXiv:1206.2056 \[hep-ex\]](#).
- [249] **ALICE** Collaboration, B. Abelev *et al.*, “Production of muons from heavy flavour decays at forward rapidity in pp and Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **109** (2012) 112301, [arXiv:1205.6443 \[hep-ex\]](#).
- [250] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of prompt J/ψ and beauty hadron production cross sections at mid-rapidity in pp collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **11** (2012) 065, [arXiv:1205.5880 \[hep-ex\]](#).

- [251] **ALICE** Collaboration, B. Abelev *et al.*, “Anisotropic flow of charged hadrons, pions and (anti-)protons measured at high transverse momentum in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV,” *Phys. Lett. B* **719** (2013) 18–28, [arXiv:1205.5761 \[nucl-ex\]](#).
- [252] **ALICE** Collaboration, B. Abelev *et al.*, “Neutral pion and η meson production in proton-proton collisions at $\sqrt{s} = 0.9$ TeV and $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **717** (2012) 162–172, [arXiv:1205.5724 \[hep-ex\]](#).
- [253] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of electrons from semileptonic heavy-flavour hadron decays in pp collisions at $\sqrt{s} = 7\text{TeV}$,” *Phys.Rev.D* **86**(2012)112007, *arXiv* : 1205.5423[*hep-ex*].
- [254] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of charm production at central rapidity in proton-proton collisions at $\sqrt{s} = 2.76$ TeV,” *JHEP* **07** (2012) 191, [arXiv:1205.4007 \[hep-ex\]](#).
- [255] **ALICE** Collaboration, B. Abelev *et al.*, “Transverse sphericity of primary charged particles in minimum bias proton-proton collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV,” *Eur. Phys. J. C* **72** (2012) 2124, [arXiv:1205.3963 \[hep-ex\]](#).
- [256] **ALICE** Collaboration, B. Abelev *et al.*, “Multi-strange baryon production in pp collisions at $\sqrt{s} = 7$ TeV with ALICE,” *Phys. Lett. B* **712** (2012) 309–318, [arXiv:1204.0282 \[nucl-ex\]](#).
- [257] **ALICE** Collaboration, B. Abelev *et al.*, “Inclusive J/ψ production in pp collisions at $\sqrt{s} = 2.76$ TeV,” *Phys. Lett. B* **718** (2012) 295–306, [arXiv:1203.3641 \[hep-ex\]](#). [Erratum: *Phys.Lett.B* 748, 472–473 (2015)].
- [258] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of the Cross Section for Electromagnetic Dissociation with Neutron Emission in Pb-Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **109** (2012) 252302, [arXiv:1203.2436 \[nucl-ex\]](#).
- [259] **ALICE** Collaboration, B. Abelev *et al.*, “Suppression of high transverse momentum D mesons in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **09** (2012) 112, [arXiv:1203.2160 \[nucl-ex\]](#).
- [260] **ALICE** Collaboration, B. Abelev *et al.*, “ J/ψ Production as a Function of Charged Particle Multiplicity in pp Collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **712** (2012) 165–175, [arXiv:1202.2816 \[hep-ex\]](#).
- [261] **ALICE** Collaboration, B. Abelev *et al.*, “ J/ψ suppression at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **109** (2012) 072301, [arXiv:1202.1383 \[hep-ex\]](#).

- [262] **ALICE** Collaboration, B. Abelev *et al.*, “Heavy flavour decay muon production at forward rapidity in proton collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **708** (2012) 265–275, [arXiv:1201.3791 \[hep-ex\]](#).
- [263] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of Event Background Fluctuations for Charged Particle Jet Reconstruction in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *JHEP* **03** (2012) 053, [arXiv:1201.2423 \[hep-ex\]](#).
- [264] **ALICE** Collaboration, B. Abelev *et al.*, “Upgrade of the ALICE Experiment: Letter Of Intent,” *J. Phys. G* **41** (2014) 087001.
- [265] **ALICE** Collaboration, B. Abelev *et al.*, “Light vector meson production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **710** (2012) 557–568, [arXiv:1112.2222 \[nucl-ex\]](#).
- [266] **ALICE** Collaboration, B. Abelev *et al.*, “Underlying Event measurements in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV with the ALICE experiment at the LHC,” *JHEP* **07** (2012) 116, [arXiv:1112.2082 \[hep-ex\]](#).
- [267] **ALICE** Collaboration, B. Abelev *et al.*, “Measurement of charm production at central rapidity in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *JHEP* **01** (2012) 128, [arXiv:1111.1553 \[hep-ex\]](#).
- [268] **ALICE** Collaboration, B. Abelev *et al.*, “ J/ψ polarization in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. Lett.* **108** (2012) 082001, [arXiv:1111.1630 \[hep-ex\]](#).
- [269] **ALICE** Collaboration, K. Aamodt *et al.*, “Particle-yield modification in jet-like azimuthal di-hadron correlations in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **108** (2012) 092301, [arXiv:1110.0121 \[nucl-ex\]](#).
- [270] **ALICE** Collaboration, K. Aamodt *et al.*, “Harmonic decomposition of two-particle angular correlations in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **708** (2012) 249–264, [arXiv:1109.2501 \[nucl-ex\]](#).
- [271] **ALICE** Collaboration, K. Aamodt *et al.*, “Higher harmonic anisotropic flow measurements of charged particles in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **107** (2011) 032301, [arXiv:1105.3865 \[nucl-ex\]](#).
- [272] **ALICE** Collaboration, K. Aamodt *et al.*, “Rapidity and transverse momentum dependence of inclusive J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Lett. B* **704** (2011) 442–455, [arXiv:1105.0380 \[hep-ex\]](#). [Erratum: *Phys. Lett. B* **718**, 692–698 (2012)].

- [273] **ALICE** Collaboration, K. Aamodt *et al.*, “Production of pions, kaons and protons in pp collisions at $\sqrt{s} = 900$ GeV with ALICE at the LHC,” *Eur. Phys. J. C* **71** (2011) 1655, [arXiv:1101.4110](#) [[hep-ex](#)].
- [274] **ALICE** Collaboration, K. Aamodt *et al.*, “Femtoscopy of pp collisions at $\sqrt{s} = 0.9$ and 7 TeV at the LHC with two-pion Bose-Einstein correlations,” *Phys. Rev. D* **84** (2011) 112004, [arXiv:1101.3665](#) [[hep-ex](#)].
- [275] **ALICE** Collaboration, K. Aamodt *et al.*, “Two-pion Bose-Einstein correlations in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **696** (2011) 328–337, [arXiv:1012.4035](#) [[nucl-ex](#)].
- [276] **ALICE** Collaboration, K. Aamodt *et al.*, “Strange particle production in proton-proton collisions at $\sqrt{s} = 0.9$ TeV with ALICE at the LHC,” *Eur. Phys. J. C* **71** (2011) 1594, [arXiv:1012.3257](#) [[hep-ex](#)].
- [277] **ALICE** Collaboration, K. Aamodt *et al.*, “Centrality dependence of the charged-particle multiplicity density at mid-rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **106** (2011) 032301, [arXiv:1012.1657](#) [[nucl-ex](#)].
- [278] **ALICE** Collaboration, K. Aamodt *et al.*, “Suppression of Charged Particle Production at Large Transverse Momentum in Central Pb-Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **696** (2011) 30–39, [arXiv:1012.1004](#) [[nucl-ex](#)].
- [279] **ALICE** Collaboration, K. Aamodt *et al.*, “Elliptic flow of charged particles in Pb-Pb collisions at 2.76 TeV,” *Phys. Rev. Lett.* **105** (2010) 252302, [arXiv:1011.3914](#) [[nucl-ex](#)].
- [280] **ALICE** Collaboration, K. Aamodt *et al.*, “Charged-particle multiplicity density at mid-rapidity in central Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. Lett.* **105** (2010) 252301, [arXiv:1011.3916](#) [[nucl-ex](#)].
- [281] **ALICE** Collaboration, K. Aamodt *et al.*, “Transverse momentum spectra of charged particles in proton-proton collisions at $\sqrt{s} = 900$ GeV with ALICE at the LHC,” *Phys. Lett. B* **693** (2010) 53–68, [arXiv:1007.0719](#) [[hep-ex](#)].
- [282] **ALICE** Collaboration, K. Aamodt *et al.*, “Two-pion Bose-Einstein correlations in pp collisions at $\sqrt{s} = 900$ GeV,” *Phys. Rev. D* **82** (2010) 052001, [arXiv:1007.0516](#) [[hep-ex](#)].
- [283] **ALICE** Collaboration, K. Aamodt *et al.*, “Midrapidity antiproton-to-proton ratio in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV measured by the ALICE experiment,” *Phys. Rev. Lett.* **105** (2010) 072002, [arXiv:1006.5432](#) [[hep-ex](#)].

- [284] **ALICE** Collaboration, K. Aamodt *et al.*, “Charged-particle multiplicity measurement in proton-proton collisions at $\sqrt{s} = 7$ TeV with ALICE at LHC,” *Eur. Phys. J. C* **68** (2010) 345–354, [arXiv:1004.3514 \[hep-ex\]](#).
- [285] **ALICE** Collaboration, K. Aamodt *et al.*, “Charged-particle multiplicity measurement in proton-proton collisions at $\sqrt{s} = 0.9$ and 2.36 TeV with ALICE at LHC,” *Eur. Phys. J. C* **68** (2010) 89–108, [arXiv:1004.3034 \[hep-ex\]](#).
- [286] **ALICE** Collaboration, K. Aamodt *et al.*, “Alignment of the ALICE Inner Tracking System with cosmic-ray tracks,” *JINST* **5** (2010) P03003, [arXiv:1001.0502 \[physics.ins-det\]](#).
- [287] **ALICE** Collaboration, K. Aamodt *et al.*, “First proton-proton collisions at the LHC as observed with the ALICE detector: Measurement of the charged particle pseudorapidity density at $s^{*}(1/2) = 900$ -GeV,” *Eur. Phys. J. C* **65** (2010) 111–125, [arXiv:0911.5430 \[hep-ex\]](#).
- [288] **PHENIX** Collaboration, A. Adare *et al.*, “Pseudorapidity Dependence of Particle Production and Elliptic Flow in Asymmetric Nuclear Collisions of p +Al, p +Au, d +Au, and ^3He +Au at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **121** no. 22, (2018) 222301, [arXiv:1807.11928 \[nucl-ex\]](#).
- [289] **PHENIX** Collaboration, C. Aidala *et al.*, “Production of π^0 and η mesons in Cu+Au collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. C* **98** no. 5, (2018) 054903, [arXiv:1805.04389 \[hep-ex\]](#).
- [290] **PHENIX** Collaboration, A. Adare *et al.*, “Beam-energy and centrality dependence of direct-photon emission from ultra-relativistic heavy-ion collisions,” *Submitted to: Phys. Rev. Lett.* (2018) , [arXiv:1805.04084 \[hep-ex\]](#).
- [291] **PHENIX** Collaboration, A. Adare *et al.*, “Low-momentum direct photon measurement in Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **98** no. 5, (2018) 054902, [arXiv:1805.04066 \[hep-ex\]](#).
- [292] **PHENIX** Collaboration, C. Aidala *et al.*, “Creating small circular, elliptical, and triangular droplets of quark-gluon plasma,” *Nature Phys.* **15** no. 3, (2019) 214–220, [arXiv:1805.02973 \[nucl-ex\]](#).
- [293] **PHENIX** Collaboration, A. Adare *et al.*, “Multiparticle azimuthal correlations for extracting event-by-event elliptic and triangular flow in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **99** no. 2, (2019) 024903, [arXiv:1804.10024 \[nucl-ex\]](#).
- [294] **PHENIX** Collaboration, A. Adare *et al.*, “Cross section and longitudinal single-spin asymmetry A_L for forward $W^\pm \rightarrow \mu^\pm \nu$ production in polarized $p+p$ collisions at $\sqrt{s} = 510$ GeV,” *Phys. Rev. D* **98** no. 3, (2018) 032007, [arXiv:1804.04181 \[hep-ex\]](#).

- [295] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of two-particle correlations with respect to second- and third-order event planes in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” [arXiv:1803.01749](#) [[hep-ex](#)].
- [296] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of emission angle anisotropy via long-range angular correlations with high p_T hadrons in d +Au and p + p collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **98** no. 1, (2018) 014912, [arXiv:1711.09003](#) [[hep-ex](#)].
- [297] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of mass-dependent azimuthal anisotropy in central p +Au, d +Au, and ^3He +Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **97** (2018) 064904, [arXiv:1710.09736](#) [[nucl-ex](#)].
- [298] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of ϕ -meson production at forward rapidity in $p + p$ collisions at $\sqrt{s} = 510$ GeV and its energy dependence from $\sqrt{s} = 200$ GeV to 7 TeV,” *Phys. Rev. D* **98** no. 9, (2018) 092006, [arXiv:1710.01656](#) [[hep-ex](#)].
- [299] **PHENIX** Collaboration, A. Adare *et al.*, “Lévy-stable two-pion Bose-Einstein correlations in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions,” *Phys. Rev. C* **97** no. 6, (2018) 064911, [arXiv:1709.05649](#) [[nucl-ex](#)].
- [300] **PHENIX** Collaboration, C. Aidala *et al.*, “Measurements of azimuthal anisotropy and charged-particle multiplicity in d +Au collisions at $\sqrt{s_{NN}} = 200, 62.4, 39,$ and 19.6 GeV,” *Phys. Rev. C* **96** no. 6, (2017) 064905, [arXiv:1708.06983](#) [[nucl-ex](#)].
- [301] **PHENIX** Collaboration, C. Aidala *et al.*, “Measurements of Multiparticle Correlations in $d + \text{Au}$ Collisions at 200, 62.4, 39, and 19.6 GeV and $p + \text{Au}$ Collisions at 200 GeV and Implications for Collective Behavior,” *Phys. Rev. Lett.* **120** no. 6, (2018) 062302, [arXiv:1707.06108](#) [[nucl-ex](#)].
- [302] **PHENIX** Collaboration, C. Aidala *et al.*, “Nuclear Dependence of the Transverse-Single-Spin Asymmetry for Forward Neutron Production in Polarized $p + A$ Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **120** no. 2, (2018) 022001, [arXiv:1703.10941](#) [[hep-ex](#)].
- [303] **PHENIX** Collaboration, C. Aidala *et al.*, “Cross section and transverse single-spin asymmetry of muons from open heavy-flavor decays in polarized p + p collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev. D* **95** no. 11, (2017) 112001, [arXiv:1703.09333](#) [[hep-ex](#)].
- [304] **PHENIX** Collaboration, C. Aidala *et al.*, “ B -meson production at forward and backward rapidity in $p + p$ and $\text{Cu} + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **96** no. 6, (2017) 064901, [arXiv:1702.01085](#) [[nucl-ex](#)].

- [305] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of e^+e^- pairs from open heavy flavor in $p+p$ and $d+A$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C96** no. 2, (2017) 024907, [arXiv:1702.01084](#) [nucl-ex].
- [306] **PHENIX** Collaboration, C. Aidala *et al.*, “Measurements of $B \rightarrow J/\psi$ at forward rapidity in $p+p$ collisions at $\sqrt{s} = 510$ GeV,” *Phys. Rev.* **D95** no. 9, (2017) 092002, [arXiv:1701.01342](#) [hep-ex].
- [307] **PHENIX** Collaboration, A. Adare *et al.*, “Angular decay coefficients of J/ψ mesons at forward rapidity from $p + p$ collisions at $\sqrt{s} = 510$ GeV,” *Phys. Rev.* **D95** (2017) 092003, [arXiv:1612.06807](#) [hep-ex].
- [308] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of the relative yields of $\psi(2S)$ to $\psi(1S)$ mesons produced at forward and backward rapidity in $p+p$, $p+Al$, $p+Au$, and ^3He+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C95** no. 3, (2017) 034904, [arXiv:1609.06550](#) [nucl-ex].
- [309] **PHENIX** Collaboration, A. Adare *et al.*, “Nonperturbative-transverse-momentum effects and evolution in dihadron and direct photon-hadron angular correlations in $p+p$ collisions at $\sqrt{s}=510$ GeV,” *Phys. Rev.* **D95** no. 7, (2017) 072002, [arXiv:1609.04769](#) [hep-ex].
- [310] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of double-helicity asymmetries in inclusive J/ψ production in longitudinally polarized $p + p$ collisions at $\sqrt{s} = 510$ GeV,” *Phys. Rev.* **D94** no. 11, (2016) 112008, [arXiv:1606.01815](#) [hep-ex].
- [311] **PHENIX** Collaboration, A. Adare *et al.*, “Inclusive cross section and double-helicity asymmetry for π^0 production at midrapidity in $p+p$ collisions at $\sqrt{s} = 510$ GeV,” *Phys. Rev.* **D93** no. 1, (2016) 011501, [arXiv:1510.02317](#) [hep-ex].
- [312] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of directed, elliptic, and triangular flow in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C94** no. 5, (2016) 054910, [arXiv:1509.07784](#) [nucl-ex].
- [313] **PHENIX** Collaboration, A. Adare *et al.*, “Azimuthally anisotropic emission of low-momentum direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C94** no. 6, (2016) 064901, [arXiv:1509.07758](#) [nucl-ex].
- [314] **PHENIX** Collaboration, A. Adare *et al.*, “Scaling properties of fractional momentum loss of high- p_T hadrons in nucleus-nucleus collisions at $\sqrt{s_{NN}}$ from 62.4 GeV to 2.76 TeV,” *Phys. Rev.* **C93** no. 2, (2016) 024911, [arXiv:1509.06735](#) [nucl-ex].

- [315] **PHENIX** Collaboration, A. Adare *et al.*, “Transverse energy production and charged-particle multiplicity at midrapidity in various systems from $\sqrt{s_{NN}} = 7.7$ to 200 GeV,” *Phys. Rev.* **C93** no. 2, (2016) 024901, [arXiv:1509.06727 \[nucl-ex\]](#).
- [316] **PHENIX** Collaboration, A. Adare *et al.*, “ ϕ meson production in the forward/backward rapidity region in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C93** no. 2, (2016) 024904, [arXiv:1509.06337 \[nucl-ex\]](#).
- [317] **PHENIX** Collaboration, A. Adare *et al.*, “Forward J/ψ production in U+U collisions at $\sqrt{s_{NN}}=193$ GeV,” *Phys. Rev.* **C93** no. 3, (2016) 034903, [arXiv:1509.05380 \[nucl-ex\]](#).
- [318] **PHENIX** Collaboration, A. Adare *et al.*, “Dielectron production in Au+Au collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev.* **C93** no. 1, (2016) 014904, [arXiv:1509.04667 \[nucl-ex\]](#).
- [319] **PHENIX** Collaboration, A. Adare *et al.*, “Single electron yields from semileptonic charm and bottom hadron decays in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C93** no. 3, (2016) 034904, [arXiv:1509.04662 \[nucl-ex\]](#).
- [320] **PHENIX** Collaboration, A. Adare *et al.*, “Centrality-dependent modification of jet-production rates in deuteron-gold collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. Lett.* **116** no. 12, (2016) 122301, [arXiv:1509.04657 \[nucl-ex\]](#).
- [321] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of elliptic and triangular flow in high-multiplicity $^3\text{He}+\text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **115** no. 14, (2015) 142301, [arXiv:1507.06273 \[nucl-ex\]](#).
- [322] **PHENIX** Collaboration, A. Adare *et al.*, “ ϕ meson production in $d+\text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C92** no. 4, (2015) 044909, [arXiv:1506.08181 \[nucl-ex\]](#).
- [323] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of higher cumulants of net-charge multiplicity distributions in Au+Au collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV,” *Phys. Rev.* **C93** no. 1, (2016) 011901, [arXiv:1506.07834 \[nucl-ex\]](#).
- [324] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of parity-violating spin asymmetries in W^\pm production at midrapidity in longitudinally polarized $p+p$ collisions,” *Phys. Rev.* **D93** no. 5, (2016) 051103, [arXiv:1504.07451 \[hep-ex\]](#).
- [325] **PHENIX** Collaboration, A. Adare *et al.*, “Systematic study of charged-pion and kaon femtoscopy in Au + Au collisions at $\sqrt{s_{NN}}=200$

- GeV,” *Phys. Rev.* **C92** no. 3, (2015) 034914, [arXiv:1504.05168 \[nucl-ex\]](#).
- [326] **PHENIX** Collaboration, A. Adare *et al.*, “An Upgrade Proposal from the PHENIX Collaboration,” [arXiv:1501.06197 \[nucl-ex\]](#).
- [327] **PHENIX** Collaboration, A. Adare *et al.*, “Systematic Study of Azimuthal Anisotropy in Cu+Cu and Au+Au Collisions at $\sqrt{s_{NN}} = 62.4$ and 200 GeV,” *Phys. Rev.* **C92** no. 3, (2015) 034913, [arXiv:1412.1043 \[nucl-ex\]](#).
- [328] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of the higher-order anisotropic flow coefficients for identified hadrons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C93** no. 5, (2016) 051902, [arXiv:1412.1038 \[nucl-ex\]](#).
- [329] **PHENIX** Collaboration, A. Adare *et al.*, “Beam-energy and system-size dependence of the space-time extent of the pion emission source produced in heavy ion collisions,” [arXiv:1410.2559 \[nucl-ex\]](#).
- [330] **PHENIX** Collaboration, A. Adare *et al.*, “Charged-pion cross sections and double-helicity asymmetries in polarized p+p collisions at $\sqrt{s}=200$ GeV,” *Phys. Rev.* **D91** no. 3, (2015) 032001, [arXiv:1409.1907 \[hep-ex\]](#).
- [331] **PHENIX** Collaboration, A. Adare *et al.*, “Search for dark photons from neutral meson decays in $p + p$ and $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C91** no. 3, (2015) 031901, [arXiv:1409.0851 \[nucl-ex\]](#).
- [332] **PHENIX** Collaboration, A. Adare *et al.*, “Cross section and transverse single-spin asymmetry of η mesons in $p^\uparrow + p$ collisions at $\sqrt{s} = 200$ GeV at forward rapidity,” *Phys. Rev.* **D90** no. 7, (2014) 072008, [arXiv:1406.3541 \[hep-ex\]](#).
- [333] **PHENIX** Collaboration, A. Adare *et al.*, “Cross section for $b\bar{b}$ production via dielectrons in d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C91** no. 1, (2015) 014907, [arXiv:1405.4004 \[nucl-ex\]](#).
- [334] **PHENIX** Collaboration, A. Adare *et al.*, “Low-mass vector-meson production at forward rapidity in $p+p$ collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D90** no. 5, (2014) 052002, [arXiv:1405.4260 \[hep-ex\]](#).
- [335] **PHENIX** Collaboration, A. Adare *et al.*, “Centrality dependence of low-momentum direct-photon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C91** no. 6, (2015) 064904, [arXiv:1405.3940 \[nucl-ex\]](#).

- [336] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of K_S^0 and K^{*0} in $p + p$, $d + Au$, and $Cu + Cu$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **90** no. 5, (2014) 054905, [arXiv:1405.3628](#) [nucl-ex].
- [337] **PHENIX** Collaboration, A. Adare *et al.*, “Heavy-quark production and elliptic flow in Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV,” *Phys. Rev. C* **91** no. 4, (2015) 044907, [arXiv:1405.3301](#) [nucl-ex].
- [338] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of long-range angular correlation and quadrupole anisotropy of pions and (anti)protons in central $d+Au$ collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. Lett.* **114** no. 19, (2015) 192301, [arXiv:1404.7461](#) [nucl-ex].
- [339] **PHENIX** Collaboration, N. N. Ajitanand *et al.*, “Comparison of the space-time extent of the emission source in $d+Au$ and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Nucl. Phys. A* **931** (2014) 1082–1087, [arXiv:1404.5291](#) [nucl-ex].
- [340] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of $\Upsilon(1S + 2S + 3S)$ production in $p + p$ and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **91** no. 2, (2015) 024913, [arXiv:1404.2246](#) [nucl-ex].
- [341] **PHENIX** Collaboration, C. Aidala *et al.*, “Nuclear matter effects on J/ψ production in asymmetric Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **90** no. 6, (2014) 064908, [arXiv:1404.1873](#) [nucl-ex].
- [342] **PHENIX** Collaboration, A. Adare *et al.*, “Inclusive double-helicity asymmetries in neutral-pion and eta-meson production in $\vec{p} + \vec{p}$ collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev. D* **90** no. 1, (2014) 012007, [arXiv:1402.6296](#) [hep-ex].
- [343] **PHENIX** Collaboration, A. Adare *et al.*, “Concept for an Electron Ion Collider (EIC) detector built around the BaBar solenoid,” [arXiv:1402.1209](#) [nucl-ex].
- [344] **PHENIX** Collaboration, A. Adare *et al.*, “Azimuthal-angle dependence of charged-pion-interferometry measurements with respect to second- and third-order event planes in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **112** no. 22, (2014) 222301, [arXiv:1401.7680](#) [nucl-ex].
- [345] **PHENIX** Collaboration, S. S. Adler *et al.*, “Transverse-energy distributions at midrapidity in $p+p$, $d+Au$, and Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ –200 GeV and implications for particle-production models,” *Phys. Rev. C* **89** no. 4, (2014) 044905, [arXiv:1312.6676](#) [nucl-ex].
- [346] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of transverse-single-spin asymmetries for midrapidity and forward-rapidity

- production of hadrons in polarized p+p collisions at $\sqrt{s}=200$ and 62.4 GeV,” *Phys. Rev. D* **90** no. 1, (2014) 012006, [arXiv:1312.1995](#) [hep-ex].
- [347] **PHENIX** Collaboration, A. Adare *et al.*, “Heavy-flavor electron-muon correlations in $p+p$ and $d+Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **89** no. 3, (2014) 034915, [arXiv:1311.1427](#) [nucl-ex].
- [348] **PHENIX** Collaboration, A. Adare *et al.*, “System-size dependence of open-heavy-flavor production in nucleus-nucleus collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. C* **90** no. 3, (2014) 034903, [arXiv:1310.8286](#) [nucl-ex].
- [349] **PHENIX** Collaboration, A. Adare *et al.*, “Centrality categorization for $R_{p(d)+A}$ in high-energy collisions,” *Phys. Rev. C* **90** no. 3, (2014) 034902, [arXiv:1310.4793](#) [nucl-ex].
- [350] **PHENIX** Collaboration, A. Adare *et al.*, “Cold-Nuclear-Matter Effects on Heavy-Quark Production at Forward and Backward Rapidity in $d+Au$ Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **112** no. 25, (2014) 252301, [arXiv:1310.1005](#) [nucl-ex].
- [351] **PHENIX** Collaboration, A. Adare *et al.*, “Azimuthal anisotropy of π^0 and η mesons in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **88** no. 6, (2013) 064910, [arXiv:1309.4437](#) [nucl-ex].
- [352] **PHENIX** Collaboration, A. Adare *et al.*, “Nuclear Modification of ψ_c and J/ψ Production in $d+Au$ Collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. Lett.* **111** no. 20, (2013) 202301, [arXiv:1305.5516](#) [nucl-ex].
- [353] **PHENIX** Collaboration, A. Adare *et al.*, “Spectra and ratios of identified particles in Au+Au and $d+Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **88** no. 2, (2013) 024906, [arXiv:1304.3410](#) [nucl-ex].
- [354] **PHENIX** Collaboration, A. Adare *et al.*, “Quadrupole Anisotropy in Dihadron Azimuthal Correlations in Central $d+Au$ Collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev. Lett.* **111** no. 21, (2013) 212301, [arXiv:1303.1794](#) [nucl-ex].
- [355] **PHENIX** Collaboration, A. Adare *et al.*, “Medium modification of jet fragmentation in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV measured in direct photon-hadron correlations,” *Phys. Rev. Lett.* **111** no. 3, (2013) 032301, [arXiv:1212.3323](#) [nucl-ex].
- [356] **PHENIX** Collaboration, A. Adare *et al.*, “ $v(1S + 2S + 3S)$ production in $d+Au$ and $p + p$ collisions at $\sqrt{s_{NN}} = 200$ GeV and cold-nuclear matter effects,” *Phys. Rev. C* **87** no. 4, (2013) 044909, [arXiv:1211.4017](#) [nucl-ex].

- [357] **PHENIX** Collaboration, A. Adare *et al.*, “Inclusive cross section and single transverse spin asymmetry for very forward neutron production in polarized p+p collisions at $\sqrt{s}=200\text{GeV}$,” *Phys. Rev.* **D88** no. 3, (2013) 032006, [arXiv:1209.3283](#) [nucl-ex].
- [358] **PHENIX** Collaboration, A. Adare *et al.*, “Double Spin Asymmetry of Electrons from Heavy Flavor Decays in $p + p$ Collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D87** no. 1, (2013) 012011, [arXiv:1209.3278](#) [hep-ex].
- [359] **PHENIX** Collaboration, A. Adare *et al.*, “Neutral pion production with respect to centrality and reaction plane in Au+Au collisions at $\sqrt{s_{NN}}=200$ GeV,” *Phys. Rev.* **C87** no. 3, (2013) 034911, [arXiv:1208.2254](#) [nucl-ex].
- [360] **PHENIX** Collaboration, A. Adare *et al.*, “ J/ψ suppression at forward rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 39$ and 62.4 GeV,” *Phys. Rev.* **C86** (2012) 064901, [arXiv:1208.2251](#) [nucl-ex].
- [361] **PHENIX** Collaboration, A. Adare *et al.*, “Cold-nuclear-matter effects on heavy-quark production in $d+\text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **109** no. 24, (2012) 242301, [arXiv:1208.1293](#) [nucl-ex].
- [362] **PHENIX** Collaboration, A. Adare *et al.*, “Direct photon production in $d+\text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C87** (2013) 054907, [arXiv:1208.1234](#) [nucl-ex].
- [363] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Measurement of Direct Photons in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **109** (2012) 152302, [arXiv:1205.5759](#) [nucl-ex].
- [364] **PHENIX** Collaboration, A. Adare *et al.*, “Direct-Photon Production in $p + p$ Collisions at $\sqrt{s} = 200$ GeV at Midrapidity,” *Phys. Rev.* **D86** (2012) 072008, [arXiv:1205.5533](#) [hep-ex].
- [365] **PHENIX** Collaboration, A. Adare *et al.*, “Evolution of π^0 suppression in Au+Au collisions from $\sqrt{s_{NN}} = 39$ to 200 GeV,” *Phys. Rev. Lett.* **109** (2012) 152301, [arXiv:1204.1526](#) [nucl-ex].
- [366] **PHENIX** Collaboration, A. Adare *et al.*, “Transverse-Momentum Dependence of the J/ψ Nuclear Modification in $d+\text{Au}$ Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C87** no. 3, (2013) 034904, [arXiv:1204.0777](#) [nucl-ex].
- [367] **PHENIX** Collaboration, A. Adare *et al.*, “Nuclear-Modification Factor for Open-Heavy-Flavor Production at Forward Rapidity in Cu+Cu Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C86** (2012) 024909, [arXiv:1204.0754](#) [nucl-ex].

- [368] **PHENIX** Collaboration, A. Adare *et al.*, “Deviation from quark-number scaling of the anisotropy parameter v_2 of pions, kaons, and protons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C85** (2012) 064914, [arXiv:1203.2644](#) [nucl-ex].
- [369] **PHENIX** Collaboration, A. Adare *et al.*, “Cross sections and double-helicity asymmetries of midrapidity inclusive charged hadrons in $p + p$ collisions at $\sqrt{s} = 62.4$ GeV,” *Phys. Rev.* **D86** (2012) 092006, [arXiv:1202.4020](#) [hep-ex].
- [370] **PHENIX** Collaboration, A. Adare *et al.*, “Suppression of back-to-back hadron pairs at forward rapidity in d +Au Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **107** (2011) 172301, [arXiv:1105.5112](#) [nucl-ex].
- [371] **PHENIX** Collaboration, A. Adare *et al.*, “Observation of direct-photon collective flow in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions,” *Phys. Rev. Lett.* **109** (2012) 122302, [arXiv:1105.4126](#) [nucl-ex].
- [372] **PHENIX** Collaboration, A. Adare *et al.*, “Measurements of Higher-Order Flow Harmonics in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **107** (2011) 252301, [arXiv:1105.3928](#) [nucl-ex].
- [373] **PHENIX** Collaboration, A. Adare *et al.*, “Production of ω mesons in $p + p$, d +Au, Cu + Cu , and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C84** (2011) 044902, [arXiv:1105.3467](#) [nucl-ex].
- [374] **PHENIX** Collaboration, A. Adare *et al.*, “Ground and excited charmonium state production in $p + p$ collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D85** (2012) 092004, [arXiv:1105.1966](#) [hep-ex].
- [375] **PHENIX** Collaboration, A. Adare *et al.*, “ J/ψ suppression at forward rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C84** (2011) 054912, [arXiv:1103.6269](#) [nucl-ex].
- [376] **PHENIX** Collaboration, A. Adare *et al.*, “Identified charged hadron production in $p + p$ collisions at $\sqrt{s} = 200$ and 62.4 GeV,” *Phys. Rev.* **C83** (2011) 064903, [arXiv:1102.0753](#) [nucl-ex].
- [377] **PHENIX** Collaboration, A. Adare *et al.*, “Azimuthal correlations of electrons from heavy-flavor decay with hadrons in p^+p and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C83** (2011) 044912, [arXiv:1011.1477](#) [nucl-ex].
- [378] **PHENIX** Collaboration, A. Adare *et al.*, “Suppression of away-side jet fragments with respect to the reaction plane in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C84** (2011) 024904, [arXiv:1010.1521](#) [nucl-ex].

- [379] **PHENIX** Collaboration, A. Adare *et al.*, “Cold Nuclear Matter Effects on J/ψ Yields as a Function of Rapidity and Nuclear Geometry in Deuteron-Gold Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **107** (2011) 142301, [arXiv:1010.1246](#) [nucl-ex].
- [380] **PHENIX** Collaboration, A. Adare *et al.*, “Cross section and double helicity asymmetry for η mesons and their comparison to neutral pion production in p+p collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D83** (2011) 032001, [arXiv:1009.6224](#) [hep-ex].
- [381] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of Transverse Single-Spin Asymmetries for J/ψ Production in Polarized $p + p$ Collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D82** (2010) 112008, [arXiv:1009.4864](#) [hep-ex]. [Erratum: *Phys. Rev.* **D86**, 099904(2012)].
- [382] **PHENIX** Collaboration, A. Adare *et al.*, “Event Structure and Double Helicity Asymmetry in Jet Production from Polarized $p + p$ Collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D84** (2011) 012006, [arXiv:1009.4921](#) [hep-ex].
- [383] **PHENIX** Collaboration, A. Adare *et al.*, “Cross Section and Parity Violating Spin Asymmetries of W^\pm Boson Production in Polarized $p + p$ Collisions at $\sqrt{s} = 500$ GeV,” *Phys. Rev. Lett.* **106** (2011) 062001, [arXiv:1009.0505](#) [hep-ex].
- [384] **PHENIX** Collaboration, A. Adare *et al.*, “Azimuthal anisotropy of neutral pion production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV: Path-length dependence of jet quenching and the role of initial geometry,” *Phys. Rev. Lett.* **105** (2010) 142301, [arXiv:1006.3740](#) [nucl-ex].
- [385] **PHENIX** Collaboration, A. Adare *et al.*, “High p_T direct photon and π^0 triggered azimuthal jet correlations and measurement of k_T for isolated direct photons in $p + p$ collisions at $\sqrt{s} = 200$ GeV,” *Phys. Rev.* **D82** (2010) 072001, [arXiv:1006.1347](#) [hep-ex].
- [386] **PHENIX** Collaboration, A. Adare *et al.*, “Transverse momentum dependence of meson suppression η suppression in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C82** (2010) 011902, [arXiv:1005.4916](#) [nucl-ex].
- [387] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of neutral mesons in p+p collisions at $\sqrt{s} = 200$ GeV and scaling properties of hadron production,” *Phys. Rev.* **D83** (2011) 052004, [arXiv:1005.3674](#) [hep-ex].
- [388] **PHENIX** Collaboration, A. Adare *et al.*, “Heavy Quark Production in $p + p$ and Energy Loss and Flow of Heavy Quarks in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev.* **C84** (2011) 044905, [arXiv:1005.1627](#) [nucl-ex].

- [389] **PHENIX** Collaboration, A. Adare *et al.*, “Nuclear modification factors of ϕ mesons in d +Au, Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200\text{GeV}$,” *Phys.Rev.C* **83**(2011)024909, *arXiv* : 1004.3532[nucl-ex].
- [390] **PHENIX** Collaboration, A. Adare *et al.*, “Elliptic and hexadecapole flow of charged hadrons in Au+Au collisions at $\sqrt{s_{NN}} = 200\text{ GeV}$,” *Phys. Rev. Lett.* **105** (2010) 062301, *arXiv*:1003.5586 [nucl-ex].
- [391] **PHENIX** Collaboration, A. Adare *et al.*, “Trends in Yield and Azimuthal Shape Modification in Dihadron Correlations in Relativistic Heavy Ion Collisions,” *Phys. Rev. Lett.* **104** (2010) 252301, *arXiv*:1002.1077 [nucl-ex].
- [392] **PHENIX** Collaboration, A. Adare *et al.*, “Transverse momentum dependence of J/psi polarization at midrapidity in p+p collisions at $\sqrt{s} = 200\text{-GeV}$,” *Phys. Rev.* **D82** (2010) 012001, *arXiv*:0912.2082 [hep-ex].
- [393] **PHENIX** Collaboration, A. Adare *et al.*, “Detailed measurement of the e^+e^- pair continuum in $p + p$ and Au+Au collisions at $\sqrt{s_{NN}} = 200\text{ GeV}$ and implications for direct photon production,” *Phys. Rev.* **C81** (2010) 034911, *arXiv*:0912.0244 [nucl-ex].
- [394] **PHENIX** Collaboration, A. Adare *et al.*, “Double-Helicity Dependence of Jet Properties from Dihadrons in Longitudinally Polarized $p + p$ Collisions at $\sqrt{s} = 200\text{ GeV}$,” *Phys. Rev.* **D81** (2010) 012002, *arXiv*:0910.1029 [hep-ex].
- [395] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Systematic Studies of Elliptic Flow Measurements in Au+Au Collisions at $\sqrt{s} = 200\text{-GeV}$,” *Phys. Rev.* **C80** (2009) 024909, *arXiv*:0905.1070 [nucl-ex].
- [396] **PHENIX** Collaboration, S. Afanasiev *et al.*, “High-pT π^0 Production with Respect to the Reaction Plane in Au + Au Collisions at $\sqrt{s_{NN}} = 200\text{-GeV}$,” *Phys. Rev.* **C80** (2009) 054907, *arXiv*:0903.4886 [nucl-ex].
- [397] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Kaon interferometric probes of space-time evolution in Au+Au collisions at $\sqrt{s_{NN}} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **103** (2009) 142301, *arXiv*:0903.4863 [nucl-ex].
- [398] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of Bottom versus Charm as a Function of Transverse Momentum with Electron-Hadron Correlations in p^+p Collisions at $\sqrt{s} = 200\text{ GeV}$,” *Phys. Rev. Lett.* **103** (2009) 082002, *arXiv*:0903.4851 [hep-ex].

- [399] **PHENIX** Collaboration, A. Adare *et al.*, “Photon-Hadron Jet Correlations in p+p and Au+Au Collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev.* **C80** (2009) 024908, [arXiv:0903.3399](#) [[nucl-ex](#)].
- [400] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Photoproduction of J/psi and of high mass e+e- in ultra-peripheral Au+Au collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Lett.* **B679** (2009) 321–329, [arXiv:0903.2041](#) [[nucl-ex](#)].
- [401] **PHENIX** Collaboration, A. Adare *et al.*, “Inclusive cross section and double helicity asymmetry for π^0 production in p^+p collisions at $\sqrt{s} = 62.4\text{ GeV}$,” *Phys. Rev.* **D79** (2009) 012003, [arXiv:0810.0701](#) [[hep-ex](#)].
- [402] **PHENIX** Collaboration, A. Adare *et al.*, “The Polarized gluon contribution to the proton spin from the double helicity asymmetry in inclusive π^0 production in polarized p + p collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **103** (2009) 012003, [arXiv:0810.0694](#) [[hep-ex](#)].
- [403] **PHENIX** Collaboration, A. Adare *et al.*, “Charged hadron multiplicity fluctuations in Au+Au and Cu+Cu collisions from $\sqrt{s_{NN}} = 22.5$ to 200 GeV,” *Phys. Rev.* **C78** (2008) 044902, [arXiv:0805.1521](#) [[nucl-ex](#)].
- [404] **PHENIX** Collaboration, D. J. Kim, “Heavy quark ‘energy loss’ and ‘flow’ in a QCD matter at RHIC,” *J. Phys.* **G35** (2008) 044024.
- [405] **PHENIX** Collaboration, A. Adare *et al.*, “Enhanced production of direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200\text{ GeV}$ and implications for the initial temperature,” *Phys. Rev. Lett.* **104** (2010) 132301, [arXiv:0804.4168](#) [[nucl-ex](#)].
- [406] **PHENIX** Collaboration, A. Adare *et al.*, “Dilepton mass spectra in p+p collisions at $s^{*}(1/2) = 200\text{-GeV}$ and the contribution from open charm,” *Phys. Lett.* **B670** (2009) 313–320, [arXiv:0802.0050](#) [[hep-ex](#)].
- [407] **PHENIX** Collaboration, A. Adare *et al.*, “Onset of π^0 Suppression Studied in Cu+Cu Collisions at $s_{NN}=22.4, 62.4$, and 200 GeV,” *Phys. Rev. Lett.* **101** (2008) 162301, [arXiv:0801.4555](#) [[nucl-ex](#)].
- [408] **PHENIX** Collaboration, A. Adare *et al.*, “Dihadron azimuthal correlations in Au+Au collisions at $s(NN)^{(1/2)} = 200\text{-GeV}$,” *Phys. Rev.* **C78** (2008) 014901, [arXiv:0801.4545](#) [[nucl-ex](#)].
- [409] **PHENIX** Collaboration, A. Adare *et al.*, “Suppression pattern of neutral pions at high transverse momentum in Au + Au collisions at $s(NN)^{(1/2)} = 200\text{-GeV}$ and constraints on medium transport coefficients,” *Phys. Rev. Lett.* **101** (2008) 232301, [arXiv:0801.4020](#) [[nucl-ex](#)].

- [410] **PHENIX** Collaboration, A. Adare *et al.*, “Quantitative Constraints on the Opacity of Hot Partonic Matter from Semi-Inclusive Single High Transverse Momentum Pion Suppression in Au+Au collisions at $\sqrt{s(NN)} = 200\text{-GeV}$,” *Phys. Rev.* **C77** (2008) 064907, [arXiv:0801.1665 \[nucl-ex\]](#).
- [411] **PHENIX** Collaboration, A. Adare *et al.*, “J/psi Production in $\sqrt{s(NN)} = 200\text{-GeV}$ Cu+Cu Collisions,” *Phys. Rev. Lett.* **101** (2008) 122301, [arXiv:0801.0220 \[nucl-ex\]](#).
- [412] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Source breakup dynamics in Au+Au Collisions at $\sqrt{s(NN)} = 200\text{-GeV}$ via three-dimensional two-pion source imaging,” *Phys. Rev. Lett.* **100** (2008) 232301, [arXiv:0712.4372 \[nucl-ex\]](#).
- [413] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Particle-species dependent modification of jet-induced correlations in Au+Au collisions at $\sqrt{s(NN)} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **101** (2008) 082301, [arXiv:0712.3033 \[nucl-ex\]](#).
- [414] **PHENIX** Collaboration, A. Adare *et al.*, “Cold Nuclear Matter Effects on J/Psi as Constrained by Deuteron-Gold Measurements at $\sqrt{s(NN)} = 200\text{-GeV}$,” *Phys. Rev.* **C77** (2008) 024912, [arXiv:0903.4845 \[nucl-ex\]](#). [Erratum: *Phys. Rev.* **C79**, 059901(2009)].
- [415] **PHENIX** Collaboration, S. S. Adler *et al.*, “Centrality dependence of charged hadron production in deuteron + gold and nucleon + gold collisions at $\sqrt{s(NN)} = 200\text{-GeV}$,” *Phys. Rev.* **C77** (2008) 014905, [arXiv:0708.2416 \[nucl-ex\]](#).
- [416] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Enhancement of the dielectron continuum in $\sqrt{s(NN)} = 200\text{-GeV}$ Au+Au collisions,” [arXiv:0706.3034 \[nucl-ex\]](#).
- [417] **PHENIX** Collaboration, A. Adare *et al.*, “Transverse momentum and centrality dependence of dihadron correlations in Au+Au collisions at $\sqrt{s(NN)} = 200\text{-GeV}$: Jet-quenching and the response of partonic matter,” *Phys. Rev.* **C77** (2008) 011901, [arXiv:0705.3238 \[nucl-ex\]](#).
- [418] **PHENIX** Collaboration, A. Adare *et al.*, “Inclusive cross-section and double helicity asymmetry for π^0 production in $p + p$ collisions at $\sqrt{s} = 200\text{-GeV}$: Implications for the polarized gluon distribution in the proton,” *Phys. Rev.* **D76** (2007) 051106, [arXiv:0704.3599 \[hep-ex\]](#).
- [419] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of density correlations in pseudorapidity via charged particle multiplicity fluctuations in Au+Au collisions at $\sqrt{s(NN)} = 200\text{-GeV}$,” *Phys. Rev.* **C76** (2007) 034903, [arXiv:0704.2894 \[nucl-ex\]](#).

- [420] **PHENIX** Collaboration, S. Afanasiev *et al.*, “Elliptic flow for phi mesons and (anti)deuterons in Au + Au collisions at $s(NN)^{1/2} = 200$ -GeV,” *Phys. Rev. Lett.* **99** (2007) 052301, [arXiv:nuc1-ex/0703024](#) [NUCL-EX].
- [421] **PHENIX** Collaboration, S. S. Adler *et al.*, “Production of omega mesons at Large Transverse Momenta in p + p and d + Au Collisions at $s^{1/2}(NN) = 200$ -GeV,” *Phys. Rev.* **C75** (2007) 051902, [arXiv:nuc1-ex/0611031](#) [nuc1-ex].
- [422] **PHENIX** Collaboration, A. Adare *et al.*, “ J/ψ Production vs Centrality, Transverse Momentum, and Rapidity in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **98** (2007) 232301, [arXiv:nuc1-ex/0611020](#) [nuc1-ex].
- [423] **PHENIX** Collaboration, A. Adare *et al.*, “System Size and Energy Dependence of Jet-Induced Hadron Pair Correlation Shapes in Cu+Cu and Au+Au Collisions at $s(NN)^{1/2} = 200$ and 62.4-GeV,” *Phys. Rev. Lett.* **98** (2007) 232302, [arXiv:nuc1-ex/0611019](#) [nuc1-ex].
- [424] **PHENIX** Collaboration, A. Adare *et al.*, “Energy Loss and Flow of Heavy Quarks in Au+Au Collisions at $s(NN)^{1/2} = 200$ -GeV,” *Phys. Rev. Lett.* **98** (2007) 172301, [arXiv:nuc1-ex/0611018](#) [nuc1-ex].
- [425] **PHENIX** Collaboration, A. Adare *et al.*, “Correlated Production of p and anti-p in Au+Au Collisions at $s(NN)^{1/2} = 200$ -GeV,” *Phys. Lett.* **B649** (2007) 359–369, [arXiv:nuc1-ex/0611016](#) [nuc1-ex].
- [426] **PHENIX** Collaboration, A. Adare *et al.*, “ J/ψ production versus transverse momentum and rapidity in p^+p collisions at $\sqrt{s} = 200$ -GeV,” *Phys. Rev. Lett.* **98** (2007) 232002, [arXiv:hep-ex/0611020](#) [hep-ex].
- [427] **PHENIX** Collaboration, S. S. Adler *et al.*, “A Detailed Study of High-p(T) Neutral Pion Suppression and Azimuthal Anisotropy in Au+Au Collisions at $s(NN)^{1/2} = 200$ -GeV,” *Phys. Rev.* **C76** (2007) 034904, [arXiv:nuc1-ex/0611007](#) [nuc1-ex].
- [428] **PHENIX** Collaboration, S. S. Adler *et al.*, “High transverse momentum η meson production in p^+p , d^+ Au and Au+Au collisions at $S(NN)^{1/2} = 200$ -GeV,” *Phys. Rev.* **C75** (2007) 024909, [arXiv:nuc1-ex/0611006](#) [nuc1-ex].
- [429] **PHENIX** Collaboration, S. S. Adler *et al.*, “Centrality dependence of pi0 and eta production at large transverse momentum in $s(NN)^{1/2} = 200$ -GeV d+Au collisions,” *Phys. Rev. Lett.* **98** (2007) 172302, [arXiv:nuc1-ex/0610036](#) [nuc1-ex].
- [430] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of Single Muons at Forward Rapidity in p+p Collisions at $s^{1/2}(1/2) = 200$ -GeV

- and Implications for Charm Production,” *Phys. Rev.* **D76** (2007) 092002, [arXiv:hep-ex/0609032](#) [hep-ex].
- [431] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of direct photon production in p + p collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **98** (2007) 012002, [arXiv:hep-ex/0609031](#) [hep-ex].
 - [432] **PHENIX** Collaboration, A. Adare *et al.*, “Measurement of high-p(T) single electrons from heavy-flavor decays in p+p collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **97** (2006) 252002, [arXiv:hep-ex/0609010](#) [hep-ex].
 - [433] **PHENIX** Collaboration, A. Adare *et al.*, “Scaling properties of azimuthal anisotropy in Au+Au and Cu+Cu collisions at $s(\text{NN}) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **98** (2007) 162301, [arXiv:nucl-ex/0608033](#) [nucl-ex].
 - [434] **PHENIX** Collaboration, S. S. Adler *et al.*, “Evidence for a long-range component in the pion emission source in Au + Au collisions at $s(\text{NN})^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **98** (2007) 132301, [arXiv:nucl-ex/0605032](#) [nucl-ex].
 - [435] **PHENIX** Collaboration, S. S. Adler *et al.*, “Jet properties from dihadron correlations in $p^{+}p$ collisions at $\sqrt{s} = 200\text{-GeV}$,” *Phys. Rev.* **D74** (2006) 072002, [arXiv:hep-ex/0605039](#) [hep-ex].
 - [436] **PHENIX** Collaboration, S. S. Adler *et al.*, “Azimuthal angle correlations for rapidity separated hadron pairs in d + Au Collisions at $s(\text{NN})^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **96** (2006) 222301, [arXiv:nucl-ex/0603017](#) [nucl-ex].
 - [437] **PHENIX** Collaboration, S. S. Adler *et al.*, “Nuclear effects on hadron production in d = Au and p + p collisions at $s(\text{NN})^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev.* **C74** (2006) 024904, [arXiv:nucl-ex/0603010](#) [nucl-ex].
 - [438] **PHENIX** Collaboration, S. S. Adler *et al.*, “Improved measurement of double helicity asymmetry in inclusive midrapidity π^0 production for polarized p+p collisions at $s^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev.* **D73** (2006) 091102, [arXiv:hep-ex/0602004](#) [hep-ex].
 - [439] **PHENIX** Collaboration, S. S. Adler *et al.*, “Common suppression pattern of eta and π^0 mesons at high transverse momentum in Au+Au collisions at $S(\text{NN})^{*}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **96** (2006) 202301, [arXiv:nucl-ex/0601037](#) [nucl-ex].
 - [440] **PHENIX** Collaboration, K. Ozawa *et al.*, “Low mass dilepton production at RHIC energies,” *Eur. Phys. J.* **C43** (2005) 421–426.

- [441] **PHENIX** Collaboration, J. Velkovska *et al.*, “Can phi mesons give an answer to the baryon puzzle at RHIC?,” *Eur. Phys. J.* **C43** (2005) 317–322.
- [442] **PHENIX** Collaboration, H. Busching *et al.*, “Medium effects on high particle production measured with the PHENIX experiment,” *Eur. Phys. J.* **C43** (2005) 303–310.
- [443] **PHENIX** Collaboration, B. A. Cole *et al.*, “Differential probes of medium-induced energy loss,” *Eur. Phys. J.* **C43** (2005) 271–280.
- [444] **PHENIX** Collaboration, O. Drapier *et al.*, “Heavy flavor production in PHENIX,” *Eur. Phys. J.* **C43** (2005) 201–208.
- [445] **PHENIX** Collaboration, M. Rosati *et al.*, “Quarkonium production from d + Au to Au + Au collisions,” *Eur. Phys. J.* **C43** (2005) 173–178.
- [446] **PHENIX** Collaboration, S. S. Adler *et al.*, “Nuclear modification of electron spectra and implications for heavy quark energy loss in Au+Au collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **96** (2006) 032301, [arXiv:nucl-ex/0510047](#) [nucl-ex].
- [447] **PHENIX** Collaboration, S. S. Adler *et al.*, “Jet structure from dihadron correlations in d+Au collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev.* **C73** (2006) 054903, [arXiv:nucl-ex/0510021](#) [nucl-ex].
- [448] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of identified π^0 and inclusive photon v_2 and implication to the direct photon production in $s(\text{NN})^{1/2} = 200\text{-GeV}$ Au+Au collisions,” *Phys. Rev. Lett.* **96** (2006) 032302, [arXiv:nucl-ex/0508019](#) [nucl-ex].
- [449] **PHENIX** Collaboration, S. S. Adler *et al.*, “Single electrons from heavy flavor decays in p+p collisions at $s^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **96** (2006) 032001, [arXiv:hep-ex/0508034](#) [hep-ex].
- [450] **PHENIX** Collaboration, S. S. Adler *et al.*, “J/psi production and nuclear effects for d+Au and p+p collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **96** (2006) 012304, [arXiv:nucl-ex/0507032](#) [nucl-ex].
- [451] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of transverse single-spin asymmetries for mid-rapidity production of neutral pions and charged hadrons in polarized p+p collisions at $s^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **95** (2005) 202001, [arXiv:hep-ex/0507073](#) [hep-ex].
- [452] **PHENIX** Collaboration, S. S. Adler *et al.*, “Dense-Medium Modifications to Jet-Induced Hadron Pair Distributions in Au+Au Collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **97** (2006) 052301, [arXiv:nucl-ex/0507004](#) [nucl-ex].

- [453] **PHENIX** Collaboration, S. S. Adler *et al.*, “Centrality dependence of direct photon production in $s(NN)^{**}(1/2) = 200\text{-GeV}$ Au + Au collisions,” *Phys. Rev. Lett.* **94** (2005) 232301, [arXiv:nucl-ex/0503003](#) [nucl-ex].
- [454] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of single electron event anisotropy in Au+Au collisions at $s(NN)^{**}(1/2) = 200\text{-GeV}$,” *Phys. Rev.* **C72** (2005) 024901, [arXiv:nucl-ex/0502009](#) [nucl-ex].
- [455] **PHENIX** Collaboration, S. S. Adler *et al.*, “Mid-rapidity direct-photon production in p^+p collisions at $\sqrt{s} = 200\text{-GeV}$,” *Phys. Rev.* **D71** (2005) 071102, [arXiv:hep-ex/0502006](#) [hep-ex].
- [456] **PHENIX** Collaboration, S. S. Adler *et al.*, “Nuclear modification factors for hadrons at forward and backward rapidities in deuteron-gold collisions at $s(NN)^{**}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **94** (2005) 082302, [arXiv:nucl-ex/0411054](#) [nucl-ex].
- [457] **PHENIX** Collaboration, S. S. Adler *et al.*, “Saturation of azimuthal anisotropy in Au + Au collisions at $s(NN)^{**}(1/2) = 62\text{-GeV}$ to 200-GeV ,” *Phys. Rev. Lett.* **94** (2005) 232302, [arXiv:nucl-ex/0411040](#) [nucl-ex].
- [458] **PHENIX** Collaboration, S. S. Adler *et al.*, “Production of phi mesons at mid-rapidity in $s(NN)^{**}(1/2) = 200\text{-GeV}$ Au+Au collisions at RHIC,” *Phys. Rev.* **C72** (2005) 014903, [arXiv:nucl-ex/0410012](#) [nucl-ex].
- [459] **PHENIX** Collaboration, K. Adcox *et al.*, “Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX collaboration,” *Nucl. Phys.* **A757** (2005) 184–283, [arXiv:nucl-ex/0410003](#) [nucl-ex].
- [460] **PHENIX** Collaboration, S. S. Adler *et al.*, “Centrality dependence of charm production from single electrons measurement in Au + Au collisions at $s(NN)^{**}(1/2) = 200\text{-GeV}$,” *Phys. Rev. Lett.* **94** (2005) 082301, [arXiv:nucl-ex/0409028](#) [nucl-ex].
- [461] **PHENIX** Collaboration, S. S. Adler *et al.*, “Systematic studies of the centrality and $s(NN)^{**}(1/2)$ dependence of the $d E(T) / d \eta$ and $d(N(\text{ch}) / d \eta$ in heavy ion collisions at mid-rapidity,” *Phys. Rev.* **C71** (2005) 034908, [arXiv:nucl-ex/0409015](#) [nucl-ex]. [Erratum: *Phys. Rev.* **C71**, 049901(2005)].
- [462] **PHENIX** Collaboration, S. S. Adler *et al.*, “Jet structure of baryon excess in Au + Au collisions at $s(NN)^{**}(1/2) = 200\text{-GeV}$,” *Phys. Rev.* **C71** (2005) 051902, [arXiv:nucl-ex/0408007](#) [nucl-ex].

- [463] **PHENIX** Collaboration, S. S. Adler *et al.*, “Deuteron and antideuteron production in Au + Au collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **94** (2005) 122302, [arXiv:nucl-ex/0406004](#) [[nucl-ex](#)].
- [464] **PHENIX** Collaboration, S. S. Adler *et al.*, “Double helicity asymmetry in inclusive mid-rapidity π^0 production for polarized p + p collisions at $s^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **93** (2004) 202002, [arXiv:hep-ex/0404027](#) [[hep-ex](#)].
- [465] **PHENIX** Collaboration, S. S. Adler *et al.*, “Bose-Einstein correlations of charged pion pairs in Au + Au collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **93** (2004) 152302, [arXiv:nucl-ex/0401003](#) [[nucl-ex](#)].
- [466] **PHENIX** Collaboration, S. S. Adler *et al.*, “Measurement of nonrandom event by event fluctuations of average transverse momentum in $s(\text{NN})^{1/2} = 200\text{-GeV}$ Au+Au and p+p collisions,” *Phys. Rev. Lett.* **93** (2004) 092301, [arXiv:nucl-ex/0310005](#) [[nucl-ex](#)].
- [467] **PHENIX** Collaboration, H. Akikawa *et al.*, “PHENIX muon arms,” *Nucl. Instrum. Meth.* **A499** (2003) 537–548.
- [468] **PHENIX** Collaboration, K. Adcox *et al.*, “PHENIX detector overview,” *Nucl. Instrum. Meth.* **A499** (2003) 469–479.
- [469] **PHENIX** Collaboration, S. S. Adler *et al.*, “High p_T charged hadron suppression in Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200\text{ GeV}$,” *Phys. Rev. C* **69** (2004) 034910, [arXiv:nucl-ex/0308006](#) [[nucl-ex](#)].
- [470] **PHENIX** Collaboration, S. S. Adler *et al.*, “Identified charged particle spectra and yields in Au+Au collisions at $S(\text{NN})^{1/2} = 200\text{-GeV}$,” *Phys. Rev. C* **69** (2004) 034909, [arXiv:nucl-ex/0307022](#) [[nucl-ex](#)].
- [471] **PHENIX** Collaboration, K. Adcox *et al.*, “Single identified hadron spectra from $s(\text{NN})^{1/2} = 130\text{-GeV}$ Au+Au collisions,” *Phys. Rev. C* **69** (2004) 024904, [arXiv:nucl-ex/0307010](#) [[nucl-ex](#)].
- [472] **PHENIX** Collaboration, S. S. Adler *et al.*, “ J/ψ production from proton proton collisions at $\sqrt{s} = 200\text{-GeV}$,” *Phys. Rev. Lett.* **92** (2004) 051802, [arXiv:hep-ex/0307019](#) [[hep-ex](#)].
- [473] **PHENIX** Collaboration, S. S. Adler *et al.*, “Absence of suppression in particle production at large transverse momentum in $S(\text{NN})^{1/2} = 200\text{-GeV}$ d + Au collisions,” *Phys. Rev. Lett.* **91** (2003) 072303, [arXiv:nucl-ex/0306021](#) [[nucl-ex](#)].
- [474] **PHENIX** Collaboration, A. Drees *et al.*, “Heavy ion collisions at collider energies: Insights from PHENIX,” *Pramana* **60** (2003) 639–650.

- [475] **PHENIX** Collaboration, S. S. Adler *et al.*, “Scaling properties of proton and anti-proton production in $s(NN)^{1/2}$ 200-GeV Au+Au collisions,” *Phys. Rev. Lett.* **91** (2003) 172301, [arXiv:nucl-ex/0305036](#) [nucl-ex].
- [476] **PHENIX** Collaboration, S. S. Adler *et al.*, “J / ψ production in Au Au collisions at $s(NN)^{1/2} = 200$ -GeV at the Relativistic Heavy Ion Collider,” *Phys. Rev.* **C69** (2004) 014901, [arXiv:nucl-ex/0305030](#) [nucl-ex].
- [477] **PHENIX** Collaboration, S. S. Adler *et al.*, “Elliptic flow of identified hadrons in Au+Au collisions at $s(NN)^{1/2} = 200$ -GeV,” *Phys. Rev. Lett.* **91** (2003) 182301, [arXiv:nucl-ex/0305013](#) [nucl-ex].
- [478] **PHENIX** Collaboration, S. S. Adler *et al.*, “Suppressed π^0 production at large transverse momentum in central Au + Au collisions at $S(NN)^{1/2} = 200$ GeV,” *Phys. Rev. Lett.* **91** (2003) 072301, [arXiv:nucl-ex/0304022](#) [nucl-ex].
- [479] **PHENIX** Collaboration, S. S. Adler *et al.*, “Mid-rapidity neutral pion production in proton proton collisions at $\sqrt{s} = 200$ -GeV,” *Phys. Rev. Lett.* **91** (2003) 241803, [arXiv:hep-ex/0304038](#) [hep-ex].
- [480] **PHENIX** Collaboration, K. Adcox *et al.*, “Centrality dependence of the high $p(T)$ charged hadron suppression in Au+Au collisions at $s(NN)^{1/2} = 130$ -GeV,” *Phys. Lett.* **B561** (2003) 82–92, [arXiv:nucl-ex/0207009](#) [nucl-ex].
- [481] **PHENIX** Collaboration, K. Adcox *et al.*, “Measurement of the Lambda and anti-Lambda particles in Au+Au collisions at $s(NN)^{1/2} = 130$ -GeV,” *Phys. Rev. Lett.* **89** (2002) 092302, [arXiv:nucl-ex/0204007](#) [nucl-ex].
- [482] **PHENIX** Collaboration, K. Adcox *et al.*, “Flow measurements via two particle azimuthal correlations in Au+Au collisions at $s(NN)^{1/2} = 130$ -GeV,” *Phys. Rev. Lett.* **89** (2002) 212301, [arXiv:nucl-ex/0204005](#) [nucl-ex].
- [483] **PHENIX** Collaboration, K. Adcox *et al.*, “Event-by-event fluctuations in mean $p(T)$ and mean $e(T)$ in $s(NN)^{1/2} = 130$ -GeV Au+Au collisions,” *Phys. Rev.* **C66** (2002) 024901, [arXiv:nucl-ex/0203015](#) [nucl-ex].
- [484] **PHENIX** Collaboration, K. Adcox *et al.*, “Net charge fluctuations in Au+Au interactions at $s^{1/2} = 130$ -GeV,” *Phys. Rev. Lett.* **89** (2002) 082301, [arXiv:nucl-ex/0203014](#) [nucl-ex].
- [485] **PHENIX** Collaboration, K. Adcox *et al.*, “Measurement of single electrons and implications for charm production in Au+Au collisions at

- $s^{**}(1/2)(NN) = 130\text{-GeV}$,” *Phys. Rev. Lett.* **88** (2002) 192303, [arXiv:nuc1-ex/0202002](#) [nuc1-ex].
- [486] **PHENIX** Collaboration, K. Adcox *et al.*, “Transverse mass dependence of two pion correlations in Au+Au collisions at $S(NN)^{**}(1/2) = 130\text{-GeV}$,” *Phys. Rev. Lett.* **88** (2002) 192302, [arXiv:nuc1-ex/0201008](#) [nuc1-ex].
- [487] **PHENIX** Collaboration, K. Adcox *et al.*, “Centrality dependence of π^+ / π^- , K^+ / K^- , p and anti-p production from $s(NN)^{**}(1/2) = 130\text{-GeV}$ Au+Au collisions at RHIC,” *Phys. Rev. Lett.* **88** (2002) 242301, [arXiv:nuc1-ex/0112006](#) [nuc1-ex].
- [488] **PHENIX** Collaboration, K. Adcox *et al.*, “First results from RHIC-PHENIX,” *Pramana* **57** (2001) 355–369.
- [489] **PHENIX** Collaboration, K. Adcox *et al.*, “Suppression of hadrons with large transverse momentum in central Au+Au collisions at $\sqrt{s_{NN}} = 130\text{-GeV}$,” *Phys. Rev. Lett.* **88** (2002) 022301, [arXiv:nuc1-ex/0109003](#) [nuc1-ex].
- [490] **PHENIX** Collaboration, W. A. Zajc *et al.*, “Overview of PHENIX results from the first RHIC run,” *Nucl. Phys.* **A698** (2002) 39–53, [arXiv:nuc1-ex/0106001](#) [nuc1-ex].
- [491] **PHENIX** Collaboration, K. Adcox *et al.*, “Measurement of the mid-rapidity transverse energy distribution from $s(NN)^{**}(1/2) = 130\text{-GeV}$ Au + Au collisions at RHIC,” *Phys. Rev. Lett.* **87** (2001) 052301, [arXiv:nuc1-ex/0104015](#) [nuc1-ex].
- [492] **PHENIX** Collaboration, K. Adcox *et al.*, “Centrality dependence of charged particle multiplicity in Au - Au collisions at $S(NN)^{**}(1/2) = 130\text{-GeV}$,” *Phys. Rev. Lett.* **86** (2001) 3500–3505, [arXiv:nuc1-ex/0012008](#) [nuc1-ex].