

several years after FDA approval, and we excluded continuation-in-part and divisional patents because they may not have identical specifications. We extracted patents listed on these drugs in the FDA's *Approved Products With Therapeutic Equivalence Evaluations* (Orange Book) from January 1, 2000, to December 31, 2021, and categorized each patent as original or continuation using the Google Patents Public Data Sets on Google BigQuery. We calculated the ratio of original and continuation patents per brand-name drug approval during each year. A similar analysis was conducted for litigated patents, based on litigation brought by either the drug manufacturer or generic competitor. Litigated patents were identified using Lex Machina, a commercial database covering 2000-2022.

Analyses were descriptive and completed in Excel version 16.

**Results** | From 2000 to 2015, the FDA approved 1421 new brand-name drugs. Manufacturers listed with the FDA 3967 distinct patents on these drugs through 2021 (2110 [53%] original; 1857 [47%] continuation). The ratio of the number of FDA-listed patents per drug increased from 1.9 for those approved in 2000 to 3.2 for those approved in 2015 (68% relative increase). While the ratio of the number of original patents per approval increased by 15% from 1.3 for drugs approved in 2000 to 1.5 for drugs approved in 2015, the ratio of continuation patents increased 200% from 0.6 for drugs approved in 2000 to 1.8 for drugs approved in 2015 (Figure, A).

There were 1936 litigated patents (985 [51%] original; 951 [49%] continuation). While the ratio of the number of litigated original patents per approval increased by 63% from 0.38 for drugs approved in 2000 to 0.62 for drugs approved in 2015, the ratio of litigated continuation patents increased 213% from 0.22 for drugs approved in 2000 to 0.69 for drugs approved in 2015 (Figure, B).

**Discussion** | Brand-name drug manufacturers listed with the FDA an increasing number of continuation patents on drugs approved from 2000 to 2015. More continuation patents mean that generic firms seeking to challenge existing protections on brand-name drugs must contest and potentially litigate more patents. Continuation patents are typically invalidated at a higher rate than patents on active ingredients.<sup>2</sup> However, lawsuits brought by brand-name firms on patents listed with the FDA can earn 30-month stays on generic drug approval even if these lawsuits eventually fail. Study limitations include that the frequency of successful challenges on litigated continuation patents was not examined.

These findings suggest that continuation patents are becoming increasingly common in drug patent thickets, likely delaying or deterring generic competition,<sup>2,5</sup> and thus potentially contributing to delays in patient access to generic medications and increases in health care spending.

S. Sean Tu, JD, PhD

Aaron S. Kesselheim, MD, JD, MPH

Kathrine Wetherbee, BA

William B. Feldman, MD, DPhil, MPH

**Author Affiliations:** West Virginia University College of Law, Morgantown (Tu); Division of Pharmacoepidemiology and Pharmacoeconomics, Brigham and Women's Hospital, Boston, Massachusetts (Kesselheim, Feldman); Sanford School of Public Policy, Duke University, Durham, North Carolina (Wetherbee).

**Accepted for Publication:** June 8, 2023.

**Corresponding Author:** S. Sean Tu, JD, PhD, West Virginia University College of Law, 101 Law School Dr, Morgantown, WV 26506 (shine.tu@mail.wvu.edu).

**Author Contributions:** Drs Tu and Feldman had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Concept and design:* Tu, Kesselheim, Feldman.

*Acquisition, analysis, or interpretation of data:* All authors.

*Drafting of the manuscript:* Tu.

*Critical revision of the manuscript for important intellectual content:* All authors.

*Statistical analysis:* Tu, Wetherbee.

*Obtained funding:* Kesselheim.

*Administrative, technical, or material support:* Tu.

*Supervision:* Tu, Kesselheim, Feldman.

**Conflict of Interest Disclosures:** Dr Kesselheim reported serving as an expert witness on behalf of a class of individual plaintiffs in a case against Gilead relating to a new formulation of tenofovir disoproxil. Dr Feldman reported serving as an expert witness in litigation against inhaler manufacturers and as a consultant to the nonprofit organization Alosa Health and as a consultant to Aetion. No other disclosures were reported.

**Funding/Support:** This work was funded by the West Virginia University Hodges Research Grant. Dr Kesselheim's and Dr Feldman's research is funded by Arnold Ventures and the Commonwealth Fund and Dr Feldman's work by the National Heart, Lung, and Blood Institute (K08HL163246).

**Role of the Funder/Sponsor:** The above listed supporters had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

**Data Sharing Statement:** See the [Supplement](#).

**Additional Contributions:** We thank Ian Wetherbee, BS, Google, for help analyzing the BigQuery data sets. He received no compensation.

1. Tu SS, Sarpatwari A. A "method of use" to prevent generic and biosimilar market entry. *N Engl J Med*. 2023;388(6):483-485. doi:10.1056/NEJMp2216020
2. Tu SS, Lemley MA. What litigators can teach the Patent Office about pharmaceutical patents. *Wash Law Rev*. 2022;99:1673-1731. doi:10.2139/ssrn.3903513
3. March 24, 2023, letter to Kathi Vidal from Jodey Arrington (R, Texas), Michael Burgess (R, Texas), Darrell Issa (R, California), Lloyd Doggett (D, Texas), and Ann Kuster (D, New Hampshire). Accessed May 16, 2023. <https://arrington.house.gov/news/documentsingle.aspx?DocumentID=956>
4. April 26, 2023, letter to Kathi Vidal from Elizabeth Warren (D, Massachusetts) and Pramila Jayapal (D, Washington). Accessed May 16, 2023. <https://www.warren.senate.gov/imo/media/doc/2023.04.26%20Letter%20to%20USPTO%20re%20patent%20abuses1.pdf>
5. Google BigQuery. Accessed May 21, 2023. <https://cloud.google.com/bigquery/>

## COMMENT & RESPONSE

### Treating Coronary Artery Disease With Treat-to-Target or High-Intensity Statin

**To the Editor** A recent article<sup>1</sup> investigated whether a treat-to-target strategy was noninferior to a strategy of high-intensity statin use for 3-year clinical outcomes in patients with coronary artery disease. We have some concerns about this study's research design.

During the 3-year follow-up, the percentage of high-intensity statin treatment increased from 47% to 56% in the treat-to-target strategy group but decreased from 100% to 89% in the high-intensity statin treatment group. This means that almost half of the patients received high-intensity

statin treatment in the treat-to-target group. While the effect of statin treatment on reducing clinical events is not solely caused by suppression of low-density lipoprotein cholesterol (LDL-C) level, and the authors adjusted for confounders, genetic factors such as apolipoprotein E gene polymorphisms may contribute to coronary artery disease and type 2 diabetes.<sup>2,3</sup> Individual variations in the response to statin treatment may be closely associated with genetic factors, and the recommendation of a treat-to-target LDL-C strategy for patients with cardiovascular disease should be made after understanding the magnitude of genetic contribution to the ability of LDL-C reduction and prevention of cardiovascular events.

Additionally, this study included substantially more men (n = 3172) than women (n = 1228), and the authors did not comment on sex difference in this study.<sup>1</sup> The majority of women in this study were likely postmenopausal, and dyslipidemia increases over a woman's life span, with adverse changes occurring around menopause.<sup>4</sup>

Tomoyuki Kawada, MD, PhD

**Author Affiliation:** Department of Hygiene and Public Health, Nippon Medical School, Tokyo, Japan.

**Corresponding Author:** Tomoyuki Kawada, MD, PhD, Department of Hygiene and Public Health, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-Ku, Tokyo 113-8602, Japan ([kawada@nms.ac.jp](mailto:kawada@nms.ac.jp)).

**Conflict of Interest Disclosures:** None reported.

1. Hong SJ, Lee YJ, Lee SJ, et al; LODESTAR Investigators. Treat-to-target or high-intensity statin in patients with coronary artery disease: a randomized clinical trial. *JAMA*. 2023;329(13):1078-1087. doi:10.1001/jama.2023.2487
2. El-Lebedy D, Raslan HM, Mohammed AM. Apolipoprotein E gene polymorphism and risk of type 2 diabetes and cardiovascular disease. *Cardiovasc Diabetol*. 2016;15:12. doi:10.1186/s12933-016-0329-1
3. Liu S, Liu J, Weng R, Gu X, Zhong Z. Apolipoprotein E gene polymorphism and the risk of cardiovascular disease and type 2 diabetes. *BMC Cardiovasc Disord*. 2019;19(1):213. doi:10.1186/s12872-019-1194-0
4. Torosyan N, Visrodia P, Torbati T, Minissian MB, Shufelt CL. Dyslipidemia in midlife women: approach and considerations during the menopausal transition. *Maturitas*. 2022;166:14-20. doi:10.1016/j.maturitas.2022.08.001

**In Reply** We appreciate Dr Kawada's comments about our study, which demonstrated that a treat-to-target LDL-C strategy of 50 to 70 mg/dL as the goal was noninferior to a high-intensity statin therapy for the 3-year composite of death, myocardial infarction, stroke, or coronary revascularization.<sup>1</sup>

First, we agree with Kawada that individual variations in the response to statin therapy may be closely related to genetic factors, and there has been accumulating evidence regarding the use of pharmacogenomics in cardiovascular medicine.<sup>2,3</sup> However, currently there is no consensus or clinical guideline about adopting pharmacogenomic biomarkers to guide selection of a drug or dose in statin therapy in daily practice.<sup>3-5</sup>

Second, regarding sex difference in our study, there was no significant interaction between sex and statin therapy strategy (treat-to-target group vs high-intensity statin group) in terms of the primary end point (8.5% vs 9.1%; difference, -0.7% [95% CI, -2.7% to 1.3%] for men and 7.3% vs 7.6%; difference, -0.3% [95% CI, -3.2% to 2.7%] for women) as well as an

achievement of LDL-C level below 70 mg/dL at 3 years (58.1% vs 60.6%; difference, -2.5% [95% CI, -6.6% to 1.6%] for men and 58.6% vs 57.2%; difference, 1.3% [95% CI, -5.2% to 7.9%] for women). Although the proportion of women was relatively low in our study,<sup>1</sup> perhaps due to sex difference in epidemiology of coronary artery disease (a higher prevalence among men), it was higher compared with the previous randomized trial regarding intensive statin therapy.<sup>6</sup> However, we agree with Kawada that the effect of sex should be cautiously considered, and further research focusing on sex and statin therapy is warranted.

Sung-Jin Hong, MD  
Yong-Joon Lee, MD  
Myeong-Ki Hong, MD, PhD

**Author Affiliations:** Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea.

**Corresponding Author:** Myeong-Ki Hong, MD, PhD, Division of Cardiology, Severance Hospital, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, 03722 Seoul, South Korea ([mkhong61@yuhs.ac](mailto:mkhong61@yuhs.ac)).

**Conflict of Interest Disclosures:** Dr M.-K. Hong reported receiving grants from Samjin Pharmaceutical and Chong Kun Dang Pharmaceutical and personal fees from Medtronic, Edwards Lifesciences, and Viatrix Korea. No other disclosures were reported.

1. Hong SJ, Lee YJ, Lee SJ, et al; LODESTAR Investigators. Treat-to-target or high-intensity statin in patients with coronary artery disease: a randomized clinical trial. *JAMA*. 2023;329(13):1078-1087. doi:10.1001/jama.2023.2487
2. El-Lebedy D, Raslan HM, Mohammed AM. Apolipoprotein E gene polymorphism and risk of type 2 diabetes and cardiovascular disease. *Cardiovasc Diabetol*. 2016;15:12. doi:10.1186/s12933-016-0329-1
3. Magavern EF, Kaski JC, Turner RM, et al. The role of pharmacogenomics in contemporary cardiovascular therapy: a position statement from the European Society of Cardiology Working Group on Cardiovascular Pharmacotherapy. *Eur Heart J Cardiovasc Pharmacother*. 2022;8(1):85-99. doi:10.1093/ehjcvp/pvab018
4. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2019;73(24):e285-e350. doi:10.1016/j.jacc.2018.11.003
5. Mach F, Baigent C, Catapano AL, et al; ESC Scientific Document Group. 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J*. 2020;41(1):111-188. doi:10.1093/eurheartj/ehz455
6. LaRosa JC, Grundy SM, Waters DD, et al; Treating to New Targets (TNT) Investigators. Intensive lipid lowering with atorvastatin in patients with stable coronary disease. *N Engl J Med*. 2005;352(14):1425-1435. doi:10.1056/NEJMoa050461

## Treatment Effects of Therapeutic-Dose Heparin in Patients Hospitalized for COVID-19

**To the Editor** A recent study<sup>1</sup> of treatment effects of therapeutic-dose heparin in patients hospitalized for COVID-19 analyzed patients on the ward (treatment-dose heparin was efficacious) separately from critically ill patients (treatment-dose heparin was not efficacious). The authors described this trial design as "fortuitous."<sup>1</sup> However, we are concerned that the no-difference result in critically ill patients may have occurred because subcutaneous depot dosing of low-molecular-weight heparin (LMWH) has poor systemic exposure in critically ill patients, particularly in those receiving vasopressors.