
Rediscovering Relativity Is Not a Single Task

What It Takes for AI to Discover a Physical Theory

Dong Zhang¹

Can an AI system rediscover general relativity? This question is increasingly discussed in the AI community, but it is often framed too narrowly. Some accounts implicitly assume that rediscovery means reproducing Einstein’s field equations; others assume that a single batch of experimental data suffices to construct a theory; still others reduce scientific discovery to one mode of inference. We argue that rediscovering a physical theory is not a single task, but a layered sequence of distinct capabilities. Using the history of relativity as a case study, we reconstruct it not as a linear path from Newton to Einstein but as a branching theory space populated by Lorentzian, Poincaréan, scalar, vector, and geometric theories. We show that theory construction and experiment design were mutually generative rather than sequential, and that at several points the available evidence could not adjudicate among live candidates. On this basis we propose a decomposition of rediscovery into distinct capabilities, and three staged tests for AI systems in theoretical physics.

1. Introduction

Can AI rediscover general relativity? This question has recently become a benchmark for evaluating the scientific capabilities of AI systems. One influential proposal is Hassabis’s “Einstein test”: train a system whose knowledge is restricted to what was known in 1911 and ask whether it can independently arrive at general relativity, just as Einstein did in 1915. Only then, the argument goes, could we claim that the system possesses genuine general intelligence.¹ One such attempt trains models from scratch on corpora containing only pre-1900 texts, aggressively removing any traces of later concepts (Hla, 2026).

The motivation behind these efforts is sound: they try to hide the answer and reduce data contamination. But they still leave the central question underspecified. What counts as “rediscovering” relativity, and how should success or failure be evaluated along the way? This is the gap this paper addresses. In our view, rediscovering relativity is not about recovering a single answer; it is about reconstructing a historical theory space.

The notion of “rediscovery” itself is ambiguous. Does it mean reproducing Einstein’s field equations? Does it mean reconstructing the entire logical path from special relativity to general relativity? Or is reproducing only one crucial step sufficient? Different answers correspond to fundamentally

different capabilities, yet current discussions often arrive at yes-or-no conclusions without first distinguishing among them.

More importantly, the framework of “existing knowledge $\rightarrow$ theory” imagines history as if all experimental facts had already been collected and were simply waiting for a theory to explain them. The actual history was not shaped in this way. Maxwell’s equations predicted electromagnetic waves and incorporated light into electromagnetism, which in turn motivated the search for the ether wind. The null result of the Michelson–Morley experiment led to Lorentz’s contraction hypothesis. The experiments carried out between 1902 and 1904 were no longer testing for the ether wind, but rather the new predictions that the contraction hypothesis itself had generated. The history of relativity was not a process of “data $\rightarrow$ theory”; it was a cycle of

theory $\rightarrow$ experiment $\rightarrow$ theory.

From Maxwell to Lorentz, from Poincaré to Einstein, and later to the competing theories of relativistic gravity, the development of relativity was a continuous process of mutual interaction rather than a one-shot inference.

We therefore decompose the question into three increasingly demanding challenges:

1. Can AI construct and revise theories as new experimental evidence arrives, as Lorentz did?
2. Can AI elevate a collection of empirical regularities into a unifying principle, as Poincaré did?
3. Can AI construct the entire space of candidate relativistic theories of gravity, derive their predictions, and

¹Affiliation, City, Country. Correspondence to: Author Name <email@example.com>.

Preprint. August 4, 2026.

¹Demis Hassabis, remarks in “Fireside Chat: Sir Demis Hassabis and Prof Govindan Rangarajan,” Indian Institute of Science, 17 February 2026, moderated by Varun Mayya. Transcript available at the IISc EECS event page.

identify the experiments that could distinguish among them?

These three challenges are not repetitions of the same capability. The first concerns adaptive modifications driven by anomalies. The second concerns the abstraction of local patches into general principles. The third requires the construction of a hypothesis space, the analysis of the empirical consequences of competing theories, and the search for experiments capable of adjudicating among them.

Yet all three challenges ultimately encounter the same boundary: theory can determine which experiments are informative, but it cannot determine the outcome of those experiments through reasoning alone. The competing theories of relativistic gravity were ultimately separated by astronomical observations, and those observations had to come from the world outside the theory itself.

Our position is that rediscovering relativity requires reconstructing a historical theory space, repeated interactions between theory and experiment, and movement across different modes of reasoning. It cannot be reduced to induction, deduction, or a single act of abduction, nor can it be completed entirely on paper.

Section 2 reconstructs this theory space. Section 3 reviews three common ways of framing the problem. Section 4 decomposes rediscovery into distinct capabilities. Section 5 proposes three staged evaluations. Section 6 explains why the validation of physical theories must ultimately come from outside the theories themselves.

2. The Historical Theory Space of Relativity

2.1. Two Nineteenth-Century Situations

- Newtonian gravity: overwhelming success, including Laplace and Neptune. One residual anomaly, Mercury's perihelion, 43''/century out of a total precession of 5557'', a relative deviation below 10^{-4} .
- The patch programme: Vulcan, Le Verrier 1859, observationally exhausted by roughly 1900; Hall's 1894 modification of the inverse-square exponent to $r^{-2+\delta}$, which fits the residual exactly and which Newcomb favoured; solar oblateness. Point to establish: from Mercury alone, the optimal data-driven answer is Hall, not Einstein.
- Maxwell's equations and the ether. $c = 1/\sqrt{\epsilon_0\mu_0}$ as a property of a medium; non-covariance under Galilean transformations was taken as evidence for the ether, not as a crisis.
- Michelson–Morley 1887: predicted 0.4 fringe shift, observed bound 0.01. Null.

2.2. Lorentz, Poincaré, Einstein

- Lorentz's contraction hypothesis: the contracting object is the moving body, not the ether; the factor is computed from Maxwell's equations plus the assumption that molecular forces are electromagnetic in nature.
- Poincaré: reverses the direction of inference. Not "bodies contract, therefore the ether is undetectable" but "undetectability is a principle, therefore contraction follows." Proves the transformations form a group; identifies the invariant; corrects Lorentz's 1904 errors; names the transformation.
- Einstein: different starting point, the magnet-conductor asymmetry, which produces no observational discrepancy; two postulates replace the electron model; t' is time.
- Empirical equivalence: the three give identical numbers for every experiment, not only in 1905 but today. Lorentzian ether theory remains empirically equivalent to special relativity. This is structural, not a limitation of period instrumentation.

2.3. Theory and Experiment as Mutually Generative

- 1887–1902: no new ether-drift experiments. The problem was regarded as solved by the contraction hypothesis.
- The 1902–04 experiments, Rayleigh, Brace, Trouton–Noble, Morley–Miller, did not test for ether drift. They tested new predictions generated by the contraction hypothesis itself: birefringence in a contracted medium, torque on a charged capacitor, material-dependence of the contraction.
- All null. Lorentz thickens the auxiliary assumptions, including local time in 1895 and deformable electron and non-electromagnetic stabilising stresses in 1904.
- Point to establish: the experiments were called into existence by the theory. A single batch of data handed to a system removes the mechanism that actually drove the sequence.

2.4. Relativistic Gravity Before General Relativity

- The shared problem: Newtonian gravity is not Lorentz covariant.
- Poincaré 1905, vector, gravitational waves, computes a Mercury correction whose stated aim is compatibility, not explanation; Minkowski 1908; Abraham 1912; Mie 1912.

- Nordström 1912–14: scalar; self-consistent; Lorentz covariant; explicitly requires $m_i = m_g$; correct gravitational redshift. Predicts $-7''/\text{century}$ for Mercury and exactly zero light deflection, with conformally flat spacetime and conformally invariant Maxwell equations.
- Einstein 1907–1912 also used a scalar theory, variable $c(x)$; the 1911 deflection of $0.83''$. The metric arrives in 1912 via Grossmann, not by selection among candidate field types.
- The field-theoretic route: a spin-2 field on a flat background, coupled to $T_{\mu\nu}$ and to its own energy, sums to the Einstein equations with $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, Gupta 1954, Kraichnan 1955, Feynman 1962, Deser 1970. Point to establish: the 1919 eclipse adjudicated the field type, spin 0 vs spin 2, not the ontology, flat background vs curved spacetime. The latter has never been adjudicated.

3. Common Framings of Rediscovery

3.1. The Whig Framing

- The framing: rediscovery means arriving at Einstein’s theory. Operationally, an evaluation scores a system on whether it outputs the Einstein field equations, or the two postulates of special relativity.
- Where it appears: informal benchmark proposals; popular framings of the question; historical narratives that follow Einstein alone and omit Nordström, Poincaré, Abraham, Mie.
- What it presupposes: that at each historical moment there was a determinate correct answer available.
- Cite specific instances.

3.2. The One-Shot Framing

- The pipeline: experimental data $\rightarrow$ AI $\rightarrow$ physical theory. Keep the existing diagram.
- Its concrete form: “give a system all of physics up to 1905 and have it produce general relativity.”
- Where it appears: public statements by AI laboratory leadership; benchmark designs that supply a complete corpus in a single prompt.
- What it presupposes: that the evidence base is fixed and prior to theorising.

3.3. The Single-Mechanism Framing

- The framing: discovery reduces to one mode of inference.
- Its three current versions: compression/induction, Schmidhuber; deduction from given axioms, the formal-mathematics line, AlphaProof; a single abductive jump requiring embodied simulation, Zahavy 2026.
- What it presupposes: that one mechanism accounts for the whole episode.

4. Rediscovery Is Not One Task

- Problem recognition. Which facts require explanation? $m_i = m_g$ had zero residual and was known for two centuries. Mercury’s $43''$ invited parameter adjustment. The magnet–conductor asymmetry produced no discrepancy at all.
- Sequential theory construction. Build from incomplete evidence; revise as observations arrive. Include the 1913 fact here: the Entwurf failed the general covariance Einstein himself required and gave $18''$ for Mercury, while Nordström’s theory was complete and self-consistent. Any search pruned on intermediate performance discards Einstein’s branch.
- Principle formation. From accumulated compensations to a single principle; the group structure as the mathematical expression of “no further patch is possible.”
- Hypothesis-space construction. Generate multiple viable theories; avoid premature collapse onto one.
- Internal screening. Lorentz covariance, conservation, stability, Newtonian limit, and the one screening that works on paper: vector gravity gives repulsion between like charges.
- Recognising empirical equivalence. Distinguish mathematical difference from observational difference. Note that adjudication at the gravity stage settled the field type, not the ontology.
- Experimental discrimination. The decisive quantity is light deflection, $1.75''$ vs 0 . Redshift does not discriminate: Nordström’s is correct. Mercury requires non-linear corrections and is not a clean criterion.
- World-coupled adjudication.

5. Three Staged Tests

- These are conceptual tests, not benchmarks ready to run. The answers are present in any modern training

corpus, so success is uninformative; only failure is informative, and only counterfactual variants, altered source symmetry or altered spacetime dimension, restore any diagnostic value.

5.1. The Lorentz Test

- Input, staged: first Maxwell plus ether plus Michelson–Morley alone; then Rayleigh/Brace, Trouton–Noble, Morley–Miller.
- Tasks: construct an explanation; derive its new predictions; on receiving the second batch, decide whether to patch, abandon, or reformulate.
- What would count as success: deriving from one’s own hypothesis the predictions that the second batch will test; recognising that those predictions have been refuted; tracking the growth in auxiliary assumptions.

5.2. The Poincaré Test

- Input: the full set of null results and Maxwell’s equations. Withheld: the relativity principle, the Lorentz group, Einstein’s postulates.
- Tasks: find a principle that explains all results at once; derive the admissible transformations; establish closure.
- What would count as success: the group property, and an explicit statement of why a principle differs from a sequence of compensations.

5.3. The Gravity Test

- Input: Minkowski spacetime; Lorentz covariance; $\partial^\mu T_{\mu\nu} = 0$; recovery of Newtonian gravity in the static weak field. Withheld: equivalence principle, Riemannian geometry, spin-2 gravity, Einstein, general relativity.
- Tasks: enumerate candidate field types with justification; write couplings and field equations; compute Newtonian limit, perihelion precession, light deflection, redshift; separate paper-eliminable from observation-requiring candidates; name the decisive observation.
- What would count as success: the enumeration is complete, spin 0, 1, 2; vector is eliminated without computation; the decisive observable is identified as light deflection with the reason why redshift and perihelion do not serve.
- Table of the three candidates and their predictions.

6. Why Physics Must Remain Coupled to the World

- Formal proof certifies mathematical validity internally; AI for mathematics can in principle close its loop.
- Physics cannot. Simulation does not supply independent evidence: it executes the theory under test. Computing 1.75'' from general relativity and then citing that number as confirmation of general relativity is circular. The 1919 measurement supplied an input from outside the theory.
- The loop: theory $\rightarrow$ prediction $\rightarrow$ experiment $\rightarrow$ observation $\rightarrow$ revision.
- Execution may be automated, including self-driving laboratories and robotic observatories. The claim is not about who operates the instrument. The claim is that the validating information must originate outside the theory being tested.

7. Discussion and Conclusion

- Restate: the question is not whether a system reaches Einstein’s answer.
- The decisive test is whether it can reconstruct the space of possibilities, recognise what remains undecidable on paper, and determine which question must next be put to nature.
- Where the division of labour falls: enumeration and paper-elimination on one side, adjudication on the other.
- Limitations: a single case; the enumeration in Section 5.3 uses representation-theoretic tools unavailable before 1939; the tests are conceptual.

References

Hla, M. *Machina mirabilis*. <https://michaelhla.com/blog/machina-mirabilis.html>, March 2026. Online research note.

A. Appendix A

- A.1 The three candidate gravity theories: couplings, field equations, and derivations of the four observables.
- A.2 Timeline of experiments and theories, 1859–1919.

Table 1. Candidate relativistic gravity theories and discriminating predictions.

CANDIDATE TYPE	FIELD	MERCURY SION	PRECES-	LIGHT DEFLECTION	GRAVITATIONAL RED- SHIFT	OUTCOME
SPIN 0						
SPIN 1						
SPIN 2						
