TOWARDS A NERVE OF TYPE THEORIES

ZACK DOOLEY

1. Introduction

(syntax/semantics for type theory/higher cat. theory, history)

(Introduce necessary syntax, say something about the choice to not study syntax directly, syntactic category of a tt, initiality?)

(clean up) Dependent type theory is an alternative foundation for mathematics in which the basic judgments are not only that a term has a type, but that a type itself may depend on a surrounding context of assumptions. In the fragment we reference throughout this paper, a typical judgment has the form

$$\Gamma \vdash A \text{ type} \qquad \text{or,} \qquad \Gamma \vdash a : A,$$

where Γ is a context, i.e. a list of typed variables $\Gamma = (x_1 : X_1, \dots, x_n : X_n)$. The basic structural rules of type theory can be extended by the inclusion of type formation rules (is this the right term?) which allow us access to various constructions within our theory such as a unit type, denoted 1, dependent pair and function types, called Σ and Π types respectively, or the type of identities on a given type. These constructions are asserted by adding rules such as the following for Id-types to our theory.

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma, x, y : A \vdash \mathrm{Id}_A(x, y) \text{ type}} \text{ ID-FORM} \qquad \frac{\Gamma \vdash A \text{ type}}{\Gamma, x : A \vdash \mathrm{refl}_A(x) : \mathrm{Id}_A(x, x)} \text{ ID-INTRO}$$

$$\frac{\Gamma, x, y : A, u : \mathrm{Id}_A(x, y) \vdash C(x, y, u) \text{ type} \qquad \Gamma, z : A \vdash d(z) : C(z, z, \mathrm{refl}_A(z))}{\Gamma, x, y : A, u : \mathrm{Id}_A(x, y) \vdash J_{z.d}(x, y, u) : C(x, y, u)} \text{ ID-ELIM}$$

$$\frac{\Gamma, x, y : A, u : \mathrm{Id}_A(x, y) \vdash C(x, y, u) \text{ type} \qquad \Gamma, z : A \vdash d(z) : C(z, z, \mathrm{refl}_A(z))}{\Gamma, x : A \vdash J_{z.d}(x, x, \mathrm{refl}_A(x)) = d(x) : C(x, x, \mathrm{refl}_A(x))} \text{ ID-COMP}$$

A presentation for the structural rules of Martin-Löf Type Theory, which is what we will consider to be the most "basic" dependent type theory, along with the rules for other common type constructors can be found in [13, Appendix A].

A central part of the value of dependent type theory is that it naturally admits a syntax-semantics relationship with category theory. On the syntactic side, contexts and substitutions naturally organize into a category whose objects are contexts and whose morphisms, from a context Γ to a context Δ, are lists of terms of the types in Δ in the context Γ.

The relationship between dependent type theory and category theory is a central theme of this paper. On the syntactic side, contexts and substitutions naturally organize into a category whose objects are contexts and whose morphisms are context morphisms. On the semantic side, the corresponding categorical structure is that of a category equipped with a distinguished class of display maps closed under pullback. When the type theory has dependent sums and extensional dependent products, this semantic structure is captured by a locally cartesian closed category (LCCC): contexts become objects, types in context become pullback-stable families of objects, dependent sums correspond to left adjoints to pullback, and extensional dependent products correspond to right adjoints to pullback together with the usual extensionality behavior.

Extensional dependent type theory is therefore closely tied to the notion of a locally cartesian closed category. For example, if $\Gamma \vdash A$ type and $\Gamma, x : A \vdash B$ type, then the dependent product $\Gamma \vdash \Pi_{x:A} B$ type is modeled by the right adjoint to the pullback functor along the projection $p_A : \Gamma.A \to \Gamma$ in an LCCC. Similarly,

dependent sums $\Sigma_{x:A} B$ are modeled by the left adjoint to the same pullback. Identity types in the extensional setting impose additional coherence conditions that are reflected semantically by a choice of diagonals and reindexing maps in the model.

Rather than studying the syntax of dependent type theory in isolation, this paper emphasizes semantic models such as contextual categories. A contextual category is a categorical abstraction of the syntactic category of a type theory: it encodes contexts, types in context, terms as sections, and substitution in a purely algebraic way. Studying these semantic models has several advantages: - it gives an invariant notion of model independent of the particular presentation of syntax; - it produces a category of models with natural morphisms between them, which is essential for comparing theories and constructing homotopies; - it makes available categorical tools such as limits, adjoints, and model structures that are hard to see directly in raw syntax; - it isolates the structural content of the theory from the bureaucratic details of derivations and proof-term representations.

In particular, contextual categories can be described as models of an essentially algebraic theory, so the collection of all such models forms a well-behaved category. This allows us to study type theories by studying their semantics directly, rather than by manipulating syntactic derivations. When we pass to homotopical questions, this semantic perspective becomes even more important, because it lets us equip the category of models with a notion of weak equivalence and to relate type-theoretic equivalence to categorical equivalence.

The remainder of the paper therefore begins by fixing the minimal syntax and notation needed for our arguments, then explains how extensional dependent type theory corresponds to locally cartesian closed structure, and finally develops the semantic framework of contextual categories as the primary objects of study. This sets the stage for the later sections, where we build homotopical models and compare them with the syntactic presentations they are meant to represent.

2. Background

2.1. **Homotopy Theory.** One of the simplest ways to realize an abstract notion of homotopy theory is through categories with weak equivalences. A **category with weak equivalences** is a category C equipped with a wide subcategory $\mathcal{W}$, whose maps we call weak equivalences. We can think of the weak equivalences as the maps which are invertible up to homotopy (whatever that may mean in this instance) but not necessarily invertible in C. We can then construct the homotopy category $\mathrm{Ho}\, C$ by freely inverting all the maps in $\mathcal{W}$. While this category correctly realizes the homotopical structure that we are interested in, it is in general difficult to work with, so we work with it indirectly by considering the pair $(C, \mathcal{W})$. We call a functor between categories with weak equivalences which preserves weak equivalences **homotopical**, and we denote the category of categories with weak equivalences and homotopical functors as **weCat**.

Remark 2.1. The definition above is sometimes used for relative categories, while categories with weak equivalences require $\mathcal{W}$ to contain all isomorphisms and be closed under 2-out-of-3. For this paper, we just use the term categories with weak equivalences to match [11], where the distinction between the two is not particularly relevant.

The category **weCat** can itself be though of as a category with weak equivalences, where the weak equivalences are given by Dwyer-Kan equivalences (DK-equivalences). A DK-equivalence is a homotopical functor $F : C \to D$, which when we take the simplicial localization $L :$ **weCat** $\to$ **sCat** (which can be implemented in a number of ways e.g. see [3]), is essentially surjective and ∞-full and faithful, i.e. for all $x, y \in LC$ the simplicial map

$$LF_{x,y} : LC(x, y) \to LD(LFx, LFy)$$

is a weak homotopy equivalence (of simplicial sets).

Ultimately, we are interested in how categories with weak equivalences can present both homotopy theories and infinity categories, so a natural model of infinity categories for us to use is that of a quasicategory.

Definition 2.2. A simplicial set X is called a **quasicategory** if it satisfies the following lifting property:

$$\begin{tikzcd} \Lambda^k_n \arrow[r] \arrow[d] & X \\ \Delta_n \arrow[ur, dashed] & \end{tikzcd},$$

for all $n > 0$ and $0 < k < n$.

We denote the full subcategory of **sSet** spanned by quasicategories by **qCat**. We write $\tau_1 : \mathbf{sSet} \to \mathbf{Cat}$ for the adjoint to the nerve functor, and we let $\mathbf{qCat}_2$ denote the 2-category of quasicategories, where the hom-categories are given by $\mathbf{qCat}_2(C, D) := \tau_1(D^C)$.

To study the homotopy theory of quasicategories, we need a notion of weak equivalences which will be given by categorical equivalences of quasicategories. To define these, we first establish the notion of an $E[1]$-homotopy, where $E[1]$ is the nerve of the contractible groupoid on two objects. Two maps $f, g : X \to Y$ of simplicial sets are $E[1]$-**homotopic**, denoted $f \sim g$, if there exists a simplicial map $H : X \times E[1] \to Y$ whose restriction to $X \times \partial\Delta_1$ is $[f, g]$. A map $f : X \to Y$ is then a **categorical equivalence** if there exists a map $g : Y \to X$ such that $fg \sim 1_Y$ and $gf \sim 1_X$.

At the core of the relationship between type theories and categories is the notion that the type formers on theory should correspond to imposing more categorical structure on the semantic side. In the case that we are interested in, the categorical structure that we expect is that of a locally Cartesian closed infinity category, a notion analogous to that of a locally Cartesian closed category, but requires some setup for our particular choice of model.

We define the **join** of simplicial sets as the unique functor $\star : \mathbf{sSet} \times \mathbf{sSet} \to \mathbf{sSet}$ with natural transformations $K \to K \star L \leftarrow L$ such that $\Delta_m \star \Delta_n \cong \Delta_{m+n+1}$, naturally in both m and n, and both functors $- \star L : \mathbf{sSet} \to L \downarrow \mathbf{sSet}$ and $K \star - : \mathbf{sSet} \to K \downarrow \mathbf{sSet}$ preserves colimits. By the adjoint functor theorem, $- \star L$ admits a right adjoint which takes a map of simplicial sets $X \to K$ to the simplicial set $K \downarrow$ defined by:

$$(K \downarrow X)_n = \{Y : \Delta_n \star L \to K \mid Y|_L = X\}.$$

Instantiating this definition with a 0-simplex $x : \Delta_0 \to C$, we obtain a notion of a **slice quasicategory**

$$(C \downarrow x)_n = \{X : \Delta_{n+1} \to C \mid X_{\Delta^{n+1}} = x\}.$$

Since the canonical map $C \downarrow x \to C$ lifts against inner horn inclusions, we get that $C \downarrow x$ is a quasicategory.

With this we can define limits in a quasicategory. For notational convenience, we will write $K^{\triangleleft}$ for $\Delta_0 \star K$.

Definition 2.3. Let C be a quasicategory and $X : K \to C$ a map of simplicial sets.

(1) A **cone** over X is a simplicial map $Y : K^{\triangleleft} \to C$ such that $Y|_K = X$.
(2) A cone $\tilde{X} : K^{\triangleleft} \to C$ is **universal** (or a **limit**) if for all $n > 0$ and all $Z : \partial\Delta_n \star K \to C$ such that $Z|_{\Delta^{\{n\}}} \star K = \tilde{X}$, there exists a lift,

$$\begin{tikzcd} \partial\Delta_n \star K \arrow[r, "Z"] \arrow[d, hook] & C \\ \Delta_n \star K \arrow[ur, dashed] & \end{tikzcd}.$$

Example 2.4. (1) A **terminal object** in quasicategory C is a limit of $\varnothing \to C$.
(2) A **pullback** in a quasicategory C is a limit of a diagram $\Lambda^2_2 \to C$.

All of these notions straightforwardly dualize to coslice categories, colimits, etc. The last definition that we need is a notion of adjoints of quasicategories.

Definition 2.5. Let $F : C \rightleftarrows D : G$ be simplicial maps between quasicategories and let $\eta : C \times \Delta_1 \to C$ be such that $\eta|_{C \times \partial\Delta_1} = [1_C, GF]$ and $\varepsilon : D \times \Delta_1 \to D$ such that $\varepsilon|_{D \times \partial\Delta_1} = [FG, 1_D]$. The quadruple $(F, G, \eta, \varepsilon)$ is an **adjunction of quasicategories** if $(F, G, \eta, \varepsilon)$ is an adjunction in the 2-category $\mathbf{qCat}_2$.

With this we can finally define what it means for a quasicategory to be locally Cartesian closed.

Definition 2.6. (1) A quasicategory C is **locally Cartesian closed** if it has finite limits and for any 1-simplex $f : x \to y$, the pullback functor $f^* : C \downarrow y \to C \downarrow x$ has a right adjoint.
(2) A simplicial map between locally Cartesian closed quasicategories is called a **locally Cartesian closed functor** if it preserves finite limits and the locally Cartesian closed structure. (?)

We denote the subcategory of **qCat** spanned by locally Cartesian closed quasicategories and locally Cartesian closed functors as $\mathbf{LCCC}_\infty$.

To study the homotopical aspects of dependent type theories, we introduce a slightly weakened version of a model category:

Definition 2.7. [11, Def. 6.6] A **left semi-model structure** on a bicomplete category $\mathcal{E}$ consists of three classes of maps: $\mathcal{W}, \mathcal{F}, \mathcal{C}$, subject to the axioms:

(1) all three classes are closed under retracts, $\mathcal{W}$ satisfies the 2-out-of-3 property,
(2) fibrations are preserved under pullback,
(3) cofibrations have the left lifting property with respect to trivial fibrations, and trivial cofibrations with cofibrant source have the left lifting property with respect to fibrations,
(4) every map can be functorially factored as a cofibration followed by trivial fibration, and every map with a cofibrant source can be functorially factored as a trivial cofibration followed by a fibration.

In many cases, one typically has slightly more structure:

Lemma 2.8. [11, Lem. 6.7] Suppose $\mathcal{E}$ is a bicomplete category, equipped with:

(1) a class of maps $\mathcal{W}$, including all identities, and closed under 2-out-of-6 and retracts, and
(2) two weak factorization systems $(\mathcal{A}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{TF})$,
(3) such that $\mathcal{TF} = \mathcal{F} \cap \mathcal{W}$, and
(4) when $A \in \mathcal{E}$ is cofibrant (i.e. the map $0 \to A$ is in $\mathcal{C}$), then a map $i : A \to B$ is in $\mathcal{A}$ if and only if it is in $\mathcal{C} \cap \mathcal{W}$.

Then the classes $(\mathcal{W}, \mathcal{C}, \mathcal{F})$ form a left semi-model structure on $\mathcal{E}$.

2.2. **Semantics of Type Theory.** The models of type theories that we are most interested in is that of contextual categories, as introduced by Cartmell in his thesis [5].

Definition 2.9. A **contextual category** is a category C equipped with:

(1) A grading on objects: $\operatorname{Ob} C = \coprod_{n\in\mathbb{N}} \operatorname{Ob}_n C$;
(2) an object $1 \in \operatorname{Ob}_0 C$;
(3) father operations: $\mathrm{ft}_n : \operatorname{Ob}_{n+1} C \to \operatorname{Ob}_n C$ (whose subscript we suppress);
(4) for each $\Gamma \in \operatorname{Ob}_{n+1} C$, a map $p_\Gamma : \Gamma \to \mathrm{ft}\,\Gamma$ (the canonical projection from Γ);
(5) for each for each $\Gamma \in \operatorname{Ob}_{n+1} C$ and $f : \Delta \to \mathrm{ft}\,\Gamma$, an object $f^*\Gamma$ together with a connecting map $f.\Gamma : f^*\Gamma \to \Gamma$;

such that:

(1) 1 is terminal in C and the unique object in $\operatorname{Ob}_0 C$;
(2) for each for each $\Gamma \in \operatorname{Ob}_{n+1} C$ and $f : \Delta \to \mathrm{ft}\,\Gamma$, we have $\mathrm{ft}(f^*\Gamma) = \Delta$, and the following square is a pullback,

$$\begin{tikzcd} f^*\Gamma \arrow[r, "f.\Gamma"] \arrow[d, "p_{f^*\Gamma}"'] \arrow[dr, phantom, "\lrcorner"] & \Gamma \arrow[d, "p_\Gamma"] \\ \Delta \arrow[r, "f"] & \mathrm{ft}\,\Gamma \end{tikzcd}$$

(3) and these canonical pullbacks are strictly functorial, i.e. for $\Gamma \in \operatorname{Ob}_{n+1} C$, $\mathrm{id}^*_{\mathrm{ft}\,\Gamma}\,\Gamma = \Gamma$ and $\mathrm{id}_{\mathrm{ft}\,\Gamma}.\Gamma = \mathrm{id}_\Gamma$; and for $\Gamma \in \operatorname{Ob}_{n+1} C$, $f : \Delta \to \mathrm{ft}\,\Gamma$ and $g : \Theta \to \Delta$, we have $(fg)^*\Gamma = g^*f^*\Gamma$ and $fg.\Gamma = f.\Gamma \circ g.f^*\Gamma$.

The structure of a contextual category seeks to axiomatize the structures found in the syntactic category of a dependent type theory. To help translating between the syntax of type theories and the semantics of contextual

categories, we identify objects $\Gamma' \in C$ such that $\mathrm{ft}(\Gamma') = \Gamma$ as types in the context Γ, the set of which we denote $\mathrm{Ty}(\Gamma)$. For a type $A \in \mathrm{Ty}(\Gamma)$, we write $\Gamma.A$ for the corresponding object in our contextual category C, and p_A for the projection $p_{\Gamma.A} : \Gamma.A \to \Gamma$. Syntactically, we think of this projection map as corresponding to the weakening of the context $\vec{x} : \Gamma, x : A$ to $\vec{x} : \Gamma$. We will also extend this notation to contexts, i.e. if $\Gamma' \in C$ such that $\mathrm{ft}^m \Gamma' = \Gamma$ for some $m \in \mathbb{N}$, we will write this object as $\Gamma.\Delta$, where we think of Δ as a context in context Γ. This again comes with a canonical projection which we denote $p_\Delta : \Gamma.\Delta \to \Delta$ obtained by composing projection maps.

Given $\Gamma \in C$ and $A \in \mathrm{Ty}(\Gamma)$, the set of terms of type A in context Γ, denoted $\mathrm{Tm}(A)$, is the set of sections $t : \Gamma \to \Gamma.A$ to the projection map p_A. To understand why we consider sections of the projection maps to be terms, take $\Gamma = x : A'$, then the projection corresponds to the judgment $x : A', y : A \vdash x : A'$, and a term of A is then given by the judgment $x : A' \vdash t : A$, so this along with the judgment $x : A' \vdash x : A'$, assembles into a context morphism $x : A'$ to $x : A', y : A$, which is a section to the projection map.

One of the benefits of using contextual categories over other potential models, is that we can describe contextual categories as models of an essentially algebraic theory.

Definition 2.10. [1, Def. 3.34] An **essentially algebraic theory** is a quadruple

$$\Gamma = (\Sigma, E, \Sigma_t, \mathrm{Def})$$

where Σ is a many-sorted signature consisting of operation symbols, E is a set of Σ-equations, $\Sigma_t \subseteq \Sigma$ is a subset of the operations in Σ which are "total", and Def is a function assigning to each operation $\sigma \in \Sigma \setminus \Sigma_t$ of type $\Pi_{i \in I} s_i \to s$, a set of Σ_t-equations involving only variables of the sorts s_i.

Essentially algebraic theories differ from algebraic theories in that they allow operations to be partially defined. This is necessary, for example, when writing a theory of categories, as the composition operation $\circ : \mathrm{Mor} \times \mathrm{Mor} \to \mathrm{Mor}$ is only defined when the source and target of our morphisms match.

Definition 2.11. A (set-theoretic) **model** of an essentially algebraic theory is then an assignment of every sort s to a set $M(s)$, and every operation symbol $\sigma : \Pi_{i \in I} s_i \to s$ to a partial function $M(\sigma) : \Pi_{i \in I} M(s_i) \to M(s)$ such that

(1) for each $\sigma \in \Sigma_t$, $M(\sigma)$ is a total function,
(2) for each $\sigma \in \Sigma \setminus \Sigma_t$, $M(\sigma)$ is defined exactly when the equations of $\mathrm{Def}(\sigma)$ are satisfied, and
(3) all the equations in E are satisfied.

The description of contextual categories as models of an essentially algebraic theory provides us with a number of useful tools for working with contextual categories, and so, indirectly with the syntax. For instance, the notion of models given provides us with a natural notion of morphism i.e. functions which preserve all of the operations. In the case of contextual categories we call these morphism **contextual functors**, and these allow us to construct the category of contextual categories, **CxlCat**, which, as a category of models of an essentially algebraic theory, must be locally presentable and thus complete and cocomplete [1, Thm. 3.36].

We wish to additionally equip our contextual categories with structures corresponding to the various forms of type constructors that appear in Martin-Löf type theory. These include:

(1) Identity types (denoted Id);
(2) unit types (1);
(3) dependent sum or Sigma types (Σ);
(4) dependent function of Pi types, with function extensionality (Π_{ext}).

The most relevant of these structures to this paper is the Id-type structure.

Definition 2.12. An Id-**type structure** on a contextual category C consists of:

(1) for each $\Gamma.A$, an object $\Gamma.A.p_A^* A.\mathrm{Id}_A$,
(2) for each $\Gamma.A$, a section $\mathrm{refl}_A : \Gamma.A \to \Gamma.A.p_A^* A.\mathrm{Id}_A$,
(3) for each $\Gamma.A.p_A^* A.\mathrm{Id}_A.C$ and $d : \Gamma.A \to \Gamma.A.p_A^* A.\mathrm{Id}_A.C$ with $p_C \circ d = \mathrm{refl}_A$, a section $J_{C,d} : \Gamma.A.p_A^* A.\mathrm{Id}_A \to \Gamma.A.p_A^* A.\mathrm{Id}_A.C$, such that $J_{C,d} \circ \mathrm{refl}_A = d$,

(4) such that for $f : \Gamma' \to \Gamma$, and all appropriate arguments above,

$$f^*(\Gamma.A.p_A^*A.\mathrm{Id}_A) = \Gamma'.f^*A.(p_{f^*A})^*(f^*A).\mathrm{Id}_{f^*A},$$
$$f^*\,\mathrm{refl}_A = \mathrm{refl}_{f^*A}, \qquad f^*\boldsymbol{J}_{C,d} = \boldsymbol{J}_{f^*C,f^*d}.$$

The first three rules of this type structure are meant to mimic the syntactic rules for identity types in our type theory, and the last rule ensures that this structure respects substitution. Similar definitions for unit, Sigma, and (extensional) Pi-type structures can be found in [13, Appendix B].

Each of these structures extend the essentially algebraic theory of contextual categories and thus also have a natural notion of morphism, preserving the corresponding structures. We denote the category of contextual categories with identity, unit, Sigma, and extensional Pi-type structures by $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$.

Following [6, Prop. 3.3.1], we can extend the notion of Id-type structures to **identity contexts**. Specifically, given $\Gamma \in C$ and a context extension $\Gamma.\Delta \in C$, we can further construct a context extension $\Gamma.\Delta.p_\Delta^*\Delta.\mathrm{Id}_\Delta$ which comes equipped with reflexivity map and an elimination term which directly generalize those of the Id-type structure on a single type. This is essentially a notational issue which helps use define a notion of homotopies within our contextual categories.

Definition 2.13. Given maps $f, g : \Gamma \to \Delta$ in a contextual category C, a **homotopy** H from f to g, denoted $H : f \sim g$, is a factorization of the map $(f, g) : \Gamma \to \Delta \times \Delta := \Delta.p_\Delta^*\Delta$ through the identity context $\Delta.p_\Delta^*\Delta.\mathrm{Id}_\Delta \to \Delta.p_\Delta^*\Delta$.

With this we can define a notion of equivalence within a contextual category to match the syntactic notion of equivalence (or one of such equivalent notions [19, Ch. 4]).

Definition 2.14. Let C be a contextual category with identity types.

A **structured equivalence** $w : \Gamma \simeq \Delta$ consists of a map $f : \Gamma \to \Delta$, along with maps $g_1, g_2 : \Delta \to \Gamma$ and homotopies $\eta : fg_1 \sim 1_\Delta$ and $\varepsilon : g_2 f \sim 1_\Gamma$.

An **equivalence** $\Gamma \xrightarrow{\sim} \Delta$ in C is a map $f : \Gamma\Delta$ for which there exists some $g_1, g_2, \eta, \varepsilon$ making it a structured equivalence.

With this definition of equivalence we can see that contextual categories have an underlying category with weak equivalences (which is in fact even a fibration category [2, Thm. 3.2.5]). Identifying this underlying structure is the first crucial step in studying the homotopical semantics of our type theories.

Our description of contextual categories as models of an essentially algebraic theory additionally provide us with the important ability to define contextual categories by generators and relations, and in particular lets us define freely generated contextual categories.

Definition 2.15. We define the following freely generated objects in $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$:

(1) $\langle \Gamma_n \rangle$ is freely generated by a context of length n.
(2) $\langle \Gamma_n \vdash A \rangle$ is freely generated by a context Γ of length n, and a type A in context Γ.
(3) $\langle \Gamma_n \vdash a : A \rangle$ is freely generated by a context Γ, and a type A and a term $a : A$ in context Γ, i.e. a section of p_A.
(4) $\langle \Gamma_n \vdash A \simeq A' \rangle$ is freely generated by a context Γ, types A, A' in context Γ, and maps $f, g_l, g_r, \alpha_l, \alpha_r$ constituting an equivalence from A to A' over Γ.
(5) $\langle \Gamma_n \vdash e : \mathrm{Id}_A(a, a') \rangle$ is freely generated by Γ and A, as in $\langle \Gamma_n \vdash A \rangle$, and a section to the projecting map $\Gamma.A.(p_A)^*A.\mathrm{Id}_A \to \Gamma.A$, giving the three terms a, a', and e.

Note that our use of the phase "freely generated" is perhaps a bit misleading, as these are not all freely generated objects of the theory of contextual categories. Since terms of the theory correspond to sections, these objects must involve some relations. Calling these objects "freely generated" emphasizes their role in relation to their corresponding syntactic objects.

With this we can define the classes of maps that will make up the left semi-model structure on $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$, which will provide us with a homotopical interpretation in service of constructing a semantics.

Definition 2.16. A map $F : C \to D$ of contextual categories with Id-types is a **type-theoretic equivalence** if it satisfies:

(1) **weak type lifting**: for any $\Gamma \in C$ and any type $A \in \mathrm{Ty}(F\Gamma)$, there exists a type $\bar{A} \in \mathrm{Ty}(\Gamma)$ along with an equivalence $F\bar{A} \simeq A$ over $F\Gamma$, and
(2) **weak term lifting**: for any $\Gamma \in C$, $A \in \mathrm{Ty}(\Gamma)$, and $a \in \mathrm{Tm}(FA)$, there exists a term $\bar{a} \in \mathrm{Tm}(A)$ along with an identity $e \in \mathrm{Tm}(\mathrm{Id}_{FA}(F\bar{a}, a)$.

These will make up the class $\mathcal{W}$ of our left semi-model structure. The remaining classes will be cofibrantly generated from the following sets of maps.

Definition 2.17. Take I and J to be the following sets of maps:

(1) I consists of the inclusions $\langle \Gamma_n \rangle \to \langle \Gamma_n \vdash A \rangle$ and $\langle \Gamma_n \vdash A \rangle \to \langle \Gamma_n \vdash a : A \rangle$, for all $n \in \mathbb{N}$, and
(2) J consists of the inclusions $\langle \Gamma_n \vdash A \rangle \to \langle \Gamma_n \vdash A \simeq A' \rangle$ and $\langle \Gamma_n \vdash a : A \rangle \to \langle \Gamma_n \vdash e : \mathrm{Id}_A(a, a') \rangle$, for all $n \in \mathbb{N}$.

Definition 2.18. We now define the **trivial fibrations**, $\mathcal{TF}$, as the right lifting class of I, the **cofibrations**, $\mathcal{C}$, as the left lifting class of $\mathcal{TF}$, the **fibrations**, $\mathcal{F}$, as the right lifting class of J, and the **anodyne maps**, $\mathcal{A}$, as the left lifting class of $\mathcal{F}$.

By the small object argument [16], we get:

Proposition 2.19. $(\mathcal{C}, \mathcal{TF})$ and $(\mathcal{A}, \mathcal{F})$ are both weak factorization systems.

Unfolding the lifting properties of $\mathcal{TF}$, we can see that a map $F : C \to D$ is a trivial fibration exactly if it strictly lifts types and terms, i.e. for any $\Gamma \in C$ and $C \in \mathrm{Ty}(F\Gamma)$, there is some term $\bar{A} \in \mathrm{Ty}(\Gamma)$ such that $F\bar{A} = A$, and similarly for terms. Since strict type and term lifting immediately implies their corresponding weak version, we get that trivial fibrations must be weak equivalences. Similarly, the lifting properties for fibrations can be seen as equivalence and equality lifting. We immediately get that every object is fibrant, since the generating anodyne maps have retracts given by the identity equivalence and reflexivity, respectively.

It follows from the construction that a typical cofibration of contextual categories is an I-cell complex, i.e. it consists of freely adding types and terms to a theory without any judgmental equalities. Cofibrant contextual categories are then type theories which are freely generated (over the constructors $\mathrm{Id}, 1, \Sigma, \Pi_{\mathrm{ext}}$) by a set of type and term formation rules with no (non-trivial) judgmental equalities. The anodyne maps, in turn, consist of freely adjoining types and terms to a theory along with equivalences and identities to a preexisting type or term.

These classes of maps satisfy the hypotheses of 2.8, providing us with homotopical structure on the category of contextual categories.

Theorem 2.20. [11, Thm. 6.9] The classes $\mathcal{W}, \mathcal{C}, \mathcal{F}$ above form a left semi-model structure on the category $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$.

Proof Sketch. We show that $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$ satisfies the hypotheses of 2.8. As noted before, we know that $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$ is bicomplete, as a category of models of an essentially algebraic theory, and as we saw above, we have two weak factorization systems, $(\mathcal{C}, \mathcal{TF})$ and $(\mathcal{A}, \mathcal{F})$, constructed with the small objects argument. Showing that the class $\mathcal{W}$ satisfies the right closure properties and that $\mathcal{TF} = \mathcal{F} \cap \mathcal{W}$ are straightforward arguments, so the bulk of the work that is required is in showing that maps with cofibrant source are anodyne exactly when they are both cofibrations and weak equivalences.

First off, the inclusion $\mathcal{A} \subseteq \mathcal{C}$ follows dually from the fact that $\mathcal{TF} \subseteq \mathcal{F}$, which is immediate from their construction.

The rest of the proof makes use of a construction that is almost a path object (examined in generality in [12], and defined in more specificity in [11, Def. 5.5]), whose specifics we will mostly black-box for the sake of brevity. Given a contextual category C with Id-types, we can construct the contextual category of **span-equivalences** C^{eqv}, which in turn allows us to define a notion of **right homotopy** between maps

$F_0, F_1 : C \to D$ in **CxlCat** when they factor through D^{eqv} in the following diagram:

$$\begin{tikzcd} & D^{\mathrm{eqv}} \arrow[d, two heads, "(P_0,P_1)"] \\ C \arrow[ur, bend left, "H"] \arrow[r, "(F_0,F_1)"] & D \times D \end{tikzcd},$$

where the maps P_0, P_1 are projections that come with the span-equivalence construction and which are trivial fibrations [11, Prop. 4.13.3, Prop. 5.12]. For C cofibrant, this notion of right homotopy, denoted $F_0 \sim_r F_1$, is an equivalence relation [11, Prop. 6.3].

Returning to the proof, if we now assume $\mathcal{A} \subseteq \mathcal{W}$ and $j : A \to B$ is anodyne with A cofibrant, then by the fibrancy of A, our lifting properties give us a retraction in the following diagram:

$$\begin{tikzcd} A \arrow[r, "1_A"] \arrow[d, tail, "j"] & A \arrow[d, two heads] \\ B \arrow[r] \arrow[ur, dashed, "r"] & 1 \end{tikzcd}.$$

But then r is also a homotopy section of j, by filling the following square:

$$\begin{tikzcd} A \arrow[r, "c_j"] \arrow[d, tail, "j"] & B^{\mathrm{eqv}} \arrow[d, two heads] \\ B \arrow[r, "(1_B, jr)"'] \arrow[ur, dashed, "H"] & B \times B \end{tikzcd},$$

where c_j is the reflexivity homotopy on j, supplied by right homotopies being an equivalence relation when the source is cofibrant. We now have that $rj = 1_A$, and $jr \sim_r 1_B$, so by 2-out-of-6 and the fact that weak equivalences are closed under right homotopies [11, Prop. 6.5], we get that $j \in \mathcal{W}$ as desired.

Now suppose that $j : A \to B$ is in $\mathcal{W}$ and C with A cofibrant. We wish to show that j lifts against all fibrations. Note that it is enough to show this for fibrations over B, as fibrations are closed under pullbacks, so assume with have the following square that we wish to find a filler for:

$$\begin{tikzcd} A \arrow[r, "h"] \arrow[d, tail, "j", "\sim"'] & Y \arrow[d, two heads, "p"] \\ B \arrow[r, "1_B"] & B \end{tikzcd}.$$

Using the reflexivity homotopy $c_h : A \to Y^{\mathrm{eqv}}$ for h, we can factor it through the pullback with the following diagram:

$$\begin{tikzcd} & (h, 1_Y)^* Y^{\mathrm{eqv}} \arrow[r] \arrow[d, two heads, "(Q_0,Q_1)"] \arrow[dr, phantom, "\lrcorner", very near start] & Y^{\mathrm{eqv}} \arrow[d, two heads, "(P_0,P_1)"] \\ A \arrow[ur, tail, "c'_h"] \arrow[r, "(1_A,h)"] \arrow[dr, tail, "1_A"'] & A \times Y \arrow[r, "(h,1_Y)"] \arrow[d, two heads, "\pi_0"] \arrow[dr, phantom, "\lrcorner", very near start] & Y \times Y \arrow[d, two heads, "\pi_0"] \\ & A \arrow[r, "h"] & Y \end{tikzcd}.$$

Since Q_0 is a pullback of the trivial fibration P_0, it is a trivial fibration as well, so by 2-out-of-3, c'_h is an equivalence since $Q_0 c'_h = 1_A$. Similarly, pQ_1 is an equivalence since $pQ_1 c'_h = ph = j$.

However, since pQ_1 is also a fibration, it is a trivial fibration, so (since j is a cofibration) we get a filler in the left square of the following diagram:

$$\begin{tikzcd}
A \arrow[r, "c'_h"'] \arrow[rr, bend left, "h"] \arrow[d, tail, "j"'] & (h, 1_Y)^* Y^{\mathrm{eqv}} \arrow[r, "Q_1"'] \arrow[d, two heads, "pQ_1", "\sim"'] & Y \arrow[d, two heads, "p"] \\
B \arrow[r, "1_B"'] \arrow[ur, dashed] & B \arrow[r, "1_B"'] & B
\end{tikzcd}.$$

Composing this lift with Q_1 then gives us the desired lift in the original square. This completes the proof that maps with cofibrant sources are anodyne exactly when they are cofibrations and weak equivalences, so we have satisfied the hypotheses of 2.8, and so constructed a left semi-model structure on $\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}}$. □

3. Conjecture

With our syntax given by contextual categories and semantics given by quasicategories, we finally have the background to start discussing the conjecture which is the primary motivation of this research. We saw previously that each contextual category has an underlying category with weak equivalences, giving us an inclusion $\mathbf{CxlCat} \to \mathbf{weCat}$. It is known that the categories **weCat** and **qCat** are DK-equivalent as categories with weak equivalences. We will call this equivalence Ho_∞ and note that it can be implemented in a number of ways (see [4, §1.6]), but, for concreteness, we can consider the following localization construction which takes a category with weak equivalences $(C, \mathcal{W})$, takes the pushout of the span

$$\coprod_{w \in \mathcal{W}} E[1] \leftarrow \coprod_{w \in \mathcal{W}} \Delta^1 \to NC$$

in **sSet** (NC is the nerve of C), and takes a fibrant replacement in the Joyal model structure. We can then define the map Cl_∞ as the following composite:

$$\mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}} \longrightarrow \mathbf{weCat} \xrightarrow{\mathrm{Ho}_\infty} \mathbf{qCat}\,.$$

[10] shows us that this functor is homotopical and valued in $\mathbf{LCCC}_\infty$.

Theorem 3.1. [10, Thm. 5.8] The functor Cl takes values in the category $\mathbf{LCCC}_\infty$. It also takes equivalences of contextual categories to categorical equivalences of quasicategories.

We can now finally state the conjecture:

Conjecture 3.2. [11, Conj. 3.7] The functor

$$\mathrm{Cl}_\infty : \ \mathbf{CxlCat}_{\mathrm{Id},1,\Sigma,\Pi_{\mathrm{ext}}} \to \mathbf{LCCC}_\infty$$

is a DK-equivalence of categories with weak equivalences.

The original conjecture additionally contained a "layer" below this, supposing that contextual categories equipped with only identity, unit, and sigma type formers should be equivalent to finitely complete quasicategories. This part of the conjecture has, however, already been proven in the most part in [9], where the authors show that the category of finitely complete quasicategories is DK-equivalent to the category of comprehension categories (another semantic model for type theories). The connection between comprehension categories and contextual categories, especially when equipped with type formers, remains not completely understood. [11] also suggests a further extension of this conjecture where contextual categories modeling homotopy type theories should be equivalent to elementary infinity toposes, however, at the time of publishing, neither of these concepts were well defined. Since then, the requisite categories have been constructed and the analogous functor has been shown to be valued in elementary infinity toposes in [15].

4. Approach

The issue with a direct approach to the conjecture is that the functor Cl_∞ is difficult to do computations with, in particular because of the fibrant replacement that is involved in constructing Ho_∞. This leads us to search for functors which are equivalent to the desired construction, but are easier to work with such as the quasicategory of frames (constructed in [18], and shown to be equivalent to Ho_∞ in [8]). Even this functor however has the problem of having no clear inverse, which would be desirable for something that we wish to show is a DK-equivalence.

Our goal then is to construct a new functor, equivalent to Cl_∞, using a nerve construction, which provides the immediate benefit of coming equipped with an adjoint that would act as a good candidate for an inverse.

Definition 4.1. Given a cosimplicial object $X : \Delta \to C$ in some cocomplete category C, we can get a map $|-| : \mathbf{sSet} \to C$, called the **(simplicial) realization**, by taking a left Kan extension of X along the Yoneda embedding $y : \Delta \to \mathbf{sSet}$. The **(simplicial) nerve** is then the induced right adjoint $N : C \to \mathbf{sSet}$. For an object $Y \in C$, the simplicial set $N(Y)$ is defined by the formula $N(Y)_n = C(X_n, Y)$.

Example 4.2. The nerve of a category is given by the inclusion $\Delta \hookrightarrow \mathbf{Cat}$, which for a category C tells us that $N(C)_n = \mathbf{Cat}([n], C)$ is the set of objects in C for $n = 0$, and composable n-chains of morphisms in C for $n > 0$.

To get the desired functor from **CxlCat** to **qCat** then, all that is necessary is to find the "correct" simplicial object in **CxlCat** which induces a nerve functor that is valued in quasicategories and equivalent to Cl_∞. There is an "obvious" cosimplicial object $\mathbb{T}$ in **CxlCat** whose theories are freely generated, for each $n \in \mathbb{N}$, by closed types A_i, for each $i \in [n]$, and terms $x : A_i \vdash f_{i+1}(x) : A_{i+1}$, for each $i \in [n-1]$,. The face maps $\delta_i : \mathbb{T}_{n-1} \to \mathbb{T}_n$ are given by sending f_i to $f_{i+1} \circ f_i$, and the degeneracy maps $\sigma_i : \mathbb{T}_{n+1} \to \mathbb{T}_n$, are defined by sending f_{i+1} to id_{A_i}.

The issue with this cosimplcial object is that it is in a sense too strict. To properly capture the structure of our type theories, we would expect the theory representing the 2-simplex, for example, to not just have a single map $x : A_0 \vdash f_2(f_1(x)) : A_2$, but rather there should be a distinct map, say f_{02} connecting the types A_0 and A_2 and a term identifying these two maps, i.e. $x : A_0 \vdash \alpha(x) : \mathrm{Id}_{A_2}(f_{02}(x), f_2(f_1(x)))$. The "simplices" in $\mathbb{T}$ contain none of the homotopical information present in our type theories i.e. they do not have any identity terms, so mapping out of them into a theory $\mathbb{S}$ would not capture any of the higher structure in $\mathbb{S}$. What this means more concretely is that $\mathbb{T}$ is not Reedy cofibrant, and so we should not expect that the mapping space from $\mathbb{T}$ to a fibrant object $\mathbb{S}$ (reminder that all theories are fibrant) would be a fibrant simplicial set in the Joyal model structure i.e. a quasicategory.

This situation is analogous to that of the homotopy coherent nerve for simplicial categories (see [14, Ch. 1] for more exposition), where the naive inclusion $\Delta \hookrightarrow \mathbf{Cat} \hookrightarrow \mathbf{sCat}$, does not produce a nerve functor that properly accounts for the homotopical data, i.e. the simplicial enrichment, on **sCat**. The solution to this is then to consider "thickened" simplices which account for the homotopical structure and are a Reedy cofibrant replacement of the naive cosimplicial object. We wish to attempt something similar in our context, but to make sure that a Reedy cofibrant replacement exists, we need some sort of Reedy left semi-model structure on $\mathbf{CxlCat}^\Delta$.

To do this, we first recall some of the basic definitions of Reedy theory.

Definition 4.3. A **Reedy category**, is a category $\mathcal{R}$ equipped with two wide subcategories $\mathcal{R}_+$ and $\mathcal{R}_-$ and a total ordering on the objects defined by a degree function $d : \mathrm{Ob}(\mathcal{R}) \to \mathbb{N}$ such that,

(1) Every non-identity morphism in $\mathcal{R}_+$ raises degree,
(2) Every non-identity morphism in $\mathcal{R}_-$ lowers degree, and
(3) Every morphism $f \in \mathcal{R}$ uniquely factors as a map in $\mathcal{R}_-$ followed by a map in $\mathcal{R}_+$.

Example 4.4. The prototypical example of a Reedy category, and the one we are interested in, is the simplex category Δ, where Δ_+ is given by the monomorphisms, and Δ_- is given by the epimorphisms.

A category of diagrams indexed by a Reedy category and valued in a model category comes equipped with a natural model structure whose (co)fibrations are defined in the following way.

Definition 4.5. Let $\mathcal{R}$ be a Reedy category, $\mathcal{M}$ a (co)complete category, X an object in $\mathcal{M}^{\mathcal{R}}$, and α an object of $\mathcal{R}$.

(1) The **latching object** of X at α is defined as $L_\alpha X := \operatorname{colim}_{\beta \underset{+}{\to} \alpha} X_\beta$, where the colimit is taken over all non-identity arrows over α in $\mathcal{R}_+$. The natural map $L_\alpha X \to X_\alpha$ is called the **latching map**.
(2) The **matching object** of X at α is defined as $M_\alpha X := \lim_{\alpha \underset{-}{\to} \beta} X_\beta$, where the limit is taken over all non-identity arrows under α in $\mathcal{R}_-$. The natural map $X_\alpha \to M_\alpha X$ is called the **matching map**.

Definition 4.6. Let $\mathcal{R}$ be a Reedy category, $\mathcal{M}$ a model category (or more generally a category with a classes of (co)fibrations), and $f : X \to Y$ a map of diagrams in $\mathcal{M}^{\mathcal{R}}$.

(1) The map f is a **Reedy cofibration** if, for every object $\alpha \in \mathcal{R}$, the **relative latching map**
$$X_\alpha \amalg_{L_\alpha X} L_\alpha Y \to Y_\alpha$$
is a cofibration in $\mathcal{M}$.
(2) The map f is a **Reedy fibration** if, for every object $\alpha \in \mathcal{R}$, the **relative matching map**
$$X_\alpha \to Y_\alpha \times_{M_\alpha Y} M_\alpha X$$
is a fibration in $\mathcal{M}$.

While the standard Reedy model structure for diagrams valued in a model category is well understood, since we only have a left semi-model structure on **CxlCat**, we wish to construct an analogous Reedy left semi-model structure on $\mathcal{M}^{\mathcal{R}}$ when $\mathcal{M}$ does not have a full model structure.

Theorem 4.7. Let $\mathcal{M}$ be a category satisfying the hypotheses of 2.8, and $\mathcal{R}$ a Reedy category. The category $\mathcal{M}^{\mathcal{R}}$ comes equipped with a left semi-model structure where weak equivalences are component-wise, and (co)fibrations are Reedy (co)fibrations.

The necessitation of $\mathcal{M}$ satisfying the slightly stronger hypotheses of 2.8, comes from the use of the following lemma, which requires the existence of the $(\mathcal{A}, \mathcal{F})$ weak factorization system.

Lemma 4.8. Suppose $\mathcal{M}$ is a category which satisfies the hypotheses of 2.8. A pushout of a trivial cofibration with cofibrant source along a cofibration in $\mathcal{M}$ is a trivial cofibration.

Proof. A trivial cofibration with cofibrant source must be in the class $\mathcal{A}$, which is the left class of a weak factorization system, and so is closed under pushouts. Since this pushout is being taken along a cofibration, the pushout is a map in $\mathcal{A}$ with a cofibrant source, and thus is a trivial cofibration. □

Proof Sketch. (Theorem 4.7) The proof mostly follows from standard inductive arguments using the Reedy structure of $\mathcal{R}$ (see [7, Ch. 15], [17, Ch. 9]).

For the inductive step of lifting a Reedy trivial cofibration with cofibrant source $A \overset{\sim}{\rightarrowtail} B$, against a Reedy fibration $X \twoheadrightarrow Y$. By [7, 15.2.11] (note: this is the lemma which motivates the definition of Reedy (co)fibrations) all that is necessary is to construct a lift in the following diagram:

$$\begin{tikzcd} A_\alpha \coprod_{L_\alpha A} L_\alpha B \arrow[r] \arrow[d, tail] & X_\alpha \arrow[d, two heads] \\ B_\alpha \arrow[r] & Y_\alpha \times_{M_\alpha Y} M_\alpha X \end{tikzcd}.$$

To get this lift, we need to know that $A_\alpha \coprod_{L_\alpha A} L_\alpha B$ is cofibrant in $\mathcal{M}$ and that the left map is a weak equivalence. Since $A \to B$ is a Reedy trivial cofibration, we get that $L_\alpha A \to L_\alpha B$ is a trivial cofibration for all α, so $A_\alpha \to A_\alpha \coprod_{L_\alpha A} L_\alpha B$ is as well by the previous lemma. Since $A_\alpha \to B_\alpha$ is a weak equivalence and can be factored as $A_\alpha \to A_\alpha \coprod_{L_\alpha A} L_\alpha B \to B_\alpha$, we get that our desired map is a weak equivalence as well by 2-out-of-3. To see that $A_\alpha \coprod_{L_\alpha A} L_\alpha B$ is cofibrant, we note that A_α is cofibrant and the map $A_\alpha \to A_\alpha \coprod_{L_\alpha A} L_\alpha B$ is a cofibration.

For the factorization of a map $f : X \to Y$, with X cofibrant, into a Reedy trivial cofibration followed by a Reedy fibration, we work degree wise by induction, assuming that f has been factored for all objects of degree less than that of α, we get an induced map $X_\alpha \coprod_{L_\alpha X} L_\alpha Z \to Y_\alpha \times_{M_\alpha Y} M_\alpha Z$, which we can apply factorization in $\mathcal{M}$ since the previous lemma tells us that the source is cofibrant. This gives us our factorization into a cofibration and a fibration, all that remains is to show that the map $X_\alpha \to Z_\alpha$ is a weak equivalence. This follows from the fact that it can be factored as $X_\alpha \to X_\alpha \coprod_{L_\alpha X} L_\alpha Z \to Z_\alpha$, where the second map is a trivial cofibration by our factorization in $\mathcal{M}$, and the first map is a trivial cofibration as a pushout of a trivial cofibration with cofibrant source along a cofibration.

The remaining parts of the proof are essentially the same as the proof for the standard Reedy model structure as that "half" of the left semi-model structure is identical to the definition of a standard model structure. □

Corollary 4.9. There exists a Reedy left semi-model structure on the category $\mathbf{CxlCat}^\Delta$.

With this we know that it is possible to take a Reedy cofibrant replacement of our cosimplicial object $\mathbb{T}$ by factoring the map $0 \to \mathbb{T}$ as a Reedy cofibration followed by a Reedy trivial fibration, but this construction is still quite difficult to work with. Even if we automatically have that the resulting nerve functor is valued in quasicategories, proving anything further, e.g. showing that these quasicategories are locally Cartesian closed, would be difficult. For this reason we wish to construct a more explicit Reedy cofibrant replacement on $\mathbf{CxlCat}^\Delta$, and this is the direction of current research.

REFERENCES

[1] Jiří Adámek and Jiří Rosický. *Locally Presentable and Accessible Categories*. Cambridge University Press, 1994.

[2] Jeremy Avigad, Krzysztof Kapulkin, and Peter Lefanu Lumsdaine. Homotopy limits in type theory. *Mathematical Structures in Computer Science*, 25(5):1040–1070, 2015.

[3] C. Barwick and D.M. Kan. A characterization of simplicial localization functors and a discussion of dk equivalences. *Indagationes Mathematicae*, 23(1):69–79, 2012.

[4] Clark Barwick. On the algebraic k-theory of higher categories. *Journal of Topology*, 9:245–347, 2016.

[5] John Cartmell. *Generalized Algebraic Thoeries and Contextual Categories*. PhD thesis, Oxford, 1978.

[6] Richard Garner. Two-dimensional models of type theory. *Math. Structures in Computer Sci.*, 2009.

[7] Philip Hirshhorn. *Model Categories and their Localizations*. American Mathematical Society, 2002.

[8] Chris Kapulkin and Karol Szumiło. Quasicategories of frames of cofibration categories. *Applied Categorical Structures*, 25:323–347, 2015.

[9] Chris Kapulkin and Karol Szumiło. Internal language of finitely complete $(\infty, 1)$-categories. *Selecta Mathematica*, 25, 09 2017.

[10] Krzysztof Kapulkin. Locally cartesian closed quasicategories from type theory. *Journal of Topology*, 10:1029–1049, 2017.

[11] Krzysztof Kapulkin and Peter LeFanu Lumsdaine. The homotopy theory of type theories. *Advances in Mathematics*, 337:1–38, 2018.

[12] Krzysztof Kapulkin and Peter LeFanu Lumsdaine. Homotopical inverse diagrams in categories with attributes. *Journal of Pure and Applied Algebra*, 225(4):106563, 2021.

[13] Krzysztof Kapulkin and Peter LeFanu Lumsdaine. The simplicial model of univalent foundations (after voevodsky). *Journal of the European Mathematical Society*, 23(6):2071–2126, March 2021.

[14] Jacob Lurie. *Higher Topos Theory*. Princeton University Press, 2009.

[15] Maximilian Petrowitsch. Elementary ∞-toposes from type theory, 2025.

[16] D. G. Quillen. *Homotopical Algebra*. Springer-Verlag, Berlin, 1967.

[17] Andrei Radulescu-Banu. Cofibrations in homotopy theory, 2009.

[18] Karol Szumiło. *Two Models for the Homotopy Theory of Cocomplete Homotopy Theories*. PhD thesis, University of Bonn, 2014.

[19] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. `https://homotopytypetheory.org/book`, Institute for Advanced Study, 2013.