The title

First Author[1] & Ernst-August Doelle[1,2]

[1] Wilhelm-Wundt-University

[2] Konstanz Business School

Author Note

Add complete departmental affiliations for each author here. Each new line herein must be indented, like this line. Enter author note here.

The authors made the following contributions. First Author: Conceptualization, Writing - Original Draft Preparation, Writing - Review & Editing; Ernst-August Doelle: Writing - Review & Editing, Supervision.

Correspondence concerning this article should be addressed to First Author, Postal address. E-mail: my@email.com

Abstract

One or two sentences providing a **basic introduction** to the field, comprehensible to a scientist in any discipline. Two to three sentences of **more detailed background**, comprehensible to scientists in related disciplines. One sentence clearly stating the **general problem** being addressed by this particular study. One sentence summarizing the main result (with the words "**here we show**" or their equivalent). Two or three sentences explaining what the **main result** reveals in direct comparison to what was thought to be the case previously, or how the main result adds to previous knowledge. One or two sentences to put the results into a more **general context**. Two or three sentences to provide a **broader perspective**, readily comprehensible to a scientist in any discipline.

Keywords: keywords

Word count: X

The title

Death is something each of us must learn to cope with, whether in healthy ways or less so. These issues may be at front of mind for many in light of the COVID-19 pandemic. Various existential philosophers and psychologists have proposed ways in which we deal with the awareness of death and the anxiety this awareness often causes. Psychoanalyst Erik Erikson (1950) proposed that during mid-life one becomes acutely aware of their oncoming death and is motivated to care for things which will outlast themselves. He called this act of caring generativity. In *The Denial of Death*, philosopher Ernest Becker (**becker1973?**) posits that humans undertake immortality projects to curb their sense of vulnerability to death. Similarly, psychiatrist Robert Jay Lifton (**lifton1979?**), a mentee of Erikson, described the awareness of death as being ever present and motivating us to create symbols, thereby allowing us to imagine ourselves as symbolically immortalized. Existential psychiatrist Irvin Yalom (**yalom2008?**) notes that many of his clients experiencing anxiety about their death take comfort in "rippling," the idea that one's lasting effects on the world will ripple out and influence the world after they have died.

Although these thinkers use different terminology, there are several common themes among their ideas. (1) Our physical death is an inevitability, and we often find our awareness of its inevitability to be aversive. This aversion may be referred to variously as angst, death-anxiety, despair, being-towards-death, terror, and so on. However, (2) we take comfort in the idea that other, non-physical parts of us continue to exist indefinitely after our biological death, through mechanisms such as the heroic archetype and symbolic self. (3) Finally, we can take action to promote these non-physical aspects of the self, such as through search for meaning, sense of immortality, care, generativity, and rippling.

One of these bodies of thought, called symbolic immortality, was originally theorized by Lifton (**lifton1979?**), who thought that awareness of death drives a fundamental human desire for a sense of continuity lasting beyond the lifespan. Essentially, humans are

meaning-seeking creatures, and throughout our lives, this search for meaning involves an evolving psychological imagery of life and death. Death, or the transient nature of life, threatens our search for meaning. Lifton thought that if we could achieve what we believe to be some form of immortality, we could overcome this loss of meaning, and the awareness of death could instead drive an inner vitality (imagery associated with connection, integrity, and movement). If this drive toward vitality is lost, we are vulnerable to a psychic numbness or death-in-life (imagery associated with separation, disintegration, and stasis). In Lifton's own words, "Death does indeed bring about biological and psychic annihilation. But life includes symbolic perceptions of connections that precede and outlast that annihilation" (**lifton1979?**).

Lifton (**lifton1979?**) proposed five modes of experience or ways of achieving symbolic immortality: The biological (or biosocial) mode in which one lives on through their genetic and sociocultural progeny, the creative mode in which one's accomplishments and contribution outlast oneself, the natural mode in which one feels they are a part of the broader universe, the spiritual mode in which one seeks to transcend the physical realm to a higher spiritual realm beyond death, and the mode of experiential transcendence in which one experiences a phenomenological state of flow. The experiential mode must occur in the context of at least one of the other four to really be considered transcendent, but it is thought to have a great capacity to bring about personal change.

Claims of how we suppress death-anxiety have been investigated experimentally, primarily through the paradigm of Terror Management Theory (TMT). Based on the theories of Ernest Becker, TMT posits that human awareness of death is always present to some degree. This awareness of our inevitable death, coupled with a strong aversion to thoughts of death, causes terror and is pushed out of our consciousness by our creation of meaning systems (Greenberg, Pyszczynski, & Solomon, 1986). TMT proposes that self-esteem, interpersonal relationships, and cultural worldview work together to buffer

against our anxiety about death. It is assumed that these buffers suppress thoughts of death by providing a sense of symbolic immortality, though little systematic research has been conducted on this construct. The results of this buffering process are not always positive. For example, experimentally priming mortality salience can lead to more positive attitudes toward in-group members but harsher negative attitudes toward out-group members (Greenberg et al., 1990).

TMT refers to a person's awareness of death as mortality salience (MS). The MS hypothesis of TMT posits than an increase in one's awareness of death causes an increase in compensatory behaviors to lower their death anxiety, either by distracting from the awareness of death or by the promotion of meaningful cultural worldviews. In the MS paradigm, experimentally priming a participant's awareness of death (for example, by having participants write about death and then complete a distraction task) is thought to cause an increase in compensatory buffers. A meta-analysis of 277 experiments found mortality salience to have a robust, moderate overall effect size: $r(276) = 0.35$, $p = .00$ (Burke, Martens, & Faucher, 2010). Altogether, these experiments provide convincing evidence for TMT and the MS hypothesis in particular.

Though some avoidance of (or buffering against) death anxiety is thought to be universal and has the potential to increase interpersonal conflict, awareness of death through symbolic immortality may also have potential as a positive force. In particular, it is thought to be an underlying motive for what Erikson referred to as generativity. Generativity is the seventh of eight proposed stages in Erikson's (1950) theory of psychosocial development, which he associated with midlife and described as "the concern in establishing and guiding the next generation" (E. H. Erikson, 1963, p. 267). Little systematic research was conducted on this subject until the 1980's. Kotre (1984) expanded on the theory and proposed that the drive for generativity was related to a motive to expand the sense of self beyond the lifetime, especially in light of the fear of death.

McAdams and de St. Aubin (1992) sought to formalize the study of generativity as a multidimensional construct. Their seven components of generativity include cultural demand, inner desire (for symbolic immortality and community), concern (for the next generation), belief (in the human species), commitment, action, and narration (of generativity within one's life story). In addition to a quantitative measure of generative concern (the Loyola Generativity Scale), they developed a system for content analysis of autobiographical episodes pertaining to generativity, and symbolic immortality is one of the five themes they found. Here they define symbolic immortality as "any reference to leaving a legacy, having an enduring influence, or leaving behind products that will outlive one's physical existence," a theme clearly related to both Lifton's and Erikson's theories (McAdams & St. Aubin, 1992, p. 1011).

These research areas depend on the construct of symbolic immortality for their theoretical frameworks, but few researchers have attempted to systematically and quantitatively assess this construct. Two attempts have been made to develop such a measurement: Drolet's (1990) Sense of Symbolic Immortality Scale and Mathews and Kling's (1988) measure of symbolic immortality, based on an original questionnaire by Mathews and Mister (1987).

Drolet (1990) developed the Sense of Symbolic Immortality Scale based on Robert J. Lifton's theory of symbolic immortality and its five modes of experience. Drolet studied 136 adults, ages 18-30 and 30-40, and hypothesized that those in their 30's (established adults) would have a greater sense of symbolic immortality than the young adults (18-30). The measure is inherently subjective, not only by the nature of self-report, but in that the scale seeks to measure what a person *believes* and how they *feel* about these subjects. The scale as a whole had a high internal consistency ($\alpha = .91$) and test-retest reliability was $r = .97$. Internal consistency of subscales for the five theoretical modes of immortality was mixed. Of the five, spiritual immortality was the most distinct from the scale as a whole and the other

subscales. Factor analysis showed three factors, mapping onto biosocial, creative, and spiritual. The transcendent and natural items may be closely related to biosocial.

Moving beyond the scale development itself, SSI correlated negatively with Templer's Death Anxiety Scale and had a strong ($r = .84$) positive relationship with Maholick's Purpose in Life Test (Drolet, 1990). In interpreting the very strong correlation, the author suggests that SSI is a broader construct than Purpose in Life and the scale itself may be less prone to social desirability effects than the PIL, although this had not been directly tested. Age group was also related, with established adults having a higher SSI, particularly in the biosocial and creative domains.

We see multiple issues with using the Symbolic Immortality Scale. First, the study was underpowered, conducting exploratory factor analysis of 67 items using a sample of 136. Second, the scale was developed in French, and we do not take for granted the psychometric properties of a translated version. Third and most fundamentally, the scale has poor face validity and appears to measure the constructs theorized to symbolically immortalize rather than a sense of symbolic immortality directly. For example, the scale includes items such as "My sex life contributes greatly to my well-being", "Intimate relationships scare me", and "I am sure of who I am." Although related to the constructs (such as interpersonal relationships and self-esteem) which theoretically help cope with death, it is unclear how these items represent the construct of symbolic immortality itself.

Mathews and Mister (1987) also developed a scale pertaining to symbolic immortality, sensation seeking, and psychic numbness in a study including 400 adults. Experiential transcendence was operationalized as similar to Zuckerman's (1979) sensation seeking, which may not fully capture the original intent (the experience of losing oneself). Items were mapped onto five factors, and the five factors largely aligned with Lifton's constructs. Although internal consistency was at least acceptable for each factor, goodness of fit statistics are not reported. Some studies have used a revised version of the scale by Mathews and

Kling (1988), who adapted it for a study on prosocial behavior in the context of nonprofit volunteer motivation. They reported similar results for their revised scale. The items on these scales seem to have more face validity than the scale by Drolet, but some factors seem more behavioral and unnecessarily specific: pertaining to one's religiosity or biological children, whereas Lifton's theory allows for a broader interpretation of these dimensions. The Nature and Creative factors seem most useful and theoretically aligned with Lifton.

Much more advanced factor analysis methods have been developed since the 1980s, but to our knowledge, these scales have not been tested with more robust tools. The goal of the present research is to develop an up-to-date symbolic immortality scale that more directly measures one's sense of symbolic immortality and which contains items more generally applicable to broad groups of participants (e.g., regardless of a person's religious beliefs and parental status). We have attempted to use current best practices for scale development and analysis.

Method

Participants

Participants were recruited from two data sources. 564 undergraduate students taking introduction to psychology were recruited for course credit at a large midwestern university. Undergraduate participants were $M = 19.62$ ($SD = 2.75$) years old at the time of the study. 240 participants were recruited from Amazon's Mechanical Turk, and these participants were $M = 35.55$ ($SD = 11.05$) years old. No other demographic information was collected in the study.

Material and Procedure

58 items were developed using previous scale development on generativity (Drolet, 1990), along with theoretical factor alignment to XXCITEXX. These items aligned with areas of biological, physical, social, spiritual, and personal immortality. The survey was

implemented in Qualtrics. First, participants entered their age in years for basic demographic information. The immortality items were separated into three blocks of 20-21 questions. These three blocks of questions were randomized across participants, and the items within each block were also randomized. In each block, one attention check item was shown (e.g., "Answer this question as somewhat agree."). At the end of the study, participants were directed back to either SONA for undergraduate course credit or Mechanical Turk for monetary compensation[ˆUnfortunately, the exact amount we paid participants is lost to time. Given our ethics application, we believe it was 1-2 USD, and the survey was completed in $M = 6.34$ ($SD = 4.85$ minutes.].

Data analysis

This study was not pre-registered. We conducted data analysis in two phases. The data were first screened for inattentive and outlier responses. The data were then randomly split into an exploratory and confirmatory dataset. We developed the XX scale by using exploratory factor analysis (EFA) with maximum likelihood estimation and an oblique (i.e., oblimin) rotation (Preacher & MacCallum, 2003). Analyses were performed with the *psych* package (Revelle, 2020). We used scree plots, parallel analysis, and theory to determine the number of factors and compared models based on fit indices to select the best model. We use the following fit indices: Root Mean Squared Error of Approximation [RMSEA; Steiger (2016)], Root Mean Squared Residual [RMSR; Jöreskog and Sörbom (1996)], Comparative Fit Index [CFI; Bentler (1990)], and the Tucker-Lewis Index [TLI; Tucker and Lewis (1973)]. Values close to one are expected for goodness of fit statistics [CFI, TLI; Bentler (1990); Hu and Bentler (1999)], while lower values close to zero are considered better models for residual statistics [RMSEA, RMSR; Steiger (2016)].

We then analyzed the best model from the exploratory stage using confirmatory factor analysis (CFA) and multigroup CFA (Brown, 2015; Kline, 2016) on the second confirmatory dataset. The models were analyzed with *lavaan* (Rosseel, 2012) and *visualizemi* (Buchanan,

2024) *R* packages. Multigroup CFA can be used to determine if the scale structure can be generalized to subsamples within the data, and thus, allowing for the same interpretation of total or subscale scores calculated from the scale. First, we modeled the overall structure from the EFA results to determine fit in a new sample. The RMSEA, CFI, and TLI were used to determine model fit along with the Standardized Root Mean Squared Residual [SRMR; Bentler (1995)]. The same interpretations were included from the exploratory stage: high fit indices close to one for goodness of fit statistics, and low fit indices close to zero for residual fit statistics. If the overall model fit well, we then modeled the subsamples of student populations and Mechanical Turk populations separately. While fit is expected to potentially degrade due to smaller sample sizes, we expect these models to converge to demonstrate that it is appropriate to combine models for multigroup testing.

For the multigroup CFA, we used Brown (2015)'s suggested terminology and steps for analysis. The *configural* model includes both subsamples combined together into one model controlling for group structure. This model supports the "configuration" or structure of the scale is at least similar within each group, but does not force each group to show similar patterns of parameter estimates. Each subsequent step constrains a set of parameters to be equal across groups to demonstrate invariance (i.e., equivalence). The configural model is then compared to the *metric* model, which sets the factor loadings for each measured item to be equal across groups. This model supports that each item is related to the underlying latent variable the same way for each group. If this assumption holds, the metric model is then compared to a *scalar* model, which sets the item intercepts (i.e., item means) to be equal across groups. Scalar model invariance suggests that each subsample shows the same item means and factor loadings across groups. Finally, we compared the scalar model to the *strict* model, which constrains item residuals (i.e., errors, variance) to be the same across groups. Invariance in model residuals suggests that the items have the same answer range/spread across groups.

To assess if models are invariant, we used Δ CFI <= .01 criterion to indicate that models were equivalent (Cheung & Rensvold, 2002). If models were not considered invariant, we investigated for the source of the invariance by using partial invariance testing (Byrne, Shavelson, & Muthén, 1989; Cheung & Rensvold, 1999). Each parameter estimate is individually relaxed to determine if that particular estimate leads to misfit in the model. For example, if the model fails scalar invariance, each item mean would be allowed to freely vary between groups (one at a time) to find the items that show group differences. If only a few items show group differences, then the model is considered "partially" invariant, and the impact on total scores can be assessed. Partially invariant models indicate areas for further investigation into group differences and ways that the final scores of the model must be carefully interpreted. Previously, Nye and Drasgow (2011) and Nye, Bradburn, Olenick, Bialko, and Drasgow (2019) proposed an effect size d_{MACS} to help quantify the size of the difference between groups in partially invariant models. However, d_{MACS} can generally only be used to understand the item means and loadings together (due to the mathematical calculation), regardless of the tyep of non-invariance. Therefore, we applied a new technique proposed by Buchanan (2024) to assess several potential effect size components of the model: 1) overall model replication likelihood, 2) individual parameter replication likelihood, and 3) effect size for individual parameters showing non-invariance.

We will use the h_{nmi} effect size as the likelihood of replication statistic for both the overall model and individual parameters. First, the data is bootstrapped with replacement n times to create similar samples to the original data. The grouping variable is then randomized to create a comparison model. The proportion of times that invariance (or non-invariance) is met is calculated for the bootstrapped and randomized group labels. h_{nmi} is Cohen's h , which is the standardized difference between two proportions for non-invariance (no measurement invariance, but the statistic can also be calculated as h_{mi} for the difference in proportions for when the model showed measurement invariance). We suggest h_{nmi} because the interpretation is a bit simpler: effect sizes close to zero indicate

that the bootstrapped data and randomized groups data show the same number of non-measurement invariance findings, indicating that they are likely the same result. The logic here is that if groups are truly equal in their measurement on a scale, then the grouping variable should not affect the results, thus, randomizing the grouping variable label does not affect the results. As h_{nmi} increases, we expect to find non-measurement invariance between groups, as the bootstrapped data is consistently different than randomized group labels. h can be interpreted much like d, but d does not technically have upper or lower limits (i.e., they are ∞), while h's range is limited to π. Therefore, we will report h_{nmi_p} which is scaled to range from -1 to +1 for easier interpretation. In addition to this effect size, we can calculate d for the differences in parameters for bootstrapped versus random group labels, thus, providing an estimate for the size of group differences on partially invariant parameter estimates.

Results

Data Screening

Undergraduate Data. Data were screened for accuracy, missing data, outliers, and multivariate parametric assumptions. The undergraduate data included $n = 564$ participants. We first screened each participant for missing data and excluded $n = 7$ participants for missing more than 5% of their data [TABACHNICK]. 105 participants included less than 5% missing data (i.e., 3 items or less), and these data points amounted to less than 2% of data missing per question. These items were replaced with a random imputation from the *mice* library [CITE MICE].

The data were then screen for attentiveness and accuracy using the framework from CITE BUCHANAN to find low quality responses. The survey items were split into three pages, and each page was screened separately using click counts, an attention check item (i.e., please answer strongly agree for this item), and number of characters on a page for reading speed. The total number of indicators for low quality data was summed for each

participant across pages (e.g., 3 indicators by 3 pages = 9 total possible items), and $n = 49$ participants showed at least 4 indicators across all pages. CITE BUCHANAN suggest a criteria of 2/5 screeners or 40%, and therefore, we used 4 items as the ~40% cut off for inclusion in the data. The final sample size for the undergraduate population was $n = 508$. We additionally screened for multivariate outliers using Mahalanobis distance across items and found $n = 43$ possible outliers, but they are included in the analyses described below.

Mechanical Turk Data. The Mechanical Turk data were screened in the same manner as the undergraduate sample. The original sample size was $n = 240$ participants. $N = 23$ participants were excluded for missing more than 5% of their data, and $n = 36$ participants had 3 or less data points estimated using the *mice* library. The same low quality data screening was performed, and $n = 20$ participants were excluded for having 40% or more low quality indicators. Last, $n = 10$ participants were flagged as potential outliers but were not included in analyses presented below.

```
## [1] 352
```

```
## [1] 353
```

```
## [1] 326
```

```
## [1] 326
```

The data was then randomly split into $n = 352$ participants for the exploratory factor analysis, and $n = 353$ for the confirmatory factor analysis. The EFA data included n undergraduate = 254 participants, and n Mechanical Turk = 98 participants. The CFA data included n undergraduate = 254 participants, and n Mechanical Turk = 99 participants.

EFA

Number of Factors. To determine the number of factors that best fit the data, we used suggested theory (5 factors), a scree plot, and a parallel analysis using the *psych* library

[CITE PSYCH]. Parallel analysis suggested 7 factors, while a visual inspection of the scree plot indicated 2 to 3 factors. We analyzed a 2-factor, 3-factor, and 5-factor model for simple structure and fit indices.

Simple Structure. The 58 items were first tested with a 2-factor fit, and 7 items were excluded for not loading on any factor or loading on multiple factors with a loading greater than .30. After removing these items, all items loaded cleanly on the two factors. The loading table for this analysis can be found at XX. This same procedure was repeated for the 3-factor fit, and 10 items were removed from the first round of analysis, and 1 item was removed from the second round of analysis, resulting in a final set of 47 items. The loading table and items can be found at XX. Last, the 5-factor model showed 15 poor items on the first round of analysis, and 10 poor items on the second round of analysis, resulting in a final set of 33 items for the final model. The items and loadings can be found at XX.

We then compared items across all factor models to reduce scale length. Items were selected for the final model if they: 1) theoretically made sense with the other items on that factor, 2) loaded on the theorized 5-factor model (i.e., items were included in the 2-factor model if they also had been included in the 5-factor model), and 3) were not face-value redundant with other items on that factor. We tested these items and found that the 2-factor and 3-factor models continued to show simple structure, but the 5-factor model required the removal of 2 more items due to cross loadings across factors. To select the best model, we examined the fit indices for each factor structure. The 5-factor model showed the best fit (RMSEA = .038, RMSR = .024, TLI = .968, CFI = .982), followed by the 3-factor model (RMSEA = .049, RMSR = .028, TLI = .962, CFI = .977), and then the 2-factor model (RMSEA = .062, RMSR = .031, TLI = .963, CFI = .977). The final factor loadings for the 5-factor scale are shown in Table 1, and the other loadings can be found at XX (2-factor) and XX (3-factor).

EXPLAIN ITEM LABELS HERE

CFA

Using the multigroup CFA procedure described above, we examined the 5-factor model for overall fit and fit across groups. Table 2 portrays the results of the assessment of all models. The model showed acceptable fit overall with lower than desired values on the TLI, and lower values on the CFI and TLI for each individual group. Model fit appeared acceptable to investigate multi-group structure when examining the configural model with good values for the residual fit statistics, and poorer values for the goodness of fit statistics. We examined for potential areas of misfit using modification indices but did not find any suggested correlated error terms (i.e., two similar items that have related residuals) or cross-loadings that fit within theoretical structure. Therefore, we did not make any modifications.

Using the ΔCFI $\leq$ .01 rule for degradation in fit, we found that the loadings (*scalar*) showed invariance across groups, while a large degradation in fit was found between the loadings and intercepts (*metric*). We individually removed the equality constraint between groups for each item, and found one item that was different between groups: *I have children or students which take after me.* The intercept for the undergraduate sample on this item was: 2.26, $SE = 0.08$, while the Mechanical Turk sample showed a higher average item score: 3.45, $SE = 0.12$. Figure 1 demonstrates a visualization of those differences with higher intercepts for the Mechanical Turk group. The overall distribution for the biological factor is also displayed in this figure.

```
## # A tibble: 37 x 47
##    free.parameter npar       fmin       chisq      df      pvalue baseline.chisq
##    <chr>          <lvn.vctr> <lvn.vctr> <lvn.vctr> <lvn.v> <lvn.> <lvn.vctr>
##  1 Q2_13 ~~ Q2_13 97         1.382986   976.3883   453     0      3650.446
##  2 Q5_12 ~~ Q5_12 97         1.382640   976.1440   453     0      3650.446
##  3 Q4_11 ~~ Q4_11 97         1.381266   975.1736   453     0      3650.446
```

```
## 4 Q4_18 ~~ Q4_18 97        1.382443   976.0046   453     0      3650.446
## 5 Q4_8 ~~ Q4_8   97        1.378295   973.0765   453     0      3650.446
## 6 Q2_19 ~~ Q2_19 97        1.378511   973.2288   453     0      3650.446
## 7 Q5_17 ~~ Q5_17 97        1.377931   972.8194   453     0      3650.446
## 8 Q5_18 ~~ Q5_18 97        1.366113   964.4759   453     0      3650.446
## 9 Q2_20 ~~ Q2_20 97        1.383588   976.8129   453     0      3650.446
## 10 Q2_9 ~~ Q2_9   97        1.383783   976.9511   453     0      3650.446
## # i 27 more rows
## # i 40 more variables: baseline.df <lvn.vctr>, baseline.pvalue <lvn.vctr>,
## #   cfi <lvn.vctr>, tli <lvn.vctr>, nnfi <lvn.vctr>, rfi <lvn.vctr>,
## #   nfi <lvn.vctr>, pnfi <lvn.vctr>, ifi <lvn.vctr>, rni <lvn.vctr>,
## #   logl <lvn.vctr>, unrestricted.logl <lvn.vctr>, aic <lvn.vctr>,
## #   bic <lvn.vctr>, ntotal <lvn.vctr>, bic2 <lvn.vctr>, rmsea <lvn.vctr>,
## #   rmsea.ci.lower <lvn.vctr>, rmsea.ci.upper <lvn.vctr>, ...
```

Discussion

References

Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, *107*(2), 238–246. https://doi.org/10.1037/0033-2909.107.2.238

Bentler, P. M. (1995). *EQS structural equations program manual*. Encino, CA.

Brown, T. A. (2015). *Confirmatory factor analysis for applied research* (Second edition). New York ; London: The Guilford Press.

Buchanan, E. M. (2024). Visualizemi: Visualization, Effect Size, and Replication of Measurement Invariance for Registered Reports. *Assessment*, 10731911241280763. https://doi.org/10.1177/10731911241280763

Burke, B. L., Martens, A., & Faucher, E. H. (2010). Two Decades of Terror Management Theory: A Meta-Analysis of Mortality Salience Research. *Personality and Social Psychology Review*, *14*(2), 155–195. https://doi.org/10.1177/1088868309352321

Byrne, B. M., Shavelson, R. J., & Muthén, B. (1989). Testing for the equivalence of factor covariance and mean structures: The issue of partial measurement invariance. *Psychological Bulletin*, *105*(3), 456–466. https://doi.org/10.1037/0033-2909.105.3.456

Cheung, G. W., & Rensvold, R. B. (1999). Testing Factorial Invariance across Groups: A Reconceptualization and Proposed New Method. *Journal of Management*, *25*(1), 1–27. https://doi.org/10.1177/014920639902500101

Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, *9*(2), 233–255. https://doi.org/10.1207/S15328007SEM0902_5

Drolet, J.-L. (1990). Transcending death during early adulthood: Symbolic immortality, death anxiety, and purpose in life. *Journal of Clinical Psychology*, *46*(2), 148–160.

Erikson, Erik H. (1950). *Childhood and society*. New York: Norton.

Erikson, E. H. (1963). *Childhood and society* (2nd ed.). New York: Norton.

Greenberg, J., Pyszczynski, T., & Solomon, S. (1986). *The Causes and Consequences of a Need for Self-Esteem: A Terror Management Theory*. New York: Springer-Verlag.

Greenberg, J., Solomon, S., Veeder, M., Pyszczynski, T., Rosenblatt, A., Kirkland, S., & Lyon, D. (1990). Evidence for Terror Management Theory II: The Effects of Mortality Salience on Reactions to Those Who Threaten or Bolster the Cultural Worldview. *Journal of Personality and Social Psychology*, *58*(2), 308–318.

Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, *6*(1), 1–55. https://doi.org/10.1080/10705519909540118

Jöreskog, K. G., & Sörbom, D. (1996). *LISREL 8: User's Reference Guide*. Scientific Software International.

Kline, R. B. (2016). *Principles and practice of structural equation modeling* (Fourth edition). New York: The Guilford Press.

Kotre, J. (1984). *Outliving the self: How we live on in future generations*. New York: W. W. Norton & Company, Inc.

Mathews, R. C., & Kling, K. J. (1988). Self-Transcendence, Time Perspective, and Prosocial Behavior. *Journal of Voluntary Action Research*, *17*(2), 4–24. https://doi.org/10.1177/089976408801700202

Mathews, R. C., & Mister, R. D. (1987). Measuring an Individual's Investment in the Future: Symbolic Immortality, Sensation Seeking, and Psychic Numbness. *OMEGA - Journal of Death and Dying*, *18*(3), 161–173. https://doi.org/10.2190/NL7H-YY22-DN6Q-5XEB

McAdams, D. P., & St. Aubin, E. de. (1992). A theory of generativity and its assessment through self-report, behavioral acts, and narrative themes in autobiography. *Journal of Personality and Social Psychology*, *62*(6), 1003–1015.

Nye, C. D., Bradburn, J., Olenick, J., Bialko, C., & Drasgow, F. (2019). How Big Are My Effects? Examining the Magnitude of Effect Sizes in Studies of Measurement Equivalence. *Organizational Research Methods*, *22*(3), 678–709. https://doi.org/10.1177/1094428118761122

Nye, C. D., & Drasgow, F. (2011). Effect size indices for analyses of measurement

equivalence: Understanding the practical importance of differences between groups. *Journal of Applied Psychology*, *96*(5), 966–980. https://doi.org/10.1037/a0022955

Preacher, K. J., & MacCallum, R. C. (2003). Repairing tom swift's electric factor analysis machine. *Understanding Statistics*, *2*(1), 13–43. https://doi.org/10.1207/S15328031US0201_02

Revelle, W. (2020). *Psych: Procedures for psychological, psychometric, and personality research*. Retrieved from https://CRAN.R-project.org/package=psych

Rosseel, Y. (2012). Lavaan: An r package for structural equation modeling. *Journal of Statistical Software*, *48*(1), 1–36. https://doi.org/10.18637/jss.v048.i02

Steiger, J. H. (2016). Notes on the Steiger–Lind (1980) Handout. *Structural Equation Modeling: A Multidisciplinary Journal*, *23*(6), 777–781. https://doi.org/10.1080/10705511.2016.1217487

Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, *38*(1), 1–10. https://doi.org/10.1007/BF02291170

Zuckerman, M., & Neeb, M. (1979). Sensation seeking and psychopathology. *Psychiatry Research*, *1*(3), 255–264. https://doi.org/10.1016/0165-1781(79)90007-6

Table 1

Spiritual	Social	Personal	Biological	Physical	Item
0.93	-0.04	0.00	0.02	-0.01	I believe in an afterlife.
0.91	0.03	0.00	-0.02	0.00	My spirit or soul will live on after I physically die.
0.83	-0.01	-0.04	-0.01	0.07	I believe in an immaterial soul or spirit.
0.82	0.01	0.04	0.04	0.00	Death is not the end.
0.51	0.15	0.07	-0.03	-0.12	The ideologies I believe in such as my religion, philo
0.01	0.79	0.08	-0.08	-0.07	I have taken part in activities or events which will i
-0.05	0.76	0.00	0.03	0.15	I have contributed to the world in a unique way.
0.02	0.73	-0.06	0.12	-0.05	I have contributed to things which will be meaningf
0.06	0.65	0.04	-0.06	0.16	My life has influenced the world.
0.12	0.59	0.08	0.10	-0.17	I feel I have taken part in something bigger than m
0.13	0.35	0.07	0.22	0.13	I believe I will be remembered for a long time after
0.00	-0.02	1.02	-0.02	0.03	Parts of my personality have rubbed off on other pe
0.03	0.08	0.69	0.11	-0.02	Through my interactions with them, others have pi
-0.06	0.29	0.35	0.17	-0.11	I have imparted knowledge and skills on to others.
0.03	-0.03	-0.04	0.76	0.04	My offspring carry parts of me with them.
0.02	0.02	0.08	0.73	0.00	I am comforted by the fact that my genes can/will
0.08	0.11	0.14	0.44	-0.10	I am comforted by knowing my loved ones will live
-0.05	0.18	-0.05	0.35	0.20	I have children or students which take after me.
-0.01	0.01	-0.01	0.05	0.75	I believe will not die.
-0.01	-0.01	0.10	0.01	0.60	I have a hard time believing I will physically die.
0.13	0.03	-0.02	-0.01	0.62	I don't entirely believe I will die.
0.07	0.04	0.07	-0.02	0.47	I believe I will be reincarnated.

Table 2

Model	Chi-Square	df	CFI	TLI	RMSEA	SRMR
Overall	503.453	76	0.901	0.885	0.066	0.065
Group Undergraduate	433.000	76	0.876	0.856	0.068	0.070
Group Mechanical Turk	331.121	76	0.899	0.883	0.082	0.094
Configural	764.120	152	0.885	0.867	0.072	0.077
Loadings	790.521	135	0.882	0.869	0.072	0.082
Intercepts (Original)	895.284	118	0.855	0.845	0.078	0.086
Intercepts (Partial)	830.685	119	0.875	0.866	0.072	0.085
Residuals	914.583	97	0.855	0.852	0.076	0.085



Figure 1. The left panel shows the loadings (slope of the regression lines), intercepts (where each line crosses the y-axis), and residuals (inset bottom right) for the non-invariant biological item. The loadings are set to be equal, but the large difference in item averages can be seen as the separation between lines. The right panel indicates the shape of the distribution of the latent variable (which includes all biological items). The Mechanical Turk group shows a normal distribution around the latent variable mean (set to zero as part of MGCFA procedures), and the undergraduate population shows a negatively skewed distribution.