

<sup>1</sup> A Meta-Analysis of Expressive Writing on Quality of Life, Posttraumatic Growth, and  
<sup>2</sup> Posttraumatic Stress

## Abstract

Emotional expression has been shown to be beneficial for promoting both positive psychological and physical health outcomes. Unfortunately, inhibiting emotions can lead to impairments in physical and psychological health. James Pennebaker showed that expressive writing is an effective form of emotional expression, and he and others have used expressive writing as an experimental manipulation to gauge its efficacy in treating a wide variety of health-related and psychological outcomes. While many studies have been conducted that examine the efficacy of expressive writing across such outcomes, a considerable amount of these studies tend to neglect necessary considerations such as power and meaningfulness of respective effect sizes. Six previous meta-analyses have been conducted that examine expressive writing's effect on psychological outcomes. However, these studies focus on the experimental versus control group effect size. Thus, our meta-analysis sought to examine the efficacy of an expressive writing task on only the experimental conditions in studies measuring posttraumatic growth, posttraumatic stress, and quality of life using random effects models. Results indicated a small overall effect size for posttraumatic stress and negligible to small effect sizes for posttraumatic growth and quality of life. Implications for future research design and interpretation of published research are discussed.

*Keywords:* meta-analysis, positive psychology, expressive writing

## A Meta-Analysis of Expressive Writing on Quality of Life, Posttraumatic Growth, and Posttraumatic Stress

### **Emotional Expression**

Emotional expression relating to negative emotions or trauma has been shown to enhance both mental and physical health outcomes (Esterling, Antoni, Kumar, & Schneiderman, 1990; Fawzy et al., 1993; Lieberman & Goldstein, 2006; Rachman, 1980; Scheff, 1979). For example, the disclosure of traumatic or stressful events has been shown to reduce stress and lead to positive health outcomes in those with diabetes (Bodor, 2002) and breast cancer (Stanton et al., 2002), among others. Inhibiting repressive thoughts or emotions, rather, may be detrimental to both physical and psychological health (H. S. Goldstein, Edelberg, Meier, & Davis, 1988; Gross & Levenson, 1997; Larson & Chastain, 1990). While some studies suggest that emotional expression in the form of “truth telling” may cause psychological harm to individuals (Brounéus, 2010), the literature presents evidence confirming the negative effects of a lack of emotional expression, such as social concerns, overall psychological dysfunction, and lack of value-congruent behaviors (Frankl, 1959; Pennebaker, 1989; Pennebaker & Beall, 1986; Schulenberg, Hutzell, Nassif, & Rogina, 2008; Wilson & DuFrene, 2009). These resulting negative outcomes may lead to detrimental effects on health (Pennebaker & Beall, 1986). Individuals having experienced a traumatic or stressful life event are significantly more likely to repress thoughts and feelings about their experience compared to individuals who have not experienced such events, thereby subjecting them to potential negative outcomes caused by a lack of emotional expression (Bodor, 2002).

### **Expressive Writing as Effective Emotional Expression**

Pennebaker and Beall (1986) first showed that emotional expression can be both experimentally manipulated and have positive benefits to participants. In their seminal study, they randomly assigned participants to several writing groups, including writing about a “stressful or traumatic” life event or a neutral event. As such, the content of the writing

likely varies widely based on contextual factors (e.g., topic, setting, sample, health concern). The group that disclosed both regarding their trauma and the emotions surrounding said trauma later showed a reduction in health visits. Pennebaker has replicated the use of expressive writing across a number of studies ranging from improved health (Pennebaker, Colder, & Sharp, 1990; Pennebaker, Kiecolt-Glaser, & Glaser, 1988) to improvements in school (Pennebaker & Francis, 1996) and work (Francis & Pennebaker, 1992). Others have expanded this work to show positive effects on mood (Schoutrop, Lange, Hanewald, Davidovich, & Salomon, 2002) and asthma (Smyth, Stone, Hurewitz, & Kaell, 1999); however, several controlled studies have shown to not replicate (Harris, Thoresen, Humphreys, & Faul, 2005) or null effects (Gidron, Peri, Connolly, & Shalev, 1996; Walker, Nail, & Croyle, 1999). This protocol, more generally, has been termed written emotional disclosure (WED).

The idea that a brief, controlled writing task can have numerous positive health and psychological benefits can certainly be controversial, given the existing literature. For example, Henry, Schlegel, Talley, Molix, and Bettencourt (2010) found that expressive writing only benefited a rural population for those individuals surviving breast cancer on physical and psychological health outcomes, while Lancaster, Klein, and Heifner (2015) found no significant evidence that expressive writing can be considered an effective approach in measuring posttraumatic growth. Additionally and as mentioned, Brounéus (2010) found that “truth telling” caused harm to individuals in a forensic setting. Regardless, the concept remains interesting due to the nature and inexpensive implementation of expressive writing. Many individuals who have experienced traumatic events do not wish to disclose their feelings regarding the events with others. Additionally, those who do not meet diagnostic criteria (e.g., subclinical symptoms) are sometimes neglected despite probable suffering (Wilson & DuFrene, 2009). However, by utilizing expressive writing as a personal method of treatment, individuals are able to effectively express their emotions while avoiding talking to another individual or clinician about the traumatic event (Smyth, 1998). Pennebaker (1993)

found that experimental conditions assigned to participate in an expressive writing task generally report more positive changes than those in control conditions. Some controversy has been observed over whether or not writing about a formerly disclosed event is more effective than writing about an undisclosed event. M. A. Greenberg and Stone (1992) conducted an experiment where they separated participants into three groups: writing about a formerly disclosed trauma, writing about an undisclosed trauma, and a control group. They found no difference between groups in efficacy. However, they did find that those who disclosed more severe traumas reportedly experienced fewer physical symptoms at follow up, which suggests that the type of trauma revealed can play a significant impact on symptom reduction and physical health. A review of current meta-analyses relative to expressive writing is presented in a subsequent section.

### **Possible Mechanisms Underlying WED Efficacy**

In order to understand why expressive writing is considered to be efficacious, one must examine the cognitive, social, and behavioral processes by which it promotes information processing. Pennebaker et al. (1990) discovered that individuals who benefited from expressive writing attributed their success from the writing task to a renewed sense of understanding. Further, Pennebaker (1993) conducted a textual analysis on expressive writing content and found that those who were more successful during the task used causation words. Pennebaker thus concluded that expressive writing was a way for individuals to effectively process the event cognitively, which may explain the aforementioned renewed sense of understanding and excess of causation-oriented words. Aside from theories related to cognitive-processing and inhibition, there are a number of other theories related to emotional disclosure that warrant mentioning. The first is the social integration model (Pennebaker & Graybeal, 2001). This model suggests emotional disclosure can have a positive impact on how people interact in their environment. This increased environmental interaction has been shown to have positive benefits on health (Frattaroli,

2006). Second, expressive writing parallels exposure therapy for phobias and Posttraumatic Stress Disorder (PTSD), which suggests that repeatedly exposing oneself to the fear or trauma can reduce the negative emotions or physical sensations associated with that fear or trauma (Meshberg-Cohen, Svikis, & McMahon, 2014). Given that exposure therapy has been shown to be effective for reducing symptoms of posttraumatic stress (PTS; Sloan, Marx, & Epstein, 2005), one would expect individuals in these studies to experience a reduction in PTS symptoms after taking part in an expressive writing task. Third, Wilson and DuFrene (2009) discussed how the nonjudgmental acceptance of emotions leads to positive health benefits by promoting value-congruent behavior, one of the main facets of Functional Contextualism theory and Logotherapy (Frankl, 1959; Schulenberg et al., 2008). Indeed, emotional expression in the form of expressive writing could be considered a form of nonjudgmental acceptance, although it may not necessarily lead to behavior change in all cases. Finally, a recently proposed theory that may help explain positive outcomes is referred to as a distance perspective (Kross & Ayduk, 2011). This theory posits that, when individuals adopt a psychologically distanced perspective, they are better able to better understand their life situation. In sum, it seems likely that there are multiple underlying mechanisms that account for the beneficial outcomes associated with expressive writing described below. Indeed, the wide range of theoretical perspectives provide further evidence which suggests that expressive writing is applicable in a variety of contexts. Previously conducted meta-analyses, however, present varying results.

### **Meta-Analytic Techniques**

Meta-analyses allow researchers the opportunity to collectively examine the efficacy of different psychological interventions/tasks on outcome variables (Borenstein, Hedges, & Rothstein, 2007; Glass, 1976; Hedges, 1982). Although many studies produced positive outcomes associated with expressive writing, some of these studies tend to neglect important questions, the most important of which is whether or not the effect sizes are meaningful

(Smyth, 1998). Meta-analyses are a technique that allows researchers to pool studies to examine an overall, weighted, population effect (Borenstein et al., 2007). Several meta-analyses of expressive writing and emotional expression have been explored that warrant explanation: Smyth (1998), Frisina, Borod, and Lepore (2004), Frattaroli (2006), Reinhold, Bürkner, and Holling (2018), Van Emmerik, Reijntjes, and Kamphuis (2013) and Mogk, Otte, Reinhold-Hurley, and Kröner-Herwig (2006). These meta-analyses have laid a foundation for exploring the effects of writing on psychological outcomes.

Smyth (1998) conducted the seminal meta-analysis regarding the efficacy of expressive writing. They included studies utilizing an expressive writing group and control group (neutral topic). This particular analysis examined the efficacy of expressive writing on psychological well-being, general health, and physical functioning. In sum, thirteen studies/effect sizes were included, and the authors found an overall medium effect size,  $d = 0.47$ , for the experimental group compared to the control group. A later meta-analysis conducted by Frisina et al. (2004) expanded these analyses. They included studies utilizing clinical samples and employing the paradigm adapted by Pennebaker and Beall (1986). This meta-analysis included nine studies in total and found an effect size of  $d = 0.19$  for health-related outcomes and  $d = 0.07$  for psychological outcomes. The next expressive writing meta-analysis was conducted by Mogk et al. (2006) and aimed to update the state of the literature on expressive writing. Similar to previously-conducted analysis, they included studies employing Pennebaker's paradigm on experimental and control groups. Additionally, they only included studies with a four-week follow up that included at least ten participants. In sum, thirty studies met their criteria. They found nonsignificant effects on somatic and psychological health outcomes and concluded that expressive wrting does not promote health-related outcomes. These findings corroborate those from Frisina et al. (2004).

Frattaroli (2006) conducted perhaps the most notable meta-analysis to date examining the efficacy of emotional disclosure on the folliwing constructs using only randomized and control conditions: psychological health, physiological functioning, reported health, health

behaviors, and general functioning/life outcomes. Additionally, their meta-analysis was the first to employ random effects models, which estimate the mean of a proposed distribution of population effect sizes. Prior meta-analyses employed fixed effects models, which assume that all studies assess the same “true” population effect size, which may be an untenable assumption across different assessment and populations (Borenstein et al., 2007). They included a wide range of studies  $N = 146$ . Individual studies were again collapsed into one publication effect size, although these effects were also examined separately by health outcome. Overall, Frattaroli (2006) found a weighted  $r$  effect size of .08 for all outcomes combined, which would be considered small. Additionally, they examined potential moderators and found larger effect sizes for the following samples: those with physical health problems, those with a history of having experienced traumatic or stressful events, samples not including college students, samples where expressive writing tasks were conducted at home and in private settings, paid participants, more male participants, and fewer participants (see Frattaroli, 2006 for a complete list of moderators). A recent analysis conducted by Van Emmerik et al. (2013) employing Pennebaker’s paradigm included six eligible studies that compared treatment to control groups. In regards to inclusion criteria, they included studies where participants had a diagnosis of Acute Stress Disorder (ASD) or PTSD. They found that those who participated in the expressive writing group experienced short-term reductions in PTS and comorbid depressive symptoms, combined Hedges’  $g = 0.81$ . The most recently published meta-analysis was conducted by Reinhold et al. (2018) and examined the effects of expressive writing on depression by randomizing participants to conditions (expressive writing vs. control). They included thirty-nine randomized controlled trials and excluded individuals with diagnoses of PTSD. This study did not support utilizing expressive writing for depression outcome measures for the specified sample, Hedges’  $g = -0.09$ . Further, they found that expressive writing did not yield any type of long-term effect on depression outcomes.



## Posttraumatic Stress

Posttraumatic Stress Disorder (PTSD) is a disorder involving re-experiencing thoughts or experiences after a traumatic event or experience. This generates a context where individuals are prone to affect-related deficiencies and maladaptive behaviors (American Psychiatric Association, 2013). DSM-5 criteria are based on twenty symptoms structured into four different subsets in those having experienced a traumatic event. These subsets are as follows: re-experiencing, avoidance, negative alterations in cognition and mood, and increased arousal (Crespo & Gomez, 2016). While the renewed DSM-5 criteria are now increasingly employed, the current meta-analysis considers studies using DSM-IV criteria. DSM-IV criteria are similar and include the following: exposure to a traumatic event, re-experiencing (intrusion), avoidance, and increased arousal (American Psychiatric Association, 2013). Further, the studies employed in the current meta-analysis are divided according to these subsets (arousal, intrusion, and avoidance). PTSD affects a wide variety of groups, a few of which are sexual assault survivors (Klump, 2008), Iraq and Afghanistan war veterans (Gentes et al., 2014), and those exposed to natural disasters (Wang et al., 2000).

Research conducted on the efficacy of expressive writing on PTSD symptoms presents intriguing results. Sloan, Marx, Epstein, and Lexington (2007) examined individuals with at least moderate PTSD symptom severity and found that individuals assigned to an emotional expression writing condition reported fewer PTSD and depression symptoms during follow up. Sloan, Marx, and Greenberg (2011) found that PTSD symptoms decreased after a written emotional disclosure task, although this decrease was not significantly different than a control group change. Di Blasio et al. (2015) recruited women who had just given birth and assessed them a few days after experiencing childbirth along with a three-month follow-up. Results showed that women who had participated in the expressive writing task had lower depression and posttraumatic stress symptoms than the group assigned to a neutral writing condition. Additionally, regression models showed that expressive writing was significantly linked to a reduction of PTSD symptoms across different dimensional levels

of symptom severity. Only 20 of the 113 women recruited for this study qualified for a diagnosis of PTSD, but those who reported mild symptomology responded better to the task than those meeting criteria for PTSD. This limitation suggests that those with moderate distress could perhaps benefit more from an expressive writing task than those diagnosed with or meeting the qualifications for PTSD. It may also explain the differences in results in comparing to Sloan et al. (2011), as they found that those with a clinical diagnosis of PTSD did not respond to an emotional disclosure writing task. Perhaps it may be more advantageous to examine effect sizes separately for diagnoses of PTSD and subclinical symptoms. Further Sloan, Marx, Bovin, Feinstein, and Gallagher (2012) adapted a writing protocol to focus primarily on the emotions, meaning, and “hot spots” associated with the trauma. They referred to this procedure as the written exposure therapy (WET) protocol, distinguishable from the paradigm adapted by Pennebaker and Beall (1986). In their seminal study examining the efficacy of WET for motor-vehicle accident related PTSD, they found that those in the WET condition experienced significant reductions in PTSD symptoms throughout the course of the study. Since then, a small number of other studies employing the WET procedure have been employed in those with PTSD. While these will be included in the current review, the newness of this protocol does not allow exclusive examination using meta-analytic techniques.

## **Posttraumatic Growth**

While the literature mostly discusses potentially harmful outcomes to traumatic events such as emotional distress, traumatic events also provide opportunities for personal growth (Aslam & Kamal, 2013). Traumatic events, either natural or human-inflicted, can lead to positive outcomes by allowing the individual to take a different perspective (Cobb, Tedeschi, Calhoun, & Cann, 2006; Taku, Calhoun, Cann, & Tedeschi, 2008). The relationship between positive growth after a traumatic event and symptom reduction is unclear, as it is a complex process. Thus, it is necessary to examine how expressive writing might influence each

variable separately, which is one of the key goals of this meta-analysis (Slavin-Spenny, Cohen, Oberleitner, & Lumley, 2011). Models receiving empirical support within the last decade suggest that traumatic events offer opportunities for both negative and positive experiences (Tedeschi & Calhoun, 1995; Weiss, 2002). Posttraumatic Growth (PTG) is a positive experience after a traumatic event (Aslam & Kamal, 2013; Yilmaz & Zara, 2016). Specifically, PTG is classified as broad cognitive benefits that are seen after a traumatic experience. These benefits can be categorized into building closer relationships, examining new possibilities, appreciating life, recognizing personal strengths, and undergoing spiritual changes (Dursun, Steger, Bentele, & Schulenberg, 2016; Tedeschi & Calhoun, 2004).

PTG is associated with a variety of desired outcomes (Dursun et al., 2016). PTG has been studied in those experiencing natural disasters, war, and other harms such as sexual assault. Finally, PTG has been studied in those experiencing medical diagnoses such as different types of cancer and diseases. Although the relationship between PTG and symptom reduction is not yet fully understood, perhaps expressive writing allows the individual to fully comprehend the event. Pennebaker and Graybeal (2001) speculated that expressive writing allows an individual to feel more connected with his or her surroundings. Although this speculation does not directly explain positive outcomes after an expressive writing task, perhaps individuals gain a better appreciation for life after gaining a better sense of connectedness with that individual's surroundings. One might expect effect sizes to be larger for those studies requiring a diagnosis of PTSD, as such growth may not be possible in those with subclinical symptomology.

## Quality of Life

Quality of Life (QOL), according to Theofilou (2013) is an evaluation of the "goodness" that an individual experiences, separated into domains of reactions to life events, disposition, life fulfillment, and satisfaction with life experiences. More generally, QOL refers to an individual's attitude towards the target life situation (Costanza et al., 2007), delineated into

objective and subjective components. Objectively, QOL refers to components outside of an individual and measurable by others, while subjective QOL is an individual's assessment of his or her own experiences (Costanza et al., 2007). The current meta-analysis will focus solely on the subjective components of QOL, as it is obtainable through questionnaires. Pennebaker and Graybeal (2001) suggested that expressive writing allows one to feel more connected with their surroundings. Further, they explain that expressive writing allows people to see things in a different way and better understand themselves. By understanding a traumatic or stressful event, one is said to see things differently and perhaps look at the situation with a more positive mindset. The changes that occur after expressive writing may also allow one to find meaning in the traumatic event, thereby increasing the QOL of that individual (Frankl, 1959). Higher QOL may be considered a type of PTG, which is why the current meta-analysis sought to examine the efficacy of studies utilizing expressive writing to improve QOL and PTG in the same study.

## Current Meta-Analysis

The purpose of the current meta-analysis is to examine studies employing expressive writing procedures using Pennebaker's paradigm (WED) and the more recent WET protocol on variables relevant to the field of positive psychology (PTG and QOL) and PTS, with effect sizes separated by the paper's indication of PTSD diagnosis when sample sizes are large enough. Based on recently published literature regarding efficacy of expressive writing for different levels of PTSD symptoms, this marker is an important facet to consider (Di Blasio et al., 2015; Reinhold et al., 2018; Sloan et al., 2011). No review has examined the effects of expressive writing on PTS separated by diagnosis. Additionally, no meta-analysis has been conducted that examines the efficacy of expressive writing on positive outcome variables such as PTG and QOL, in line with the field of positive psychology. The meta-analyses described sequentially above also focused on experimental versus control group effect sizes or *p*-values, rather than emphasizing change for the expressive writing group. This focus is likely because

of the analyses provided in these publications, especially when using randomized controlled trial research designs. While this design is the gold standard for medicine, the current meta-analysis sought to examine the magnitude of change for participants who experienced an expressive writing task. For example, a comparison group may increase their quality of life scores by two points in a controlled study, while the experimental group increases their quality of life scores by four points; thus, creating a significant difference in change between the two groups. This information is valuable, but it does not tell the reader the magnitude of the change for the writing group, wherein four points might only be a small effect when examined within the group who received the writing task.

This analysis will also focus on changes across time for groups who received the expressive writing task to determine what size of effects one might expect given a specific measurement schedule (i.e., one to three months, three months to six months, etc.). This analysis should present researchers with a renewed examination of the efficacy of expressive writing on the aforementioned variables using newer meta-analytic techniques. Newer methods of meta-analysis, including *p*-curve (Simonsohn, Nelson, & Simmons, 2014; Simonsohn, Simmons, & Nelson, 2015), *p*-uniform (Aert, Wicherts, & Van Assen, 2016), PET-PEESE (Stanley & Doucouliagos, 2014), selection models (Vevea & Hedges, 1995), and trim and fill methods (Carter & McCullough, 2014) allow for better estimation of meta-analytic effect sizes. These analyses would be best performed by examining each potential effect separately, rather than averaging effects of each publication into one study effect size (a common trend in the previously mentioned meta-analysis). In addition to an estimate of overall effect sizes using updated techniques, the current meta-analysis estimates power for effects on writing groups, as research has shown a consistent underpowering of psychological studies, combined with a misunderstanding of the sample size needed for adequately powering one's work (Bakker, Hartgerink, Wicherts, & Maas, 2016).

## Method

### Data Collection

Studies were collected through online databases, such as PsycINFO and Google Scholar, using the following search terms and their combinations: *Posttraumatic Growth*, *PTG*, *Quality of Life*, *QOL*, *Posttraumatic Stress*, *PTS*, *Expressive Writing*, *Emotional Disclosure*, *Written Emotional Disclosure (WED)*, *Written Exposure Therapy (WET)*. Within these articles, the change in outcome variables (PTS, PTG, QOL) from pre- to post-test was the dependent variable of interest. Generally, groups were separated into an experimental and control group and then examined at different time points. For purposes of this meta-analysis, only participants assigned to the experimental condition were examined due to having received the expressive writing task. If a study included multiple assessment time points, then these measurements were examined sequentially (i.e., time 1 to time 2, time 2 to time 3) to determine change across time for the dependent variable.

264 citations focusing on PTS, PTG, and QOL were identified through the literature search and previous meta-analyses. After screening these studies, 53 articles were retained for containing the appropriate information for this meta-analysis. This manuscript was written with *papaja* in *R* (???) with the analyses inline with the text. The complete set of data, excluded article list with reasoning, and other relevant information can be found at: <https://osf.io/4mjqt>. Generally, studies were included if they utilized WED or WET, included relevant numbers to compute an effect size, and included the relevant outcome variables. After having two reviewers independently code articles, 223 effect sizes were calculated. On average, each study represented  $M = 4.21$ ,  $SD = 3.31$  effects, ranging from 1 to 16 effects. 165 effects were calculated for PTS, 21 for PTG, and 37 for QOL. Studies were coded for PTSD diagnosis as no (not mentioned or not included), mixed (mentioned number of participants but all included), and yes (included as criteria). After examining the number of effects in each of these categories for each variable, only the PTS results will be split by PTSD diagnosis with 88 no mention, 32 in the mixed category, and PTSD\_diag[2,3] yeses.

### Calculations for Effect Size, Variance, and Confidence Intervals

For our purposes, we used Cohen's (1988) standards for nomenclature for small (0.20), medium (0.50), and large (0.80)  $d$  values, although it is important to note that Cohen himself suggested that these values should be based on the area of study. Generally, however, these effect size criteria are used within the social sciences. Each study implemented a pre-test to post-test style repeated measures design, usually with paired  $t$ -tests, ANOVA, or regression analyses. The means, standard deviations, and  $N$  values were collected from each study. In general, Cohen's  $d$  values were calculated using the following formula for paired  $t$  using means and standard deviations:

$$d_{av} = \frac{M_1 - M_2}{\frac{SD_1 + SD_2}{2}}$$

This equation is described in detail in Cumming (2012) as an alternative to the traditional calculation of  $d$  for paired samples  $t$ , wherein the denominator is the standard deviation of the difference scores:

$$d_z = \frac{M_1 - M_2}{SD_{diff}}$$

This equation for  $d_{av}$  not only allows for calculations from published articles that do not include  $SD_{diff}$  (i.e., most articles included), but also has been shown to be less upwardly biased than  $d_z$ . Alternative formulas include controlling for  $r$  between paired levels, as described in Lakens (2013); however, these values were not available in the selected articles, and Lakens also recommends  $d_{av}$  as an effect size for paired designs. When only mean differences and standard deviation of the difference scores were available, the second equation for  $d_z$  was used.

We planned to use traditional and newer methods of meta-analysis, following guidelines from Cooper, Hedges, and Valentine (2009) and Borenstein et al. (2007), as well as Aert et al. (2016). Sampling variance of the effect sizes were estimated using the *escalc()* function

from the *metafor* package in *R* (Viechtbauer, 2010). The variance formula was originally published in S. B. Morris and DeShon (2002) and is shown below:

$$v = \frac{1}{n} \left( \frac{n-1}{n-3} \right) (1 + n * d^2) - \frac{d^2}{[c(n-1)]^2}$$

In this formula,  $n$  is the number of paired observations,  $d$  is the calculated effect size, and  $c$  is a correction factor, wherein  $df$  are  $n - 1$  (Hedges, 1982):

$$c = 1 - \frac{3}{4 * df - 1}$$

We used the *metagen()* function in the *metafor* package to calculate both fixed and random effects models, which uses standard error of the effect to calculate overall estimates of an effect and their confidence intervals. Thus, we took the square root of the variance estimate for standard error. Given these calculations, the goal of this analysis was to calculate a combined effect size, along with a confidence interval for study planning and an assessment of the literature. A fixed effects model requires the assumption that there is a true population effect size across all studies. By including multiple measures of psychological outcomes, this assumption may be tenuous, and therefore, a random effects model was also calculated. In random effects models, the true effect is assumed to vary across studies (Borenstein et al., 2007). For a fixed effects model, the effect sizes are weighted by their inverse variance ( $v$ ; Sánchez-Meca & Marín-Martínez, 2008), which is calculated automatically in *metafor* by:

$$w_i^{FE} = \frac{1}{v}$$

The advantage to this procedure is that analyses are weighted by their precision, that is, that studies with more information (often, larger samples), are given larger weights in the overall estimated effect size (Borenstein et al., 2007). Random effects models are also weighted by inverse variance, with an additional correction for variance between studies,  $\tau_{DL}^2$ , as described by DerSimonian and Laird (1986):



$$w_i^{RE} = \frac{1}{v + \tau_{DL}^2}$$

Confidence intervals were calculated in two ways for this study. Cumming (2012), Kelley (2007), and Smithson (2001) have shown that the distribution of  $d$  values are non-normal, and thus, CIs should be estimated using the non-centrality parameter and a non-normal distribution. These values were calculated using the functions in the *MOTE* library which iteratively estimates the appropriate non-centrality parameter and converts back to  $d$  values (i.e., non-centrality parameter divided by the square root of  $n$ ; Buchanan, Valentine, & Scofield, 2017; Smithson, 2001, 2003). However, the *metafor* package in *R* uses central distributions to estimate CIs for each study and overall effect sizes. Therefore, we present both sets of values for the interested reader, as meta-analytic procedures have not implemented non-central distributions of effect sizes.

### Additional Meta-Analytic Techniques

**p-Curve and p-Uniform.** We used *p-curve.com* to conduct a *p-curve* analysis (Simonsohn et al., 2014). The purpose of this type of analysis is to detect true effects. Specifically, *p-curve* is used to reveal possible *p-hacking* in published literature in order to decipher whether or not a true effect exists. Broadly, *p-hacking* occurs when researchers use questionable research practices to create significant results by manipulating dependent variables or covariates. Additionally, authors may add participants if the initial findings are not significant (Bruns & Ioannidis, 2016). Researchers may also decide to exclude participants for final analyses if that exclusion leads to a significant difference (L. K. John, Loewenstein, & Prelec, 2012). Thus, it is necessary to distinguish between true and false effects in order to effectively interpret effect sizes corresponding to those *p-values*. *p-curve* accomplishes this task by examining the distributions of the published *p-values*. If an effect exists, or rather the results should be interpreted as presented, the distribution of *p-values* will be positively skewed (Simonsohn et al., 2014). If, however, no effect exists, then the

distribution of  $p$ -values will be flat.  $p$ -curve analyses ultimately provide evidence of  $p$ -hacking in groups of studies and has become an important tool for interpreting meta-analyses. In order to accurately estimate effect sizes because of scrutiny associated with effect size estimation of  $p$ -curve, we also conducted  $p$ -uniform.  $p$ -uniform analyses, too, are interpreted by examining the distribution of  $p$ -values in a set of studies (Aert et al., 2016). However, it is assumed that the population effect size equals the effect size from the dataset. Because of this assumption, the population effect size is referred to as uniform. This analysis also examines for publication bias and presents the researcher with a corrected effect size. Publication bias occurs when only select studies are published, usually only significant studies, although many factors can bias a study's publication (McShane, Böckenholt, & Hansen, 2016).  $p$ -uniform was calculated from code provided by Van Aert (2017) on GitHub.

**PET-PEESE.** Originally, meta-analyses relied on the calculation of Egger's regression test which examined the relationship of the standard error (predictor) to the effect size estimates (criterion). In this regression, the intercept values were used to determine if effect size measures were different than zero, by providing a meta-analytic estimate (Egger, Davey Smith, Schneider, & Minder, 1997; Stanley, 2005). PET-PEESE analyses examine for publication bias by adapting parts from Egger's traditional regression tests: PET (Precision Effect Test) and PEESE (Precision Effect Estimate with Standard Error, Carter & McCullough, 2014). PET is a more reliable test of publication bias with effect size estimates of zero,  $b_0 = 0$ , while PEESE is more accurate with non-zero effect size estimates,  $b_0 \neq 0$  (Stanley & Doucouliagos, 2014). PET-PEESE was calculated using Hilgard's (2016) code provided on GitHub.

**Selection Models.** Selection model analyses provide the researcher with a test of publication bias and effect size estimates using maximum likelihood estimation (Vevea & Hedges, 1995; Vevea & Woods, 2005). Using selection models, researchers are able to discover effect size estimates as well as evidence of publication bias (McShane et al., 2016) by using a mixed general linear model to estimate these values. Selection models were

calculated with the *weightr* package in *R* (Coburn & Vevea, 2017).

**Trim and Fill.** Trim and Fill analyses, in contrast to PET-PEESE, regress standard error (criterion) and effect size estimates (predictor). Specifically, the purpose of Trim and Fill techniques is to examine whether or not publication bias may influence the regression equation (Carter & McCullough, 2014). Effect sizes and standard error terms are graphically displayed on x and y-axes, respectively, in a funnel plot. If this graphical representation indicates asymmetry, considered a gap of missing data points in the lower center area of the plot, the study set can be assumed to have studies that are both non-significant and small in sample size (Van Assen, Van Aert, & Wicherts, 2015). This funnel is then trimmed until symmetry is achieved. Missing studies from the symmetrical graph are imputed (filled) while maintaining the given symmetry (Duval & Tweedie, 2000). The meta-analytic effect size is then estimated from the trimmed and filled funnel plot. Trim and fill analyses, as well as funnel plots included below, were calculated with the *metafor* package.

## Results

### PTS

**Overall Effect Size.** As described above, both fixed effects and random effects models with centralized confidence intervals are presented in Table 1. Studies were examined for potential outliers using the *metafor* package in *R*. This package calculates traditional regression influence values, such as Cook's and hat values (J. Cohen, 1988). These values indicate change in overall meta-analytic model with and without the effect; thus, determining their impact on the pooled effect size (Viechtbauer, 2010). Because published studies likely represent the range of the sampling distribution of effect sizes, we included the analyses with and without outliers to present evidence for both paths a researcher might take when examining an overall effect.

3 outliers were detected with this procedure, all showing very large effect sizes, average  $d = 2.35$ . The fixed and random effects estimates without these points are also included in

Table 1. Figures 1, 2, 3, and 4 portray the effect sizes for PTS studies, separated by intrusions, avoidance, hyperarousal, and total scores for easier viewing (i.e., over 100+ effect sizes did not fit easily on one combined graph). Although these categories are not reflective of updated DSM-5 criteria, researchers have not yet conducted enough studies using expressive writing on PTS with updated PTSD criteria to warrant a meta-analysis. Name acronym coding can be found in the data online. This forest plot includes the non-centralized confidence interval calculated from the *MOTE* library (Buchanan et al., 2017). Shape size indicates study weight, and these values were taken from the overall random effects meta-analysis and normalized by dividing by the mean weight. The dashed lines indicate the average non-weighted lower and upper confidence interval limit for the non-centralized estimates. Overall, PTS studies include a small effect size that appears to be significantly greater than zero across all estimate types (fixed, random, with or without outliers).

We further calculated the overall effect sizes by PTSD diagnosis category. Studies only including individuals with a diagnosis of PTSD exhibited a medium to large effect size (before and after outlier exclusion), while studies not requiring (or listing) a PTSD diagnosis showed a small to medium effect size. Similarly, the mixed category showed a small to medium effect size. Complete estimates are included in Table (LIST OUR SUPER SUPPLEMENTAL TABLE LINK HERE).

**Homogeneity.** A prerequisite for newer meta-analytic techniques includes the assessment of homogeneity of the effects (Aert et al., 2016). Using the *metafor* package in *R*, we calculated the  $Q$ -statistic and the  $I^2$  index (Cochran, 1954; Huedo-Medina, Sánchez-Meca, Marín-Martínez, & Botella, 2006). Significant values imply inconsistencies across the variable or variables of interest and are represented by  $Q$ . In contrast,  $I^2$  indicates the percentage of heterogeneity along with a 95% CI. Both can, however, be biased with a small number of experiments included for analyses (Higgins, Thompson, Deeks, & Altman, 2003; Huedo-Medina et al., 2006). Thus, we sought to calculate an overall level of heterogeneity after examining each variable separately before and after excluding outliers.

For PTS studies including outliers, we found significant heterogeneity,  $Q(164) = 780.46$ ,  $p < .001$  and  $I^2 = 79.0$ , 95% CI[75.8 - 81.8]. These values were reduced slightly with the exclusion of outliers,  $Q(161) = 642.72$ ,  $p < .001$  and  $I^2 = 75.0$ , 95% CI[70.9 - 78.5].

**Power.** Power was calculated in two different ways using the *pwr* package in *R* (Champely, 2016). *Post hoc* power was first calculated using sample size and effect size statistics from each individual study. Additionally, we calculated power using the study sample size and estimated overall effect size from the random effects model with and without outliers, as explained by G. Francis (2012) and G. Francis (2014). The first estimate indicates the likelihood of finding an effect from our sample statistics, while the second indicates the likelihood of finding the true population effect size. If each study had been conducted on only the change in the experimental group, 46.1% of studies would have been considered significant at  $\alpha < .05$ . The average power of these studies based on their original study characteristics was .48 ( $SD = .36$ ). Power for the random-effects meta-analytic effect size with outliers was .52 ( $SD = .25$ ) and without outliers was .47 ( $SD = .24$ ). Therefore, power consistently was around 40-50% for studies examining PTS, regardless of outlier effects. In these studies, only 28.5% achieved recommended 80% power for their found effect size, a smaller 23.6% for the random-effect outlier effect size, and even smaller 17.6% for power calculations on the random-effect size without the outliers.

**Other Meta-Analytic Estimates.** As noted in Aert et al. (2016), *p*-curve and *p*-uniform analyses are upwardly biased when heterogeneity is high. Therefore, we use caution when interpreting these analyses on PTS outcomes. As seen in Table 1, the estimates for *p*-uniform were higher than other techniques, likely because of the focus on significant *p*-values and the great degree of heterogeneity described earlier. *P*-curve pictures can be found at <https://osf.io/4mjqt/> online, and this analysis indicated evidentiary value at  $p < .001$ . Additionally, the *p*-uniform analysis indicated that there was likely no publication bias present,  $Z = -5.76$ ,  $p = 1.000$ . When examining the PET analysis, we found that the intercept was significant, which indicated that PEESE was likely a better estimator of the

meta-analytic effect size. PEESE estimates were lower than the original meta-analytic estimate, but confidence intervals indicated that the effect is small to medium, and still larger than zero. Selection models indicated a larger effect size, especially with the random-effects models, and these effects were influenced by the outliers found in the published studies. Trim and fill models are shown in Table 1, and figures are included online. Nineteen missing studies were imputed for both models with and without outliers. Across all these effect size estimates, we found that expressive writing was likely to decrease PTS symptoms in a small to moderate way. The correlation of effect size with time between measurement times was  $r = -.01$ , 95% CI  $[-.17, .14]$ ,  $t(163) = -0.17$ ,  $p = .865$ , and  $r = -.07$ , 95% CI  $[-.22, .09]$ ,  $t(160) = -0.89$ ,  $p = .377$  without outliers. This result indicated that the effect of expressive writing slightly decreased across time.

## PTG

**Overall Effect Size.** Both fixed and random effects models with centralized confidence intervals for PTG are presented in Table 2. When examining expressive writing on PTG, no outliers were detected. Fixed and random effects estimates are included in Table 2, while Figure 5 shows effect sizes for PTG studies where shape size indicates the normalized weight of the study. Dashed lines indicate non-weighted lower and upper confidence intervals for non-centralized estimates. Overall, PTG studies indicated a negligible to small effect size across both random and fixed effects models, and the non-centralized confidence intervals indicated an effect that crossed zero.

**Homogeneity.** Using the *metafor* package in *R*, we calculated both a  $Q$  statistic and  $I^2$  index. Since PTG studies did not contain any outliers, we did not calculate two separate analyses to examine heterogeneity both with and without outliers. We did not find significant heterogeneity across PTG studies,  $Q(20) = 14.18$ ,  $p = .82$  and  $I^2 = 0.0$ , 95% CI  $[0.0 - 25.3]$ .

**Power.** First, we calculated *post hoc* power using both sample and effect size statistics from individual studies. Individual studies examining change in experimental

groups showed that 9.5% of studies would have been considered significant at  $\alpha < .05$ . Average power of PTG studies was .15 ( $SD = .16$ ). 0.0% achieved recommended 80% power for their found effect size. Additionally, we calculated power using study sample size and estimated effect size from our random effects model. Power for the true effect size was .08 ( $SD = .02$ ). Again, 0.0% achieved recommended 80% power.

**Other Meta-Analytic Estimates.** Due to no heterogeneity across PTG studies, we can use both  $p$ -curve and  $p$ -uniform analyses with more confidence. A pictorial representation of  $p$ -curve can be found at <https://osf.io/4mjqt/>. This analysis did not indicate evidentiary value,  $p = .75$ , as only two of the results would be considered significant at  $\alpha < .05$ .  $p$ -uniform estimates are presented in Table 2. Specifically, these analyses indicated that there was no publication bias present,  $Z = 0.70$ ,  $p = .243$ . The  $p$ -uniform estimates of the effect size for PTG were negative, in contrast to the fixed and random effects overall model. The confidence interval for this analysis indicates a wide range of possible effects. In examining PET-PEESE analyses, we did not find a significant intercept, indicating that PET is most likely a better effect size estimator. PET analyses indicated that the effect size is negligible to small, with our confidence interval crossing zero. These results corroborated our original effect size calculations. Selection models indicated negligible to small effect sizes, again wherein the confidence interval includes zero effect. Trim and fill models are shown in Table 2, and figures are included online. Zero studies were imputed for our model, and thus, the effect size estimate is the same as the overall model. Across techniques, we found that expressive writing has little to no effect on PTG. The correlation of effect size across measurement times in PTG studies at subsequent time points was  $r = .09$ , 95% CI  $[-.36, .50]$ ,  $t(19) = 0.38$ ,  $p = .707$ , and no change over time was found.

## QOL

**Overall Effect Size.** Finally, for QOL, both fixed and random effects models with centralized confidence intervals are presented in Table 3. Two outliers were detected with

this procedure, average  $d = -0.07$ . While the average effect of these outliers indicates a small number, it is important to note that these two outliers were the largest positive and negative effects found from the Possemato, Ouimette, and Geller (2010) study. Fixed and random effects estimates without these points are also included in Table 3, while Figure 6 shows effect sizes for QOL studies. Overall, QOL studies indicated a negligible to small effect that showed a non-significant decrease in quality of life as a result of expressive writing.

**Homogeneity.** For QOL studies including outliers, we found significant heterogeneity from our random effects model,  $Q(36) = 200.09$ ,  $p < .001$  and  $I^2 = 82.0$ , 95% CI[75.9 - 86.5]. After excluding outliers, our random effects model still indicated heterogeneity,  $Q(34) = 93.18$ ,  $p < .001$  and  $I^2 = 63.5$ , 95% CI[47.6 - 74.6].

**Power.** In conducting *post hoc* power using sample and effect size statistics from individual studies, we found that 21.6% of studies would have been considered significant at  $\alpha < .05$ . Average power based on actual study characteristics was .33 ( $SD = .32$ ). Power for the random effects meta-analytic effect size with outliers was .05 ( $SD = .00$ ) and without outliers was .05 ( $SD = .00$ ). Unfortunately, power was around 5% for both random effects models with and without outliers. In these studies, 18.9% achieved adequate power of 80% on their found effect size, while 0.0% achieved 80% power for our random effects model with outliers. Finally, without outliers, 0.0% achieved 80% power.

**Other Meta-Analytic Estimates.** We exert caution in interpreting  $p$ -curve and  $p$ -uniform analyses on QOL outcomes with and without outliers due to heterogeneity. As seen in Table 1,  $p$ -uniform estimates were stronger and positive than other techniques because of the high degree of heterogeneity recently described.  $p$ -curve pictures can be found at the following OSF Link: <https://osf.io/4mjqt>. Eight studies were significant at  $\alpha < .05$ , and the studies indicated evidentiary value,  $p = .004$ .  $p$ -uniform analyses did not indicate publication bias,  $Z = -2.75$ ,  $p = .997$ . In PET-PEESE analyses, we found that the intercept was not significant, and therefore, PET was a better estimator of the meta-analytic effect. Table 1 indicates that both of these analyses estimate the effect size around zero, with a



confidence interval that includes zero. Selection models correspondingly show small effects crossing zero, except for random effects models with outliers, that appear to be heavily influenced by the outliers. Trim and fill models are shown in Table 3, and figures are included online. No studies were imputed for these analyses, and therefore, the effect size estimates match the original meta-analysis. Overall, these results appear to point to no effects, ranging across zero with several negative estimates. Interestingly, the correlation of effect sizes across measurement times with outliers was  $r = -.37$ , 95% CI  $[-.62, -.05]$ ,  $t(35) = -2.33$ ,  $p = .026$  and  $r = -.64$ , 95% CI  $[-.80, -.39]$ ,  $t(33) = -4.75$ ,  $p < .001$  without outliers. The effect of expressive writing appears to be positive at short time intervals and decreases into negative effects at longer time intervals.

## Discussion

In examining pre- to post-test comparisons across each variable separately, we found that PTS studies indicated a small effect size across all meta-analytic estimates. Interestingly, those studies requiring a diagnosis of PTSD for inclusion resulted in a medium to large effect size (before and after outlier exclusion), while those studies not requiring a PTSD diagnosis resulted in a small to medium effect size. These results suggest that those with clinical symptoms of PTSD may benefit more from expressive writing interventions. Further, these results are in contrast to recently-conducted studies, which suggest that those with subclinical symptoms benefit more from expressive writing tasks (???; Sloan et al., 2011). Both QOL and PTG studies indicated a negligible to small effect size using random effects models. Although the PTG effect in our overall meta-analysis estimate was significant, other methods indicate this small effect is likely not different from zero. We also examined the relationship of time between measurements of the dependent variables and the corresponding effect size to determine if effects change over time. For both PTS and PTG, there was no relationship between effect size and time; yet, PTS indicated a small negative correlation. This correlation was not, however, significant. For QOL studies, a medium to

large negative correlation was found. A negative relationship between time and effect size implies that writing tasks were more effective in the initial time points, and effects decreased over longer time spans.

The authors note several limitations. Generally, ineffective emotional expression may be a contributing factor. If participants/clients are not deeply engaged with the material, an expressive writing task may not be effective, as Pennebaker and Graybeal (2001) imply that connectedness is an important factor for the task. However, it may be difficult to implement a check for engagement in these types of research designs. Doing so may also set a context that will inhibit emotional processing and general responses. Research on expressive writing has found a wide range of outcomes for different variables (Fratraroli, 2006), and these various results may explain the large heterogeneity found in this study. Encouragingly, we did not find much evidence of publication bias, and therefore, these estimates may represent a true population effect size. Regardless, methodology of expressive writing studies is variable, as it is applied in different forms across different contexts. Ideally, it would be possible to control for these varied instructions and protocols. However, this is simply not feasible, as most studies do not use measures that examine how engaged an individual is with the material. As such, this current meta-analysis sought to provide readers with a global effect of expressive writing on the aforementioned outcome variables. More studies are needed to examine potential moderating effects of participant engagement. The authors also note limitations in regards to the specific outcome variables. The nature of the construct of PTG makes it difficult to analyze rigorously. For example, on the Posttraumatic Growth Inventory (commonly used to study PTG), one could respond 0 to the item “I have a greater appreciation for the value in my own life” because they already had a high level of appreciation in their life (i.e. ceiling effect). This conceptual issue may account for the non-effect of expressive writing on PTG. Logically, it would be difficult to determine whether or not an individual experiences growth from trauma without having experienced trauma. In conducting the literature search for the present meta-analysis, an insufficient number of

studies requiring a diagnosis of PTSD employed PTG as an outcome variable. Thus, it was difficult to determine whether participants in the studies employed had experienced trauma in line with DSM criteria. For PTS, studies not specifying whether or not participants had a diagnosis of PTSD were included. It is possible that studies included in the subclinical symptom category did in fact include participants without PTSD diagnosis (perhaps it was simply not assessed by means of a structured clinical interview). It is also crucial to consider mainstream issues not specific to expressive writing and the outcome variables utilized in the present study.

The psychological scientific community has shifted focus to reproducibility and research design in the last several years (Nelson, Simmons, & Simonsohn, 2018), and much of this discussion has focused on adequately powering studies for publication (Bakker et al., 2016; S. E. Maxwell, Lau, & Howard, 2015). S. E. Maxwell et al. (2015) and Open Science Collaboration (2015) have shown that the “replication crisis” may be attributed to low power in published studies. The power found in the current meta-analysis was very poor, with very few studies reaching the suggested 80% criterion to adequately power their study. This result was the same when considering individual study characteristics or the estimate true population effect size. Research by Bakker et al. (2016) indicates that researchers’ intuitions about power are particularly poor, and many studies could benefit from more informed power analyses. Anderson, Kelley, and Maxwell (2017) recently published a primer on power, with an online application to help with sample size planning for many types of research designs. Additionally, we encourage researchers to report power analyses of studies in order to better understand methodology for replication and reproducibility.

Meta-analyses, while useful tools to pool for population effect sizes, contain various limitations to their usefulness (Van Elk et al., 2015). As mentioned previously, these analyses can be affected by high heterogeneity, which was found in this study (Aert et al., 2016). Selection models have been criticized when using a smaller number of studies (Van Assen et al., 2015), and trim and fill analyses may not always estimate accurate confidence

intervals and funnel plots may be biased with heterogeneity (Terrin, Schmid, Lau, & Olkin, 2003). When focusing on improving the psychological sciences, Van Elk et al. (2015) suggest that the reliability and size of effects may be best elucidated by conducting large preregistered studies. This suggestion will also improve the outlook for power in published studies, and projects such as Many Labs can aide in subsidizing large samples (R. A. Klein et al., 2014). Even with limitations, meta-analyses allow researchers to examine the state of a research area, and we find potential with expressive writing on reducing PTS symptoms, and an overall need for better sample size and power planning for studies.

## References

- Aert, R. C. M. van, Wicherts, J. M., & Van Assen, M. A. L. M. (2016). Conducting meta-analyses based on p-values: Reservations and recommendations for applying p-uniform and p-curve. *Perspectives on Psychological Science*, 11(5), 713–729. doi:10.1017/CBO9781107415324.004
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*. doi:10.1176/appi.books.9780890425596.744053
- Anderson, S. F., Kelley, K., & Maxwell, S. E. (2017). Sample-size planning for more accurate statistical power: A method adjusting sample effect sizes for publication bias and uncertainty. *Psychological Science*, 28(11), 1547–1562. doi:10.1177/0956797617723724
- Aslam, N., & Kamal, A. (2013). Gender differences in distress responses, rumination patterns, perceived social support and posttraumatic growth among flood affected individuals. *Journal of Pakistan Psychiatric Society*, 10, 86–90.
- Bakker, M., Hartgerink, C. H. J., Wicherts, J. M., & Maas, H. L. J. van der. (2016). Researchers' intuitions about power in psychological research. *Psychological Science*, 27(8), 1069–1077. doi:10.1177/0956797616647519
- Bodor, N. Z. (2002). *The health effects of emotional disclosure for individuals with Type 1 diabetes* (PhD thesis No. 10-B). Retrieved from <http://ezproxy.lib.utexas.edu/login?url=http://search.ebscohost.com/login.aspx?direct=true{\&}db>
- Borenstein, M., Hedges, L. V., & Rothstein, H. (2007). Meta-analysis fixed effect vs. random effects. Retrieved from <https://www.meta-analysis.com/downloads/Meta-analysis-fixed-effect-vs-random-effects-072607.pdf>
- Brounéus, K. (2010). The trauma of truth telling: Effects of witnessing in the Rwandan Gacaca Courts on psychological health. *Journal of Conflict Resolution*, 54(3), 408–437. doi:10.1177/0022002709360322
- Bruns, S. B., & Ioannidis, J. P. A. (2016). P-curve and p-hacking in observational research.

*PLoS ONE*, 11(2). doi:[10.1371/journal.pone.0149144](https://doi.org/10.1371/journal.pone.0149144)

Buchanan, E. M., Valentine, K. D., & Scofield, J. E. (2017). MOTE. Retrieved from

<https://github.com/doomlab/MOTE>

Carter, E. C., & McCullough, M. E. (2014). Publication bias and the limited strength model

of self-control: Has the evidence for ego depletion been overestimated? *Frontiers in*

*Psychology*, 5(July), 1–11. doi:[10.3389/fpsyg.2014.00823](https://doi.org/10.3389/fpsyg.2014.00823)

Champely, S. (2016). pwr: Basic functions for power analysis. R package version 1.2-0.

Retrieved from <https://cran.r-project.org/package=pwr>

Cobb, A. R., Tedeschi, R. G., Calhoun, L. G., & Cann, A. (2006). Correlates of

posttraumatic growth in survivors of intimate partner violence. *Journal of Traumatic*

*Stress*, 19(6), 895–903. doi:[10.1002/jts.20171](https://doi.org/10.1002/jts.20171)

Coburn, K. M., & Vevea, J. L. (2017). Weightr. Retrieved from

<https://cran.r-project.org/web/packages/weightr/index.html>

Cochran, W. G. (1954). Some methods for strengthening the common  $\chi^2$  tests. *Biometrics*,

10(4), 417–451. doi:[10.2307/3001616](https://doi.org/10.2307/3001616)

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale,

NJ: Earlbaum.

Cooper, H., Hedges, L. V., & Valentine, J. (2009). *The handbook of research synthesis and*

*meta-analysis* (2nd ed.). New York, NY: Russell Sage Foundation.

Costanza, R., Fisher, B., Ali, S., Beer, C., Bond, L., Boumans, R., . . . Snapp, R. (2007).

Quality of life: An approach integrating opportunities, human needs, and subjective

well-being. *Ecological Economics*, 61(2-3), 267–276.

doi:[10.1016/j.ecolecon.2006.02.023](https://doi.org/10.1016/j.ecolecon.2006.02.023)

Crespo, M., & Gomez, M. M. (2016). Diagnostic concordance of DSM-IV and DSM-5

posttraumatic stress disorder (PTSD) in a clinical sample. *Psicothema*, 28(2),

161–166. doi:[10.7334/psicothema2015.213](https://doi.org/10.7334/psicothema2015.213)

Cumming, G. (2012). *Understanding the new statistics: Effect sizes, confidence intervals,*

- and meta-analysis. New York, NY: Routledge.
- DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3), 177–188. doi:[10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)
- Di Blasio, P., Camisasca, E., Caravita, S. C. S., Ionio, C., Milani, L., Valtolina, G. G., . . . Valtolina, G. G. (2015). The effects of expressive writing on postpartum depression and posttraumatic stress symptoms. *Psychological Reports*, 117(3), 856–882. doi:[10.2466/02.13.PR0.117c29z3](https://doi.org/10.2466/02.13.PR0.117c29z3)
- Dursun, P., Steger, M. F., Bentele, C., & Schulenberg, S. E. (2016). Meaning and posttraumatic growth among survivors of the September 2013 Colorado floods. *Journal of Clinical Psychology*, 72(12), 1247–1263. doi:[10.1002/jclp.22344](https://doi.org/10.1002/jclp.22344)
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56(2), 455–463. doi:[10.1111/j.0006-341X.2000.00455.x](https://doi.org/10.1111/j.0006-341X.2000.00455.x)
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315(7109), 629–634. doi:[10.1136/bmj.316.7129.469](https://doi.org/10.1136/bmj.316.7129.469)
- Esterling, B. A., Antoni, M. H., Kumar, M., & Schneiderman, N. (1990). Emotional repression, stress disclosure responses, and Epstein-Barr viral capsid antigen titers. *Psychosomatic Medicine*, 52, 397–410. doi:[10.1097/00006842-199007000-00002](https://doi.org/10.1097/00006842-199007000-00002)
- Fawzy, N. W., Fawzy, N. W., Hyun, C. S., Elashoff, R., Guthrie, D., Fahey, J. L., & Morton, D. L. (1993). Malignant melanoma. Effects of an early structured psychiatric intervention, coping, and affective state on recurrence and survival 6 years later. *Archives of General Psychiatry*, 50(9), 681–689. doi:[10.1001/archpsyc.1993.01820210015002](https://doi.org/10.1001/archpsyc.1993.01820210015002)
- Francis, G. (2012). Publication bias and the failure of replication in experimental psychology. *Psychonomic Bulletin & Review*, 19(6), 975–991. doi:[10.3758/s13423-012-0322-y](https://doi.org/10.3758/s13423-012-0322-y)
- Francis, G. (2014). The frequency of excess success for articles in Psychological Science.

- Psychonomic Bulletin & Review*, 21(5), 1180–1187. doi:[10.3758/s13423-014-0601-x](https://doi.org/10.3758/s13423-014-0601-x)
- Francis, M. E., & Pennebaker, J. W. (1992). Putting stress into words: The impact of writing on physiological, absentee, and self-reported emotional well-being measures. *American Journal of Health Promotion*, 6(4), 280–287. doi:[10.4278/0890-1171-6.4.280](https://doi.org/10.4278/0890-1171-6.4.280)
- Frankl, V. (1959). *Man's search for meaning* (3rd ed.). Boston, MA: Beacon Press.
- Frattaroli, J. (2006). Experimental disclosure and its moderators: A meta-analysis. *Psychological Bulletin*, 132(6), 823–865. doi:[10.1037/0033-2909.132.6.823](https://doi.org/10.1037/0033-2909.132.6.823)
- Frisina, P. G., Borod, J. C., & Lepore, S. J. (2004). A meta-analysis of the effects of written emotional disclosure on the health outcomes of clinical populations. *The Journal of Nervous and Mental Disease*, 192(9), 629–634. doi:[10.1097/01.nmd.0000138317.30764.63](https://doi.org/10.1097/01.nmd.0000138317.30764.63)
- Gentes, E. L., Dennis, P. A., Kimbrel, N. A., Rissling, M. B., Beckham, J. C., & Calhoun, P. S. (2014). DSM-5 posttraumatic stress disorder: Factor structure and rates of diagnosis. *Journal of Psychiatric Research*, 59(1), 60–67. doi:[10.1016/j.jpsychires.2014.08.014](https://doi.org/10.1016/j.jpsychires.2014.08.014)
- Gidron, Y., Peri, T., Connolly, J. F., & Shalev, A. Y. (1996). Written disclosure in posttraumatic stress disorder: Is it beneficial for the patient? *The Journal of Nervous and Mental Disease*, 184(8), 505–506. doi:[10.1097/00005053-199608000-00009](https://doi.org/10.1097/00005053-199608000-00009)
- Glass, G. V. (1976). Primary, secondary, and meta-analysis of research. *Educational Researcher*, 5(10), 3–8. doi:[10.3102/0013189X005010003](https://doi.org/10.3102/0013189X005010003)
- Goldstein, H. S., Edelberg, R., Meier, C. F., & Davis, L. (1988). Relationship of resting blood pressure and heart rate to experienced anger and expressed anger. *Psychosomatic Medicine*, 50(4), 321–329. doi:[10.1097/00006842-198807000-00001](https://doi.org/10.1097/00006842-198807000-00001)
- Greenberg, M. A., & Stone, A. A. (1992). Emotional disclosure about traumas and its relation to health: Effects of previous disclosure and trauma severity. *Journal of Personality and Social Psychology*, 63, 75–84. doi:[10.1037/h0090372](https://doi.org/10.1037/h0090372)
- Gross, J. J., & Levenson, R. W. (1997). Hiding feelings: The acute effects of inhibiting



negative and positive emotion. *Journal of Abnormal Psychology*, 106(1), 95–103.

doi:[10.1037/0021-843X.106.1.95](https://doi.org/10.1037/0021-843X.106.1.95)

Harris, A. H. S., Thoresen, C. E., Humphreys, K., & Faul, J. (2005). Does writing affect asthma? A randomized trial. *Psychosomatic Medicine*, 67(1), 130–136.

doi:[10.1097/01.psy.0000146345.73510.d5](https://doi.org/10.1097/01.psy.0000146345.73510.d5)

Hedges, L. V. (1982). Estimation of effect size from a series of independent experiments.

*Psychological Bulletin*, 92(2), 490–499. doi:[10.1037/0033-2909.92.2.490](https://doi.org/10.1037/0033-2909.92.2.490)

Henry, E. A., Schlegel, R. J., Talley, A. E., Molix, L. A., & Bettencourt, B. A. (2010). The feasibility and effectiveness of expressive writing for rural and urban breast cancer

survivors. *Oncology Nursing Forum*, 37(6), 749–757. doi:[10.1188/10.ONF.749-757](https://doi.org/10.1188/10.ONF.749-757)

Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *British Medical Journal*, 327(7414), 557–560.

doi:[10.1136/bmj.327.7414.557](https://doi.org/10.1136/bmj.327.7414.557)

Hilgard, J. (2016). PETPEESE. GitHub. Retrieved from

<https://github.com/Joe-Hilgard/PETPEESE>

Huedo-Medina, T. B., Sánchez-Meca, J., Marín-Martínez, F., & Botella, J. (2006). Assessing heterogeneity in meta-analysis: Q statistic or I<sup>2</sup> index? *Psychological Methods*, 11(2),

193–206. doi:[10.1037/1082-989X.11.2.193](https://doi.org/10.1037/1082-989X.11.2.193)

John, L. K., Loewenstein, G., & Prelec, D. (2012). Measuring the prevalence of questionable research practices with incentives for truth telling. *Psychological Science*, 23(5),

524–532. doi:[10.1177/0956797611430953](https://doi.org/10.1177/0956797611430953)

Kelley, K. (2007). Confidence intervals for standardized effect sizes. *Journal of Statistical Software*, 20(8), 1–24. doi:[10.18637/jss.v020.i08](https://doi.org/10.18637/jss.v020.i08)

Klein, R. A., Ratliff, K. A., Vianello, M., Adams, R. B., Bahník, Š., Bernstein, M. J., . . .

Nosek, B. A. (2014). Investigating variation in replicability. *Social Psychology*, 45(3), 142–152. doi:[10.1027/1864-9335/a000178](https://doi.org/10.1027/1864-9335/a000178)

Klump, M. C. (2008). Posttraumatic stress disorder and sexual assault in women. *Journal of*

*College Student Development*, 8225 (May 2014), 37–41. doi:[10.1300/J035v21n02](https://doi.org/10.1300/J035v21n02)

Kross, E., & Ayduk, O. (2011). Making meaning out of negative experiences by self-distancing. *Current Directions in Psychological Science*, 20(3), 187–191. doi:[10.1177/0963721411408883](https://doi.org/10.1177/0963721411408883)

Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: A practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, 4. doi:[10.3389/fpsyg.2013.00863](https://doi.org/10.3389/fpsyg.2013.00863)

Lancaster, S. L., Klein, K. P., & Heifner, A. (2015). The validity of self-reported growth after expressive writing. *Traumatology*, 21(4), 293–298. doi:[10.1037/trm0000052](https://doi.org/10.1037/trm0000052)

Larson, D. G., & Chastain, R. L. (1990). Self-concealment: Conceptualization, measurement, and health implications. *Journal of Social and Clinical Psychology*, 9(4), 439–455. doi:[10.1521/jscp.1990.9.4.439](https://doi.org/10.1521/jscp.1990.9.4.439)

Lieberman, M. A., & Goldstein, B. A. (2006). Not all negative emotions are equal: The role of emotional expression in online support groups for women with breast cancer. *Psycho-Oncology*, 15(2), 160–168. doi:[10.1002/pon.932](https://doi.org/10.1002/pon.932)

Maxwell, S. E., Lau, M. Y., & Howard, G. S. (2015). Is psychology suffering from a replication crisis? What does “failure to replicate” really mean? *American Psychologist*, 70(6), 487–498. doi:[10.1037/a0039400](https://doi.org/10.1037/a0039400)

McShane, B. B., Böckenholt, U., & Hansen, K. T. (2016). Adjusting for publication bias in meta-analysis. *Perspectives on Psychological Science*, 11(5), 730–749. doi:[10.1177/1745691616662243](https://doi.org/10.1177/1745691616662243)

Meshberg-Cohen, S., Svikis, D., & McMahon, T. J. (2014). Expressive writing as a therapeutic process for drug-dependent women. *Substance Abuse*, 35(1), 80–88. doi:[10.1080/08897077.2013.805181](https://doi.org/10.1080/08897077.2013.805181)

Mogk, C., Otte, S., Reinhold-Hurley, B., & Kröner-Herwig, B. (2006). Health effects of expressive writing on stressful or traumatic experiences - a meta-analysis.

838 *Psychosocial Medicine*, 3, Doc06.

839 Morris, S. B., & DeShon, R. P. (2002). Combining effect size estimates in meta-analysis with  
840 repeated measures and independent-groups designs. *Psychological Methods*, 7(1),  
841 105–125. doi:[10.1037/1082-989X.7.1.105](https://doi.org/10.1037/1082-989X.7.1.105)

842 Nelson, L. D., Simmons, J., & Simonsohn, U. (2018). Psychology's renaissance. *Annual*  
843 *Review of Psychology*, 69(1), 511–534. doi:[10.1146/annurev-psych-122216-011836](https://doi.org/10.1146/annurev-psych-122216-011836)

844 Open Science Collaboration. (2015). Estimating the reproducibility of psychological science.  
845 *Science*, 349(6251), aac4716–aac4716. doi:[10.1126/science.aac4716](https://doi.org/10.1126/science.aac4716)

846 Pennebaker, J. W. (1989). Confession, inhibition, and disease. In L. Berkowitz (Ed.),  
847 *Advances in experimental social psychology* (Vol. 22, pp. 211–244). Academic Press.  
848 doi:[10.1016/S0065-2601\(08\)60309-3](https://doi.org/10.1016/S0065-2601(08)60309-3)

849 Pennebaker, J. W. (1993). Putting stress into words: Health, linguistic, and therapeutic  
850 implications. *Behaviour Research and Therapy*, 31(6), 539–548.  
851 doi:[10.1016/0005-7967\(93\)90105-4](https://doi.org/10.1016/0005-7967(93)90105-4)

852 Pennebaker, J. W., & Beall, S. K. (1986). Confronting a traumatic event: Toward an  
853 understanding of inhibition and disease. *Journal of Abnormal Psychology*, 95(3),  
854 274–281. doi:[10.1037//0021-843X.95.3.274](https://doi.org/10.1037//0021-843X.95.3.274)

855 Pennebaker, J. W., & Francis, M. E. (1996). Cognitive, emotional, and language processes in  
856 disclosure. *Cognition & Emotion*, 10(6), 601–626. doi:[10.1080/026999396380079](https://doi.org/10.1080/026999396380079)

857 Pennebaker, J. W., & Graybeal, A. (2001). Patterns of natural language use: Disclosure,  
858 personality, and social integration. *Current Directions in Psychological Science*, 10(3),  
859 90–93. doi:[10.1111/1467-8721.00123](https://doi.org/10.1111/1467-8721.00123)

860 Pennebaker, J. W., Colder, M., & Sharp, L. K. (1990). Accelerating the coping process.  
861 *Journal of Personality and Social Psychology*, 58(3), 528–537.  
862 doi:[10.1037//0022-3514.58.3.528](https://doi.org/10.1037//0022-3514.58.3.528)

863 Pennebaker, J. W., Kiecolt-Glaser, J. K., & Glaser, R. (1988). Disclosure of traumas and  
864 immune function: Health implications for psychotherapy. *Journal of Consulting and*

- Clinical Psychology*, 56(2), 239–245. doi:[10.1037/0022-006X.56.2.239](https://doi.org/10.1037/0022-006X.56.2.239)
- Possemato, K., Ouimette, P., & Geller, P. (2010). Internet-based expressive writing for kidney transplant recipients: Effects on posttraumatic stress and quality of life. *Traumatology*, 16(1), 49–54. doi:[10.1177/1534765609347545](https://doi.org/10.1177/1534765609347545)
- Rachman, S. (1980). Emotional processing. *Behaviour Research and Therapy*, 18(1), 51–60. doi:[10.1016/0005-7967\(80\)90069-8](https://doi.org/10.1016/0005-7967(80)90069-8)
- Reinhold, M., Bürkner, P. C., & Holling, H. (2018). Effects of expressive writing on depressive symptoms—A meta-analysis. *Clinical Psychology: Science and Practice*, 25(1). doi:[10.1111/cpsp.12224](https://doi.org/10.1111/cpsp.12224)
- Sánchez-Meca, J., & Marín-Martínez, F. (2008). Confidence intervals for the overall effect size in random-effects meta-analysis. *Psychological Methods*, 13(1), 31–48. doi:[10.1037/1082-989X.13.1.31](https://doi.org/10.1037/1082-989X.13.1.31)
- Scheff, T. J. (1979). *Catharsis in healing, ritual, and drama*. Los Angeles: University of California Press.
- Schoutrop, M. J. A., Lange, A., Hanewald, G., Davidovich, U., & Salomon, H. H. (2002). Structured writing and processing major stressful events: A controlled trial. *Psychotherapy and Psychosomatics*, 71(3), 151–157. doi:[10.1159/000056282](https://doi.org/10.1159/000056282)
- Schulenberg, S. E., Hutzell, R. R., Nassif, C., & Rogina, J. M. (2008). Logotherapy for clinical practice. *Psychotherapy*, 45(4), 447–463. doi:[10.1037/a0014331](https://doi.org/10.1037/a0014331)
- Simonsohn, U., Nelson, L. D., & Simmons, J. P. (2014). p-curve: A key to the file-drawer. *Journal of Experimental Psychology: General*, 143(2), 534–547. doi:[10.1037/a0033242](https://doi.org/10.1037/a0033242)
- Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2015). Better p-curves: Making p-curve analysis more robust to errors, fraud, and ambitious p-hacking, a reply to Ulrich and Miller (2015). *Journal of Experimental Psychology: General*, 144(6), 1146–1152. doi:[10.1037/xge0000104](https://doi.org/10.1037/xge0000104)
- Slavin-Spenny, O. M., Cohen, J. L., Oberleitner, L. M., & Lumley, M. A. (2011). The effects of different methods of emotional disclosure: Differentiating posttraumatic growth

892 from stress symptoms. *Journal of Clinical Psychology*, 67(10), 993–1007.

893 doi:[10.1002/jclp.20750](https://doi.org/10.1002/jclp.20750)

894 Sloan, D. M., Marx, B. P., & Epstein, E. M. (2005). Further examination of the exposure  
895 model underlying the efficacy of written emotional disclosure. *Journal of Consulting  
896 and Clinical Psychology*, 73(3), 549–554. doi:[10.1037/0022-006X.73.3.549](https://doi.org/10.1037/0022-006X.73.3.549)

897 Sloan, D. M., Marx, B. P., & Greenberg, E. M. (2011). A test of written emotional  
898 disclosure as an intervention for posttraumatic stress disorder. *Behaviour Research  
899 and Therapy*, 49(4), 299–304. doi:[10.1016/j.brat.2011.02.001](https://doi.org/10.1016/j.brat.2011.02.001)

900 Sloan, D. M., Marx, B. P., Bovin, M. J., Feinstein, B. A., & Gallagher, M. W. (2012).  
901 Written exposure as an intervention for PTSD: A randomized clinical trial with motor  
902 vehicle accident survivors. *Behaviour Research and Therapy*, 50(10), 627–635.  
903 doi:[10.1016/j.brat.2012.07.001](https://doi.org/10.1016/j.brat.2012.07.001)

904 Sloan, D. M., Marx, B. P., Epstein, E. M., & Lexington, J. M. (2007). Does altering the  
905 writing instructions influence outcome associated with written disclosure? *Behavior  
906 Therapy*, 38(2), 155–168. doi:[10.1016/j.beth.2006.06.005](https://doi.org/10.1016/j.beth.2006.06.005)

907 Smithson, M. (2001). Correct confidence intervals for various regression effect sizes and  
908 parameters: The importance of noncentral distributions in computing intervals.  
909 *Educational and Psychological Measurement*, 61(4), 605–632.  
910 doi:[10.1177/00131640121971392](https://doi.org/10.1177/00131640121971392)

911 Smithson, M. (2003). *Confidence intervals*. Thousand Oaks, CA: Sage.

912 Smyth, J. M. (1998). Written emotional expression: Effect sizes , outcome types, and  
913 moderating variables. *Journal of Consulting and Clinical Psychology*, 66(1), 174–184.  
914 doi:[10.1037/0022-006X.66.1.174](https://doi.org/10.1037/0022-006X.66.1.174)

915 Smyth, J. M., Stone, A. A., Hurewitz, A., & Kaell, A. (1999). Effects of writing about  
916 stressful experiences on symptom reduction in patients with asthma or rheumatoid  
917 arthritis: A randomized trial. *JAMA: The Journal of the American Medical*

- Association, 281(14), 1304–1309. doi:[10.1001/jama.281.14.1304](https://doi.org/10.1001/jama.281.14.1304)
- Stanley, T. D. (2005). Beyond publication bias. *Journal of Economic Surveys*, 19(3), 309–345. doi:[10.1111/j.0950-0804.2005.00250.x](https://doi.org/10.1111/j.0950-0804.2005.00250.x)
- Stanley, T. D., & Doucouliagos, H. (2014). Meta-regression approximations to reduce publication selection bias. *Research Synthesis Methods*, 5(1), 60–78. doi:[10.1002/jrsm.1095](https://doi.org/10.1002/jrsm.1095)
- Stanton, A. L., Danoff-Burg, S., Sworowski, L. A., Collins, C. A., Branstetter, A. D., Rodriguez-Hanley, A., . . . Austenfeld, J. L. (2002). Randomized, controlled trial of written emotional expression and benefit finding in breast cancer patients. *Journal of Clinical Oncology*, 20(20), 4160–4168. doi:[10.1200/JCO.2002.08.521](https://doi.org/10.1200/JCO.2002.08.521)
- Taku, K., Calhoun, L. G., Cann, A., & Tedeschi, R. G. (2008). The role of rumination in the coexistence of distress and posttraumatic growth among bereaved Japanese University students. *Death Studies*, 32(5), 428–444. doi:[10.1080/07481180801974745](https://doi.org/10.1080/07481180801974745)
- Tedeschi, R. G., & Calhoun, L. G. (1995). *Trauma & transformation: Growing in the aftermath of suffering*. Thousand Oaks, CA: Sage Publications.
- Tedeschi, R. G., & Calhoun, L. G. (2004). Posttraumatic growth: Conceptual foundations and empirical evidence. *Psychological Inquiry*, 15(1), 1–18. doi:[10.1207/s15327965pli1501](https://doi.org/10.1207/s15327965pli1501)
- Terrin, N., Schmid, C. H., Lau, J., & Olkin, I. (2003). Adjusting for publication bias in the presence of heterogeneity. *Statistics in Medicine*, 22(13), 2113–2126. doi:[10.1002/sim.1461](https://doi.org/10.1002/sim.1461)
- Theofilou, P. (2013). Quality of life: Definition and measurement. *Europe's Journal of Psychology*, 9(1), 150–162. doi:[10.5964/ejop.v9i1.337](https://doi.org/10.5964/ejop.v9i1.337)
- Van Aert, R. C. M. (2017). P-uniform. GitHub. Retrieved from <https://github.com/RobbievanAert/puniform>
- Van Assen, M. A. L. M., Van Aert, R. C. M., & Wicherts, J. M. (2015). Meta-analysis using effect size distributions of only statistically significant studies. *Psychological Methods*,

20(3), 293–309. doi:<http://dx.doi.org/10.1037/met0000025>

Van Elk, M., Matzke, D., Gronau, Q. F., Guan, M., Vandekerckhove, J., & Wagenmakers, E.-J. (2015). Meta-analyses are no substitute for registered replications: A skeptical perspective on religious priming. *Frontiers in Psychology*, 6, 1365. doi:[10.3389/fpsyg.2015.01365](https://doi.org/10.3389/fpsyg.2015.01365)

Van Emmerik, A. A. P., Reijntjes, A., & Kamphuis, J. H. (2013). Writing therapy for posttraumatic stress: A meta-analysis. *Psychotherapy and Psychosomatics*, 82(2), 82–88. doi:[10.1159/000343131](https://doi.org/10.1159/000343131)

Vevea, J. L., & Hedges, L. V. (1995). A general linear model for estimating effect size in the presence of publication bias. *Psychometrika*, 60(3), 419–435. doi:[10.1007/BF02294384](https://doi.org/10.1007/BF02294384)

Vevea, J. L., & Woods, C. M. (2005). Publication bias in research synthesis: Sensitivity analysis using a priori weight functions. *Psychological Methods*, 10(4), 428–443. doi:[10.1037/1082-989X.10.4.428](https://doi.org/10.1037/1082-989X.10.4.428)

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36(3), 1–48. doi:[10.18637/jss.v036.i03](https://doi.org/10.18637/jss.v036.i03)

Walker, B. L., Nail, L. M., & Croyle, R. T. (1999). Does emotional expression make a difference in reactions to breast cancer? *Oncology Nursing Forum*, 26(6), 1025–1032.

Wang, X., Gao, L., Shinfuku, N., Zhang, H., Zhao, C., & Shen, Y. (2000). Longitudinal study of earthquake-related PTSD in a randomly selected community sample in North China. *American Journal of Psychiatry*, 157(8), 1260–1266. doi:[10.1176/appi.ajp.157.8.1260](https://doi.org/10.1176/appi.ajp.157.8.1260)

Weiss, T. (2002). Posttraumatic growth in women with breast cancer and their husbands – An intersubjective validation study. *Journal of Psychosocial Oncology*, 20(2), 65–80. doi:[10.1300/J077v20n02\\_04](https://doi.org/10.1300/J077v20n02_04)

Wilson, K. G., & DuFrene, T. (2009). *Mindfulness for two: An acceptance and commitment Therapy approach to mindfulness in psychotherapy*. Oakland, CA: New Harbinger

971 Publications.

972 Yilmaz, M., & Zara, A. (2016). Traumatic loss and posttraumatic growth: The effect of  
973 traumatic loss related factors on posttraumatic growth. *Anatolian Journal of*  
974 *Psychiatry*, 17(1), 5–11. doi:[10.5455/apd.188311](https://doi.org/10.5455/apd.188311)



Table 1

*Effect Size Estimates for PTS Results*

Model	Fixed Effects	Random Effects	Fixed No Outliers	Random No Outliers
Overall Effects	0.36 [0.33, 0.39]	0.42 [0.35, 0.48]	0.34 [0.32, 0.37]	0.39 [0.32, 0.45]
Z Values	24.61, $p < .001$	12.38, $p < .001$	23.33, $p < .001$	12.42, $p < .001$
$p$ -Uniform	0.63 [0.54, 0.72]	-	0.60 [0.51, 0.69]	-
PET	0.09 [0.01, 0.17]	-	0.13 [0.04, 0.21]	-
PEESE	0.24 [0.19, 0.29]	-	0.25 [0.20, 0.30]	-
Selection Models	0.33 [0.28, 0.37]	0.45 [0.33, 0.57]	0.29 [0.24, 0.33]	0.39 [0.27, 0.50]
Trim and Fill	0.28 [0.25, 0.31]	0.28 [0.20, 0.35]	0.28 [0.25, 0.31]	0.28 [0.21, 0.35]

*Note.* [] indicates the 95 percent confidence interval for each effect size estimate.

Table 2

*Effect Size Estimates for PTG Results*

Model	Fixed Effects	Random Effects
Overall Effects	0.10 [0.02, 0.17]	0.10 [0.02, 0.17]
Z Values	2.45, $p = .014$	2.45, $p = .014$
$p$ -Uniform	-0.11 [-1.43, 0.42]	-
PET	0.06 [-0.20, 0.32]	-
PEESE	0.08 [-0.04, 0.20]	-
Selection Models	0.09 [-0.01, 0.18]	0.09 [-0.03, 0.20]
Trim and Fill	0.10 [0.02, 0.17]	0.10 [0.02, 0.17]

*Note.* [] indicates the 95 percent confidence interval for each effect size estimate.

Table 3

*Effect Size Estimates for QOL Results*

Model	Fixed Effects	Random Effects	Fixed No Outliers	Random No Outliers
Overall Effects	-0.01 [-0.07, 0.05]	-0.01 [-0.16, 0.13]	-0.01 [-0.07, 0.05]	-0.01 [-0.11, 0.09]
Z Values	-0.33, $p = .745$	-0.18, $p = .860$	-0.25, $p = .805$	-0.20, $p = .838$
$p$ -Uniform	0.79 [0.33, 1.61]	-	0.62 [0.10, 0.96]	-
PET	0.05 [-0.26, 0.36]	-	0.05 [-0.29, 0.38]	-
PEESE	0.00 [-0.17, 0.17]	-	0.00 [-0.19, 0.19]	-
Selection Models	-0.06 [-0.12, 0.01]	0.51 [-0.09, 1.12]	-0.04 [-0.11, 0.03]	0.05 [-0.15, 0.24]
Trim and Fill	-0.01 [-0.07, 0.05]	-0.01 [-0.16, 0.13]	-0.01 [-0.07, 0.05]	-0.01 [-0.11, 0.09]

*Note.* [] indicates the 95 percent confidence interval for each effect size estimate.





Figure 1. Effect sizes and their non-centralized confidence interval for PTS total scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.



*Figure 2.* Effect sizes and their non-centralized confidence interval for PTS Hyperarousal. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 3. Effect sizes and their non-centralized confidence interval for PTS Intrusion scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 4. Effect sizes and their non-centralized confidence interval for PTS Avoidance Scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 5. Effect sizes and their non-centralized confidence interval for PTG outcome variables. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.






Figure 6. Effect sizes and their non-centralized confidence interval for QOL outcome variables. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.