

1 A Meta-Analysis of Expressive Writing on Positive Psychology Variables and Traumatic
2 Stress

Abstract

Emotional expression has been shown to be beneficial for promoting both positive psychological and physical health outcomes. Unfortunately, inhibiting emotions can lead to impairments in physical and psychological health. James Pennebaker showed that expressive writing is an effective form of emotional expression, and he and others have used expressive writing as an experimental manipulation to gauge its effectiveness in treating a wide variety of health-related and psychological outcomes. While many studies have been conducted that examine the effectiveness of expressive writing across such outcomes, a considerable amount of these studies tend to neglect necessary considerations such as power and meaningfulness of respective effect sizes. Four previous meta-analyses have been conducted that examine expressive writing's effect on psychological outcomes. However, these studies focus on the experimental versus control group effect size. Thus, our meta-analysis sought to examine the effectiveness of an expressive writing intervention on only the experimental conditions in studies measuring posttraumatic growth, posttraumatic stress, and quality of life using random effects models. Results indicated a small overall effect size for posttraumatic stress and negligible to small effect sizes for posttraumatic growth and quality of life. Implications for future research design and interpretation of published research are discussed.

Keywords: meta-analysis, positive psychology, expressive writing

A Meta-Analysis of Expressive Writing on Positive Psychology Variables and Traumatic Stress

Emotional expression, especially focusing on negative emotions or trauma, has been shown to increase both mental and physical health (Esterling, Antoni, Kumar, & Schneiderman, 1990; Fawzy et al., 1993; Lieberman & Goldstein, 2006; Rachman, 1980; Scheff, 1979). In contrast, inhibiting repressive thoughts or emotions can be detrimental to both physical and psychological health (H. S. Goldstein, Edelberg, Meier, & Davis, 1988; Gross & Levenson, 1997; Larson & Chastain, 1990). Additionally, repressing or avoiding certain emotions may prevent an individual from living in accordance with his or her values (Wilson & DuFrene, 2009). Value-congruent behavior allows individuals to foster a sense of meaning (Frankl, 1959; Schulenberg, Hutzell, Nassif, & Rogina, 2008). Further, individuals experiencing traumatic events are more likely to repress thoughts and feelings about a given traumatic experience (Bodor, 2002). Generally, preventing the disclosure of harmful thoughts and feelings can be harmful to individuals, while disclosing these events can reduce stress and lead to various positive health outcomes, such as with diabetes (Bodor, 2002), breast cancer related illnesses (Stanton et al., 2002), and other physical health conditions.

As a caveat, however, the disclosure or acceptance of harmful thoughts or emotions can sometimes lead to negative outcomes. Thus, caution should be used when interpreting results from interventions based on emotional expression (Wilson & DuFrene, 2009). For example, Brounéus (2010) found that truth telling, a form of emotional expression, caused harm to individuals. However, in general, these repressive maneuvers have the capability to lead to social concerns, overall psychological dysfunction, and inhibit people from living in accordance with their values (Frankl, 1959; Pennebaker, 1989; Pennebaker & Beall, 1986; Schulenberg et al., 2008; Wilson & DuFrene, 2009). Psychological dysfunction can obviously lead to detrimental effects on health, including unhealthy everyday life habits, which could lead to biological problems, especially immune system and neurotransmitter deficiencies (Pennebaker & Beall, 1986). By preventing an individual from living in accordance with his

or her values, emotional inhibition may prohibit individuals from gaining access to values that are intrinsically reinforcing. Additionally, living in accordance with personal values allows the individual to foster a sense of meaning despite debilitating life circumstances (Frankl, 1959; Schulenberg et al., 2008). Therefore, it is important to identify and foster ways in which individuals can effectively express emotions, thereby improving both physical and psychological health.

Pennebaker and Beall (1986) first showed that emotional expression can be both experimentally manipulated and have positive benefits to participants. In their seminal study, they randomly assigned participants to several writing groups including writing about an experienced trauma or a neutral event. The group that disclosed both about their trauma and the emotions surrounding that trauma later showed a reduction in health visits. Pennebaker has replicated the use of expressive writing (i.e., a paradigm in which one writes about emotions) across a number of studies ranging from improving health (Pennebaker, Colder, & Sharp, 1990; Pennebaker, Kiecolt-Glaser, & Glaser, 1988) to improvements in school (Pennebaker & Francis, 1996) and work (Francis & Pennebaker, 1992). Others have expanded this work to show positive effects on mood (Schoutrop, Lange, Hanewald, Davidovich, & Salomon, 2002) and asthma (Smyth, Stone, Hurewitz, & Kaell, 1999); however, several controlled studies have shown to not replicate (Harris, Thoresen, Humphreys, & Faul, 2005) or null effects (Gidron, Peri, Connolly, & Shalev, 1996; Walker, Nail, & Croyle, 1999).

The idea that a brief, controlled writing intervention can have numerous positive health and psychological benefits can certainly be controversial, especially when recent studies show contradicting results. For example, Henry, Schlegel, Talley, Molix, and Bettencourt (2010) found that expressive writing only benefited a rural population for those individuals surviving breast cancer, while Lancaster, Klein, and Heifner (2015) found no significant evidence that expressive writing can be considered an effective approach. Additionally, Brounéus (2010) found that truth telling caused harm to individuals in a

forensic setting. Regardless, the concept remains interesting due to the nature and inexpensive implementation of expressive writing. Many individuals who have experienced traumatic events do not wish to disclose their feelings regarding the events with others. Additionally, those who do not meet diagnostic criteria are sometimes neglected despite probable suffering (Wilson & DuFrene, 2009). However, by utilizing expressive writing as a personal method of treatment, individuals are able to effectively express their emotions while avoiding talking to another individual or clinician about the traumatic event (Smyth, 1998). Pennebaker (1993) found that experimental conditions assigned to participate in an expressive writing task generally report more positive changes than those in control conditions. Some controversy has been observed over whether or not writing about a formerly disclosed event is more effective than writing about an undisclosed event. M. A. Greenberg and Stone (1992) conducted an experiment where they separated participants into three groups: writing about a formerly disclosed trauma, writing about an undisclosed trauma, and a control group. They found no difference between groups in effectiveness. However, they did find that those who disclosed more severe traumas reportedly experienced fewer physical symptoms at follow up, which suggests that the type of trauma revealed can play a significant impact on symptom reduction and physical health. From a meta-analytic perspective, Mogk, Otte, Reinhold-Hurley, and Kröner-Herwig (2006) found a non-meaningful effect size when examining the effect of expressive writing on somatic health symptoms, psychological health, and miscellaneous outcomes, such as grades and self-efficacy. These results were in contrast to the meta-analysis conducted by Smyth (1998), which found a medium overall effect size ($d = 0.47$).

In order to understand why expressive writing is considered to be an effective, intervention-based approach, one must examine the cognitive and social processes by which it allows an individual to process information. Pennebaker et al. (1990) discovered that individuals who had benefited from expressive writing attributed their success to ways in which the intervention allowed them to understand what had happened to them.

Furthermore, in an additional study, Pennebaker (1993) conducted a textual analysis on expressive writing content and found that those who were more successful during the intervention used words that can be categorized as causation words. Pennebaker attributed these results as individuals effectively processing the event in their own minds. Aside from cognitive-processing and inhibition theories, there are a number of other theories that researchers have used to explain emotional disclosure. The first theory that warrants explanation is the social integration model (Pennebaker & Graybeal, 2001). This model discusses how emotional disclosure can have a positive impact on how people interact in their environment. This increased environmental interaction has been shown to have positive benefits on health (Frattaroli, 2006). Finally, expressive writing parallels exposure therapy for a variety of phobias and post-traumatic stress disorder, which suggests that repeatedly exposing oneself to the seemingly anxious thought or trauma can reduce the anxiety, fear, or stress associated with that event (Meshberg-Cohen, Svikis, & McMahon, 2014). Given that exposure therapy has been shown to be effective for reducing symptoms of post-traumatic stress (PTS; Sloan, Marx, & Epstein, 2005), one would expect individuals in these studies to experience a reduction in PTS symptoms after taking part in an expressive writing intervention. Additionally, Wilson and DuFrene (2009) discussed how the nonjudgmental acceptance of emotions leads to positive health benefits by promoting value-congruent behavior, one of the main facets of Acceptance and Commitment Therapy. Engaging in value-congruent in the presence of inevitable human suffering is also the foundation of Logotherapy. Further, these value-based approaches can be integrated with orientations dedicated to symptom reduction (Frankl, 1959; Schulenberg et al., 2008). Enhancing life circumstances through valued action is important, as it has built the foundation for modern psychotherapy. Sometimes, however, emotional inhibition prevents an individual from engaging in such action.

Meta-Analytic Techniques

Meta-analyses allow researchers the opportunity to collectively examine the effectiveness of different psychological interventions on outcome variables (Borenstein, Hedges, & Rothstein, 2007; Glass, 1976; Hedges, 1982). Although many studies produced positive outcomes associated with expressive writing, some of these studies tend to neglect important questions, the most important of which is whether or not the effect sizes are meaningful (Smyth, 1998). Meta-analyses are a technique that allow researchers to pool studies to examine an overall, weighted, population effect (Borenstein et al., 2007). Several meta-analyses of expressive writing and emotional expression have been explored: Smyth (1998), Frisina, Borod, and Lepore (2004), Frattaroli (2006), and Mogk et al. (2006). These meta-analyses have laid a foundation for exploring the effects of writing on psychological outcomes. For our purposes, we used Cohen's (1988) standards for nomenclature for small (0.20), medium (0.50), and large (0.80) d values, although it is important to note that Cohen himself suggested that these values should be based on the area of study. Generally, however, these effect size criteria are used within the social sciences.

The meta-analysis conducted by Smyth (1998) found an overall medium effect size, $d = 0.47$, for the experimental group compared to the control group. This particular analysis examined the effectiveness of expressive writing on psychological well-being, general health, and physical functioning. Frisina et al. (2004) expanded these analyses and found that expressive writing had a small to no effect on health outcomes, weighted $d = 0.07$ to $d = 0.21$. The meta-analyses conducted by Mogk et al. (2006) corroborated findings from Frisina et al. (2004). Specifically, Mogk et al. (2006) examined the effectiveness of expressive writing on somatic health symptoms, psychological outcomes, grades, and self-efficacy. The meta-analyses conducted by Smyth (1998) and Frisina et al. (2004) were relatively small in nature, using only 12-14 studies. The meta-analysis conducted by Mogk et al. (2006) included 30 studies. Newer methods of meta-analysis, including p -curve (Simonsohn, Nelson, & Simmons, 2014; Simonsohn, Simmons, & Nelson, 2015), p -uniform (van Aert, Wicherts, &

van Assen, 2016), PET-PEESE (Stanley & Doucouliagos, 2014), selection models (Vevea & Hedges, 1995), and trim and fill methods (Carter & McCullough, 2014) allow for better estimation of meta-analytic effect sizes. These analyses would be best performed by examining each potential effect separately, rather than averaging effects of each publication into one study effect size.

Additionally, Frattaroli (2006) conducted a meta-analysis to examine the effects of emotional disclosure on a variety of variables such as psychological health, physiological functioning, reported health, health behaviors, subjective impact of intervention, and general functioning/life outcomes. This meta-analysis is different from the meta conducted by Smyth (1998) in that it utilizes random effects modeling in order to calculate effect sizes. The current meta-analysis includes both random and fixed effects models for comparison. Fixed effects models assume that all studies assess the same “true” population effect size, which may be an untenable assumption across different assessments and populations (Borenstein et al., 2007). Random effects models estimate the mean of a proposed distribution of population effect sizes, which may vary by subject type or research design. Overall, Frattaroli (2006) found a weighted r effect size of .08 for all outcomes combined, which would be considered small. This meta-analysis included a very large range of studies, $N = 146$, but individual studies were again collapsed into one publication effect size, although these effects were also examined separately by health outcome.

The previous meta-analyses often focused on experimental versus control group effect sizes or p -values, rather than emphasizing change for an intervention group. This focus is likely because of the analyses provided in these publications, especially when using randomized controlled trial research designs. While this design is the gold standard for medicine, we sought to examine the magnitude of change for participants who experienced an expressive writing intervention. For example, a comparison group may increase their quality of life scores by two points in a controlled study, while the experimental group increases their quality of life scores by four points; thus, creating a significant difference in change between

the two groups. This information is valuable, but it does not tell us the magnitude of the change for the intervention group, wherein four points might only be a small effect when examined within the group who received the intervention. This study will focus on changes across time for groups who received the expressive writing task to determine what size of effects one might expect given a specific measurement schedule (i.e., one to three months, three months to six months, etc.). Additionally, we will examine the change in effects across measurement time to determine whether or not effects decrease over time.

Expressive writing tasks fit well within the framework of different psychological interventions and can be adapted for treatment, which is why the literature includes a multitude of different studies looking at a wide variety of outcomes. However, it is important to focus on individual variables in order to determine the effectiveness of expressive writing for specific diagnoses and psychopathology. As previously mentioned, some studies have found long-term benefits of expressive writing on psychological well-being (Park & Blumberg, 2002). However, other studies, such as the research completed by Lancaster et al. (2015), found no evidence supporting the utilization of expressive writing as an effective therapeutic approach. Thus, it is necessary to evaluate the effectiveness of expressive writing on specific outcome variables, and we chose to focus specifically on post-traumatic stress (PTS), post-traumatic growth (PTG), and quality of life (QOL), in line with the current positive psychology trends. By focusing on specific outcome variables, rather than specific patient or study characteristics (which are covered extensively in Frattaroli, 2006), we can examine how effective writing can be for changing stress and positive psychology phenomena as described below. With this knowledge, we can decide whether or not to incorporate this type of intervention into protocols for certain diagnoses.

Posttraumatic Stress

Post-traumatic Stress Disorder (PTSD) is a disorder involving reoccurring thoughts or experiences after a traumatic event or experience (American Psychiatric Association, 2013).

The diagnosis is based on 20 symptoms structured into four different subsets. These subsets are as follows: re-experiencing, avoidance, negative alterations in cognition and mood, and arousal (Crespo & Gomez, 2016). PTSD is a concerning disorder that affects a wide variety of groups, a few of which are sexual assault survivors (Klump, 2008), Iraq and Afghanistan war veterans (Gentes et al., 2014), and those exposed to natural disasters (Wang et al., 2000). Research conducted on the effectiveness of expressive writing on PTSD symptoms has been less successful and shows outcomes that are not as effective as other studies. While PTSD symptoms decreased after a written emotional disclosure intervention, this decrease was not significantly different than a control group change (Sloan, Marx, & Greenberg, 2011). Additionally, the emotional disclosure group showed greater emotional and heart rate responding, leading the researchers to conclude that writing may not be effective for treating PTSD.

Di Blasio et al. (2015) suggested that those meeting the criteria for moderate PTSD benefit more from expressive writing interventions in comparison to those with greater PTSD symptoms. They recruited women who had just given birth and assessed them a few days after experiencing childbirth along with a three-month follow-up. Results showed that women who had participated in the expressive writing task had lower depression and post-traumatic stress symptoms than the group assigned to a neutral writing condition. Additionally, regression models showed that expressive writing was significantly linked to a reduction of PTSD symptoms at all baseline levels. Contradicting the work conducted by Sloan et al. (2011), these results suggest that expressive writing may be an effective tool for women managing postpartum distress. However, it is important to note that only 20 of the 113 women recruited for this study qualified for a diagnosis of PTSD. This limitation suggests that those with moderate distress could perhaps benefit more from an expressive writing intervention than those diagnosed with or meeting the qualifications for PTSD. It may also explain the differences in results between the two studies, as Sloan et al. (2011) found that those with a clinical diagnosis of PTSD did not respond to an emotional

disclosure writing task. However, Di Blasio et al. (2015) found that those who reported mild symptomology better responded to the intervention than those meeting the criteria for a diagnosis of PTSD. As previously mentioned, expressive writing mirrors exposure therapy in that it encourages individuals to express avoided emotions related to the aversive event. Thus, given this observation, one would expect PTS scores to decrease in those taking part in an expressive writing intervention.

Posttraumatic Growth

While the literature mostly discusses potentially harmful outcomes to traumatic events such as emotional distress, traumatic events also provide opportunities for personal growth (Aslam & Kamal, 2013). Traumatic events, either natural or man-inflicted, can lead to positive outcomes by allowing the individual to take a different perspective (Cobb, Tedeschi, Calhoun, & Cann, 2006; Taku, Calhoun, Cann, & Tedeschi, 2008). The relationship between positive growth after a traumatic event and symptom reduction is unclear, as it is a complex process. Thus, it is necessary to examine how expressive writing might influence each variable separately, which is one of the key goals of this meta-analysis (Slavin-Spenny, Cohen, Oberleitner, & Lumley, 2011). Models receiving empirical support within the last decade suggest that traumatic events offer opportunities for both negative and positive experiences (Tedeschi & Calhoun, 1995; Weiss, 2002). Post-traumatic Growth is a positive experience after a traumatic event (Aslam & Kamal, 2013; Yilmaz & Zara, 2016). Specifically, PTG is classified as broad cognitive benefits that are seen after a traumatic experience. These benefits can be categorized into building closer relationships, examining new possibilities, appreciating life, recognizing personal strengths, and undergoing spiritual changes. (Dursun, Steger, Bentele, & Schulenberg, 2016; Tedeschi & Calhoun, 2004).

Simply, PTG is associated with a variety of desired outcomes (Dursun et al., 2016). PTG has been studied in those experiencing natural disasters, war, and other harms such as sexual assault. Finally, PTG has been studied in those experiencing medical diagnoses such

as different types of cancer and diseases. Although the relationship between PTG and symptom reduction is not yet fully understood, perhaps expressive writing allows the individual to fully comprehend the event. Expressive writing has been shown to be an effective method for reducing psychological distress among those suffering from trauma (Sloan, Marx, Epstein, & Lexington, 2007). Thus, it only makes sense to examine positive growth in response to a traumatic event after exposing participants to an expressive writing intervention. Pennebaker and Graybeal (2001) speculated that expressive writing allows an individual to feel more connected with his or her surroundings. Although this speculation does not directly explain positive outcomes after an expressive writing intervention, perhaps individuals gain a better appreciation for life after gaining a better sense of connectedness with that individual's surroundings.

Quality of Life

QOL is another positive outcome variable that is worth examining with expressive writing interventions. QOL is described as a concept comprised of multiple domains, both subjective and objective. Objectively, QOL is a measure of the extent to which an individual's needs are met. Subjectively, QOL measures an individual's attitude towards their given situation (Costanza et al., 2007). Pennebaker and Graybeal (2001) suggested that expressive writing allows one to feel more connected with their surroundings. Furthermore, they explain that expressive writing allows people to see things in a different way and better understand themselves. By understanding a traumatic event, one is able to see things differently and perhaps look at the situation with a more positive mindset. The changes that occur after expressive writing may also allow one to find meaning in the traumatic event, thereby increasing the QOL of that individual (Frankl, 1959). Higher QOL may be considered a type of PTG, which is why we thought to examine the effectiveness of studies utilizing expressive writing to improve QOL and PTG in the same study.

Current Meta-Analysis

The purpose of this meta-analysis was to examine studies utilizing expressive writing on positive outcome variables (PTG and QOL) and PTS. Due to inconsistent results in current studies published and outdated meta-analyses, it is important to clarify the effectiveness of expressive writing on promoting positive change after a traumatic event, improving overall quality of life, and reducing post-traumatic stress. This meta-analysis will provide researchers with a collected look at the use of expressive writing to promote increased PTG, increased QOL, and decreased PTS. This particular meta-analysis examines studies of patients with different types of psychopathology and medical diagnoses on PTG, QOL, and PTS. The main focus of this study is to examine PTG, QOL, and PTS by estimating effect sizes of experimental groups assigned to participate in the expressive writing intervention using newer techniques that have not been implemented in previous research. In addition to an estimate of overall effect sizes, we estimate power for effects on intervention groups, as research has shown a consistent under-powering of psychological studies, combined with a misunderstanding of the sample size needed for adequately powering one's work (Bakker, Hartgerink, Wicherts, & van der Maas, 2016).

Method

Data Collection

Studies were collected through online databases, such as PsycINFO and Google Scholar, using the following search terms and their combinations: *Posttraumatic Growth*, *PTG*, *Quality of Life*, *QOL*, *Posttraumatic Stress*, *PTS*, *Expressive Writing*, *Emotional Disclosure*. Within these articles, the change in outcome variables (PTS, PTG, QOL) from pre- to post-test was the dependent variable of interest. Generally, groups were separated into an experimental and control group and then examined at different time points. For purposes of this meta-analysis, only participants assigned to the experimental condition were examined due to having received the expressive writing intervention. If a study included multiple

assessment time points, then these measurements were examined sequentially (i.e., time 1 to time 2, time 2 to time 3) to determine change across time for the dependent variable.

220 citations focusing on PTS, PTG, and QOL were identified through the literature search and previous meta-analyses. After screening these studies, forty-five articles were retained for containing the appropriate information for this meta-analysis. A complete list of excluded articles can be found at <https://osf.io/4mjqt>, along with reasons why they were excluded. After coding articles, 202 effect sizes were calculated from the forty-five studies. On average, each study represented $M = 4.49$ ($SD = 3.50$) effects, ranging from 1 to 16 effects. 144 effects were calculated for PTS, 21 for PTG, and 37 for QOL.

Calculations for Effect Size, Variance, and Confidence Intervals

Each study implemented a pre-test to post-test style repeated measures design, usually with paired t -tests, ANOVA, or regression analyses. The means, standard deviations, and N values were collected from each study. In general, Cohen's d values were calculated using the following formula for paired t using means and standard deviations:

$$d_{av} = \frac{M_1 - M_2}{\frac{SD_1 + SD_2}{2}}$$

This equation is described in detail in Cumming (2012) as an alternative to the traditional calculation of d for paired samples t , wherein the denominator is the standard deviation of the difference scores:

$$d_z = \frac{M_1 - M_2}{SD_{diff}}$$

This equation for d_{av} not only allows for calculations from published articles that do not include SD_{diff} (i.e., most articles included), but also has been shown to be less upwardly biased than d_z . Alternative formulas include controlling for r between paired levels, as described in Lakens (2013); however, these values were not available in the selected articles, and Lakens also recommends d_{av} as an effect size for paired designs. When only mean

differences and standard deviation of the difference scores were available, the second equation for d_z was used.

We planned to use traditional and newer methods of meta-analysis, following guidelines from Cooper, Hedges, and Valentine (2009) and Borenstein et al. (2007), as well as van Aert et al. (2016). Sampling variance of the effect sizes were estimated using the *escalc()* function from the *metafor* package in *R* (Viechtbauer, 2010). The variance formula was originally published in S. B. Morris and DeShon (2002) and is shown below:

$$v = \frac{1}{n} \left(\frac{n-1}{n-3} \right) (1 + n * d^2) - \frac{d^2}{[c(n-1)]^2}$$

In this formula, n is the number of paired observations, d is the calculated effect size, and c is a correction factor, wherein df are $n - 1$ (Hedges, 1982):

$$c = 1 - \frac{3}{4 * df - 1}$$

We used the *metagen()* function in the *metafor* package to calculate both fixed and random effects models, which uses standard error of the effect to calculate overall estimates of an effect and their confidence intervals. Thus, we took the square root of the variance estimate for standard error. Given these calculations, the goal of this analysis was to calculate a combined effect size, along with a confidence interval for study planning and an assessment of the literature. A fixed effects model requires the assumption that there is a true population effect size across all studies. By including multiple measures of psychological outcomes, this assumption may be tenuous, and therefore, a random effects model was also calculated. In random effects models, the true effect is assumed to vary across studies (Borenstein et al., 2007). For a fixed effects model, the effect sizes are weighted by their inverse variance (v ; Sánchez-Meca & Marín-Martínez, 2008), which is calculated automatically in *metafor* by:

$$w_i^{FE} = \frac{1}{v}$$

The advantage to this procedure is that analyses are weighted by their precision, that is, that studies with more information (often, larger samples), are given larger weights in the overall estimated effect size (Borenstein et al., 2007). Random effects models are also weighted by inverse variance, with an additional correction for variance between studies, τ_{DL}^2 , as described by DerSimonian and Laird (1986):

$$w_i^{RE} = \frac{1}{v + \tau_{DL}^2}$$

Confidence intervals were calculated in two ways for this study. Cumming (2012), Kelley (2007), and Smithson (2001) have shown that the distribution of d values are non-normal, and thus, CIs should be estimated using the non-centrality parameter and a non-normal distribution. These values were calculated using the functions in the *MOTE* library which iteratively estimates the appropriate non-centrality parameter and converts back to d values (i.e., non-centrality parameter divided by the square root of n ; Buchanan, Valentine, & Scofield, 2017; Smithson, 2001, 2003). However, the *metafor* package in *R* uses central distributions to estimate CIs for each study and overall effect sizes. Therefore, we present both sets of values for the interested reader, as meta-analytic procedures have not implemented non-central distributions of effect sizes.

Additional Meta-Analytic Techniques

p-Curve and p-Uniform. We used *p-curve.com* to conduct a *p-curve* analysis (Simonsohn et al., 2014). The purpose of this type of analysis is to detect true effects. Specifically, *p-curve* is used to reveal possible *p-hacking* in published literature in order to decipher whether or not a true effect exists. Broadly, *p-hacking* occurs when researchers use questionable research practices to create significant results by manipulating dependent variables or covariates. Additionally, authors may add participants if the initial findings are not significant (Bruns & Ioannidis, 2016). Researchers may also decide to exclude participants for final analyses if that exclusion leads to a significant difference (L. K. John,

Loewenstein, & Prelec, 2012). Thus, it is necessary to distinguish between true and false effects in order to effectively interpret effect sizes corresponding to those p -values. p -curve accomplishes this task by examining the distributions of the published p -values. If an effect exists, or rather the results should be interpreted as presented, the distribution of p -values will be positively skewed (Simonsohn et al., 2014). If, however, no effect exists, then the distribution of p -values will be flat. p -curve analyses ultimately provide evidence of p -hacking in groups of studies and has become an important tool for interpreting meta-analyses. In order to accurately estimate effect sizes because of scrutiny associated with effect size estimation of p -curve, we also conducted p -uniform. p -uniform analyses, too, are interpreted by examining the distribution of p -values in a set of studies (van Aert et al., 2016). However, it is assumed that the population effect size equals the effect size from the dataset. Because of this assumption, the population effect size is referred to as uniform. This analysis also examines for publication bias and presents the researcher with a corrected effect size. Publication bias occurs when only select studies are published, usually only significant studies, although many factors can bias a study's publication (McShane, Böckenholt, & Hansen, 2016). p -uniform was calculated from code provided by van Aert (2017) on GitHub.

PET-PEESE. Originally, meta-analyses relied on the calculation of Egger's regression test which examined the relationship of the standard error (predictor) to the effect size estimates (criterion). In this regression, the intercept values were used to determine if effect size measures were different than zero, by providing a meta-analytic estimate (Egger, Davey Smith, Schneider, & Minder, 1997; Stanley, 2005). PET-PEESE analyses examine for publication bias by adapting parts from Egger's traditional regression tests: PET (Precision Effect Test) and PEESE (Precision Effect Estimate with Standard Error, Carter & McCullough, 2014). PET is a more reliable test of publication bias with effect size estimates of zero, $b_0 = 0$, while PEESE is more accurate with non-zero effect size estimates, $b_0 \neq 0$ (Stanley & Doucouliagos, 2014). PET-PEESE was calculated using Hilgard's (2016) code provided on GitHub.

Selection Models. Selection model analyses provide the researcher with a test of publication bias and effect size estimates using maximum likelihood estimation (Vevea & Hedges, 1995; Vevea & Woods, 2005). Using selection models, researchers are able to discover effect size estimates as well as evidence of publication bias (McShane et al., 2016) by using a mixed general linear model to estimate these values. Selection models were calculated with the *weightr* package in *R* (Coburn & Vevea, 2017).

Trim and Fill. Trim and Fill analyses, in contrast to PET-PEESE, regress standard error (criterion) and effect size estimates (predictor). Specifically, the purpose of Trim and Fill techniques is to examine whether or not publication bias may influence the regression equation (Carter & McCullough, 2014). Effect sizes and standard error terms are graphically displayed on x and y-axes, respectively, in a funnel plot. If this graphical representation indicates asymmetry, considered a gap of missing data points in the lower center area of the plot, the study set can be assumed to have studies that are both non-significant and small in sample size (van Assen, van Aert, & Wicherts, 2015). This funnel is then trimmed until symmetry is achieved. Missing studies from the symmetrical graph are imputed (filled) while maintaining the given symmetry (Duval & Tweedie, 2000). The meta-analytic effect size is then estimated from the trimmed and filled funnel plot. Trim and fill analyses, as well as funnel plots included below, were calculated with the *metafor* package.

Results

PTS

Overall Effect Size. As described above, both fixed effects and random effects models with centralized confidence intervals are presented in Table 1. Studies were examined for potential outliers using the *metafor* package in *R*. This package calculates traditional regression influence values, such as Cook's and hat values (J. Cohen, 1988). These values indicate change in overall meta-analytic model with and without the effect; thus, determining their impact on the pooled effect size (Viechtbauer, 2010). Because published

studies likely represent the range of the sampling distribution of effect sizes, we included the analyses with and without outliers to present evidence for both paths a researcher might take when examining an overall effect.

Three outliers were detected with this procedure, all showing very large effect sizes, average $d = 1.63$. The fixed and random effects estimates without these points are also included in Table 1. Figures 1, 2, 3, and 4 portray the effect sizes for PTS studies, separated by intrusions, avoidance, hyperarousal, and total scores for easier viewing (i.e., over 100+ effect sizes did not fit easily on one combined graph). Although these categories are not reflective of updated DSM-V criteria, researchers have not yet conducted enough studies using expressive writing on PTS with updated PTSD criteria to warrant a meta-analysis. Name acronym coding can be found in the data online. This forest plot includes the non-centralized confidence interval calculated from the *MOTE* library (Buchanan et al., 2017). Shape size indicates study weight, and these values were taken from the overall random effects meta-analysis and normalized by dividing by the mean weight. The dashed lines indicate the average non-weighted lower and upper confidence interval limit for the non-centralized estimates. Overall, PTS studies include a small effect size that appears to be significantly greater than zero across all estimate types (fixed, random, with or without outliers).

Homogeneity. A prerequisite for newer meta-analytic techniques includes the assessment of homogeneity of the effects (van Aert et al., 2016). Using the *metafor* package in *R*, we calculated the Q -statistic and the I^2 index (Cochran, 1954; Huedo-Medina, Sánchez-Meca, Marín-Martínez, & Botella, 2006). Significant values imply inconsistencies across the variable or variables of interest and are represented by Q . In contrast, I^2 indicates the percentage of heterogeneity along with a 95% CI. Both can, however, be biased with a small number of experiments included for analyses (Higgins, Thompson, Deeks, & Altman, 2003; Huedo-Medina et al., 2006). Thus, we sought to calculate an overall level of heterogeneity after examining each variable separately before and after excluding outliers.

For PTS studies including outliers, we found significant heterogeneity, $Q(143) = 639.98$, $p < .001$ and $I^2 = 77.7$, 95% CI[73.9 - 80.9]. These values were reduced slightly with the exclusion of outliers, $Q(140) = 519.75$, $p < .001$ and $I^2 = 73.1$, 95% CI[68.2 - 77.2].

Power. Power was calculated in two different ways using the *pwr* package in *R* (Champely, 2016). *Post hoc* power was first calculated using sample size and effect size statistics from each individual study. Additionally, we calculated power using the study sample size and estimated overall effect size from the random effects model with and without outliers, as explained by G. Francis (2012) and G. Francis (2014). The first estimate indicates the likelihood of finding an effect from our sample statistics, while the second indicates the likelihood of finding the true population effect size. If each study had been conducted on only the change in the experimental group, 45.1% of studies would have been considered significant at $\alpha < .05$. The average power of these studies based on their original study characteristics was .46 ($SD = .36$). Power for the random-effects meta-analytic effect size with outliers was .47 ($SD = .24$) and without outliers was .42 ($SD = .23$). Therefore, power consistently was around 40-50% for studies examining PTS, regardless of outlier effects. In these studies, only 26.4% achieved recommended 80% power for their found effect size, a smaller 16.7% for the random-effect outlier effect size, and even smaller 6.9% for power calculations on the random-effect size without the outliers.

Other Meta-Analytic Estimates. As noted in van Aert et al. (2016), *p*-curve and *p*-uniform analyses are upwardly biased when heterogeneity is high. Therefore, we use caution when interpreting these analyses on PTS outcomes. As seen in Table 1, the estimates for *p*-uniform were higher than other techniques, likely because of the focus on significant *p*-values and the great degree of heterogeneity described earlier. *P*-curve pictures can be found at <https://osf.io/4mjqt/> online, and this analysis indicated evidentiary value at $p < .001$. Additionally, the *p*-uniform analysis indicated that there was likely no publication bias present, $Z = -5.02$, $p = 1.000$. When examining the PET analysis, we found that the intercept was significant, which indicated that PEESE was likely a better estimator of the

meta-analytic effect size. PEESE estimates were lower than the original meta-analytic estimate, but confidence intervals indicated that the effect is small to medium, and still larger than zero. Selection models indicated a larger effect size, especially with the random-effects models, and these effects were influenced by the outliers found in the published studies. Trim and fill models are shown in Table 1, and figures are included online. Nineteen missing studies were imputed for both models with and without outliers. Across all these effect size estimates, we found that expressive writing was likely to decrease PTS symptoms in a small to moderate way. The correlation of effect size with time between measurement times was $r = -.16$, 95% CI $[-.32, .00]$, $t(142) = -1.99$, $p = .049$, and $r = -.15$, 95% CI $[-.30, .02]$, $t(139) = -1.75$, $p = .082$ without outliers. This result indicated that the effect of expressive writing slightly decreased across time.

PTG

Overall Effect Size. Both fixed and random effects models with centralized confidence intervals for PTG are presented in Table 2. When examining expressive writing on PTG, no outliers were detected. Fixed and random effects estimates are included in Table 2, while Figure 5 shows effect sizes for PTG studies where shape size indicates the normalized weight of the study. Dashed lines indicate non-weighted lower and upper confidence intervals for non-centralized estimates. Overall, PTG studies indicated a negligible to small effect size across both random and fixed effects models, and the non-centralized confidence intervals indicated an effect that crossed zero.

Homogeneity. Using the *metafor* package in *R*, we calculated both a Q statistic and I^2 index. Since PTG studies did not contain any outliers, we did not calculate two separate analyses to examine heterogeneity both with and without outliers. We did not find significant heterogeneity across PTG studies, $Q(20) = 14.18$, $p = .82$ and $I^2 = 0.0$, 95% CI $[0.0 - 25.3]$.

Power. First, we calculated *post hoc* power using both sample and effect size statistics from individual studies. Individual studies examining change in experimental groups showed that 9.5% of studies would have been considered significant at $\alpha < .05$. Average power of PTG studies was .15 ($SD = .16$). 0.0% achieved recommended 80% power for their found effect size. Additionally, we calculated power using study sample size and estimated effect size from our random effects model. Power for the true effect size was .08 ($SD = .02$). Again, 0.0% achieved recommended 80% power.

Other Meta-Analytic Estimates. Due to no heterogeneity across PTG studies, we can use both *p*-curve and *p*-uniform analyses with more confidence. A pictorial representation of *p*-curve can be found at <https://osf.io/4mjqt/>. This analysis did not indicate evidentiary value, $p = .75$, as only two of the results would be considered significant at $\alpha < .05$. *p*-uniform estimates are presented in Table 2. Specifically, these analyses indicated that there was no publication bias present, $Z = 0.70$, $p = .243$. The *p*-uniform estimates of the effect size for PTG were negative, in contrast to the fixed and random effects overall model. The confidence interval for this analysis indicates a wide range of possible effects. In examining PET-PEESE analyses, we did not find a significant intercept, indicating that PET is most likely a better effect size estimator. PET analyses indicated that the effect size is negligible to small, with our confidence interval crossing zero. These results corroborated our original effect size calculations. Selection models indicated negligible to small effect sizes, again wherein the confidence interval includes zero effect. Trim and fill models are shown in Table 2, and figures are included online. Zero studies were imputed for our model, and thus, the effect size estimate is the same as the overall model. Across techniques, we found that expressive writing has little to no effect on PTG. The correlation of effect size across measurement times in PTG studies at subsequent time points was $r = .09$, 95% CI $[-.36, .50]$, $t(19) = 0.38$, $p = .707$, and no change over time was found.

QOL

Overall Effect Size. Finally, for QOL, both fixed and random effects models with centralized confidence intervals are presented in Table 3. Two outliers were detected with this procedure, average $d = -0.07$. While the average effect of these outliers indicates a small number, it is important to note that these two outliers were the largest positive and negative effects found from the Possemato, Ouimette, and Geller (2010) study. Fixed and random effects estimates without these points are also included in Table 3, while Figure 6 shows effect sizes for QOL studies. Overall, QOL studies indicated a negligible to small effect that showed a non-significant decrease in quality of life as a result of expressive writing.

Homogeneity. For QOL studies including outliers, we found significant heterogeneity from our random effects model, $Q(36) = 200.09$, $p < .001$ and $I^2 = 82.0$, 95% CI[75.9 - 86.5]. After excluding outliers, our random effects model still indicated heterogeneity, $Q(34) = 93.18$, $p < .001$ and $I^2 = 63.5$, 95% CI[47.6 - 74.6].

Power. In conducting *post hoc* power using sample and effect size statistics from individual studies, we found that 21.6% of studies would have been considered significant at $\alpha < .05$. Average power based on actual study characteristics was .33 ($SD = .32$). Power for the random effects meta-analytic effect size with outliers was .05 ($SD = .00$) and without outliers was .05 ($SD = .00$). Unfortunately, power was around 5% for both random effects models with and without outliers. In these studies, 18.9% achieved adequate power of 80% on their found effect size, while 0.0% achieved 80% power for our random effects model with outliers. Finally, without outliers, 0.0% achieved 80% power.

Other Meta-Analytic Estimates. We exert caution in interpreting p -curve and p -uniform analyses on QOL outcomes with and without outliers due to heterogeneity. As seen in Table 1, p -uniform estimates were stronger and positive than other techniques because of the high degree of heterogeneity recently described. p -curve pictures can be found at the following OSF Link: <https://osf.io/4mjqt>. Eight studies were significant at $\alpha < .05$, and the studies indicated evidentiary value, $p = .004$. p -uniform analyses did not indicate publication bias, $Z = -2.75$, $p = .997$. In PET-PEESE analyses, we found that the intercept

was not significant, and therefore, PET was a better estimator of the meta-analytic effect. Table 1 indicates that both of these analyses estimate the effect size around zero, with a confidence interval that includes zero. Selection models correspondingly show small effects crossing zero, except for random effects models with outliers, that appear to be heavily influenced by the outliers. Trim and fill models are shown in Table 3, and figures are included online. No studies were imputed for these analyses, and therefore, the effect size estimates match the original meta-analysis. Overall, these results appear to point to no effects, ranging across zero with several negative estimates. Interestingly, the correlation of effect sizes across measurement times with outliers was $r = -.37$, 95% CI $[-.62, -.05]$, $t(35) = -2.33$, $p = .026$ and $r = -.64$, 95% CI $[-.80, -.39]$, $t(33) = -4.75$, $p < .001$ without outliers. The effect of expressive writing appears to be positive at short time intervals and decreases into negative effects at longer time intervals.

Discussion

In examining pre- to post-test comparisons across each variable separately, we found that PTS studies indicated a small effect size across all meta-analytic estimates. Both QOL and PTG studies indicated a negligible to small effect size using random effects models. Although the PTG effect in our overall meta-analysis estimate was significant, other methods indicate this small effect is likely not different from zero. QOL was not different from zero, which suggests no effect of expressive writing on quality of life. Interestingly, these results are in contrast to Sloan et al. (2011), which suggested that only certain groups may respond to these interventions. Potentially, the high heterogeneity may be due to the mixed levels of PTSD in these studies, as Di Blasio et al. (2015) indicates that only certain levels of PTSD are responsive to an expressive writing condition.

Expressive writing does not appear to play an important role in influencing positive growth or improved quality of life post intervention. Ineffective emotional expression may be a contributing factor. Additionally, future research might examine specific methodology for

these types of studies. If participants/clients are not deeply engaged with the material, an expressive writing intervention may not be effective, as Pennebaker and Graybeal (2001) imply that connectedness is an important factor for the intervention. However, it may be difficult to implement a check for engagement in these types of research designs. Doing so may also set a context that will inhibit emotional processing and general responses. Research on expressive writing has found a wide range of outcomes for different variables (Frattaroli, 2006), and these various results may explain the large heterogeneity found in this study. Encouragingly, however, we did not find much evidence of publication bias, and therefore, these estimates may represent a true population effect size.

We also examined the relationship of time between measurements of the dependent variables and the corresponding effect size to determine if effects change over time. For both PTS and PTG, there was no relationship between effect size and time; yet, PTS indicated a small negative correlation. For QOL studies, a medium to large negative correlation was found. A negative relationship between time and effect size implies that interventions were more effective in the initial time points, and effects decreased over longer time spans.

The psychological scientific community has shifted focus to reproducibility and research design in the last several years (Nelson, Simmons, & Simonsohn, 2018), and much of this discussion has focused on adequately powering studies for publication (Bakker et al., 2016; S. E. Maxwell, Lau, & Howard, 2015). S. E. Maxwell et al. (2015) and Open Science Collaboration (2015) have shown that the “replication crisis” may be attributed to low power in published studies. The power found in the current meta-analysis was very poor, with very few studies reaching the suggested 80% criterion to adequately power their study. This result was the same when considering individual study characteristics or the estimate true population effect size. Research by Bakker et al. (2016) indicates that researchers’ intuitions about power are particularly poor, and many studies could benefit from more informed power analyses. Anderson, Kelley, and Maxwell (2017) recently published a primer on power, with an online application to help with sample size planning for many types of research

617 designs. Additionally, we encourage researchers to report power analyses of studies in order
618 to better understand methodology for replication and reproducibility.

619 Meta-analyses, while useful tools to pool for population effect sizes, contain various
620 limitations to their usefulness (van Elk et al., 2015). As mentioned previously, these analyses
621 can be affected by high heterogeneity, which was found in this study (van Aert et al., 2016).
622 Selection models have been criticized when using a smaller number of studies (van Assen et
623 al., 2015), and trim and fill analyses may not always estimate accurate confidence intervals
624 and funnel plots may be biased with heterogeneity (Terrin, Schmid, Lau, & Olkin, 2003).
625 When focusing on improving the psychological sciences, van Elk et al. (2015) suggest that
626 the reliability and size of effects may be best elucidated by conducting large preregistered
627 studies. This suggestion will also improve the outlook for power in published studies, and
628 projects such as Many Labs can aide in subsidizing large samples (R. A. Klein et al., 2014).
629 Even with limitations, meta-analyses allow researchers to examine the state of a research
630 area, and we find potential with expressive writing on reducing PTS symptoms, and an
631 overall need for better sample size and power planning for studies.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*. doi:[10.1176/appi.books.9780890425596.744053](https://doi.org/10.1176/appi.books.9780890425596.744053)
- Anderson, S. F., Kelley, K., & Maxwell, S. E. (2017). Sample-size planning for more accurate statistical power: A method adjusting sample effect sizes for publication bias and uncertainty. *Psychological Science*, 28(11), 1547–1562. doi:[10.1177/0956797617723724](https://doi.org/10.1177/0956797617723724)
- Aslam, N., & Kamal, A. (2013). Gender differences in distress responses, rumination patterns, perceived social support and posttraumatic growth among flood affected individuals. *Journal of Pakistan Psychiatric Society*, 10, 86–90.
- Bakker, M., Hartgerink, C. H. J., Wicherts, J. M., & van der Maas, H. L. J. (2016). Researchers' intuitions about power in psychological research. *Psychological Science*, 27(8), 1069–1077. doi:[10.1177/0956797616647519](https://doi.org/10.1177/0956797616647519)
- Bodor, N. (2002). The health effects of emotional disclosure for individuals with Type I Diabetes. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 64(10-B), 5207.
- Borenstein, M., Hedges, L. V., & Rothstein, H. (2007). Meta-analysis fixed effect vs. random effects. Retrieved from <https://www.meta-analysis.com/downloads/Meta-analysis-fixed-effect-vs-random-effects-072607.pdf>
- Brounéus, K. (2010). The trauma of truth telling: Effects of witnessing in the Rwandan Gacaca Courts on psychological health. *Journal of Conflict Resolution*, 54(3), 408–437. doi:[10.1177/0022002709360322](https://doi.org/10.1177/0022002709360322)
- Bruns, S. B., & Ioannidis, J. P. A. (2016). p-curve and p-hacking in observational research. *PLoS ONE*, 11(2). doi:[10.1371/journal.pone.0149144](https://doi.org/10.1371/journal.pone.0149144)
- Buchanan, E. M., Valentine, K. D., & Scofield, J. E. (2017). MOTE. Retrieved from <https://github.com/doomlab/MOTE>
- Carter, E. C., & McCullough, M. E. (2014). Publication bias and the limited strength model of self-control: Has the evidence for ego depletion been overestimated? *Frontiers in*

Psychology, 5(July), 1–11. doi:[10.3389/fpsyg.2014.00823](https://doi.org/10.3389/fpsyg.2014.00823)

Champely, S. (2016). pwr: Basic functions for power analysis. R package version 1.2-0.

Retrieved from <https://cran.r-project.org/package=pwr>

Cobb, A. R., Tedeschi, R. G., Calhoun, L. G., & Cann, A. (2006). Correlates of posttraumatic growth in survivors of intimate partner violence. *Journal of Traumatic Stress*, 19(6), 895–903. doi:[10.1002/jts.20171](https://doi.org/10.1002/jts.20171)

Coburn, K. M., & Vevea, J. L. (2017). Weightr. Retrieved from

<https://cran.r-project.org/web/packages/weightr/index.html>

Cochran, W. G. (1954). Some methods for strengthening the common χ^2 tests. *Biometrics*, 10(4), 417–451. doi:[10.2307/3001616](https://doi.org/10.2307/3001616)

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Earlbaum.

Cooper, H., Hedges, L. V., & Valentine, J. (2009). *The handbook of research synthesis and meta-analysis* (2nd ed.). New York, NY: Russell Sage Foundation.

Costanza, R., Fisher, B., Ali, S., Beer, C., Bond, L., Boumans, R., . . . Snapp, R. (2007).

Quality of life: An approach integrating opportunities, human needs, and subjective well-being. *Ecological Economics*, 61(2-3), 267–276.

doi:[10.1016/j.ecolecon.2006.02.023](https://doi.org/10.1016/j.ecolecon.2006.02.023)

Crespo, M., & Gomez, M. M. (2016). Diagnostic concordance of DSM-IV and DSM-5 posttraumatic stress disorder (PTSD) in a clinical sample. *Psicothema*, 28(2), 161–166. doi:[10.7334/psicothema2015.213](https://doi.org/10.7334/psicothema2015.213)

Cumming, G. (2012). *Understanding the new statistics: Effect sizes, confidence intervals, and meta-analysis*. New York, NY: Routledge.

DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3), 177–188. doi:[10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2)

Di Blasio, P., Camisasca, E., Caravita, S. C. S., Ionio, C., Milani, L., Valtolina, G. G., . . . Valtolina, G. G. (2015). The effects of expressive writing on postpartum depression

and posttraumatic stress symptoms. *Psychological Reports*, 117(3), 856–882.

doi:[10.2466/02.13.PR0.117c29z3](https://doi.org/10.2466/02.13.PR0.117c29z3)

Dursun, P., Steger, M. F., Bentele, C., & Schulenberg, S. E. (2016). Meaning and posttraumatic growth among survivors of the September 2013 Colorado floods. *Journal of Clinical Psychology*, 72(12), 1247–1263. doi:[10.1002/jclp.22344](https://doi.org/10.1002/jclp.22344)

Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56(2), 455–463. doi:[10.1111/j.0006-341X.2000.00455.x](https://doi.org/10.1111/j.0006-341X.2000.00455.x)

Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315(7109), 629–634. doi:[10.1136/bmj.316.7129.469](https://doi.org/10.1136/bmj.316.7129.469)

Esterling, B. A., Antoni, M. H., Kumar, M., & Schneiderman, N. (1990). Emotional repression, stress disclosure responses, and Epstein-Barr viral capsid antigen titers. *Psychosomatic Medicine*, 52, 397–410. doi:[10.1097/00006842-199007000-00002](https://doi.org/10.1097/00006842-199007000-00002)

Fawzy, N. W., Fawzy, N. W., Hyun, C. S., Elashoff, R., Guthrie, D., Fahey, J. L., & Morton, D. L. (1993). Malignant melanoma. Effects of an early structured psychiatric intervention, coping, and affective state on recurrence and survival 6 years later. *Archives of General Psychiatry*, 50(9), 681–689. doi:[10.1001/archpsyc.1993.01820210015002](https://doi.org/10.1001/archpsyc.1993.01820210015002)

Francis, G. (2012). Publication bias and the failure of replication in experimental psychology. *Psychonomic Bulletin & Review*, 19(6), 975–991. doi:[10.3758/s13423-012-0322-y](https://doi.org/10.3758/s13423-012-0322-y)

Francis, G. (2014). The frequency of excess success for articles in Psychological Science. *Psychonomic Bulletin & Review*, 21(5), 1180–1187. doi:[10.3758/s13423-014-0601-x](https://doi.org/10.3758/s13423-014-0601-x)

Francis, M. E., & Pennebaker, J. W. (1992). Putting stress into words: The impact of writing on physiological, absentee, and self-reported emotional well-being measures.

- American Journal of Health Promotion*, 6(4), 280–287. doi:[10.4278/0890-1171-6.4.280](https://doi.org/10.4278/0890-1171-6.4.280)
- Frankl, V. (1959). *Man's search for meaning*. Boston, MA: Beacon Press.
- Frattaroli, J. (2006). Experimental disclosure and its moderators: A meta-analysis. *Psychological Bulletin*, 132(6), 823–865. doi:[10.1037/0033-2909.132.6.823](https://doi.org/10.1037/0033-2909.132.6.823)
- Frisina, P. G., Borod, J. C., & Lepore, S. J. (2004). A meta-analysis of the effects of written emotional disclosure on the health outcomes of clinical populations. *The Journal of Nervous and Mental Disease*, 192(9), 629–634. doi:[10.1097/01.nmd.0000138317.30764.63](https://doi.org/10.1097/01.nmd.0000138317.30764.63)
- Gentes, E. L., Dennis, P. A., Kimbrel, N. A., Rissling, M. B., Beckham, J. C., & Calhoun, P. S. (2014). DSM-5 posttraumatic stress disorder: Factor structure and rates of diagnosis. *Journal of Psychiatric Research*, 59(1), 60–67. doi:[10.1016/j.jpsychires.2014.08.014](https://doi.org/10.1016/j.jpsychires.2014.08.014)
- Gidron, Y., Peri, T., Connolly, J. F., & Shalev, A. Y. (1996). Written disclosure in posttraumatic stress disorder: Is it beneficial for the patient? *The Journal of Nervous and Mental Disease*, 184(8), 505–506. doi:[10.1097/00005053-199608000-00009](https://doi.org/10.1097/00005053-199608000-00009)
- Glass, G. V. (1976). Primary, secondary, and meta-analysis of research. *Educational Researcher*, 5(10), 3–8. doi:[10.3102/0013189X005010003](https://doi.org/10.3102/0013189X005010003)
- Goldstein, H. S., Edelberg, R., Meier, C. F., & Davis, L. (1988). Relationship of resting blood pressure and heart rate to experienced anger and expressed anger. *Psychosomatic Medicine*, 50(4), 321–329. doi:[10.1097/00006842-198807000-00001](https://doi.org/10.1097/00006842-198807000-00001)
- Greenberg, M. A., & Stone, A. A. (1992). Emotional disclosure about traumas and its relation to health: Effects of previous disclosure and trauma severity. *Journal of Personality and Social Psychology*, 63, 75–84. doi:[10.1037/h0090372](https://doi.org/10.1037/h0090372)
- Gross, J. J., & Levenson, R. W. (1997). Hiding feelings: The acute effects of inhibiting negative and positive emotion. *Journal of Abnormal Psychology*, 106(1), 95–103. doi:[10.1037/0021-843X.106.1.95](https://doi.org/10.1037/0021-843X.106.1.95)
- Harris, A. H. S., Thoresen, C. E., Humphreys, K., & Faul, J. (2005). Does writing affect

asthma? A randomized trial. *Psychosomatic Medicine*, 67(1), 130–136.

doi:[10.1097/01.psy.0000146345.73510.d5](https://doi.org/10.1097/01.psy.0000146345.73510.d5)

Hedges, L. V. (1982). Estimation of effect size from a series of independent experiments.

Psychological Bulletin, 92(2), 490–499. doi:[10.1037/0033-2909.92.2.490](https://doi.org/10.1037/0033-2909.92.2.490)

Henry, E. A., Schlegel, R. J., Talley, A. E., Molix, L. A., & Bettencourt, B. A. (2010). The

feasibility and effectiveness of expressive writing for rural and urban breast cancer

survivors. *Oncology Nursing Forum*, 37(6), 749–757. doi:[10.1188/10.ONF.749-757](https://doi.org/10.1188/10.ONF.749-757)

Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring

inconsistency in meta-analyses. *British Medical Journal*, 327(7414), 557–560.

doi:[10.1136/bmj.327.7414.557](https://doi.org/10.1136/bmj.327.7414.557)

Hilgard, J. (2016). PETPEESE. GitHub. Retrieved from

<https://github.com/Joe-Hilgard/PETPEESE>

Huedo-Medina, T. B., Sánchez-Meca, J., Marín-Martínez, F., & Botella, J. (2006). Assessing

heterogeneity in meta-analysis: Q statistic or I² index? *Psychological Methods*, 11(2),

193–206. doi:[10.1037/1082-989X.11.2.193](https://doi.org/10.1037/1082-989X.11.2.193)

John, L. K., Loewenstein, G., & Prelec, D. (2012). Measuring the prevalence of questionable

research practices with incentives for truth telling. *Psychological Science*, 23(5),

524–532. doi:[10.1177/0956797611430953](https://doi.org/10.1177/0956797611430953)

Kelley, K. (2007). Confidence intervals for standardized effect sizes. *Journal of Statistical*

Software, 20(8), 1–24. doi:[10.18637/jss.v020.i08](https://doi.org/10.18637/jss.v020.i08)

Klein, R. A., Ratliff, K. A., Vianello, M., Adams, R. B., Bahník, Š., Bernstein, M. J., . . .

Nosek, B. A. (2014). Investigating variation in replicability. *Social Psychology*, 45(3),

142–152. doi:[10.1027/1864-9335/a000178](https://doi.org/10.1027/1864-9335/a000178)

Klump, M. C. (2008). Posttraumatic stress disorder and sexual assault in women. *Journal of*

College Student Development, 8225(May 2014), 37–41. doi:[10.1300/J035v21n02](https://doi.org/10.1300/J035v21n02)

Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: A

practical primer for t-tests and ANOVAs. *Frontiers in Psychology*, 4.

doi:[10.3389/fpsyg.2013.00863](https://doi.org/10.3389/fpsyg.2013.00863)

Lancaster, S. L., Klein, K. P., & Heifner, A. (2015). The validity of self-reported growth after expressive writing. *Traumatology*, 21(4), 293–298. doi:[10.1037/trm0000052](https://doi.org/10.1037/trm0000052)

Larson, D. G., & Chastain, R. L. (1990). Self-concealment: Conceptualization, measurement, and health implications. *Journal of Social and Clinical Psychology*, 9(4), 439–455. doi:[10.1521/jscp.1990.9.4.439](https://doi.org/10.1521/jscp.1990.9.4.439)

Lieberman, M. A., & Goldstein, B. A. (2006). Not all negative emotions are equal: The role of emotional expression in online support groups for women with breast cancer. *Psycho-Oncology*, 15(2), 160–168. doi:[10.1002/pon.932](https://doi.org/10.1002/pon.932)

Maxwell, S. E., Lau, M. Y., & Howard, G. S. (2015). Is psychology suffering from a replication crisis? What does “failure to replicate” really mean? *American Psychologist*, 70(6), 487–498. doi:[10.1037/a0039400](https://doi.org/10.1037/a0039400)

McShane, B. B., Böckenholt, U., & Hansen, K. T. (2016). Adjusting for publication bias in meta-analysis. *Perspectives on Psychological Science*, 11(5), 730–749. doi:[10.1177/1745691616662243](https://doi.org/10.1177/1745691616662243)

Meshberg-Cohen, S., Svikis, D., & McMahon, T. J. (2014). Expressive writing as a therapeutic process for drug-dependent women. *Substance Abuse*, 35(1), 80–88. doi:[10.1080/08897077.2013.805181](https://doi.org/10.1080/08897077.2013.805181)

Mogk, C., Otte, S., Reinhold-Hurley, B., & Kröner-Herwig, B. (2006). Health effects of expressive writing on stressful or traumatic experiences - a meta-analysis. *Psychosocial Medicine*, 3, Doc06.

Morris, S. B., & DeShon, R. P. (2002). Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychological Methods*, 7(1), 105–125. doi:[10.1037/1082-989X.7.1.105](https://doi.org/10.1037/1082-989X.7.1.105)

Nelson, L. D., Simmons, J., & Simonsohn, U. (2018). Psychology’s renaissance. *Annual Review of Psychology*, 69(1), 511–534. doi:[10.1146/annurev-psych-122216-011836](https://doi.org/10.1146/annurev-psych-122216-011836)

Open Science Collaboration. (2015). Estimating the reproducibility of psychological science.

Science, 349(6251), aac4716–aac4716. doi:[10.1126/science.aac4716](https://doi.org/10.1126/science.aac4716)

Park, C. L., & Blumberg, C. J. (2002). Disclosing trauma through writing: Testing the meaning-making hypothesis. *Cognitive Therapy and Research*, 26(5), 597–616.

doi:[10.1023/A:1020353109229](https://doi.org/10.1023/A:1020353109229)

Pennebaker, J. W. (1989). Confession, inhibition, and disease. In L. Berkowitz (Ed.), *Advances in experimental social psychology* (Vol. 22, pp. 211–244). Academic Press.

doi:[10.1016/S0065-2601\(08\)60309-3](https://doi.org/10.1016/S0065-2601(08)60309-3)

Pennebaker, J. W. (1993). Putting stress into words: Health, linguistic, and therapeutic implications. *Behaviour Research and Therapy*, 31(6), 539–548.

doi:[10.1016/0005-7967\(93\)90105-4](https://doi.org/10.1016/0005-7967(93)90105-4)

Pennebaker, J. W., & Beall, S. K. (1986). Confronting a traumatic event: Toward an understanding of inhibition and disease. *Journal of Abnormal Psychology*, 95(3), 274–281.

doi:[10.1037//0021-843X.95.3.274](https://doi.org/10.1037//0021-843X.95.3.274)

Pennebaker, J. W., & Francis, M. E. (1996). Cognitive, emotional, and language processes in disclosure. *Cognition & Emotion*, 10(6), 601–626. doi:[10.1080/026999396380079](https://doi.org/10.1080/026999396380079)

Pennebaker, J. W., & Graybeal, A. (2001). Patterns of natural language use: Disclosure, personality, and social integration. *Current Directions in Psychological Science*, 10(3), 90–93.

doi:[10.1111/1467-8721.00123](https://doi.org/10.1111/1467-8721.00123)

Pennebaker, J. W., Colder, M., & Sharp, L. K. (1990). Accelerating the coping process. *Journal of Personality and Social Psychology*, 58(3), 528–537.

doi:[10.1037//0022-3514.58.3.528](https://doi.org/10.1037//0022-3514.58.3.528)

Pennebaker, J. W., Kiecolt-Glaser, J. K., & Glaser, R. (1988). Disclosure of traumas and immune function: Health implications for psychotherapy. *Journal of Consulting and Clinical Psychology*, 56(2), 239–245.

doi:[10.1037/0022-006X.56.2.239](https://doi.org/10.1037/0022-006X.56.2.239)

Possemato, K., Ouimette, P., & Geller, P. (2010). Internet-based expressive writing for kidney transplant recipients: Effects on posttraumatic stress and quality of life.

Traumatology, 16(1), 49–54. doi:[10.1177/1534765609347545](https://doi.org/10.1177/1534765609347545)

Rachman, S. (1980). Emotional processing. *Behaviour Research and Therapy*, 18(1), 51–60.

doi:[10.1016/0005-7967\(80\)90069-8](https://doi.org/10.1016/0005-7967(80)90069-8)

Sánchez-Meca, J., & Marín-Martínez, F. (2008). Confidence intervals for the overall effect size in random-effects meta-analysis. *Psychological Methods*, 13(1), 31–48.

doi:[10.1037/1082-989X.13.1.31](https://doi.org/10.1037/1082-989X.13.1.31)

Scheff, T. J. (1979). *Catharsis in healing, ritual, and drama*. Los Angeles: University of California Press.

Schoutrop, M. J. A., Lange, A., Hanewald, G., Davidovich, U., & Salomon, H. H. (2002). Structured writing and processing major stressful events: A controlled trial.

Psychotherapy and Psychosomatics, 71(3), 151–157. doi:[10.1159/000056282](https://doi.org/10.1159/000056282)

Schulenberg, S. E., Hutzell, R. R., Nassif, C., & Rogina, J. M. (2008). Logotherapy for clinical practice. *Psychotherapy*, 45(4), 447–463. doi:[10.1037/a0014331](https://doi.org/10.1037/a0014331)

Simonsohn, U., Nelson, L. D., & Simmons, J. P. (2014). p-curve: A key to the file-drawer.

Journal of Experimental Psychology: General, 143(2), 534–547. doi:[10.1037/a0033242](https://doi.org/10.1037/a0033242)

Simonsohn, U., Simmons, J. P., & Nelson, L. D. (2015). Better p-curves: Making p-curve analysis more robust to errors, fraud, and ambitious p-hacking, a reply to Ulrich and Miller (2015). *Journal of Experimental Psychology: General*, 144(6), 1146–1152.

doi:[10.1037/xge0000104](https://doi.org/10.1037/xge0000104)

Slavin-Spenny, O. M., Cohen, J. L., Oberleitner, L. M., & Lumley, M. A. (2011). The effects of different methods of emotional disclosure: Differentiating posttraumatic growth from stress symptoms. *Journal of Clinical Psychology*, 67(10), 993–1007.

doi:[10.1002/jclp.20750](https://doi.org/10.1002/jclp.20750)

Sloan, D. M., Marx, B. P., & Epstein, E. M. (2005). Further examination of the exposure model underlying the efficacy of written emotional disclosure. *Journal of Consulting and Clinical Psychology*, 73(3), 549–554. doi:[10.1037/0022-006X.73.3.549](https://doi.org/10.1037/0022-006X.73.3.549)

Sloan, D. M., Marx, B. P., & Greenberg, E. M. (2011). A test of written emotional

- disclosure as an intervention for posttraumatic stress disorder. *Behaviour Research and Therapy*, 49(4), 299–304. doi:[10.1016/j.brat.2011.02.001](https://doi.org/10.1016/j.brat.2011.02.001)
- Sloan, D. M., Marx, B. P., Epstein, E. M., & Lexington, J. M. (2007). Does altering the writing instructions influence outcome associated with written disclosure? *Behavior Therapy*, 38(2), 155–168. doi:[10.1016/j.beth.2006.06.005](https://doi.org/10.1016/j.beth.2006.06.005)
- Smithson, M. (2001). Correct confidence intervals for various regression effect sizes and parameters: The importance of noncentral distributions in computing intervals. *Educational and Psychological Measurement*, 61(4), 605–632. doi:[10.1177/00131640121971392](https://doi.org/10.1177/00131640121971392)
- Smithson, M. (2003). *Confidence intervals*. Thousand Oaks, CA: Sage.
- Smyth, J. M. (1998). Written emotional expression: Effect sizes, outcome types, and moderating variables. *Journal of Consulting and Clinical Psychology*, 66(1), 174–184. doi:[10.1037/0022-006X.66.1.174](https://doi.org/10.1037/0022-006X.66.1.174)
- Smyth, J. M., Stone, A. A., Hurewitz, A., & Kaell, A. (1999). Effects of writing about stressful experiences on symptom reduction in patients with asthma or rheumatoid arthritis: A randomized trial. *JAMA: The Journal of the American Medical Association*, 281(14), 1304–1309. doi:[10.1001/jama.281.14.1304](https://doi.org/10.1001/jama.281.14.1304)
- Stanley, T. D. (2005). Beyond publication bias. *Journal of Economic Surveys*, 19(3), 309–345. doi:[10.1111/j.0950-0804.2005.00250.x](https://doi.org/10.1111/j.0950-0804.2005.00250.x)
- Stanley, T. D., & Doucouliagos, H. (2014). Meta-regression approximations to reduce publication selection bias. *Research Synthesis Methods*, 5(1), 60–78. doi:[10.1002/jrsm.1095](https://doi.org/10.1002/jrsm.1095)
- Stanton, A. L., Danoff-Burg, S., Sworowski, L. A., Collins, C. A., Branstetter, A. D., Rodriguez-Hanley, A., . . . Austenfeld, J. L. (2002). Randomized, controlled trial of written emotional expression and benefit finding in breast cancer patients. *Journal of Clinical Oncology*, 20(20), 4160–4168. doi:[10.1200/JCO.2002.08.521](https://doi.org/10.1200/JCO.2002.08.521)
- Taku, K., Calhoun, L. G., Cann, A., & Tedeschi, R. G. (2008). The role of rumination in the

- coexistence of distress and posttraumatic growth among bereaved Japanese
University students. *Death Studies*, 32(5), 428–444. doi:[10.1080/07481180801974745](https://doi.org/10.1080/07481180801974745)
- Tedeschi, R. G., & Calhoun, L. G. (1995). *Trauma & transformation: Growing in the aftermath of suffering*. Thousand Oaks, CA: Sage Publications.
- Tedeschi, R. G., & Calhoun, L. G. (2004). Posttraumatic growth: Conceptual foundations and empirical evidence. *Psychological Inquiry*, 15(1), 1–18.
doi:[10.1207/s15327965pli1501](https://doi.org/10.1207/s15327965pli1501)
- Terrin, N., Schmid, C. H., Lau, J., & Olkin, I. (2003). Adjusting for publication bias in the presence of heterogeneity. *Statistics in Medicine*, 22(13), 2113–2126.
doi:[10.1002/sim.1461](https://doi.org/10.1002/sim.1461)
- van Aert, R. C. M. (2017). p-uniform. GitHub. Retrieved from
<https://github.com/RobbievanAert/puniform>
- van Aert, R. C. M., Wicherts, J. M., & van Assen, M. A. L. M. (2016). Conducting meta-analyses based on p-values: Reservations and recommendations for applying p-uniform and p-curve. *Perspectives on Psychological Science*, 11(5), 713–729.
doi:[10.1017/CBO9781107415324.004](https://doi.org/10.1017/CBO9781107415324.004)
- van Assen, M. A. L. M., van Aert, R. C. M., & Wicherts, J. M. (2015). Meta-analysis using effect size distributions of only statistically significant studies. *Psychological Methods*, 20(3), 293–309. doi:<http://dx.doi.org/10.1037/met0000025>
- van Elk, M., Matzke, D., Gronau, Q. F., Guan, M., Vandekerckhove, J., & Wagenmakers, E.-J. (2015). Meta-analyses are no substitute for registered replications: A skeptical perspective on religious priming. *Frontiers in Psychology*, 6, 1365.
doi:[10.3389/fpsyg.2015.01365](https://doi.org/10.3389/fpsyg.2015.01365)
- Vevea, J. L., & Hedges, L. V. (1995). A general linear model for estimating effect size in the presence of publication bias. *Psychometrika*, 60(3), 419–435. doi:[10.1007/BF02294384](https://doi.org/10.1007/BF02294384)
- Vevea, J. L., & Woods, C. M. (2005). Publication bias in research synthesis: Sensitivity analysis using a priori weight functions. *Psychological Methods*, 10(4), 428–443.

doi:[10.1037/1082-989X.10.4.428](https://doi.org/10.1037/1082-989X.10.4.428)

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36(3), 1–48. doi:[10.18637/jss.v036.i03](https://doi.org/10.18637/jss.v036.i03)

Walker, B. L., Nail, L. M., & Croyle, R. T. (1999). Does emotional expression make a difference in reactions to breast cancer? *Oncology Nursing Forum*, 26(6), 1025–1032.

Wang, X., Gao, L., Shinfuku, N., Zhang, H., Zhao, C., & Shen, Y. (2000). Longitudinal study of earthquake-related PTSD in a randomly selected community sample in North China. *American Journal of Psychiatry*, 157(8), 1260–1266.

doi:[10.1176/appi.ajp.157.8.1260](https://doi.org/10.1176/appi.ajp.157.8.1260)

Weiss, T. (2002). Posttraumatic growth in women with breast cancer and their husbands – An intersubjective validation study. *Journal of Psychosocial Oncology*, 20(2), 65–80.

doi:[10.1300/J077v20n02_04](https://doi.org/10.1300/J077v20n02_04)

Wilson, K. G., & DuFrene, T. (2009). *Mindfulness for two: An acceptance and commitment Therapy approach to mindfulness in psychotherapy*. Oakland, CA: New Harbinger Publications.

Yilmaz, M., & Zara, A. (2016). Traumatic loss and posttraumatic growth: The effect of traumatic loss related factors on posttraumatic growth. *Anatolian Journal of Psychiatry*, 17(1), 5–11. doi:[10.5455/apd.188311](https://doi.org/10.5455/apd.188311)

Table 1

Effect Size Estimates for PTS Results

Model	Fixed Effects	Random Effects	Fixed No Outliers	Random No Outliers
Overall Effects	0.34 [0.31, 0.37]	0.39 [0.32, 0.46]	0.32 [0.29, 0.35]	0.36 [0.29, 0.42]
\$Z\$ Values	21.75, \$p\$ < .001	11.06, \$p\$ < .001	20.00, \$p\$ < .001	11.03, \$p\$ < .001
\$p\$-Uniform	0.60 [0.50, 0.71]	-	0.57 [0.47, 0.67]	-
PET	0.12 [0.03, 0.21]	-	0.11 [0.02, 0.20]	-
PEESE	0.25 [0.20, 0.30]	-	0.23 [0.18, 0.28]	-
Selection Models	0.33 [0.28, 0.37]	0.45 [0.33, 0.57]	0.29 [0.24, 0.33]	0.39 [0.27, 0.50]
Trim and Fill	0.26 [0.23, 0.29]	0.26 [0.18, 0.34]	0.25 [0.22, 0.28]	0.25 [0.18, 0.32]

Note. [] indicates the 95 percent confidence interval for each effect size estimate.

Table 2

Effect Size Estimates for PTG Results

Model	Fixed Effects	Random Effects
Overall Effects	0.10 [0.02, 0.17]	0.10 [0.02, 0.17]
\$Z\$ Values	2.45, \$p\$ = .014	2.45, \$p\$ = .014
\$p\$-Uniform	-0.11 [-1.43, 0.42]	-
PET	0.06 [-0.20, 0.32]	-
PEESE	0.08 [-0.04, 0.20]	-
Selection Models	0.09 [-0.01, 0.18]	0.09 [-0.03, 0.20]
Trim and Fill	0.10 [0.02, 0.17]	0.10 [0.02, 0.17]

Note. [] indicates the 95 percent confidence interval for each effect size estimate.

Table 3

Effect Size Estimates for QOL Results

Model	Fixed Effects	Random Effects	Fixed No Outliers	Random No Outliers
Overall Effects	-0.01 [-0.07, 0.05]	-0.01 [-0.16, 0.13]	-0.01 [-0.07, 0.05]	-0.01 [-0.11, 0.09]
\$Z\$ Values	-0.33, \$p\$ = .745	-0.18, \$p\$ = .860	-0.25, \$p\$ = .805	-0.20, \$p\$ = .838
\$p\$-Uniform	0.79 [0.33, 1.61]	-	0.62 [0.10, 0.96]	-
PET	0.05 [-0.26, 0.36]	-	0.05 [-0.29, 0.38]	-
PEESE	0.00 [-0.17, 0.17]	-	0.00 [-0.19, 0.19]	-
Selection Models	-0.06 [-0.12, 0.01]	0.51 [-0.09, 1.12]	-0.04 [-0.11, 0.03]	0.05 [-0.15, 0.24]
Trim and Fill	-0.01 [-0.07, 0.05]	-0.01 [-0.16, 0.13]	-0.01 [-0.07, 0.05]	-0.01 [-0.11, 0.09]

Note. [] indicates the 95 percent confidence interval for each effect size estimate.




Figure 1. Effect sizes and their non-centralized confidence interval for PTS total scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 2. Effect sizes and their non-centralized confidence interval for PTS Hyperarousal. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 3. Effect sizes and their non-centralized confidence interval for PTS Intrusion scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 4. Effect sizes and their non-centralized confidence interval for PTS Avoidance Scores. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 5. Effect sizes and their non-centralized confidence interval for PTG outcome variables. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.




Figure 6. Effect sizes and their non-centralized confidence interval for QOL outcome variables. Dashed lines indicated average non-weighted lower and upper confidence interval limits. Diamond size indicates normalized study weight from a random effects model.