


# Hexagulation numbers: magic numbers on the gyroid surfaces

Kana Yamamoto and Tomonari Dotera\*


Department of Physics, Kindai University, Higashiosaka, Osaka 577-8502, Japan

\*email: dotera@phys.kindai.ac.jp

Regular structures of equal spheres on the gyroid minimal surfaces have been investigated as obtained through Monte Carlo simulations of hard spheres undergoing the Alder transition<sup>1,2</sup>. Remarkably, there exist magic numbers (Figure 1) producing regular structures, which are simply explained in terms of *hexagulation numbers* defined as  $H = h^2 + k^2 - hk$ , in analogy with the Caspar and Klug's *triangulation numbers*  $T = h^2 + k^2 + hk$  for icosahedral viruses, where  $h$  and  $k$  are nonnegative integers<sup>3</sup>. The total number of spheres per cubic unit cell  $N$  is represented by  $N = 16H$ . Here we extend our simulations up to  $H = 21$ . When  $H$  is a multiple of three, we find that the space group  $Ia\bar{3}d$  is broken into  $I\bar{4}3d$ , which is chiral with right- and left-handed version. These arrangements are analyzed in terms of the space groups, equivalent positions (Wyckoff positions), and polygonal-tiling representations (Figure 2). We also present bilayer regular structures composed of both right- and left-handed chiral layers having multiple of three  $H$  numbers.



**Figure 1** (a) Caspar and Klug's  $T$ -diagram for counting the triangulation number  $T$  and (b)  $H$ -diagram for counting the hexagulation number  $H$ . Lattice points denoted as  $(h, k)$  with  $h, k$  nonnegative integers in two oblique coordinate systems indicate circled numbers  $T = h^2 + k^2 + hk$  and  $H = h^2 + k^2 - hk$ :  $T$  is half the number of lattice points inside a triangle (solid line), while  $H$  is the number of dashed hexagons inside a hexagon (solid line). Shown here are  $T=13$  with  $(3,1)$  and  $H=7$  with  $(3,1)$ .



**Figure 2** Top: Regular structures of hard spheres on the gyroid surface ( $Ia\bar{3}d$ ). Bottom: their polygonal-tiling representations.  $H$  is the Hexagulation number and  $N$  is the number of spheres in a cubic unit cell. Colors indicate Wyckoff positions. Structures for  $H = 12$  and  $H = 21$  are chiral ( $I\bar{4}3d$ ) having right- and left-handed version, both of which have been obtained by simulations.

The key is that only a limited number of efficient physical designs are possible even on negatively curved triply periodic minimal surfaces<sup>4</sup> like in icosahedral viruses. Future applications of such regular arrangements are to construct complex assemblies using the concepts of bijels, coloidosomes, polymersomes, and DNA-origami self-assemblies<sup>5</sup>.

[1] T. Dotera, M. Kimoto, J. Matsuzawa, *Interface Focus* **2**, 575 (2012).

[2] T. Dotera, H. Tanaka, Y. Takahashi, *Struct. Chem.* **28**, 105 (2017).

[3] D. L. Caspar, A. Klug, *Cold Spring Harbor Symp. Quant. Biol.* **27**, 1 (1962).

[4] M. C. Pedersen, S. T. Hyde, S. Ramsden, J. J. K. Kirkensgaard, *Soft Matter* **19**, 1586 (2023).

[5] C. Duque, D. Hall, B. Tyukodi, M. Hagan, C. Santangelo, G. Grason, *PNAS* **121**, e231548212 (2024).