
a Morphological ordination of S. lautus ecotypes
PC2 (25.6% variance)
PC1 (36.5% variance)
Index width
N indents
Circularity
Dune (N)
Headland (N)
Headland (S)
Alpine (E)
Alpine (P)
Brigalow
Lacustrinus
Depressicola
Lanceolatus
Mallee
Sea Bird Colony
Tableland
Woodland
b Phenotypic divergence (proxy: θ12)
Phenotypic divergence Δμ (standardised Euclidean)
Eastern clade
Southeastern clade
P1 DN0 HN0
P2 DN1 HN1
P3 DN3 HN2
P4 DN4 HN5
P5 DN12 HN14
P6 DN23 HS21
P7 DN32 HN12
P8 DN14 HN15
c Environmental divergence (proxy: ΔH)
Environmental divergence ΔE (standardised Euclidean)
d Environmental vs. phenotypic divergence per pair
(prediction: ΔE ∝ Δμ ∝ θ12 ⇒ Igeom ≈ k θ12²)
Phenotypic divergence Δμ (proxy for θ12)
Environmental divergence ΔE (proxy for ΔH)
Pair 4 (DN4/HN5)
Pair 7 (DN32/HN12)
Pair 1 (DN0/HN0)
Pair 5 (DN12/HN14)
Pair 3 (DN3/HN2)
Pair 2 (DN1/HN1)
Pair 8 (DN14/HN15)
r = 0.01, p = 0.983 (n = 8 pairs)
Large ΔE, large Δμ
→ predicted high Igeom