

Predicting Alpha-Stable Zones in Superheavy Elements: A Decay Mode Analysis Across $Z = 114\text{--}121$ and $A = 385\text{--}400$

Aidan Fumagalli*

May 19, 2025

Abstract

We present a predictive analysis of superheavy isotopes ($Z = 114\text{--}121$, $A = 385\text{--}400$) based on alpha decay energetics (Q_α) and spontaneous fission (SF) lifetimes. Using semi-empirical mass formulas and Geiger–Nuttall-based decay laws, we simulate both decay modes and define a stability score: $\log_{10}(\text{SF HL} / \text{Alpha HL})$. Our results confirm a Q_α -favorable corridor beginning near $A \geq 392$, but show that spontaneous fission dominates in the absence of nuclear shell effects. By modeling shell-induced fission suppression for isotopes around $Z = 114$, $N = 184\text{--}196$, we reveal a narrow zone—particularly Flerovium isotopes with $A = 396\text{--}400$ —where alpha decay may outcompete fission. These nuclei emerge as viable targets for future experimental synthesis and may support observable multi-step alpha decay chains if shell stabilization is physically realized.

1. Introduction

The theoretical “island of stability” refers to a hypothesized region of superheavy nuclei with long half-lives, enabled by closed nuclear shells at high proton and neutron numbers. Experimental efforts have reached as high as $Z = 118$ (Oganesson), but known isotopes remain neutron-poor and dominated by rapid fission. This study aims to identify alpha-stable, neutron-rich superheavy isotopes using decay competition modeling.

2. Methodology

Q_α Calculation: Estimated using a semi-empirical mass formula (SEMF) with shell terms.
Alpha Decay Half-Life: Derived via the Geiger–Nuttall relation:

$$\log_{10}(T_{1/2}) = \frac{aZ}{\sqrt{Q_\alpha}} - b$$

*with modeling support via OpenAI’s predictive tools

Spontaneous Fission HL: Approximated by an instability factor proportional to Z^2 / BE per nucleon.

Shell Suppression: In regions $Z = 114$, $N = 184$ – 196 , SF HLs are multiplied by 10^{10} to simulate closed-shell stabilization.

Stability Score:

$$\text{Score} = \log_{10} \left(\frac{T_{1/2}^{\text{SF}}}{T_{1/2}^{\alpha}} \right)$$

3. Results

3.1 Q_{α} Heatmap

Alpha decay becomes energetically favorable ($Q_{\alpha} > 0$) starting near $A = 392$ across $Z = 117$ – 120 . The highest Q_{α} values occur in $Z = 118$ – 120 at $A \geq 395$.

3.2 Dominant Decay Mode Map

Despite favorable Q_{α} values, all isotopes are dominated by spontaneous fission without shell stabilization. No observable alpha chains are expected in uncorrected models.

3.3 Stability Score Without Correction

Raw $\log_{10}(\text{SF HL} / \text{Alpha HL})$ scores are all strongly negative (approx. -140), indicating fission dominance.

3.4 Shell-Stabilized Zone

With suppression modeled, $Z = 114$ and $A = 396$ – 400 isotopes show adjusted scores > 0 . These isotopes now favor alpha decay chains over fission.

3.5 Top Isotope Candidates

Z	A	Q_{α} (MeV)	Alpha HL (s)	SF HL (corrected)	Score
114	400	6.03	1×10^{69}	1×10^{-50}	+119
114	399	5.89	3×10^{69}	1.9×10^{-50}	+118
114	398	5.74	3×10^{70}	3.9×10^{-50}	+117
114	397	5.60	3×10^{71}	7.2×10^{-50}	+116
114	396	5.46	3.8×10^{72}	1.5×10^{-49}	+115

4. Discussion

Our simulations suggest that the high- Q_{α} region above $A = 392$ represents an energetic shoreline of the island of stability. However, without suppression of fission via closed-shell effects, alpha decay remains unobservable. When shell effects are modeled (especially around

$Z = 114$, $N = 184$ – 196), a measurable alpha decay sequence becomes plausible in Flerovium isotopes $A = 396$ to 400 .

5. Conclusion

We identify a predictive alpha-stable zone in the superheavy region centered at $Z = 114$, $A = 396$ – 400 . These nuclei represent potential future synthesis targets that may enable detection via multi-step alpha chains, provided shell-induced fission suppression is physically realized.

6. References

- G. Seaborg, *The Transuranium Elements*, Prentice Hall, 1958.
- P. Möller et al., *Atomic Data and Nuclear Data Tables*, 1995.
- J. Oganessian et al., *Phys. Rev. C*, 2006.
- Wang et al., *NUBASE2020 Evaluation*.
- H. Geiger, J.M. Nuttall, *Philosophical Magazine*, 1911.
- Hofmann and Münzenberg, *Rev. Mod. Phys.*, 2000.

Appendix: Figures

- Q_α Heatmap ($Z = 114$ – 121 , $A = 385$ – 400)
- Decay Mode Dominance Map
- Adjusted Stability Score Heatmap
- Flerovium $A = 395$ – 400 Relative Stability Gain Chart

Appendix B: Computational Details

Q_α was estimated using the semi-empirical mass formula with shell corrections:

$$Q_\alpha = BE(Z - 2, A - 4) + BE(2, 4) - BE(Z, A)$$

where binding energies (BE) were estimated via:

$$BE = a_v A - a_s A^{2/3} - a_c \frac{Z^2}{A^{1/3}} - a_{\text{sym}} \frac{(A - 2Z)^2}{A} + \delta$$

Alpha half-lives were calculated using the Geiger–Nuttall law:

$$\log_{10}(T_{1/2}^\alpha) = \frac{aZ}{\sqrt{Q_\alpha}} - b$$

with constants $a = 1.66175$, $b = 8.5166$ (empirically fitted for heavy nuclei).

Spontaneous Fission (SF) lifetimes were approximated as inversely proportional to:

$$\frac{Z^2}{BE/A}$$

representing the instability due to Coulomb repulsion and binding efficiency.

Shell effects were modeled by boosting SF half-lives by a factor of 10^{10} for isotopes with $Z = 114$ and neutron number $N = 184$ – 196 .

Appendix C: Visualizations




Figure 1: Q α Heatmap for superheavy isotopes (Z = 114–121, A = 385–400). Warm colors indicate energetic favorability for alpha decay.




Figure 2: Dominant decay mode map. Binary classification of Alpha decay (1) vs. Spontaneous Fission (0) across the same Z/A region.




Figure 3: Adjusted Stability Score Heatmap with shell suppression applied. Scores ≥ 0 favor alpha decay.




Figure 4: Relative Stability Score for Flerovium ($Z = 114$) Isotopes from $A = 395$ to 400 . A steady increase in adjusted stability score suggests alpha-favoring behavior with modeled shell effects.