

GENE638 - Homework 2

Daniel Osorio - dcosorih@tamu.edu

Department of Veterinary Integrative Biosciences

Texas A&M University

1. For this system of equations:

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 5 \\ 20 \\ 25 \end{bmatrix}$$

- (a) Determine the rank of the coefficient matrix

```
> as.numeric(Matrix::rankMatrix(coefficientMatrix))  
[1] 2
```

- (b) Express any linearly dependent columns in the matrix form $\begin{bmatrix} A_{12} \\ A_{22} \end{bmatrix} = \begin{bmatrix} A_{11} \\ A'_{12} \end{bmatrix} L$

- (c) Find a generalized inverse of the coefficient matrix. Prove that it is a generalized inverse

```
> gInverse <- MASS::ginv(coefficientMatrix)  
> gInverse  
      [,1]      [,2]      [,3]  
[1,] 0.5000000 -3.333333e-01 1.666667e-01  
[2,] -0.3333333  3.333333e-01 1.942890e-16  
[3,] 0.1666667  6.938894e-17 1.666667e-01  
> round(gInverse %*% coefficientMatrix)  
      [,1] [,2] [,3]  
[1,]    1    0    0  
[2,]    0    1    0  
[3,]    0    0    1  
> round(coefficientMatrix %*% gInverse %*% coefficientMatrix)  
      [,1] [,2] [,3]  
[1,]    1    0    1  
[2,]    0    2    2  
[3,]    1    2    3
```

- (d) Using the inverse from part (c), solve for x . Prove your solution satisfies the equations

```
> x <- gInverse %*% c(5, 20, 25)  
> round(x)  
      [,1]  
[1,]    0  
[2,]    5  
[3,]    5  
> coefficientMatrix %*% x
```

```

      [,1]
[1,]    5
[2,]   20
[3,]   25

```

(e) What do your solutions estimate?

(f) Based in what you did in part (e): Can you estimate x_1 ? Can you estimate $x_1 - x_2$?

2. Using the partitioned matrix inverse procedure on page 14 in the notes

(a) Find the inverse of: $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 3 \end{bmatrix}$

(b) Prove that your answer in (a) is an inverse

(c) Solve $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$

(d) What is the solution in (c) an estimate of?

(e) Can you estimate x_3 ? With what? Can you estimate $6x_1 + 4x_2 - 8x_3$? With what?

3. Assuming the linear model $y_i = \mu + \epsilon_i$ where y_i is and observed first lactation milkfat production, μ is the population mean milkfat production, and ϵ_i is the deviation of an individual cow's production from the mean. Write this model as

$$\underline{y} = 1\mu + \underline{\epsilon}$$

Cow	1 st lactation milkfat (lb)
1	300
2	290
3	405
4	360
5	315

(a) Calculate:

i. $X'X$

```

> X <- rep(1,length(y))
> t(X) %*% X

```

```

      [,1]
[1,]    5

```

ii. $(X'X)^{-1}$

```

> solve(t(X) %*% X)

```

```

      [,1]
[1,]  0.2

```

iii. $X'y$

```

> t(X) %*% y

```

```

      [,1]
[1,] 1670

```

(b) Solve $\hat{\mu} = (X'X)^{-1}X'y$

```

> muHat <- as.numeric(solve(t(X) %*% X) %*% X %*% y)
> muHat

```

```
[1] 334
```

(c) Show that $\hat{\mu} = \bar{y}$

```
> meanY <- mean(y)
> meanY
```

```
[1] 334
```

```
> all.equal(muHat, meanY)
```

```
[1] TRUE
```

(d) Find the predicted deviations $\hat{\epsilon} = \underline{y} - \underline{1}\hat{\mu}$

```
> epsilonHat <- (y - muHat)
> epsilonHat
```

```
[1] -34 -44 71 26 -19
```

(e) Find $\frac{\hat{\epsilon}'\hat{\epsilon}}{n-\text{rank}(X)}$ and show that this is s^2 the sample variance.

```
> s2Hat <- as.numeric((t(epsilonHat) %*% epsilonHat) / (length(y) - 1))
> s2Hat
```

```
[1] 2292.5
```

```
> var(y)
```

```
[1] 2292.5
```

```
> all.equal(s2Hat, var(y))
```

```
[1] TRUE
```