

1 Neuroprotective effects of tibolone during astrocytic metabolic inflammation: a network based approach

Abstract:

1.1. Introduction

Astrocyte-Neuron Metabolic Relationships

Astrocytes are the most abundant cells in the human brain and play important roles in the central nervous system (CNS) [1]. They are highly associated to several homeostatic functions such as glutamate, ion, and water homeostasis, energy storage in the form of glycogen, synapse formation and remodeling, defense against oxidative stress, scar formation, tissue repair and modulation of synaptic activity via the release of gliotransmitters [2]. Astrocytes metabolize glucose in anaerobic way to produce lactate, which is released to neurons through monocarboxylate transporters [3]. Lactate is used in neurons as an energy substrate after its conversion to pyruvate and subsequently to ATP via oxidative phosphorylation [4]. Astrocytes play an important role in glutamate mediated synaptic activity [5]; according to the astrocyte–neuron lactate shuttle model, astrocytes respond to glutamate induced activation by increasing their rate of glucose uptake and the release of lactate into the extracellular space, increasing the lactate available to be used by neurons to supply their energetic needs [6]. Glutamate is uptaken by astrocytes through the glutamate aspartate transporter and glial glutamate transporter-1, inducing events that involves the activation of $\text{Na}^+\text{--K}^+\text{--ATPase}$ and maintaining extracellular glutamate at homeostatic levels [7]. Part of incorporated glutamate is converted to glutamine through glutamine synthetase, which is only associated to glial cells and released to neurons using electroneutral systems-N transporters coupled to Na^+ and H^+ [8]. In neurons glutaminase enzyme converts glutamine back into glutamate which can be used again for neurotransmission or metabolized into the neuronal Krebs cycle [9]. Astrocytes release many other substances related to synaptic transmission [10]. However D-serine, a neurotransmitter that act as a coagonist with glutamate at NMDA receptors is

one of the most important [5]. Due only glial cells can synthesize serine, all available D-serine at synapsis is associated to be primarily produced and secreted by astrocytes [8]. D-serine is synthesized in astrocytes by serine racemase from L-serine [11]. Additionally to these energy and synaptic support associated functions, astrocytes also play an important role in the reduced glutathione (GSH) metabolism of the brain [12]. GSH is the major cellular antioxidant and plays an important neuroprotective role [13]. Cellular GSH levels are closely correlated with cell survival under adverse conditions [14]. GSH is synthesized from glutamate, cysteine, and glycine and release directly from Astrocytes through GSH transporters ion-independent, net transport is concentration-gradient dependent [15].

This strong metabolic cooperation between astrocytes and neurons allows to predict that even an small astrocytic dysfunction might cause and/or contribute neurodegenerative processes.

Inflammation

Inflammation is a complex biological response to injuries, metabolic disorders or infections and its dysregulation induce many complex diseases [13, 16, 17]. In brain, inflammatory response acts as a defense mechanism against any threat to homeostatic state inducing changes in glucose metabolism and release of proinflammatory factors [14]. Inflammation responses in CNS are mediated by glial cells that acquire reactive phenotypes to participate in repair mechanisms [1, 13, 18]. Astrocytes, as glial cells are highly sensitive cells to inflammatory mediators, they respond to inflammation through a complex reaction named astrogliosis [19]. During astrogliosis, glial cells generally associated to several beneficial activities in the CNS, also act as a source of inflammatory mediators and as generators of reactive oxidant species (ROS) that have the potential to damage neurons [20]. Astrogliosis is characterized by a low regulation of mitochondrial dynamics that result in mitochondrial failure [21]. Mitochondrial failure induces the deregulation of Ca^{2+} homeostasis and increased ROS generation, both of which are linked to neurotoxicity [2]. At metabolic level, inflammatory process has been associated to an increase of free saturated fatty acid in comparison with healthy conditions in some brain tissues [22]. The increase of free saturated fatty acid induce metabolic inflammation, a response associated with the induction of diverse intracellular stresses, such as mitochondrial oxidative stress, ER stress, and autophagy defects [13]. Lipid excess in metabolic inflammation activates hypothalamic $\text{IKK}\beta$ and $\text{NF-}\kappa\beta$ signaling pathways, which ultimately impairs hypothalamic leptin and insulin signaling and further triggers the synthesis and release of increased amounts of ROS and proinflammatory cytokines ($\text{TNF-}\alpha$ and IL-6) from glial cells to sustain the neuroinflammatory state [23]. Enhanced ROS generation by reactive glial cells trigger mitochondria dysfunction in neuron, which induces neuronal apoptosis, the prerequisite for a diverse number of neurodegenerative conditions [24].

SysBio Inflammation

Tibolone

1.2. Material and Methods

Tissue Specific Model Construction

The tissue specific model construction process started with the identification of all enzyme-coding genes expressed over the mean in at least 50 % of samples for healthy human astrocytes indexed in the GEO database [25] as GSE73721 [26]. Gene identifiers conversion from GeneCards[27] to ENTREZ [28] was performed through ‘UniProt.ws’ R Package [29]. Reactions associated with the identified genes were mapped from the Human Genome Scale Metabolic Reconstruction RECON 2.04 downloaded from the VMH Lab (<https://vmh.uni.lu>) [30]. The R package ‘g2f’ [31] was used to identify and fill the gaps using all no gene associated reactions included in RECON 2.04, as well as to identify and remove all blocked reactions from the reconstruction. All reactions involved in the conversion of extracellular glutamate, glycine, cysteine and glucose to extracellular glutamine, glycine, serine-D, reduced glutathione, lactate and ATP respectively were added. Exchange reactions were limited to components of the Dulbecco’s Modified Eagle Medium (DMEM) as input and gliotransmitters (glutamine, D-serine, ATP, glutamate), reduced glutathione, lactate, glucose, nitric oxide, prostaglandins and leukotrienes as output. Finally, syntax, mass-charge validation and creation of SBML files were carried out through the ‘minval’ R Package [32]. Reaction limits (upper and lower bounds) were constrained proportional to the mean gene expression reported for genes included in Gene-Protein-Reaction (GPR) [33] associated to each reaction in samples of 47 to 63 years old using ‘exp2flux’ R package [34]. All analysis were done by the ‘sybil’ [35] R Package running under R 3.3.1 [36].

Flux Balance Analysis

Flux Balance Analysis (FBA) is a linear optimization method for simulating metabolism that allows to identify the set of reactions involved in the production of a biological response within a metabolic model [37]. The metabolic reactions are represented internally as a stoichiometric matrix (S), of size $m * n$, where m represents the compounds and n the reactions; the entries in the matrix are the stoichiometric coefficients of the metabolites participating in a reaction [38]. The flux through all of the reactions in a network is represented by the vector v , which has a length of n . The concentrations of all metabolites are represented by the vector x , with length m . The systems of mass balance equations at steady state, $\frac{d_x}{d_t} = 0$ or $S * v = 0$. FBA seeks to maximize or minimize an objective function which can be any linear combination fluxes, to obtain a flux for each reaction, indicating how much each reaction contributes to the objective function [37]. FBA for healthy, inflamed and medicated scenarios was

resolved using GLPK 4.60, setting the generic human biomass reaction included in RECON 2.04 and each one of reactions described in table **1-1** as objective functions. Models were analyzed by comparing fluxes between scenarios, metabolites production rate and sensitivity analysis.

Table 1-1: Main metabolic capabilities associated to astrocytes represented as the set of objective functions used to evaluate neuroprotective effects of Tibolone under inflamed scenarios

ID	FORMULA REACTION	DESCRIPTION
Glu2Gln	$1 \text{ glu_L[e]} \Rightarrow 1 \text{ gln_L[e]}$	Glutamate - Glutamine Cycle
Gly2SerD	$1 \text{ gly[e]} \Rightarrow 1 \text{ ser_D[e]}$	Glycine to D-serine conversion
Glc2Lac	$1 \text{ glc_D[e]} \Rightarrow 2 \text{ lac_L[e]}$	Lactate production from Glucose
Glc2ATP	$1 \text{ glc_D[e]} \Rightarrow 36 \text{ atp[e]}$	ATP production from Glucose
Cys2GTHRD	$1 \text{ cys_L[e]} + 1 \text{ glu_L[c]} + 1 \text{ gly[c]} \Rightarrow 1 \text{ gthrd[e]}$	Catch of Cysteine to produce reduced Glutathione

Metabolic Scenarios

To test neuroprotective effects of tibolone during astrocytic metabolic inflammation we define three different metabolic scenarios. A ‘healthy’ scenario, where palmitate uptake rate was freely set by optimizer; an ‘inflamed’ scenario, where uptake rate of palmitate was forced to be stable in the mean of the half maximal inhibitory concentration (IC50) value for all objective functions included in table **1-1**. IC50 values were calculated through a robustness analysis performed using uptake of palmitate (‘EX_hdca(e)’ in RECON 2.04) as control reaction and a 1000 points in the range from 0 to 1 mMgDW⁻¹h⁻¹ for each objective function. Uptake value where each objective function reached IC50 was selected and subsequently averaged. Finally, a medicated scenario, defined as an inflamed scenario that include 279 reactions associated with tibolone and estradiol-derived compounds metabolism. Ten specific reactions described in table **1-2** associated to specific Tibolone action mechanism non included in RECON 2.04 were added to medicated scenario.

Metabolic Changes

Metabolic changes across metabolic scenarios were measured through two different approximations. Flux differences for each reaction between optimized scenarios were measured using

Table 1-2: Set of reactions associated to tibolone specific action mechanism in brain reported by Kloosterboer, H. J. (2004) added to medicated scenario model.

ID	FORMULA REACTION	DESCRIPTION
T1	tibolone[e] \Leftrightarrow	Tibolone exchange reaction
T2	tibolone[e] \Leftrightarrow a3OHTibolone[e]	3 α hydroxytibolone interconversion
T3	tibolone[e] \Leftrightarrow b3OHTibolone[e]	3 β hydroxytibolone interconversion
T4	tibolone[e] \Rightarrow d4tibolone[e]	Tibolone Δ^4 isomer formation
T5	b3OHTibolone[e] \Rightarrow d4tibolone[e]	Tibolone Δ^4 isomer formation from 3 β -hydroxytibolone
T6	a3OHTibolone[e] \Rightarrow estradiol[c]	Estradiol receptor agonist action mechanism of 3 α -hydroxytibolone
T7	b3OHTibolone[e] \Rightarrow estradiol[c]	Estradiol receptor agonist action mechanism of 3 β -hydroxytibolone
T8	d4tibolone[e] \Rightarrow prgstrn[c] + tststerone[c]	Progesterone and androgen receptor activation by tibolone Δ^4 isomer
T9	a3OHTibolone[e] \Leftrightarrow a3SOTibolone[e]	3 α hydroxytibolone interconversion to sulfated inactive compounds
T10	a3SOTibolone[e] \Rightarrow	Tibolone inactive form in blood

the fold change calculated as described in equation 1-1.

$$foldChange = \frac{valueModel2 - valueModel1}{|valueModel1|} \quad (1-1)$$

Additionally, to obtain a full perspective about inflammation effects in metabolites production, the production of each metabolite was set as objective function in each metabolic scenario and differences were evaluated as well as flux differences.

Proinflammatory, Antiinflammatory and Tibolone Action Mechanism Associated Enzymes

Identification of enzymes involved in proinflammatory and antiinflammatory responses as well as in the tibolone action mechanism were identified through several sensitivity analysis as follows: Proinflammatory enzymes, are those that catalyze reactions that being knocked out allows an increase of objective function value. Antiinflammatory enzymes, are those associated to reactions that being knocked out reduce even more the objective function value. Tibolone action mechanism associated enzymes are those that catalyze reactions that being knocked out inhibit entirely the metabolic effect of tibolone.

1.3. Results

1.4. Conclusion

Bibliography

- [1] Kazuhiro Takuma, Akemichi Baba, and Toshio Matsuda. Astrocyte apoptosis: Implications for neuroprotection. *Progress in Neurobiology*, 72:111–127, 2004.
- [2] Sofie C. Lange, Lasse K. Bak, Helle S. Waagepetersen, Arne Schousboe, and Michael D. Norenberg. Primary cultures of astrocytes: Their value in understanding astrocytes in health and disease. *Neurochemical Research*, 37:2569–2588, 2012.
- [3] Harold K Kimelberg. Functions of Mature Mammalian Astrocytes: A Current View. *The Neuroscientist*, 16(1):79–106, feb 2010.
- [4] Nicola J Allen and Ben Barres. Neuroscience: Glia - more than just brain glue. *Nature*, 457(7230):675–677, 2009.
- [5] Michael M Halassa and Philip G Haydon. Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annual review of physiology*, 72(2):335–355, 2010.
- [6] Christian Giaume, Annette Koulakoff, Lisa Roux, David Holcman, and Nathalie Rouach. Astroglial networks: a step further in neuroglial and gliovascular interactions. *Nature reviews. Neuroscience*, 11(fEbRuARy):87–99, 2010.
- [7] Cora H. Nijboer, Cobi J. Heijnen, Vincent Degos, Hanneke L M Willemsen, Pierre Gresens, and Annemieke Kavelaars. Astrocyte GRK2 as a novel regulator of glutamate transport and brain damage. *Neurobiology of Disease*, 54:206–215, 2013.
- [8] Ben Barres. The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron*, 60(3):430–40, nov 2008.
- [9] Jun Shen. Modeling the glutamate-glutamine neurotransmitter cycle. *Frontiers in Neuroenergetics*, 5(JAN):1–13, 2013.
- [10] Francesco Petrelli and Paola Bezzi. Novel insights into gliotransmitters. *Current Opinion in Pharmacology*, 26(Table 1):138–145, 2016.
- [11] Andrea R Durrant and Uriel Heresco-levy. D-Serine in Neuropsychiatric Disorders : New Advances. 2014, 2014.

-
- [12] Stephen P. Raps, James C K Lai, Leif Hertz, and Arthur J L Cooper. Glutathione is present in high concentrations in cultured astrocytes but not in cultured neurons. *Brain Research*, 493(2):398–401, 1989.
 - [13] Mithilesh Kumar Jha, Dong Ho Park, Hyun Kook, In-Kyu Lee, Won-Ha Lee, and Kyoungso Suk. Metabolic Control of Glia-Mediated Neuroinflammation. *Current Alzheimer research*, 13(4):387–402, 2016.
 - [14] Igor Allaman, Mireille Bélanger, and Pierre J. Magistretti. Astrocyte–neuron metabolic relationships: for better and for worse. *Trends in Neurosciences*, 34(2):76–87, feb 2011.
 - [15] Xue Feng Wang and Max S. Cynader. Astrocytes provide cysteine to neurons by releasing glutathione. *Journal of Neurochemistry*, 74(4):1434–1442, 2000.
 - [16] Brent E Masel and Douglas S DeWitt. Traumatic brain injury: a disease process, not an event. *Journal of neurotrauma*, 27(8):1529–1540, 2010.
 - [17] Qing Yan. *Systems biology in drug discovery and development*. 2013.
 - [18] Michael T. Fitch and Jerry Silver. CNS injury, glial scars, and inflammation: Inhibitory extracellular matrices and regeneration failure. *Experimental Neurology*, 209(2):294–301, 2008.
 - [19] James a. Dowell, Jeffrey a. Johnson, and Lingjun Li. Identification of astrocyte secreted proteins with a combination of shotgun proteomics and bioinformatics. *Journal of Proteome Research*, 8(8):4135–4143, 2009.
 - [20] Anna V. Molofsk, Robert Krenick, Erik Ullian, Hui Hsin Tsai, Benjamin Deneen, William D. Richardson, Ben. Barres, and David H. Rowitch. Astrocytes and disease: A neurodevelopmental perspective. *Genes and Development*, 26:891–907, 2012.
 - [21] Marta Sidoryk-Wegrzynowicz and Michael Aschner. Role of astrocytes in manganese mediated neurotoxicity. *BMC pharmacology & toxicology*, 14:23, 2013.
 - [22] Sunita Gupta, Alecia G. Knight, Shruti Gupta, Jeffrey N. Keller, and Annadora J. Bruce-Keller. Saturated long-chain fatty acids activate inflammatory signaling in astrocytes. *Journal of Neurochemistry*, 120(6):1060–1071, 2012.
 - [23] Sudarshana Purkayastha and Dongsheng Cai. Neuroinflammatory basis of metabolic syndrome. *Molecular Metabolism*, 2(4):356–363, 2015.
 - [24] Suk K. Proteomics-based discovery of biomarkers and therapeutic targets in neurodegenerative diseases: Perspective of microglia and neuroinflammation. *Expert Opinion on Therapeutic Patents*, 16(3):237–247, 2006.

- [25] Ron Edgar, Michael Domrachev, and Alex E Lash. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res*, 30(1):207–210, 2002.
- [26] Ye Zhang, Steven A. Sloan, Laura E. Clarke, Christine Caneda, Colton A. Plaza, Paul D. Blumenthal, Hannes Vogel, Gary K. Steinberg, Michael S B Edwards, Gordon Li, John A. Duncan, Samuel H. Cheshier, Lawrence M. Shuer, Edward F. Chang, Gerald A. Grant, Melanie G Hayden Gephart, and Ben A. Barres. Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron*, 89(1):37–53, 2016.
- [27] Michael Rebhan, Vered Chalifa-Caspi, Jaime Prilusky, and Doron Lancet. GeneCards: integrating information about genes, proteins and diseases. *Trends in Genetics*, 13(4):163, 1997.
- [28] Donna Maglott, Jim Ostell, Kim D Pruitt, and Tatiana Tatusova. Entrez Gene: gene-centered information at NCBI. *Nucleic acids research*, 33(suppl 1):D54—D58, 2005.
- [29] Marc Carlson. *UniProt.ws: R Interface to UniProt Web Services*, 2016.
- [30] Ines Thiele, Neil Swainston, Ronan M T Fleming, Andreas Hoppe, Swagatika Sahoo, Maike K Aurich, Hulda Haraldsdottir, Monica L Mo, Ottar Rolfsson, Miranda D Stobbe, and Others. A community-driven global reconstruction of human metabolism. *Nature biotechnology*, 31(5):419–425, 2013.
- [31] Kelly Botero, Daniel Osorio, Janneth Gonzalez, and Andres Pinzon. *g2f: Find and Fill Gaps in Metabolic Networks*, 2016.
- [32] Daniel Osorio, Janneth Gonzalez, and Andres Pinzon. minval: MINimal VALidation for Stoichiometric Reactions, 2016.
- [33] Ines Thiele and Bernhard Ø Palsson. A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nature Protocols*, 5(1):93–121, jan 2010.
- [34] Daniel Osorio, Kelly Botero, Janneth Gonzalez, and Andres Pinzon. *exp2flux: Convert Gene EXPression Data to FBA FLUXes*, 2016.
- [35] Gabriel Gelius-dietrich, Abdelmoneim Amer Desouki, Claus Jonathan Fritzemeier, and Martin J Lercher. sybil – Efficient constraint-based modelling in R. 2013.
- [36] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2016.
- [37] Jeffrey D Orth, Ines Thiele, and B O Palsson. What is flux balance analysis? *Nature Biotechnology*, 28(3):245–248, 2010.

- [38] Karthik Raman and Nagasuma Chandra. Flux balance analysis of biological systems: Applications and challenges. *Briefings in Bioinformatics*, 10(4):435–449, 2009.