

# 1 Neuroprotective effects of tibolone 2 during astrocytic metabolic 3 inflammation: a network based 4 approach

## 5 Abstract:

6 d

## 7 1.1. Introduction

## 8 1.2. Material and Methods

### 9 1.2.1. Tissue Specific Model Construction

10 The tissue specific model construction process started with the identification of all enzyme-  
11 coding genes expressed over the mean in at least 50 % of samples for healthy human astrocy-  
12 tes indexed in the GEO database [3] as GSE73721 [14]. Gene identifiers conversion from  
13 GeneCards[11] to ENTREZ [5] was performed through ‘UniProt.ws’ R Package [2]. Reac-  
14 tions associated with the identified genes were mapped from the Human Genome Scale Me-  
15 tabolic Reconstruction RECON 2.04 downloaded from the VMH Lab (<https://vmh.uni.lu>)  
16 [13]. The R package ‘g2f’ [1] was used to identify and fill the gaps using all no gene as-  
17 sociated reactions included in RECON 2.04, as well as to identify and remove all blocked  
18 reactions from the reconstruction. All reactions involved in the conversion of extracellular  
19 glutamate, glycine, cysteine and glucose to extracellular glutamine, glycine, serine-D, redu-  
20 ced glutathione, lactate and ATP respectively were added. Exchange reactions were limited  
21 to components of the Dulbecco’s Modified Eagle Medium (DMEM) as input and gliotrans-  
22 mitters () as output. Finally, syntax, mass-charge validation and creation of SBML files were  
23 carried out through the ‘minval’ R Package [8]. Reaction limits (upper and lower bounds)  
24 were constrained proportional to the mean gene expression reported for genes included in  
25 Gene-Protein-Reaction (GPR) [12] associated to each reaction in samples of 47 to 63 years

old using ‘exp2flux’ R package [7]. All analysis were done by the ‘sybil’ [4] R Package running  
under R 3.3.1 [9].

## 1.2.2. Flux Balance Analysis

Flux Balance Analysis (FBA) is a linear optimization method for simulating metabolism that  
allows to identify the set of reactions involved in the production of a biological response within  
a metabolic model [6]. The metabolic reactions are represented internally as a stoichiometric  
matrix ( $S$ ), of size  $m * n$ , where  $m$  represents the compounds and  $n$  the reactions; the entries  
in the matrix are the stoichiometric coefficients of the metabolites participating in a reaction  
[10]. The flux through all of the reactions in a network is represented by the vector  $v$ , which  
has a length of  $n$ . The concentrations of all metabolites are represented by the vector  $x$ , with  
length  $m$ . The systems of mass balance equations at steady state,  $\frac{dx}{dt} = 0$  or  $S * v = 0$ . FBA  
seeks to maximize or minimize an objective function which can be any linear combination  
fluxes, to obtain a flux for each reaction, indicating how much each reaction contributes  
to the objective function [6]. FBA for healthy, inflamed and medicated scenarios was  
resolved using GLPK 4.60, setting the generic human biomass reaction included in RECON  
2.04 and each one of functions described in table 1-1 as objective function. Models were  
analyzed by comparing fluxes between scenarios, metabolites production rate and sensibility  
analysis.

**Table 1-1:** Objective functions used to evaluate astrocytes metabolic capabilities

ID	FORMULA REACTION	DESCRIPTION
Glu2Gln	1 glu[e] $\rightarrow$ 1 gln[e]	C
Gly2SerD	1 gly[e] $\rightarrow$ 1 serD[e]	C
Glc2Lac	$\rightarrow$	C
Glc2ATP	$\rightarrow$	C
Cys2GTHRD	$\rightarrow$	C

44 **1.2.3. Inflamed Scenario**

45 **1.2.4. Medicated Scenario**

46 **1.3. Results**

47 **1.4. Conclusion**

48 **1.5. Bibliography**

## Bibliografía

- [1] Kelly Botero, Daniel Osorio, Janneth Gonzalez, and Andres Pinzon. *g2f: Find and Fill Gaps in Metabolic Networks*, 2016.
- [2] Marc Carlson. *UniProt.ws: R Interface to UniProt Web Services*, 2016.
- [3] Ron Edgar, Michael Domrachev, and Alex E Lash. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res*, 30(1):207–210, 2002.
- [4] Gabriel Gelius-Dietrich, Abdelmoneim Desouki, Claus Fritzemeier, and Martin J Lercher. sybil – Efficient constraint-based modelling in R. *BMC Systems Biology*, 7(1):125, 2013.
- [5] Donna Maglott, Jim Ostell, Kim D Pruitt, and Tatiana Tatusova. Entrez Gene: gene-centered information at NCBI. *Nucleic acids research*, 33(suppl 1):D54—D58, 2005.
- [6] Jeffrey D Orth, Ines Thiele, and B O Palsson. What is flux balance analysis? *Nature Biotechnology*, 28(3):245–248, 2010.
- [7] Daniel Osorio, Kelly Botero, Janneth Gonzalez, and Andres Pinzon. *exp2flux: Convert Gene EXPression Data to FBA FLUXes*, 2016.
- [8] Daniel Osorio, Janneth Gonzalez, and Andres Pinzon. minval: MINimal VALidation for Stoichiometric Reactions, 2016.
- [9] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2016.
- [10] Karthik Raman and Nagasuma Chandra. Flux balance analysis of biological systems: Applications and challenges. *Briefings in Bioinformatics*, 10(4):435–449, 2009.
- [11] Michael Rebhan, Vered Chalifa-Caspi, Jaime Prilusky, and Doron Lancet. GeneCards: integrating information about genes, proteins and diseases. *Trends in Genetics*, 13(4):163, 1997.
- [12] Ines Thiele and Bernhard Ø Palsson. A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nature Protocols*, 5(1):93–121, jan 2010.

- 
- 76 [13] Ines Thiele, Neil Swainston, Ronan M T Fleming, Andreas Hoppe, Swagatika Sahoo,  
77 Maike K Aurich, Hulda Haraldsdottir, Monica L Mo, Ottar Rolfsson, Miranda D Stobbe,  
78 and Others. A community-driven global reconstruction of human metabolism. *Nature*  
79 *biotechnology*, 31(5):419–425, 2013.
- 80 [14] Ye Zhang, Steven A. Sloan, Laura E. Clarke, Christine Caneda, Colton A. Plaza,  
81 Paul D. Blumenthal, Hannes Vogel, Gary K. Steinberg, Michael S B Edwards, Gor-  
82 don Li, John A. Duncan, Samuel H. Cheshier, Lawrence M. Shuer, Edward F. Chang,  
83 Gerald A. Grant, Melanie G Hayden Gephart, and Ben A. Barres. Purification and  
84 Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional  
85 and Functional Differences with Mouse. *Neuron*, 89(1):37–53, 2016.