




# Black hole

A **black hole** is an astronomical body so compact that its gravity prevents anything from escaping, even light. Albert Einstein's theory of general relativity predicts that a sufficiently compact mass will form a black hole.<sup>[4]</sup> The boundary of no escape is called the event horizon. In general relativity, a black hole's event horizon seals an object's fate but produces no locally detectable change when crossed.<sup>[5]</sup> In many ways, a black hole acts like an ideal black body, as it reflects no light.<sup>[6][7]</sup> Quantum field theory in curved spacetime predicts that event horizons emit Hawking radiation, with the same spectrum as a black body of a temperature inversely proportional to its mass. This temperature is of the order of billionths of a kelvin for stellar black holes, making it essentially impossible to observe directly.

Objects whose gravitational fields are too strong for light to escape were first considered in the 18th century by John Michell and Pierre-Simon Laplace. In 1916, Karl Schwarzschild found the first modern solution of general relativity that would characterise a black hole. Due to his influential research, the Schwarzschild metric is named after him. David Finkelstein, in 1958, first published the interpretation of "black hole" as a region of space from which nothing can escape. Black holes were long considered a mathematical curiosity; it was not until the 1960s that theoretical work showed they were a generic prediction of general relativity. The first black hole known was Cygnus X-1, identified by several researchers independently in 1971.<sup>[8][9]</sup>

Black holes typically form when massive stars collapse at the end of their life cycle. After a black hole has formed, it can grow by absorbing mass from its surroundings. Supermassive black holes of millions of solar masses may form by absorbing other stars and merging with other black holes, or via direct collapse of gas clouds. There is consensus that supermassive black holes exist in the centres of most galaxies.



An image of the core region of Messier 87, a supermassive black hole, processed from a sparse array of radio telescopes known as the EHT with colors indicating brightness temperature.<sup>[1][2]</sup>



Simulated view of a Schwarzschild black hole in front of the Large Magellanic Cloud. Note the gravitational lensing effect, which produces two enlarged but highly distorted views of the Cloud. Across the top, the Milky Way disk appears distorted into an arc.<sup>[3]</sup>

The presence of a black hole can be inferred through its interaction with other matter and with electromagnetic radiation such as visible light. Matter falling toward a black hole can form an accretion disk of infalling plasma, heated by friction and emitting light. In extreme cases, this creates a quasar, some of the brightest objects in the universe. Stars passing too close to a supermassive black hole can be shredded into streamers that shine very brightly before being "swallowed".<sup>[10]</sup> If other stars are orbiting a black hole, their orbits can be used to determine the black hole's mass and location. Such observations can be used to exclude possible alternatives such as neutron stars. In this way, astronomers have identified numerous stellar black hole candidates in binary systems and established that the radio source known as Sagittarius A\*, at the core of the Milky Way galaxy, contains a supermassive black hole of about 4.3 million solar masses.

## History

---

The idea of a body so massive that even light could not escape was briefly proposed by English astronomical pioneer and clergyman John Michell and independently by French scientist Pierre-Simon Laplace. Both scholars proposed very large stars, analogous to modern models of supermassive black holes.<sup>[11]</sup>

Michell's idea, in a short part of a letter published in 1784, calculated that a star with the same density but 500 times the radius of the sun would not let any emitted light escape; the surface escape velocity would exceed the speed of light. Michell correctly noted that such supermassive but non-radiating bodies might be detectable through their gravitational effects on nearby visible bodies.<sup>[11][12][13]</sup>

In 1796, Laplace mentioned that a star could be invisible if it were sufficiently large while speculating on the origin of the Solar System in his book *Exposition du Système du Monde*. Franz Xaver von Zach asked Laplace for a mathematical analysis, which Laplace provided and published in a journal edited by von Zach.<sup>[11]</sup>

Scholars of the time were initially excited by the proposal that giant but invisible 'dark stars' might be hiding in plain view, but enthusiasm dampened in the early 19th century, when light was discovered to be wavelike.<sup>[14]</sup> Since light was understood as a wave rather than a particle, it was unclear what, if any, influence gravity would have on escaping light waves.<sup>[11][13]</sup>

## General relativity

In 1911, Albert Einstein published a paper about the properties of acceleration in special relativity, noting that an object accelerating through space outside of a gravitational field would be physically indistinguishable from an object in a static gravitational field. The paper also predicted the deflection of light by massive bodies.<sup>[15]</sup> In 1915, Einstein refined these ideas into his general theory of relativity, which explained how matter affects spacetime, which in turn affects the motion of other matter.<sup>[16][17][18]</sup> This theory formed the basis for black hole physics, although Einstein himself would later try, and fail, to refute the idea that black holes could exist.<sup>[19][20][21]</sup>

Only a few months after Einstein published the field equations describing general relativity, Karl Schwarzschild found a solution describing the gravitational field of a point mass and a spherical mass.<sup>[22][23]</sup> A few months after Schwarzschild, Johannes Droste, a student of Hendrik Lorentz, independently gave the same solution for the point mass and wrote more extensively about its

properties.<sup>[24][25]</sup> At a certain radius from the center of the mass, the Schwarzschild solution became singular, meaning that some of the terms in the Einstein equations became infinite. The nature of this radius, which later became known as the Schwarzschild radius, was not understood at the time.<sup>[26][27]</sup>

In 1924, Arthur Eddington showed that the singularity disappeared after a change of coordinates. In 1933, Georges Lemaître realised that this meant the singularity at the Schwarzschild radius was a non-physical coordinate singularity.<sup>[27]</sup> Arthur Eddington commented on the possibility of a star with mass compressed to the Schwarzschild radius in a 1926 book, noting that Einstein's theory allows us to rule out overly large densities for visible stars like Betelgeuse because the extreme force of gravity would make light unable to escape from the star, redshift the star's entire light spectrum out of existence, and "produce so much curvature of the spacetime metric that space would close up around the star, leaving us outside (i.e., nowhere)."<sup>[28][29]</sup>

In 1931, using special relativity, Subrahmanyan Chandrasekhar calculated that a non-rotating body of electron-degenerate matter above a certain limiting mass (now called the Chandrasekhar limit at  $1.4 M_{\odot}$ ) has no stable solutions.<sup>[30]</sup> His arguments were opposed by many of his contemporaries like Eddington and Lev Landau, who argued that some yet unknown mechanism would stop the collapse.<sup>[31]</sup> They were partially correct: a white dwarf slightly more massive than the Chandrasekhar limit will collapse into a neutron star, which is itself stable.<sup>[32]</sup>

In 1939, based on Chandrasekhar's reasoning, Robert Oppenheimer and George Volkoff predicted that neutron stars above another mass limit, now known as the Tolman–Oppenheimer–Volkoff limit, would collapse further, concluding that no law of physics was likely to intervene and stop at least some stars from collapsing to black holes.<sup>[33]</sup> Their original calculations, based on the Pauli exclusion principle, gave it as  $0.7 M_{\odot}$ . Subsequent consideration of neutron-neutron repulsion mediated by the strong force raised the estimate to approximately  $1.5 M_{\odot}$  to  $3.0 M_{\odot}$ .<sup>[34]</sup> Observations of the neutron star merger GW170817, which is thought to have generated a black hole shortly afterward, have refined the TOV limit estimate to  $\sim 2.17 M_{\odot}$ .<sup>[35][36][37][38][39]</sup>

Oppenheimer and his co-authors interpreted the singularity at the boundary of the Schwarzschild radius as indicating that this was the boundary of a bubble in which time stopped.<sup>[33]:380–381</sup> This is a valid point of view for external observers, but not for infalling observers.<sup>[40]</sup> John Wheeler later described black holes viewed from an external reference frame as "frozen stars" because an outside observer would see the surface of the star frozen in time due to gravitational time dilation, never to fully collapse.<sup>[40]</sup>

Also in 1939, Einstein attempted to prove that black holes were impossible in his publication "On a Stationary System with Spherical Symmetry Consisting of Many Gravitating Masses", using his theory of general relativity to defend his argument.<sup>[41]</sup> Months later, Oppenheimer and his student Hartland Snyder provided the Oppenheimer–Snyder model in their paper "On Continued Gravitational Contraction",<sup>[42]</sup> which predicted the existence of black holes. In the paper, which made no reference to Einstein's recent publication, Oppenheimer and Snyder used Einstein's own theory of general relativity to show the conditions on how a black hole could develop, for the first time in contemporary physics.<sup>[41]</sup>

In 1958, David Finkelstein identified the Schwarzschild surface as an event horizon, calling it "a perfect unidirectional membrane: causal influences can cross it in only one direction". In this sense, events that occur inside of the black hole cannot affect events that occur outside of the black hole.<sup>[43]</sup> Finkelstein created a new reference frame to include the point of view of infalling observers.<sup>[44]:103</sup> Finkelstein's

solution extended the Schwarzschild solution for the future of observers falling into a black hole. A similar concept had already been found by Martin Kruskal, but its significance had not been fully understood at the time.<sup>[44]:103</sup>

## Golden age


The era from the mid-1960s to the mid-1970s was the "golden age of black hole research", when general relativity and black holes became mainstream subjects of research.<sup>[47][48]:258</sup>

In this period, more general black hole solutions were found. In 1963, Roy Kerr found the exact solution for a rotating black hole.<sup>[49][50]</sup> Two years later, Ezra Newman found the cylindrically symmetric solution for a black hole that is both rotating and electrically charged.<sup>[51]</sup>

In 1967, Werner Israel found that the Schwarzschild solution was the only possible solution for a nonspinning, uncharged black hole, and couldn't have any additional parameters. In that sense, a Schwarzschild black hole would be defined by its mass alone, and any two Schwarzschild black holes with the same mass would be identical.<sup>[52]</sup> Israel later found that Reissner-Nordstrom black holes were only defined by their mass and electric charge, while Brandon Carter discovered that Kerr black holes only had two degrees of freedom, mass and spin.<sup>[53][54]</sup> Together, these findings became known as the no-hair theorem, which states that a stationary black hole is completely described by the three parameters of the Kerr–Newman metric: mass, angular momentum, and electric charge.<sup>[55]</sup>

At first, it was suspected that the strange mathematical singularities found in each of the black hole solutions only appeared due to the assumption that a black hole would be perfectly spherically symmetric, and therefore the singularities would not appear in generic situations where black holes would not necessarily be symmetric. This view was held in particular by Vladimir Belinski, Isaak Khalatnikov, and Evgeny Lifshitz, who tried to prove that no singularities appear in generic solutions, although they would later reverse their positions.<sup>[56]</sup> However, in 1965, Roger Penrose proved that general relativity without quantum mechanics requires that singularities appear in all black holes.<sup>[57][58]</sup> Shortly afterwards, Hawking generalized Penrose's solution to find that in all but a few physically infeasible scenarios, a cosmological Big Bang singularity is inevitable unless quantum gravity intervenes.<sup>[59]</sup> For his work, Penrose received half of the 2020 Nobel Prize in Physics, Hawking having died in 2018.<sup>[60]</sup>

Astronomical observations also made great strides during this era. In 1967, Antony Hewish and Jocelyn Bell Burnell discovered pulsars.<sup>[61][62]</sup> and by 1969, these were shown to be rapidly rotating neutron stars.<sup>[63]</sup> Until that time, neutron stars, like black holes, were regarded as just theoretical curiosities, but the discovery of pulsars showed their physical relevance and spurred a further interest in all types of compact objects that might be formed by gravitational collapse.<sup>[64]</sup> Based on observations in Greenwich and Toronto in the early 1970s, Cygnus X-1, a galactic X-ray source discovered in 1964, became the first astronomical object commonly accepted to be a black hole.<sup>[65][66]</sup>



The first simulated image of a black hole, created by Jean-Pierre Luminet in 1978 and featuring the characteristic shadow, photon sphere, and lensed accretion disk. The disk is brighter on one side due to the Doppler beaming.<sup>[45][46]</sup>

Work by James Bardeen, Jacob Bekenstein, Carter, and Hawking in the early 1970s led to the formulation of black hole thermodynamics.<sup>[67]</sup> These laws describe the behaviour of a black hole in close analogy to the laws of thermodynamics by relating mass to energy, area to entropy, and surface gravity to temperature. The analogy was completed when Hawking, in 1974, showed that quantum field theory implies that black holes should radiate like a black body with a temperature proportional to the surface gravity of the black hole, predicting the effect now known as Hawking radiation.<sup>[68]</sup>

## Modern research and observation

The first strong evidence for black holes came from combined X-ray and optical observations of Cygnus X-1 in 1972.<sup>[69]</sup> The x-ray source, located in the Cygnus constellation, was discovered through a survey by two suborbital rockets, as the blocking of x-rays by Earth's atmosphere makes it difficult to detect them from the ground.<sup>[70][71][72]</sup> Unlike stars or pulsars, Cygnus X-1 was not associated with any prominent radio or optical source.<sup>[72][73]</sup> In 1972, Louise Webster, Paul Murdin, and, independently, Charles Thomas Bolton, found that Cygnus X-1 was actually in a binary system with the supergiant star HDE 226868. Using the emission patterns of the visible star, both research teams found that the mass of Cygnus X-1 was likely too large to be a white dwarf or neutron star, indicating that it was probably a black hole.<sup>[74][75]</sup> Further research strengthened their hypothesis.<sup>[76][77]</sup>

While Cygnus X-1, a stellar-mass black hole, was generally accepted by the scientific community as a black hole by the end of 1973,<sup>[76]</sup> it would be decades before a supermassive black hole would gain the same broad recognition. Although, as early as the 1960s, physicists such as Donald Lynden-Bell and Martin Rees had suggested that powerful quasars in the center of galaxies were powered by accreting supermassive black holes, little observational proof existed at the time.<sup>[78][79]</sup> However, the Hubble Space Telescope, launched decades later, found that supermassive black holes were not only present in these active galactic nuclei, but that supermassive black holes in the center of galaxies were ubiquitous: Almost every galaxy had a supermassive black hole at its center, many of which were quiescent.<sup>[80][81]</sup> In 1999, David Merritt proposed the M–sigma relation, which related the dispersion of the velocity of matter in the center bulge of a galaxy to the mass of the supermassive black hole at its core.<sup>[82]</sup> Subsequent studies confirmed this correlation.<sup>[83][84][85]</sup> Around the same time, based on telescope observations of the velocities of stars at the center of the Milky Way galaxy, independent work groups led by Andrea Ghez and Reinhard Genzel concluded that the compact radio source in the center of the galaxy, Sagittarius A\*, was likely a supermassive black hole.<sup>[86][87]</sup> In 2020, Ghez and Genzel won the Nobel Prize in Physics for their prediction.<sup>[88][89]</sup>

On 11 February 2016, the LIGO Scientific Collaboration and Virgo Collaboration announced the first direct detection of gravitational waves, named GW150914, representing the first observation of a black hole merger.<sup>[90]</sup> At the time of the merger, the black holes were approximately 1.4 billion light-years away from Earth and had masses of 30 and 35 solar masses.<sup>[91]:6</sup> The mass of the resulting black hole was approximately 62 solar masses, with an additional three solar masses radiated away as gravitational waves.<sup>[91][92]</sup> The Laser Interferometer Gravitational-Wave Observatory (LIGO) detected the gravitational waves by using two mirrors spaced four kilometers apart to measure microscopic changes in length.<sup>[93]</sup> In 2017, Rainer Weiss, Kip Thorne, and Barry Barish, who had spearheaded the project, were awarded the Nobel Prize in Physics for their work.<sup>[94]</sup> Since the initial discovery in 2015, hundreds more gravitational waves have been observed by LIGO and another interferometer, Virgo.<sup>[95]</sup>

On 10 April 2019, the first direct image of a black hole and its vicinity was published, following observations made by the Event Horizon Telescope (EHT) in 2017 of the supermassive black hole in Messier 87's galactic centre.<sup>[96][97][98]</sup> The observations were carried out by eight observatories in six geographical locations across four days and totaled five petabytes of data.<sup>[99][100][101]</sup> In 2022, the Event Horizon Telescope collaboration released an image of the black hole in the center of the Milky Way galaxy, Sagittarius A\*; The data had been collected in 2017.<sup>[102]</sup> Detailed analysis of the motion of stars recorded by the Gaia mission produced evidence in 2022<sup>[103]</sup> and 2023<sup>[104]</sup> of a black hole named Gaia BH1 in a binary with a Sun-like star about 1,560 light-years (480 parsecs) away. Gaia BH1 is currently the closest known black hole to Earth.<sup>[105][106]</sup> Two more black holes have since been found from Gaia data, one in a binary with a red giant<sup>[107]</sup> and the other in a binary with a G-type star.<sup>[108]</sup>


## Etymology

In December 1967, a student reportedly suggested the phrase "black hole" at a lecture by John Wheeler; Wheeler adopted the term for its brevity and "advertising value", and Wheeler's stature in the field ensured it quickly caught on,<sup>[44][109]</sup> leading some to credit Wheeler with coining the phrase.<sup>[110]</sup>

However, the term was used by others around that time. Science writer Marcia Bartusiak traces the term "black hole" to physicist Robert H. Dicke, who in the early 1960s reportedly compared the phenomenon to the Black Hole of Calcutta, notorious as a prison where people entered but never left alive. The term *black hole* was used in print by Life and Science News magazines in 1963, and by science journalist Ann Ewing in her article "'Black Holes' in Space", dated 18 January 1964, which was a report on a meeting of the American Association for the Advancement of Science held in Cleveland, Ohio.<sup>[44]</sup>

## Properties and structure

A black hole is generally defined as a region of spacetime for which no information-carrying signals or objects can escape.<sup>[111][112]</sup> However, physicists do not have a precisely-agreed-upon definition of a black hole.<sup>[113]</sup> From a local perspective, a black hole can be viewed as a region with a gravitational pull, or, equivalently, spacetime curvature so intense that nothing can escape.<sup>[112][114]</sup> A black hole may also be defined as a region where the escape velocity is greater than the speed of light, or where space is falling inwards faster than the speed of light.<sup>[115]:27[116]</sup> The formula for escape velocity is  $V = \sqrt{2MG/R}$ , where  $M$  represents mass,  $G$  is the gravitational constant, and  $R$  is the radius from the center of the mass.<sup>[115]</sup> From a global perspective, a black hole can be understood as a region of the universe for which no causal influences can reach the outside.<sup>[113][111]</sup>



The first detection of gravitational waves, imaged by LIGO observatories in Hanford Site, Washington and Livingston, Louisiana




Image by the Event Horizon Telescope of the supermassive black hole in the center of Messier 87



The no-hair theorem postulates that, once it achieves a stable condition after formation, a black hole has only three independent physical properties: mass, electric charge, and angular momentum; the black hole is otherwise featureless. If the conjecture is true, any two black holes that share the same values for these properties, or parameters, are indistinguishable from one another. The degree to which the conjecture is true for real black holes is currently an unsolved problem.<sup>[55]</sup>

These properties are unique because they are visible from outside a black hole. For example, a charged black hole repels other like charges just like any other charged object.<sup>[117]</sup> Similarly, the total mass of a black hole can be estimated by analyzing the motion of objects near the black hole, such as stars or gas.<sup>[81]</sup> A black hole's angular momentum (or spin) can be measured from far away by analyzing the electromagnetic spectrum of the accretion disk—The faster the black hole is spinning, the closer matter can get to the event horizon without falling in, and the redder that matter appears due to gravitational redshift.<sup>[118][119]</sup> Properties of black holes can also be detected from the gravitational waves they emit.<sup>[118][120][121][122]</sup>



An artistic depiction of a black hole and its features


Because a black hole eventually achieves a stable state with only three parameters, there is no way to avoid losing information about the initial conditions: the gravitational and electric fields of a black hole give very little information about what went in, and every quantity that cannot be measured far away from the black hole horizon is lost. Additionally, because the Hawking radiation emitted from a black hole is thermal, it cannot carry any information about what fell into the black hole, causing the information to seemingly vanish as the black hole radiates away. Since this seemingly violates the law of conservation of energy, it has been called the black hole information loss paradox.<sup>[123][124][125]</sup>

## Physical parameters

The simplest static black holes have mass but neither electric charge nor angular momentum. These black holes are often referred to as Schwarzschild black holes after Karl Schwarzschild, who discovered the solution in 1916.<sup>[23]</sup> According to Birkhoff's theorem, it is the only vacuum solution that is spherically symmetric.<sup>[126]</sup> This means there is no observable difference at a distance between the gravitational field

of such a black hole and that of any other spherical object of the same mass. Contrary to the popular notion of a black hole "sucking in everything" in its surroundings, from far away, the external gravitational field of a black hole is identical to that of any other body of the same mass.<sup>[127]</sup>

Solutions describing more general black holes also exist. Non-rotating charged black holes are described by the Reissner–Nordström metric, while the Kerr metric describes a non-charged rotating black hole. The most general stationary black hole solution known is the Kerr–Newman metric, which describes a black hole with both charge and angular momentum.<sup>[128]</sup>



Radii for shadow and photon sphere relative to event horizon

While the mass of a black hole can theoretically take any positive value, the charge and angular momentum are constrained by the mass. The total electric charge  $Q$  and the total angular momentum  $J$  are expected to satisfy the inequality

$$\frac{Q^2}{4\pi\epsilon_0} + \frac{c^2 J^2}{GM^2} \leq GM^2$$

for a black hole of mass  $M$ . Black holes with the minimum possible mass satisfying this inequality are called extremal. Solutions of Einstein's equations that violate this inequality exist, but they do not possess an event horizon. These solutions have so-called naked singularities that can be observed from the outside.<sup>[129]</sup> Because these singularities make the universe inherently unpredictable, many physicists believe they could not exist.<sup>[130]</sup> The weak cosmic censorship hypothesis, proposed by Sir Roger Penrose, rules out the formation of such singularities, when they are created through the gravitational collapse of realistic matter. However, this theory has not yet been proven, and some physicists believe that naked singularities could exist.<sup>[131]</sup> It is also unknown whether black holes could even become extremal, forming naked singularities, since natural processes counteract increasing spin and charge when a black hole becomes near-extremal.<sup>[131][132][133]</sup>

## Charge

Most black holes are believed to have an approximately neutral charge. For example, Michal Zajaček, Arman Tursunov, Andreas Eckart, and Silke Britzen found the electric charge of Sagittarius A\*, the black hole at the center of the Milky Way galaxy, to be at least ten orders of magnitude below the theoretical maximum.<sup>[134]</sup> If a black hole were to become charged, particles with an opposite sign of charge would be pulled in by the extra electromagnetic force, while particles with the same sign of charge would be repelled, neutralizing the black hole. This effect may not be as strong if the black hole is also spinning.<sup>[135]</sup> The presence of charge can reduce the diameter of the black hole by up to 38% and moves the innermost stable circular orbit inwards, regardless whether the particles in the ISCO are electrically charged or electrically neutral.<sup>[134][136]</sup>



The charge Q for a nonspinning black hole is bounded by

$$Q \leq \sqrt{GM},$$

where G is the gravitational constant and M is the black hole's mass.<sup>[137]</sup>

Spin

Unlike charge, rotation is expected to be a universal feature of compact astrophysical objects, with many black holes spinning at near the maximum rate.<sup>[138][139][140]</sup> For example, the Milky Way's central black hole Sagittarius A\* rotates at about 90% of the maximum rate.<sup>[139][141]</sup> The supermassive black hole in the center of the M87 galaxy appears to have an angular momentum very close to the maximum allowed value.<sup>[139][142][143]</sup> That uncharged limit is<sup>[144]</sup>

$$J \leq \frac{GM^2}{c},$$

allowing definition of a dimensionless spin parameter such that<sup>[144]</sup>

$$0 \leq \frac{cJ}{GM^2} \leq 1.$$

<sup>[144][Note 1]</sup>

Mass

Black holes can have a wide range of masses. The minimum mass of a black hole formed by stellar gravitational collapse is governed by the maximum mass of a neutron star and is believed to be approximately two-to-four solar masses.<sup>[146][147][148]</sup> However, theoretical primordial black holes, believed to have formed soon after the Big Bang, could be far smaller, with masses as little as 10<sup>-5</sup> grams at formation.<sup>[149]</sup> These very small black holes are sometimes called micro black holes.<sup>[150][151]</sup>

Black hole classifications

| Class                               | Approx. mass                                     | Approx. radius                          |
|-------------------------------------|--|---|
| <u>Ultramassive black hole</u>      | 10 <sup>9</sup> –10 <sup>11</sup> M <sub>☉</sub> | >1,000 AU                               |
| <u>Supermassive black hole</u>      | 10 <sup>6</sup> –10 <sup>9</sup> M <sub>☉</sub>  | 0.001–400 AU                            |
| <u>Intermediate-mass black hole</u> | 10 <sup>2</sup> –10 <sup>5</sup> M <sub>☉</sub>  | 10 <sup>3</sup> km ≈ R <sub>Earth</sub> |
| <u>Stellar black hole</u>           | 2–150 M <sub>☉</sub>                             | 30 km                                   |
| <u>Micro black hole</u>             | up to M <sub>Moon</sub>                          | up to 0.1 mm                            |

Black holes formed by stellar collapse are called stellar black holes. Estimates of their maximum mass at formation vary, but generally range from 10 to 100 solar masses, with higher estimates for black holes progenated by low-metallicity stars.<sup>[152][153][154][155][156]</sup> These black holes can also gain mass via accretion of nearby matter.<sup>[157]</sup> Stellar black holes are often found in binaries with stars.<sup>[158][159][160]</sup> These binaries can be categorized as either *low-mass* or *high-mass*; This classification is based on the mass of the companion star, not the compact object itself.<sup>[158]</sup> Stellar black holes are also sometimes found in binaries with other compact objects, such as white dwarfs,<sup>[158]</sup> neutron stars,<sup>[161][162]</sup> and other black holes.<sup>[163][164]</sup>

Black holes that are larger than stellar black holes but smaller than supermassive black holes are called intermediate-mass black holes, with masses of approximately 10<sup>2</sup> to 10<sup>5</sup> solar masses. These black holes seem to be rarer than their stellar and supermassive counterparts, with relatively few candidates having been observed.<sup>[165][156][166]</sup> Physicists have speculated that such black holes may form from collisions in

globular and star clusters or at the center of low-mass galaxies.<sup>[167][168][169][170][171]</sup> They may also form as the result of mergers of smaller black holes, with several LIGO observations finding merged black holes within the 110-350 solar mass range.<sup>[172][173]</sup>

The black holes with the largest masses are called supermassive black holes, with masses more than  $10^6$  times that of the Sun.<sup>[165][174][175]</sup> These black holes are believed to exist at the centers of almost every large galaxy, including the Milky Way.<sup>[80][81][176][177]</sup> Scientists have speculated that these black holes formed via the collapse of large population III stars, direct collapse of large amounts of matter, or the merging of many smaller black holes.<sup>[178][179][179]</sup> Some scientists have proposed a subcategory of even larger black holes, called ultramassive black holes, with masses greater than  $10^9$ - $10^{10}$  solar masses.<sup>[179][180][181]</sup> It is unlikely that black holes with masses greater than 50-100 billion times that of the Sun could exist now, as black hole growth is limited by the age of the universe.<sup>[182][183][184][185]</sup>

## Radius

For a nonspinning, uncharged black hole, the radius of the event horizon, or Schwarzschild radius, is proportional to the mass,  $M$ , through

$$r_s = \frac{2GM}{c^2} \approx 2.95 \frac{M}{M_\odot} \text{ km},$$

where  $r_s$  is the Schwarzschild radius and  $M_\odot$  is the mass of the Sun.<sup>[186]</sup> For a black hole with nonzero spin or electric charge, the radius is smaller,<sup>[Note 2]</sup> until an extremal black hole could have an event horizon close to

$$r_+ = \frac{GM}{c^2},$$

half the radius of a nonspinning, uncharged black hole of the same mass.<sup>[187]</sup>

## External geometry

### Ergosphere

General relativity predicts that any rotating mass will slightly "drag" along the spacetime immediately surrounding it. This will cause the spacetime around the mass to rotate around it, similar to a vortex. The rotating spacetime will then drag any matter and light in it into rotation around the spinning mass as well. This effect, called *frame dragging*, gets stronger closer to the spinning mass.<sup>[189]</sup>

Near a rotating black hole, this effect becomes very strong. In fact, rotating black holes are surrounded by a region of spacetime in which it is impossible to stay still, called the ergosphere. This is because frame dragging is so strong near the event horizon that an object would have to move faster than the speed of light in the opposite direction to just stay still.<sup>[190][189]</sup>

The ergosphere of a black hole is a volume bounded by the black hole's event horizon and the *ergosurface*, which coincides with the event horizon at the poles but is at a much greater distance around the equator.<sup>[188]</sup>


Matter and radiation can escape from the ergosphere. Through the Penrose process, objects can emerge from the ergosphere with more energy than they entered with. The extra energy is taken from the rotational energy of the black hole, slowing down the rotation of the black hole.<sup>[191]</sup> A variation of the Penrose process in the presence of strong magnetic fields, the Blandford–Znajek process, is considered a likely mechanism for the enormous luminosity and relativistic jets of quasars and other active galactic nuclei.<sup>[192]</sup>

## Accretion disk


Due to conservation of angular momentum, gas falling into the gravitational well created by a massive object will typically form a disk-like structure around the object.<sup>[195]</sup> As the disk's angular momentum is transferred outward due to internal processes, its matter falls farther inward, converting its gravitational energy into heat and releasing a large flux of x-rays.<sup>[196][197][198][199]</sup> The temperature of these disks can range from thousands to millions of Kelvin, and temperatures can differ throughout a single accretion disk.<sup>[200][201]</sup> Accretion disks can also emit in other parts of the electromagnetic spectrum, depending on the disk's turbulence and magnetization and the black hole's mass and angular momentum.<sup>[199][202][203]</sup>

Accretion disks can be defined as geometrically thin or geometrically thick. Geometrically thin disks are mostly confined to the black hole's equatorial plane and have a well-defined edge at the innermost stable circular orbit (ISCO), while geometrically thick disks are supported by internal pressure and temperature and can extend inside the ISCO. Disks with high rates of electron scattering and absorption, appearing bright and opaque, are called *optically thick*; *Optically thin* disks are more translucent and produce fainter images when viewed from afar.<sup>[204]</sup> Accretion disks of black holes accreting beyond the Eddington limit are often referred to as "polish donuts" due to their thick, toroidal shape that resembles that of a donut.<sup>[205][206][207]</sup>

Quasar accretion disks are expected to generally appear blue in color.<sup>[208]</sup> The disk for a stellar black hole, on the other hand, would likely look orange, yellow, or red, with its inner regions being the brightest.<sup>[209]</sup> Theoretical research suggests that the hotter a disk is, the bluer it should be, although this is not always supported by observations of real astronomical objects.<sup>[210]</sup> Accretion disk colors may also be altered by the Doppler effect, with the part of the disk travelling towards an observer appearing bluer and brighter and the part of the disk travelling away from the observer appearing redder and dimmer.<sup>[211][212][213]</sup>



The ergosphere is a region outside of the event horizon, where objects cannot remain in place.<sup>[188]</sup>



A black hole with an orange accretion disk. The parts of the disk circling over and under the hole are actually gravitationally lensed from the back side of the black hole.<sup>[193][194]</sup>

## Quasi-periodic oscillations

The X-ray emissions from the disks of accreting black holes sometimes flicker at certain frequencies. These signals are called quasi-periodic oscillations and are thought to be caused by material moving along the inner edge of the accretion disk (the innermost stable circular orbit).<sup>[214][215]</sup> Some scientists also suggest that these oscillations may be caused by the black hole's axis of rotation being out of alignment with the binary system's axis of rotation.<sup>[215]</sup> Since the frequency of quasi-periodic oscillations is correlated with the mass and rotation rate of the compact object, it can be used as an alternative way to determine the properties of candidate black holes.<sup>[214][215][216]</sup>

## Relativistic jets

Some black holes have relativistic jets--thin streams of plasma travelling away from the black hole at more than one-tenth of the speed of light.<sup>[217][Note 3]</sup> These jets can extend as far as millions of parsecs from the black hole itself.<sup>[219]</sup>

Black holes of any mass can have jets.<sup>[220]</sup> However, they are typically observed around spinning black holes with strongly-magnetized accretion disks.<sup>[221][222]</sup> Relativistic jets were more common in the early universe, when galaxies and their corresponding supermassive black holes were rapidly gaining mass.<sup>[221][223]</sup> All black holes with jets also have an accretion disk, but the jets are usually brighter than the disk.<sup>[217][224]</sup> Supermassive black holes with jets are often called quasars, and their stellar-mass companions are referred to as microquasars.<sup>[225]</sup>



Relativistic jets from the supermassive black hole in Centaurus A extend perpendicularly from the galaxy.

The mechanism of formation of jets is not yet known,<sup>[220]</sup> but several options have been proposed. One method proposed to fuel these jets is the Blandford-Znajek process, which suggests that the dragging of magnetic field lines by a black hole's rotation could launch jets of matter into space.<sup>[192][226]</sup> The Penrose process, which involves extraction of a black hole's rotational energy, has also been proposed as a potential mechanism of jet propulsion.<sup>[227][228]</sup>

## Innermost stable circular orbit (ISCO)

In Newtonian gravity, test particles can stably orbit at arbitrary distances from a central object. In general relativity, however, there exists a smallest possible radius for which a massive particle can orbit stably. Any infinitesimal inward perturbations to this orbit will lead to the particle spiraling into the black hole, and any outward perturbations will, depending on the energy, cause the particle to spiral in, move to a stable orbit further from the black hole, or escape to infinity. This orbit is called the **innermost stable circular orbit**, or ISCO.<sup>[230][231]</sup> The location of the ISCO depends on the spin of the black hole and the spin of the particle itself. In the case of a Schwarzschild black hole (spin zero) and a particle without spin, the location of the ISCO is:

$$r_{\text{ISCO}} = 3r_s = \frac{6GM}{c^2}.$$

<sup>[232]</sup> The radius of this orbit changes slightly based on particle spin.<sup>[233][234]</sup> The ISCO moves inward for

charged black holes.<sup>[233]</sup> For spinning black holes, the ISCO is moved inwards for particles orbiting in the same direction that the black hole is spinning (prograde) and outwards for particles orbiting in the opposite direction (retrograde).<sup>[231]</sup> For example, the ISCO for a particle orbiting retrograde can be as far out as about  $9r_s$ , while the ISCO for a particle orbiting prograde can be as close as at the event horizon itself.<sup>[231][235]</sup>


## Plunging region

The final observable region of spacetime around a black hole is called the plunging region. In this area it is no longer possible for matter to follow circular orbits or to stop a final descent into the black hole. Instead, it will rapidly plunge toward the black hole at close to the speed of light, growing increasingly hot and producing a characteristic, detectable thermal emission.<sup>[236][237][238]</sup> However, light and thermal radiation emitted from this region can still escape from the black hole's gravitational pull.<sup>[239]</sup>


## Photon sphere

The photon sphere is a spherical boundary where photons that move on tangents to that sphere would be trapped in an unstable, circular orbit around the black hole.<sup>[240]</sup> For Schwarzschild black holes, the photon sphere has a radius 1.5 times the Schwarzschild radius.<sup>[241][242]</sup> Non-Schwarzschild black holes have a photon sphere radius at least 1.5 times that of the event horizon, but the photon sphere's radius is never larger than the radius of the black hole's shadow.<sup>[241][243]</sup> Their orbits would likely be unstable, so any small perturbation, such as a particle of infalling matter, would cause an instability that would grow over time. This would either set the photon on an outward trajectory, causing it to escape from the black hole's orbit, or on an inward spiral, where it would eventually cross the event horizon.<sup>[241][242][244]</sup>

For a rotating, uncharged black hole, the radius of the photon sphere depends on the spin parameter and whether the photon is orbiting prograde or retrograde.<sup>[232]</sup> For a photon orbiting prograde, the photon sphere will be 1-3 Schwarzschild radii from the center of the black hole, while for a photon orbiting retrograde, the photon sphere will be between 3-5 Schwarzschild radii from the center of the black hole. The exact location of the photon sphere depends on the magnitude of the black hole's rotation.<sup>[245]</sup> For a charged, nonrotating black hole, there will only be one photon sphere, and the radius of the photon sphere will decrease for increasing black hole charge.<sup>[246]</sup> For non-extremal, charged, rotating black holes, there will always be two photon spheres, with the exact radii depending on the parameters of the black hole.<sup>[247]</sup>



Since particles in a black hole's accretion disk must orbit at or outside the ISCO, astronomers can observe the properties of accretion disks to determine black hole spins.<sup>[229]</sup>



Video of a photon being captured by a Schwarzschild black hole



While light can still escape from the photon sphere, any light that crosses the photon sphere on an inbound trajectory will be captured by the black hole. Therefore, any light that reaches an outside observer from the photon sphere must have been emitted by objects between the photon sphere and the event horizon.<sup>[244]</sup>

## Event horizon


The defining feature of a black hole is the existence of an event horizon, a boundary in spacetime through which matter and light can pass only inward towards the center of the black hole. Nothing, not even light, can escape from inside the event horizon.<sup>[249][250]</sup> The event horizon is referred to as such because if an event occurs within the boundary, information from that event cannot reach or affect an outside observer, making it impossible to determine whether such an event occurred.<sup>[251]</sup>

As predicted by general relativity, the presence of a mass deforms spacetime in such a way that the paths taken by particles bend towards the mass.<sup>[252]</sup> At the event horizon of a black hole, this deformation becomes so strong that there are no paths that lead away from the black hole.<sup>[253]</sup>

To a distant observer, a clock near a black hole would appear to tick more slowly than one further from the black hole.<sup>[254]</sup> This effect, known as gravitational time dilation, would also cause an object falling into a black hole to appear to slow as it approached the event horizon, never quite reaching the horizon from the perspective of an outside observer.<sup>[255]</sup> All processes on this object would appear to slow down, and any light emitted by the object to appear redder and dimmer, an effect known as gravitational redshift.<sup>[256]</sup> The falling object would fade away until it could no longer be seen, disappearing from view within less than a second.<sup>[257]</sup>

On the other hand, an observer falling into a black hole would not notice any of these effects as they cross the event horizon. Their own clocks appear to them to tick normally, and they cross the event horizon after a finite time without noting any singular behaviour. In general relativity, it is impossible to determine the location of the event horizon from local observations, due to Einstein's equivalence principle.<sup>[258][259]</sup>

For non-rotating black holes, the geometry of the event horizon is precisely spherical, while for rotating black holes, the event horizon is oblate.<sup>[260][261][262]</sup>



## Internal geometry

### Cauchy horizon

Black holes that are rotating and/or charged have an inner horizon, often called the Cauchy horizon, inside of the black hole.<sup>[263][264]</sup> The inner horizon is divided up into two segments: an ingoing section and an outgoing section.<sup>[265]</sup>

At the ingoing section of the Cauchy horizon, radiation and matter that fall into the black hole would build up at the horizon, causing the curvature of spacetime to go to infinity. This would cause an observer falling in to experience tidal forces.<sup>[263][264][265]</sup> This phenomenon is often called mass inflation, since it is associated with a parameter dictating the black hole's internal mass growing exponentially,<sup>[264][266]</sup> and the buildup of tidal forces is called the mass-inflation singularity<sup>[267][265]</sup> or Cauchy horizon singularity.<sup>[268][269]</sup> Some physicists have argued that in realistic black holes, accretion and Hawking radiation would stop mass inflation from occurring.<sup>[270][271]</sup>

At the outgoing section of the inner horizon, infalling radiation would backscatter off of the black hole's spacetime curvature and travel outward, building up at the outgoing Cauchy horizon. This would cause an infalling observer to experience a gravitational shock wave and tidal forces as the spacetime curvature at the horizon grew to infinity. This buildup of tidal forces is called the shock singularity.<sup>[266][265]</sup>

Both of these singularities are weak, meaning that an object crossing them would only be deformed a finite amount by tidal forces, even though the spacetime curvature would still be infinite at the singularity. This is as opposed to a strong singularity, where an object hitting the singularity would be stretched and squeezed by an infinite amount.<sup>[263][267][266]</sup> They are also null singularities, meaning that a photon could travel parallel to them without ever being intercepted.<sup>[265]</sup>

### Singularity

Mathematical models of black holes based on general relativity have singularities at their centers—points where the curvature of spacetime becomes infinite, and geodesics terminate within a finite proper time. However, it is unknown whether these singularities truly exist in real black holes.<sup>[272]</sup> Some physicists believe that singularities do not exist, and that their existence, which would make spacetime unpredictable, signals a breakdown of general relativity and a need for a more complete understanding of quantum gravity.<sup>[273][274][275]</sup> Others believe that such singularities could be resolved within the current framework of physics, without having to introduce quantum gravity.<sup>[272]</sup> There are also physicists, including Kip Thorne<sup>[211]</sup> and Charles Misner,<sup>[276]</sup> who believe that not all singularities can be resolved, and that some likely still exist in the real universe despite the effects of quantum gravity.<sup>[272][277]</sup> Finally, still others believe that singularities do not exist, and that their existence in general relativity does not matter, since general relativity is already believed to be an incomplete theory.<sup>[272]</sup>


According to general relativity, every black hole has a singularity inside.<sup>[278][279]</sup> For a non-rotating black hole, this region takes the shape of a single point; for a rotating black hole it is smeared out to form a ring singularity that lies in the plane of rotation.<sup>[280]</sup> In both cases, the singular region has zero volume. All of the mass of the black hole ends up in the singularity.<sup>[281]</sup> Since the singularity has nonzero mass in an infinitely small space, it can be thought of as having infinite density.<sup>[282]</sup>

Observers falling into a Schwarzschild black hole (i.e., non-rotating and not charged) cannot avoid being carried into the singularity once they cross the event horizon.<sup>[283][284]</sup> As they fall further into the black hole, they will be torn apart by the growing tidal forces in a process sometimes referred to as spaghettification or the "noodle effect". Eventually, they will reach the singularity and be crushed into an infinitely small point.<sup>[285]</sup>

Before the 1970s, most physicists believed that the interior of a Schwarzschild black hole curved inwards towards a sharp point at the singularity. However, in the late 1960s, Soviet physicists Vladimir Belinskii, Isaak Khalatnikov, and Evgeny Lifshitz discovered that this model was only true when the spacetime inside the black hole had not been perturbed. Any perturbations, such as those caused by matter or radiation falling in, would cause space to oscillate chaotically near the singularity. Any matter falling in would experience intense tidal forces rapidly changing in direction, all while being compressed into an increasingly small volume. Physicists termed these oscillations "Mixmaster dynamics", after a brand of mixer that was popular at the time that Belinskii, Khalatnikov, and Lifshitz made their discovery, because they have a similar effect on matter near a singularity as an electric mixer would have on dough.<sup>[286][211][287]</sup>

In the case of a charged (Reissner–Nordström) or rotating (Kerr) black hole, it is possible to avoid the singularity. Extending these solutions as far as possible reveals the hypothetical possibility of exiting the black hole into a different spacetime with the black hole acting as a wormhole.<sup>[288]</sup> The possibility of travelling to another universe is, however, only theoretical, since any perturbation would destroy this possibility.<sup>[289]</sup> It also appears to be possible to follow closed timelike curves (returning to one's own past) around the Kerr singularity, which leads to problems with causality like the grandfather paradox.<sup>[290][291]</sup> However, processes inside the black hole, such as quantum gravity effects or mass inflation, might prevent closed timelike curves from arising.<sup>[291]</sup>

Some models of gravity seeking to solve technical issues with general relativity do not include black hole singularities. These theoretical black holes without singularities are called *regular*, or *nonsingular*, black holes.<sup>[292][293]</sup> For example, the fuzzball model, based on string theory, states that black holes are actually made up of quantum microstates and need not have a singularity or an event horizon.<sup>[294][295]</sup> The theory of loop quantum gravity proposes that the curvature and density at the center of a black hole is large, but not infinite.<sup>[296]</sup>



Chaotic oscillations of spacetime experienced by an object approaching a gravitational singularity



A Mixmaster electric mixer

# Formation

---


Black holes are formed by gravitational collapse of massive stars, either by direct collapse or during a supernova explosion in a process called "fallback".<sup>[297]</sup> Black holes can result from the merger of two neutron stars or a neutron star and a black hole.<sup>[298]</sup> Other more speculative mechanisms include primordial black holes created from density fluctuations in the early universe, the collapse of dark stars, a hypothetical object powered by annihilation of dark matter, or from hypothetical self-interacting dark matter.<sup>[299]</sup>

## Gravitational collapse

Gravitational collapse occurs when an object's internal pressure is insufficient to resist the object's own gravity. At the end of a star's life, it will run out of hydrogen to fuse, and will start fusing more and more massive elements, until it gets to iron. Since the fusion of any heavier elements would require more energy than they would release, the star can no longer perform nuclear fusion. If the iron core of the star is too massive, the star will no longer be able to support itself and will undergo gravitational collapse.<sup>[301][302]</sup>

The collapse may be stopped by the degeneracy pressure of the star's constituents, allowing the condensation of matter into an exotic denser state. Degeneracy pressure occurs from the Pauli exclusion principle—Particles will resist being in the same place as each other. Smaller progenitor stars, with masses less than about  $8 M_{\odot}$ , will be held together by the degeneracy pressure of electrons and will become a white dwarf. For more massive progenitor stars, electron degeneracy pressure is no longer strong enough to resist the force of gravity and the star will be held together by neutron degeneracy pressure, which can occur at much higher densities, forming a neutron star. If the star is still too massive, even neutron degeneracy pressure will not be able to resist the force of gravity and the star will collapse into a black hole.<sup>[303][304]</sup> Which type forms depends on the mass of the remnant of the original star left if the outer layers have been blown away (for example, in a Type II supernova). The mass of the remnant, the collapsed object that survives the explosion, can be substantially less than that of the original star. Remnants exceeding  $5 M_{\odot}$  are produced by stars that were over  $20 M_{\odot}$  before the collapse.<sup>[304]</sup>

The gravitational collapse of heavy stars is assumed to be responsible for the formation of stellar mass black holes. Star formation in the early universe may have resulted in very massive stars, which upon their collapse would have produced black holes of up to  $10^3 M_{\odot}$ . These black holes could be the seeds of the supermassive black holes found in the centres of most galaxies.<sup>[305]</sup> It has further been suggested that massive black holes with typical masses of  $\sim 10^5 M_{\odot}$  could have formed from the direct collapse of gas clouds in the young universe.<sup>[306]</sup> These massive objects have been proposed as the seeds that eventually



Gas cloud being ripped apart by black hole at the centre of the Milky Way (observations from 2006, 2010 and 2013 are shown in blue, green and red, respectively)<sup>[300]</sup>

formed the earliest quasars observed already at redshift  $z \sim 7$ , less than approximately one billion years after the Big Bang.<sup>[307][308]</sup> Some candidates for such objects have been found in observations of the young universe.<sup>[306]</sup>

While most of the energy released during gravitational collapse is emitted very quickly, an outside observer does not actually see the end of this process. Even though the collapse takes a finite amount of time from the reference frame of infalling matter, a distant observer would see the infalling material slow and halt just above the event horizon, due to gravitational time dilation. Light from the collapsing material takes longer and longer to reach the observer, with the light emitted just before the event horizon forms delayed an infinite amount of time. Thus the external observer never sees the formation of the event horizon; instead, the collapsing material seems to become dimmer and increasingly red-shifted, eventually fading away.<sup>[309]</sup>

## Primordial black holes and the Big Bang

In the current epoch of the universe, conditions needed to form black holes are rare and are mostly only found in stars. However, in the early universe, conditions may have allowed for black hole formations via other means. Fluctuations of spacetime soon after the Big Bang may have formed areas that were denser than their surroundings. Initially, these regions would not have been compact enough to form a black hole, but eventually, the curvature of spacetime in the regions become large enough to cause them to collapse into a black hole.<sup>[310][311]</sup> Different models for the early universe vary widely in their predictions of the scale of these fluctuations. Various models predict the creation of primordial black holes ranging in size from a Planck mass ( $\sim 2.2 \times 10^{-8}$  kg) to hundreds of thousands of solar masses.<sup>[312]</sup> Primordial black holes with masses less than  $10^{15}$  g would have evaporated by now due to Hawking radiation.<sup>[313]</sup>

Despite the early universe being extremely dense, it did not re-collapse into a black hole during the Big Bang, since the universe was expanding rapidly and did not have the gravitational differential necessary for black hole formation. Models for the gravitational collapse of objects of relatively constant size, such as stars, do not necessarily apply in the same way to rapidly expanding space such as the Big Bang.<sup>[314][315][316]</sup>

## High-energy collisions

In principle, black holes could be formed in high-energy particle collisions that achieve sufficient density, although no such events have been detected.<sup>[317][318]</sup> These hypothetical micro black holes, which could form from the collision of cosmic rays and Earth's atmosphere or in particle accelerators like the Large Hadron Collider, would not be able to aggregate additional mass.<sup>[319]</sup> Instead, they would evaporate in about  $10^{-25}$  seconds, posing no threat to the Earth.<sup>[320]</sup>




# Evolution

---

## Growth

Once a black hole has formed, it can continue to grow by absorbing additional matter. Any black hole will continually absorb gas and interstellar dust from its surroundings. This growth process is one possible way through which some supermassive black holes may have been formed, although the formation of supermassive black holes is still an open field of research.<sup>[305]</sup> A similar process has been suggested for the formation of intermediate-mass black holes found in globular clusters.<sup>[321]</sup> Black holes can also merge with other objects such as stars or even other black holes. This is thought to have been important, especially in the early growth of supermassive black holes, which could have formed from the aggregation of many smaller objects.<sup>[305]</sup> The process has also been proposed as the origin of some intermediate-mass black holes.<sup>[322][323]</sup>




Simulation of two black holes colliding

Restrictions have been proposed to limit the growth rate of black holes. In theory, at a certain rate of accretion, the outward radiation pressure will become as strong as the inward gravitational force, and the black hole will be unable to accrete any faster. This limit is called the Eddington limit. However, in practicality, many black holes accrete beyond this rate due to their non-spherical geometry or instabilities in the accretion disk. Accretion beyond the limit is called Super-Eddington accretion and may have been commonplace in the early universe.<sup>[324][325][326]</sup> Additionally, mergers of supermassive black holes may take a long time: As a binary of supermassive black holes approach each other, most nearby stars are ejected, leaving little for the remaining black holes to gravitationally interact with that would allow them to get closer to each other. This phenomenon has been called the final parsec problem, as the distance at which this happens is usually around one parsec.<sup>[327][328]</sup>

## Accretion of matter

When a black hole accretes matter, the gas in the inner accretion disk orbits at very high speeds because of its proximity to the black hole. The resulting friction is so significant that it heats the inner disk to temperatures at which it emits vast amounts of electromagnetic radiation (mainly X-rays). These bright X-ray sources may be detected by telescopes. By the time the matter of the disk reaches the ISCO, it will have given off a significant amount of energy: Between 5.7% and 42% of its mass will have been converted to energy, depending on the black hole's spin. Most of this energy (about 90%) is released in a relatively small area, within about 20 black hole radii.<sup>[196]</sup> In many cases, accretion disks are accompanied by relativistic jets that are emitted



The active galactic nucleus of galaxy Centaurus A in X-ray light, believed to be powered by a supermassive black hole (centre) and surrounded by x-ray binaries (blue dots).

along the black hole's poles, which carry away much of the energy. The mechanism for the creation of these jets is currently not well understood, in part due to insufficient data.<sup>[329]</sup>


As such, many of the universe's most energetic phenomena have been attributed to the accretion of matter on black holes. In particular, active galactic nuclei and quasars are believed to be the accretion disks of supermassive black holes.<sup>[330]</sup> Similarly, X-ray binaries are generally accepted to be binary star systems in which one of the two stars is a compact object accreting matter from its companion.<sup>[330]</sup> It has also been suggested that some ultraluminous X-ray sources may be the accretion disks of intermediate-mass black holes.<sup>[331]</sup>

Stars have been observed to get torn apart by tidal forces in the immediate vicinity of supermassive black holes in galaxy nuclei, in what is known as a tidal disruption event (TDE). Some of the material from the disrupted star forms an accretion disk around the black hole, which emits observable electromagnetic radiation.<sup>[332][333][334]</sup>

## Evaporation

In 1974, Stephen Hawking predicted that black holes emit small amounts of thermal radiation at a temperature of  $\frac{\hbar c^3}{8\pi G M k_B}$ , where  $\hbar$  is the reduced Planck constant,  $c$  is the speed of light,  $G$  is the gravitational constant,  $M$  is the mass of the black hole and  $k_B$  is the Boltzmann constant.<sup>[68]</sup> This effect has become known as Hawking radiation. By applying quantum field theory to black holes, Hawking determined that a black hole should continuously emit thermal blackbody radiation. This theory was supported by previous work by Jacob Bekenstein, who theorized that black holes should have a finite entropy proportional to their surface area, and therefore should also have a temperature.<sup>[335]</sup> It is also analogous to a special relativistic effect called the Unruh effect, which predicts that an accelerating observer should detect a higher temperature of their surroundings than a nonaccelerating observer. The temperature experienced by an accelerating observer is the same as an observer near the horizon of a black hole with an equivalent surface gravity. This result is in accordance with the equivalence principle, which states that the effects of acceleration in flat spacetime should be the same as the effects of the equivalent acceleration due to gravity in curved spacetime.<sup>[336]</sup>

Since Hawking's publication, many others have mathematically verified the result through different approaches.<sup>[335]</sup> If Hawking's theory of black hole radiation is correct, then black holes are expected to shrink and evaporate over time as they lose mass by the emission of photons and other particles.<sup>[68]</sup> The temperature of this thermal spectrum (Hawking temperature) is proportional to the surface gravity of the black hole, which is inversely proportional to the mass. Hence, large black holes emit less radiation than small black holes.<sup>[337][338]</sup>



An artist's impression (top) of a supermassive black hole tidally deforming a star based on observations from the Chandra X-ray observatory and the European Southern Observatory.

A stellar black hole of  $1 M_{\odot}$  has a Hawking temperature of 62 nanokelvins.<sup>[339]</sup> This is far less than the 2.7 K temperature of the cosmic microwave background radiation. Stellar-mass or larger black holes receive more mass from the cosmic microwave background than they emit through Hawking radiation and thus will grow instead of shrinking.<sup>[340]</sup> To have a Hawking temperature larger than 2.7 K (and be able to evaporate), a black hole would need a mass less than the Moon. Such a black hole would have a diameter of less than a tenth of a millimetre.<sup>[341]</sup>

If a black hole is very small, the radiation effects are expected to become very strong. A black hole with the mass of a car would have a diameter of about  $10^{-24}$  m and take a nanosecond to evaporate, during which time it would briefly have a luminosity of more than 200 times that of the Sun. Lower-mass black holes are expected to evaporate even faster. For a very small black hole, quantum gravity effects are expected to play an important role and could hypothetically make the black hole stable, although current developments in quantum gravity do not indicate this is the case.<sup>[342][343]</sup>

The Hawking radiation for an astrophysical black hole is predicted to be very weak and would thus be exceedingly difficult to detect from Earth. A possible exception is the burst of gamma rays emitted in the last stage of the evaporation of primordial black holes. Searches for such flashes have proven unsuccessful and provide stringent limits on the possibility of existence of low mass primordial black holes, with modern research predicting that primordial black holes must make up less than a fraction of  $10^{-7}$  of the universe's total mass.<sup>[344][345]</sup> NASA's Fermi Gamma-ray Space Telescope, launched in 2008, has searched for these flashes, but has not yet found any.<sup>[346][347]</sup>

If black holes evaporate via Hawking radiation, a non-accreting solar mass black hole will evaporate (beginning once the temperature of the cosmic microwave background drops below that of the black hole) over a period of  $10^{64}$  years.<sup>[348]</sup> A supermassive black hole with a mass of  $10^{11} M_{\odot}$  will evaporate in around  $2 \times 10^{100}$  years.<sup>[349]</sup> During the collapse of a supercluster of galaxies, supermassive black holes are predicted to grow to up to  $10^{14} M_{\odot}$ . Even these would evaporate over a timescale of up to  $10^{106}$  years.<sup>[348]</sup> It is unknown exactly what would happen at the end of a black hole's evaporation. Some physicists theorize that it would leave behind a remnant, such as a naked singularity.<sup>[350][351][352]</sup>

## Observational evidence

---

Millions of black holes with around 30 solar masses derived from stellar collapse are expected to exist in the Milky Way. Even a dwarf galaxy like Draco should have hundreds.<sup>[353]</sup> Only a few of these have been detected. By nature, black holes do not themselves emit any electromagnetic radiation other than the hypothetical Hawking radiation, so astrophysicists searching for black holes must generally rely on indirect observations.

### Direct interferometry


The Event Horizon Telescope (EHT) is an active program that directly observes the immediate environment of black holes' event horizons, such as the black hole at the centre of the Milky Way. In April 2017, EHT began observing the black hole at the centre of Messier 87.<sup>[354][355]</sup> Using petabytes of data from eight different radio observatories over a ten-day observation period, the EHT team created a composite image of the black hole, which they debuted in April 2019.<sup>[356][357][358][359]</sup> The black hole's shadow appears as a dark circle in the centre of the image, bordered by the orange-red ring of its

accretion disk.<sup>[360]</sup> The bottom half of the disk is brighter than the top due to Doppler beaming: Material at the bottom of the disk, which is travelling towards the viewer at relativistic speeds, appears brighter than the material at the top of the disk, which is travelling away from the viewer.<sup>[361][360]</sup> In April 2023, the EHT team presented an image of the shadow of the Messier 87 black hole and its high-energy jet, viewed together for the first time.<sup>[362][363]</sup>


On 12 May 2022, the EHT released the first image of Sagittarius A\*, the supermassive black hole at the centre of the Milky Way galaxy. The EHT team had previously detected magnetic field lines around the black hole, confirming theoretical predictions of magnetic fields around black holes.<sup>[364][365]</sup> The imaging of Sagittarius A\* was done concurrently with the imaging of the Messier 87 (M87\*) black hole. Like M87\*, Sagittarius A\*'s shadow and accretion disk can be seen in the EHT image, with the size of the shadow matching theoretical projections.<sup>[357][366]</sup> Although the image of Sagittarius A\* was created through the same process as for M87\*, it was significantly more complex to image Sagittarius A\* because of the instability of its surroundings. Because Sagittarius A\* is one thousand times less massive as M87\*, its accretion disk has a much shorter orbital period, so the environment around Sagittarius A\* was rapidly changing as the EHT team was trying to image it.<sup>[367]</sup> Additionally, turbulent plasma lies between Sagittarius A\* and Earth, preventing resolution of the image at longer wavelengths.<sup>[368]</sup>

## Detection of gravitational waves from merging black holes


On 14 September 2015, the LIGO gravitational wave observatory made the first-ever successful direct observation of gravitational waves.<sup>[90][369]</sup> The signal was consistent with theoretical predictions for the gravitational waves produced by the merger of two black holes: one with about 36 solar masses, and the other around 29 solar masses.<sup>[90][370]</sup> Because the two objects were only 350 km apart just before the merger, yet were more massive than possible for a neutron star, the LIGO team concluded that the gravitational waves must have come from a merger of two black holes.<sup>[90]</sup> The signal observed by LIGO also included the start of the post-merger ringdown, the signal produced as the newly formed



An image of supermassive black hole M87\* with lines drawn over it to indicate the degree and direction of light polarization



The M87\* black hole and its relativistic jet



Sagittarius A\*, the supermassive black hole in the center of the Milky Way

compact object settles down to a stationary state.<sup>[371]</sup> From the ringdown, the LIGO team was able to determine that the resulting merged black hole was spinning at 67% of the maximum rate and had a mass of 62 solar masses, having lost three solar masses as gravitational waves during the merger.<sup>[90][370]</sup>

The observation also provides the first observational evidence for the existence of stellar-mass black hole binaries. Furthermore, it is the first observational evidence of stellar-mass black holes weighing 25 solar masses or more.<sup>[372]</sup>

Since then, many more gravitational wave events have been observed.<sup>[373]</sup>

## Stars orbiting Sagittarius A\*

The proper motions of stars near the centre of the Milky Way provide strong observational evidence that these stars are orbiting a supermassive black hole.<sup>[374]</sup> Since 1995, astronomers have tracked the motions of 90 stars orbiting an invisible object coincident with the radio source Sagittarius A\*. In 1998, by fitting the motions of the stars to Keplerian orbits, the astronomers were able to infer that Sagittarius A\* must be a  $2.6 \times 10^6 M_{\odot}$  object must be contained within a radius of 0.02 light-years.<sup>[375]</sup>

Since then, one of the stars—called S2—has completed a full orbit. From the orbital data, astronomers were able to refine the calculations of the mass of Sagittarius A\* to  $4.3 \times 10^6 M_{\odot}$ , with a radius of less than 0.002 light-years.<sup>[374]</sup>


The upper limit on Sagittarius A\*'s size is still too large to test whether it is smaller than its Schwarzschild radius. Nevertheless, these observations strongly suggest that the

central object is a supermassive black hole as there are no other plausible scenarios for confining so much invisible mass into such a small volume.<sup>[375]</sup> Additionally, there is some observational evidence that this object might possess an event horizon, a feature unique to black holes.<sup>[376]</sup> The Event Horizon Telescope image of Sagittarius A\*, released in 2022, provided further confirmation that it is indeed a black hole.<sup>[377]</sup>

In 2020, Andrea Ghez and Reinhard Genzel shared one-half of the Nobel Prize in Physics for their discovery that Sagittarius A\* was a supermassive black hole.<sup>[378][379]</sup>

## X-ray binaries

X-ray binaries are binary star systems that emit a majority of their radiation in the X-ray part of the electromagnetic spectrum. These X-ray emissions result when a compact object accretes matter from an ordinary star.<sup>[380]</sup> The presence of an ordinary star in such a system provides an opportunity for studying



LIGO measurement of the gravitational waves at the Livingston (right) and Hanford (left) detectors, compared with the theoretical predicted values



Stars moving around Sagittarius A\* as seen in 2021



the central object and to determine if it might be a black hole. By measuring the orbital period of the binary, the distance to the binary from Earth, and the mass of the companion star, scientists can estimate the mass of the compact object.<sup>[381]</sup> The Tolman-Oppenheimer-Volkoff limit (TOV limit) dictates the largest mass a nonrotating neutron star can be, and is estimated to be about two solar masses. While a rotating neutron star can be slightly more massive, if the compact object is much more massive than the TOV limit, it cannot be a neutron star and is generally expected to be a black hole.<sup>[330][382]</sup>


The first strong candidate for a black hole, Cygnus X-1, was discovered in this way by Charles Thomas Bolton,<sup>[9]</sup> Louise Webster, and Paul Murdin<sup>[8]</sup> in 1972.<sup>[383][66]</sup> Observations of rotation broadening of the optical star reported in 1986 lead to a compact object mass estimate of 16 solar masses, with 7 solar masses as the lower bound.<sup>[330]</sup> In 2011, this estimate was updated to  $14.1 \pm 1.0 M_{\odot}$  for the black hole and  $19.2 \pm 1.9 M_{\odot}$  for the optical stellar companion.<sup>[384]</sup>

In a class of X-ray binaries called soft X-ray transients, the companion star is of relatively low mass, allowing for more accurate estimates of the black hole mass. These systems actively emit X-rays for only several months once every 10–50 years. During the period of low X-ray emission, called quiescence, the accretion disk is extremely faint, allowing detailed observation of the companion star.<sup>[330]</sup> Numerous black hole candidates have been measured by this method.<sup>[385]</sup>

## Galactic nuclei

Astronomers use the term "active galaxy" to describe galaxies with unusual characteristics, such as unusual spectral line emission and very strong radio emission. Theoretical and observational studies have shown that the high levels of activity in the centers of these galaxies, regions called active galactic nuclei (AGN), may be explained by accretion onto supermassive black holes. These AGN consist of a central black hole that may be millions or billions of times more massive than the Sun, a disk of interstellar gas and dust called an accretion disk, and two jets perpendicular to the accretion disk.<sup>[387][388][389]</sup>

Although supermassive black holes are expected to be found in most AGN, only some galaxies' nuclei have been more carefully studied in attempts to both identify and measure the actual masses of the central supermassive black hole candidates. Some of the most notable galaxies with supermassive black hole candidates include the Andromeda Galaxy, Messier 32, Messier 87, the Sombrero Galaxy, and the Milky Way itself.<sup>[390][391]</sup>



A Chandra X-Ray Observatory image of Cygnus X-1, which was the first strong black hole candidate discovered



Detection of unusually bright X-ray flare from Sagittarius A\*, a black hole in the centre of the Milky Way galaxy on 5 January 2015<sup>[386]</sup>

It is now widely accepted that the centre of nearly every galaxy, not just active ones, contains a supermassive black hole.<sup>[392]</sup> The close observational correlation between the mass of this hole and the velocity dispersion of the host galaxy's bulge, known as the M–sigma relation, strongly suggests a connection between the formation of the black hole and that of the galaxy itself.<sup>[393]</sup>

## Microlensing

Another way the black hole nature of an object may be tested is through observation of effects caused by a strong gravitational field in their vicinity. One such effect is gravitational lensing: The deformation of spacetime around a massive object causes light rays to be deflected, such as light passing through an optic lens. Observations have been made of weak gravitational lensing, in which light rays are deflected by only a few arcseconds. Microlensing occurs when the sources are unresolved and the observer sees a small brightening. The turn of the millennium saw the first 3 candidate detections of black holes in this way,<sup>[394][395]</sup> and in January 2022, astronomers reported the first confirmed detection of a microlensing event from an isolated black hole.<sup>[396]</sup>

Another possibility for observing gravitational lensing by a black hole would be to observe stars orbiting the black hole. There are several candidates for such an observation in orbit around Sagittarius A\*.<sup>[397]</sup>

## Alternatives

---

While there is a strong case for supermassive black holes, the model for stellar-mass black holes assumes of an upper limit for the mass of a neutron star: objects observed to have more mass are assumed to be black holes. However, the properties of extremely dense matter are poorly understood. New exotic phases of matter could allow other kinds of massive objects.<sup>[330]</sup> A phase of free quarks at high density might allow the existence of dense quark stars,<sup>[398]</sup> and some supersymmetric models predict the existence of Q stars.<sup>[399]</sup> Some extensions of the Standard Model posit the existence of preons as fundamental building blocks of quarks and leptons, which could hypothetically form preon stars.<sup>[400]</sup> These hypothetical models could potentially explain a number of observations of stellar black hole candidates. However, it can be shown from arguments in general relativity that any such object will have a maximum mass.<sup>[330]</sup>

Since the average density of a black hole inside its Schwarzschild radius is inversely proportional to the square of its mass, supermassive black holes are much less dense than stellar black holes. The average density of a  $10^8 M_{\odot}$  black hole is comparable to that of water.<sup>[330]</sup> Consequently, the physics of matter forming a supermassive black hole is much better understood and the possible alternative explanations for supermassive black hole observations are much more mundane. For example, a supermassive black hole could be modelled by a large cluster of very dark objects. However, such alternatives are typically not stable enough to explain the supermassive black hole candidates.<sup>[330]</sup>

A few theoretical objects have been conjectured to match observations of astronomical black hole candidates identically or near-identically,<sup>[401]</sup> but which function via a different mechanism. These include the gravastar,<sup>[402]</sup> the black star,<sup>[403]</sup> related nestar<sup>[404]</sup> and the dark-energy star.<sup>[405]</sup>

# Open questions

---

## Entropy and thermodynamics

In 1971, Hawking showed under general conditions<sup>[Note 5]</sup> that the total area of the event horizons of any collection of classical black holes can never decrease, even if they collide and merge.<sup>[406]</sup> This result, now known as the second law of black hole mechanics, is remarkably similar to the second law of thermodynamics, which states that the total entropy of an isolated system can never decrease. As with classical objects at absolute zero temperature, it was assumed that black holes had zero entropy. If this were the case, the second law of thermodynamics would be violated by entropy-laden matter entering a black hole, resulting in a decrease in the total entropy of the universe. Therefore, Bekenstein proposed that a black hole should have an entropy, and that it should be proportional to its horizon area.<sup>[407]</sup>

$$S = \frac{1}{4} \frac{c^3 k}{G \hbar} A$$

The formula for the Bekenstein–Hawking entropy ( $S$ ) of a black hole, which depends on the area of the black hole ( $A$ ). The constants are the speed of light ( $c$ ), the Boltzmann constant ( $k$ ), Newton's constant ( $G$ ), and the reduced Planck constant ( $\hbar$ ). In Planck units, this reduces to  $S = \frac{A}{4}$ .

The link with the laws of thermodynamics was further strengthened by Hawking's discovery in 1974 that quantum field theory predicts that a black hole radiates blackbody radiation at a constant temperature. This seemingly causes a violation of the second law of black hole mechanics, since the radiation will carry away energy from the black hole causing it to shrink. The radiation also carries away entropy, and it can be proven under general assumptions that the sum of the entropy of the matter surrounding a black hole and one quarter of the area of the horizon as measured in Planck units is in fact always increasing. This allows the formulation of the first law of black hole mechanics as an analogue of the first law of thermodynamics, with the mass acting as energy, the surface gravity as temperature and the area as entropy.<sup>[407]</sup>

One puzzling feature is that the entropy of a black hole scales with its area rather than with its volume, since entropy is normally an extensive quantity that scales linearly with the volume of the system. This odd property led Gerard 't Hooft and Leonard Susskind to propose the holographic principle, which suggests that anything that happens in a volume of spacetime can be described by data on the boundary of that volume.<sup>[408]</sup>

Although general relativity can be used to perform a semiclassical calculation of black hole entropy, this situation is theoretically unsatisfying. In statistical mechanics, entropy is understood as counting the number of microscopic configurations of a system that have the same macroscopic qualities, such as mass, charge, and pressure. Without a satisfactory theory of quantum gravity, one cannot perform such a computation for black holes.<sup>[409]</sup> Some progress has been made in various approaches to quantum gravity. In 1995, Andrew Strominger and Cumrun Vafa showed that counting the microstates of a specific supersymmetric black hole in string theory reproduced the Bekenstein–Hawking entropy.<sup>[410]</sup> Since then, similar results have been reported for different black holes both in string theory and in other approaches to quantum gravity like loop quantum gravity.<sup>[411]</sup>

## Information loss paradox

### Unsolved problem in physics



*Is physical information lost in black holes?*

[More unsolved problems in physics](#)

Because, according to the no-hair theorem, a black hole has only a few internal parameters, most of the information about the matter that went into forming the black hole is lost. Regardless of the type of matter which goes into a black hole, it appears that only information concerning the total mass, charge, and angular momentum are conserved. When black holes were thought to persist forever, this information loss was not problematic, as the information can be thought of as existing inside the black hole, inaccessible from the outside, but represented on the event horizon in accordance with the holographic principle. However, black holes slowly evaporate by emitting Hawking radiation. This radiation does not appear to carry any additional information about the matter that formed the black hole, meaning that this information appears to be gone forever.<sup>[412]</sup>

The question whether information is truly lost in black holes (the black hole information paradox) has divided the theoretical physics community. In quantum mechanics, loss of information corresponds to the violation of a property called unitarity, and it has been argued that loss of unitarity would also imply violation of conservation of energy,<sup>[413]</sup> though this has also been disputed.<sup>[414]</sup> Over recent years evidence has been building that indeed information and unitarity are preserved in a full quantum gravitational treatment of the problem.<sup>[415]</sup>

One attempt to resolve the black hole information paradox is known as black hole complementarity. In 2012, the "firewall paradox" was introduced with the goal of demonstrating that black hole complementarity fails to solve the information paradox. According to quantum field theory in curved spacetime, a single emission of Hawking radiation involves two mutually entangled particles. The outgoing particle escapes and is emitted as a quantum of Hawking radiation; the infalling particle is swallowed by the black hole. Assume a black hole formed a finite time in the past and will fully evaporate away in some finite time in the future. Then, it will emit only a finite amount of information encoded within its Hawking radiation. According to research by physicists like Don Page<sup>[416][417]</sup> and Leonard Susskind, there will eventually be a time by which an outgoing particle must be entangled with all the Hawking radiation the black hole has previously emitted.

This seemingly creates a paradox: a principle called "monogamy of entanglement" requires that, like any quantum system, the outgoing particle cannot be fully entangled with two other systems at the same time; yet here the outgoing particle appears to be entangled both with the infalling particle and, independently, with past Hawking radiation.<sup>[418]</sup> In order to resolve this contradiction, physicists may eventually be forced to give up one of three time-tested principles: Einstein's equivalence principle, unitarity, or local quantum field theory. One possible solution, which violates the equivalence principle, is that a "firewall" destroys incoming particles at the event horizon.<sup>[419]</sup> In general, which—if any—of these assumptions should be abandoned remains a topic of debate.<sup>[414]</sup>

## In science fiction

---

The concept of black holes became popular in science and fiction alike in the 1960s. Authors quickly seized upon the relativistic effect of gravitational time dilation, whereby time passes more slowly closer to a black hole due to its immense gravitational field. Black holes also became a popular means of space travel in science fiction, especially when the notion of wormholes emerged as a relatively plausible way to achieve faster-than-light travel. In this concept, a black hole is connected to its theoretical opposite, a so-called white hole, and as such acts as a gateway to another point in space which might be very distant from the point of entry. More exotically, the point of emergence is occasionally portrayed as another point in time—thus enabling time travel—or even an entirely different universe.

More fanciful depictions of black holes that do not correspond to their known or predicted properties also appear. As nothing inside the event horizon—the distance away from the black hole where the escape velocity exceeds the speed of light—can be observed from the outside, authors have been free to employ artistic license when depicting the interiors of black holes. A small number of works also portray black holes as being sentient.

Besides stellar-mass black holes, supermassive and especially micro black holes also make occasional appearances. Supermassive black holes are a common feature of modern space opera. Recurring themes in stories depicting micro black holes include spaceship propulsion, threatening or causing the destruction of the Earth, and serving as a source of gravity in outer-space settlements.

## See also

---



***Astronomy portal***



***History of science portal***



***Mathematics portal***



***Physics portal***



***Stars portal***

- Black brane or Black string
- Black Hole Initiative
- Black hole starship
- Black holes in fiction
- Blanet
- BTZ black hole
- Dark star (dark matter)
- Golden binary
- Hypothetical black hole (disambiguation)
- Kugelblitz (astrophysics)
- List of black holes
- List of nearest black holes
- Outline of black holes
- Planck star
- Sonic black hole
- Susskind-Hawking battle
- Timeline of black hole physics
- Virtual black hole
- White hole



## Notes

---

1. The value of  $cJ/GM^2$  can exceed 1 for objects other than black holes. The largest value known for a neutron star is  $\leq 0.4$ , and commonly used equations of state would limit that value to  $< 0.7$ .<sup>[145]</sup>
2. The (outer) event horizon radius scales as:  $M + \sqrt{M^2 - (J/M)^2 - Q^2}$ .
3. Relativistic jets do not come from matter from inside of the black hole, but from matter close to the black hole.<sup>[218]</sup>
4. The set of possible paths, or more accurately the future light cone containing all possible world lines (in this diagram the light cone is represented by the V-shaped region bounded by arrows representing light ray world lines), is tilted in this way in Eddington–Finkelstein coordinates (the diagram is a "cartoon" version of an Eddington–Finkelstein coordinate diagram), but in other coordinates the light cones are not tilted in this way, for example in Schwarzschild coordinates they narrow without tilting as one approaches the event horizon, and in Kruskal–Szekeres coordinates the light cones do not change shape or orientation at all.<sup>[248]</sup>
5. In particular, he assumed that all matter satisfies the weak energy condition.

## References

---

1. The Event Horizon Telescope Collaboration; Akiyama, Kazunori; Alberdi, Antxon; Alef, Walter; Asada, Keiichi; Azulay, Rebecca; Bacsko, Anne-Kathrin; Ball, David; Baloković, Mislav; Barrett, John; Bintley, Dan; Blackburn, Lindy; Boland, Wilfred; Bouman, Katherine L.; Bower, Geoffrey C. (10 April 2019). "First M87 Event Horizon Telescope Results. IV. Imaging the Central Supermassive Black Hole" (<https://doi.org/10.3847%2F2041-8213%2Fab0e85>). *The Astrophysical Journal Letters*. **875** (1): L4. arXiv:1906.11241 (<https://arxiv.org/abs/1906.11241>). Bibcode:2019ApJ...875L...4E (<https://ui.adsabs.harvard.edu/abs/2019ApJ...875L...4E>). doi:10.3847/2041-8213/ab0e85 (<https://doi.org/10.3847%2F2041-8213%2Fab0e85>). ISSN 2041-8205 (<https://search.worldcat.org/issn/2041-8205>).
2. "Astronomers capture first image of a black hole" (<https://new.nsf.gov/news/astronomers-capture-first-image-black-hole#image-caption-credit-block>). *new.nsf.gov*. 10 April 2019. Retrieved 28 January 2025.
3. Riazuelo, Alain (2019). "Seeing relativity -- I. Ray tracing in a Schwarzschild metric to explore the maximal analytic extension of the metric and making a proper rendering of the stars". *International Journal of Modern Physics D*. **28** (2): 1950042. arXiv:1511.06025 (<https://arxiv.org/abs/1511.06025>). Bibcode:2019IJMPD..2850042R (<https://ui.adsabs.harvard.edu/abs/2019IJMPD..2850042R>). doi:10.1142/S0218271819500421 (<https://doi.org/10.1142%2FS0218271819500421>). S2CID 54548877 (<https://api.semanticscholar.org/CorpusID:54548877>).
4. Overbye, Dennis (8 June 2015). "Black Hole Hunters" (<https://www.nytimes.com/2015/06/09/science/black-hole-event-horizon-telescope.html>). *NASA*. Archived (<https://web.archive.org/web/20150609023631/http://www.nytimes.com/2015/06/09/science/black-hole-event-horizon-telescope.html>) from the original on 9 June 2015. Retrieved 8 June 2015.
5. Hamilton, A. "Journey into a Schwarzschild black hole" (<http://jila.colorado.edu/~ajsh/insidebh/schw.html>). *jila.colorado.edu*. Archived (<https://web.archive.org/web/20190903235853/http://jila.colorado.edu/~ajsh/insidebh/schw.html>) from the original on 3 September 2019. Retrieved 28 June 2020.

6. Schutz, Bernard F. (2003). *Gravity from the ground up* ([https://books.google.com/books?id=P\\_T0xxhDcslC](https://books.google.com/books?id=P_T0xxhDcslC)). Cambridge University Press. p. 110. ISBN 978-0-521-45506-0. Archived ([https://web.archive.org/web/20161202222711/https://books.google.com/books?id=P\\_T0xxhDcslC](https://web.archive.org/web/20161202222711/https://books.google.com/books?id=P_T0xxhDcslC)) from the original on 2 December 2016.
7. Davies, P. C. W. (1978). "Thermodynamics of Black Holes" (<https://web.archive.org/web/20130510184530/http://cosmos.asu.edu/publications/papers/ThermodynamicTheoryofBlackHoles%2034.pdf>) (PDF). *Reports on Progress in Physics* (Review). **41** (8): 1313–1355. Bibcode:1978RPPh...41.1313D (<https://ui.adsabs.harvard.edu/abs/1978RPPh...41.1313D>). doi:10.1088/0034-4885/41/8/004 (<https://doi.org/10.1088/0034-4885/41/8/004>). S2CID 250916407 (<https://api.semanticscholar.org/CorpusID:250916407>). Archived from the original (<http://cosmos.asu.edu/publications/papers/ThermodynamicTheoryofBlackHoles%2034.pdf>) (PDF) on 10 May 2013.
8. Webster, B. Louise; Murdin, Paul (1972), "Cygnus X-1—a Spectroscopic Binary with a Heavy Companion?", *Nature* (Letter), **235** (5332): 37–38, Bibcode:1972Natur.235...37W (<https://ui.adsabs.harvard.edu/abs/1972Natur.235...37W>), doi:10.1038/235037a0 (<https://doi.org/10.1038/235037a0>), S2CID 4195462 (<https://api.semanticscholar.org/CorpusID:4195462>)
9. Bolton, C. T. (1972), "Identification of Cygnus X-1 with HDE 226868", *Nature* (Letter), **235** (5336): 271–273, Bibcode:1972Natur.235..271B (<https://ui.adsabs.harvard.edu/abs/1972Natur.235..271B>), doi:10.1038/235271b0 (<https://doi.org/10.1038/235271b0>), S2CID 4222070 (<https://api.semanticscholar.org/CorpusID:4222070>)
10. Clery D (2020). "Black holes caught in the act of swallowing stars" (<https://doi.org/10.1126/2Fscience.367.6477.495>). *Science*. **367** (6477): 495. Bibcode:2020Sci...367..495C (<https://ui.adsabs.harvard.edu/abs/2020Sci...367..495C>). doi:10.1126/science.367.6477.495 (<https://doi.org/10.1126/science.367.6477.495>). PMID 32001633 (<https://pubmed.ncbi.nlm.nih.gov/32001633>). S2CID 210984462 (<https://api.semanticscholar.org/CorpusID:210984462>).
11. Montgomery, Colin; Orchiston, Wayne; Whittingham, Ian (2009) [Available online 18 April 2023]. "Michell, Laplace and the origin of the black hole concept" ([https://researchonline.jcu.edu.au/9892/1/Microsoft\\_Word\\_-\\_Paper\\_Black\\_Hole\\_Concept\\_Final\\_.pdf](https://researchonline.jcu.edu.au/9892/1/Microsoft_Word_-_Paper_Black_Hole_Concept_Final_.pdf)) (PDF). *Journal of Astronomical History and Heritage* (Research article). **12** (2): 90–96. Bibcode:2009JAHH...12...90M (<https://ui.adsabs.harvard.edu/abs/2009JAHH...12...90M>). doi:10.3724/SP.J.1440-2807.2009.02.01 (<https://doi.org/10.3724/SP.J.1440-2807.2009.02.01>). S2CID 55890996 (<https://api.semanticscholar.org/CorpusID:55890996>).
12. Michell, J. (1784). "On the Means of Discovering the Distance, Magnitude, &c. of the Fixed Stars, in Consequence of the Diminution of the Velocity of Their Light, in Case Such a Diminution Should be Found to Take Place in any of Them, and Such Other Data Should be Procured from Observations, as Would be Farther Necessary for That Purpose" (<https://doi.org/10.1098/2Frstl.1784.0008>). *Philosophical Transactions of the Royal Society*. **74**: 35–57. Bibcode:1784RSPT...74...35M (<https://ui.adsabs.harvard.edu/abs/1784RSPT...74...35M>). doi:10.1098/rstl.1784.0008 (<https://doi.org/10.1098/2Frstl.1784.0008>). JSTOR 106576 (<https://www.jstor.org/stable/106576>).
13. Thorne 1994, pp. 123–124
14. Slayter, Elizabeth M.; Slayter, Henry S. (1992). *Light and Electron Microscopy* (<https://books.google.com/books?id=LlePVS9oq7MC>). Cambridge University Press. ISBN 978-0-521-33948-3. Archived (<https://web.archive.org/web/20171130151044/https://books.google.com/books?id=LlePVS9oq7MC>) from the original on 30 November 2017.
15. Einstein, Albert (1911), "Einfluss der Schwerkraft auf die Ausbreitung des Lichtes (On the Influence of Gravity on the Propagation of Light)" (<https://zenodo.org/record/1424213>), *Annalen der Physik*, **35** (10): 898–908, Bibcode:1911AnP...340..898E (<https://ui.adsabs.harvard.edu/abs/1911AnP...340..898E>), doi:10.1002/andp.19113401005 (<https://doi.org/10.1002/andp.19113401005>) (also in *Collected Papers* Vol. 3, document 23)
16. Einstein, Albert (1915). "Feldgleichungen der Gravitation" [Field Equations of Gravitation]. *Preussische Akademie der Wissenschaften, Sitzungsberichte*: 844–847.

17. Janssen, Michel; Renn, Jürgen (2015). "Arch and scaffold: How Einstein found his field equations" (<https://doi.org/10.1063%2FPT.3.2979>). *Physics Today* (Feature article). **68** (11): 30–36. Bibcode:2015PhT....68k..30J (<https://ui.adsabs.harvard.edu/abs/2015PhT....68k..30J>). doi:10.1063/PT.3.2979 (<https://doi.org/10.1063%2FPT.3.2979>). hdl:11858/00-001M-0000-002A-8ED7-1 (<https://hdl.handle.net/11858%2F00-001M-0000-002A-8ED7-1>).
18. O'Connor, J J; Robertson, E F (May 1996). "General Relativity" ([https://mathshistory.st-andrews.ac.uk/HistTopics/General\\_relativity/](https://mathshistory.st-andrews.ac.uk/HistTopics/General_relativity/)). *MacTutor*. School of Mathematics and Statistics, University of St Andrews, Scotland. Archived ([https://web.archive.org/web/20200712062639/https://mathshistory.st-andrews.ac.uk/HistTopics/General\\_relativity/](https://web.archive.org/web/20200712062639/https://mathshistory.st-andrews.ac.uk/HistTopics/General_relativity/)) from the original on 12 July 2020. Retrieved 22 October 2025.
19. Fraknoi, Andrew; Morrison, David; Wolff, Sidney C. (2022). "24.5 Black Holes". *Astronomy 2e* (<https://assets.openstax.org/oscms-prodcms/media/documents/Astronomy2e-WEB.pdf>) (PDF) (2e ed.). OpenStax. pp. 839–846. ISBN 978-1-951693-50-3. OCLC 1322188620 (<http://search.worldcat.org/oclc/1322188620>).
20. O'Neill, Ian (10 April 2019). "Black Holes Were Such an Extreme Concept, Even Einstein Had His Doubts" (<https://www.history.com/articles/black-holes-albert-einstein-theory-relativity-space-time>). *History* (published 28 May 2025). Archived (<https://web.archive.org/web/20250813095832/https://www.history.com/articles/black-holes-albert-einstein-theory-relativity-space-time>) from the original on 13 August 2025. Retrieved 22 October 2025.
21. Einstein, Albert (10 May 1939). "On a Stationary System With Spherical Symmetry Consisting of Many Gravitating Masses". *Annals of Mathematics*. **40** (4): 922–936. doi:10.2307/1968902 (<https://doi.org/10.2307%2F1968902>). JSTOR 1968902 (<https://www.jstor.org/stable/1968902>).
22. Levy, Adam (11 January 2021). "How black holes morphed from theory to reality" (<https://knowablemagazine.org/article/physical-world/2021/how-black-holes-morphed-theory-reality>). *Knowable Magazine*. doi:10.1146/knowable-010921-1 (<https://doi.org/10.1146%2Fknowable-010921-1>). Retrieved 25 March 2022.
23. Schwarzschild, K. (1916). "Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie" (<https://archive.org/stream/sitzungsberichte1916deutsch#page/188/mode/2up>) [On the gravitational field of a mass point according to Einstein's theory]. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*. **7**: 189–196. Bibcode:1916SPAW.....189S (<https://ui.adsabs.harvard.edu/abs/1916SPAW.....189S>) – via Internet Archive.
  - Translation: Antoci, S.; Loinger, A. (12 May 1999). "On the gravitational field of a mass point according to Einstein's theory". arXiv:physics/9905030 (<https://arxiv.org/abs/physics/9905030>). and Schwarzschild, K. (1916). "Über das Gravitationsfeld einer Kugel aus inkompressibler Flüssigkeit nach der Einsteinschen Theorie" (<https://archive.org/stream/sitzungsberichte1916deutsch#page/424/mode/2up>). *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*. **18**: 424–434. Bibcode:1916skpa.conf..424S (<https://ui.adsabs.harvard.edu/abs/1916skpa.conf..424S>).
  - Translation: Antoci, S. (1999). "On the gravitational field of a sphere of incompressible fluid according to Einstein's theory". arXiv:physics/9912033 (<https://arxiv.org/abs/physics/9912033>).
24. Droste, J. (1917). "The field of a single centre in Einstein's theory of gravitation, and the motion of a particle in that field" (<http://www.dwc.knaw.nl/DL/publications/PU00012325.pdf>) (PDF). *Physics. Proceedings of the Section of Sciences*. **19** (1). Koninklijke Akademie van Wetenschappen: 197–215. Bibcode:1917KNAB...19..197D (<https://ui.adsabs.harvard.edu/abs/1917KNAB...19..197D>). Archived (<https://web.archive.org/web/20130518034708/http://www.dwc.knaw.nl/DL/publications/PU00012325.pdf>) (PDF) from the original on 18 May 2013. Retrieved 16 September 2012.

25. Kox, A. J. (1992). "General Relativity in the Netherlands: 1915–1920" ([https://books.google.com/books?id=vDHCF\\_3vIhUC&pg=PA41](https://books.google.com/books?id=vDHCF_3vIhUC&pg=PA41)). In Eisenstaedt, Jean; Kox, A. J. (eds.). *Studies in the history of general relativity*. Birkhäuser. p. 41. ISBN 978-0-8176-3479-7. Archived ([https://web.archive.org/web/20160810215219/https://books.google.com/books?id=vDHCF\\_3vIhUC&pg=PA41](https://web.archive.org/web/20160810215219/https://books.google.com/books?id=vDHCF_3vIhUC&pg=PA41)) from the original on 10 August 2016. Retrieved 23 February 2016.
26. 't Hooft, G. (2009). "Introduction to the Theory of Black Holes" ([http://www.phys.uu.nl/~thoof/lectures/blackholes/BH\\_lecturenotes.pdf](http://www.phys.uu.nl/~thoof/lectures/blackholes/BH_lecturenotes.pdf)) (PDF). Institute for Theoretical Physics / Spinoza Institute. pp. 47–48. Archived ([https://web.archive.org/web/20090521082736/http://www.phys.uu.nl/~thoof/lectures/blackholes/BH\\_lecturenotes.pdf](https://web.archive.org/web/20090521082736/http://www.phys.uu.nl/~thoof/lectures/blackholes/BH_lecturenotes.pdf)) (PDF) from the original on 21 May 2009. Retrieved 24 June 2010.
27. Petrescu, Rely Victoria; Aversa, Raffaella; Apicella, Antonio; Petrescu, Florian Ion (11 June 2018). "Nasa Has Found the Most Distant Black Hole" (<https://doi.org/10.3844%2Fjastsp.2018.31.39>). *Journal of Aircraft and Spacecraft Technology* (Review). **2** (1): 31–39. doi:10.3844/jastsp.2018.31.39 (<https://doi.org/10.3844%2Fjastsp.2018.31.39>). eISSN 2523-1200 (<https://search.worldcat.org/issn/2523-1200>). SSRN 3184263 ([https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=3184263](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3184263)).
28. Eddington, Arthur (1926). *The Internal Constitution of the Stars* (<https://books.google.com/books?id=RjC9DpnWFbkC&pg=PA6>). Science. Vol. 52. Cambridge University Press. pp. 233–40. Bibcode:1920Sci....52..233E (<https://ui.adsabs.harvard.edu/abs/1920Sci....52..233E>). doi:10.1126/science.52.1341.233 (<https://doi.org/10.1126%2Fscience.52.1341.233>). PMID 17747682 (<https://pubmed.ncbi.nlm.nih.gov/17747682>). Archived (<https://web.archive.org/web/20160811034409/https://books.google.com/books?id=RjC9DpnWFbkC&pg=PP1&pg=PA6>) from the original on 11 August 2016. ISBN 978-0-521-33708-3
29. Thorne, Kip S.; Hawking, Stephen (1994). Agrawal, Milan (ed.). *Black Holes and Time Warps: Einstein's Outrageous Legacy* (<https://archive.org/details/blackholestimewa0000thor/page/134>) (1st ed.). W. W. Norton & Company. pp. 134 (<https://archive.org/details/blackholestimewa0000thor/page/134>)–135. ISBN 978-0-393-31276-8. Retrieved 12 April 2019. "The first conclusion was the Newtonian version of light not escaping; the second was a semi-accurate, relativistic description; and the third was typical Eddingtonian hyperbole [...] when a star is as small as the critical circumference, the curvature is strong but not infinite, and space is definitely not wrapped around the star. Eddington may have known this, but his description made a good story, and it captured in a whimsical way the spirit of Schwarzschild's spacetime curvature." "
30. Venkataraman, G. (1992). *Chandrasekhar and his limit* (<https://books.google.com/books?id=HNSdDFOJ4wkC&pg=PA89>). Universities Press. p. 89. ISBN 978-81-7371-035-3. Archived (<https://web.archive.org/web/20160811035848/https://books.google.com/books?id=HNSdDFOJ4wkC&pg=PA89>) from the original on 11 August 2016.
31. Detweiler, S. (1981). "Resource letter BH-1: Black holes". *American Journal of Physics* (Paper). **49** (5): 394–400. Bibcode:1981AmJPh..49..394D (<https://ui.adsabs.harvard.edu/abs/1981AmJPh..49..394D>). doi:10.1119/1.12686 (<https://doi.org/10.1119%2F1.12686>).
32. Harpaz, A. (1994). *Stellar evolution* (<https://books.google.com/books?id=kd4VEZv8oo0C&pg=PA105>). A K Peters. p. 105. ISBN 978-1-56881-012-6. Archived (<https://web.archive.org/web/20160811025449/https://books.google.com/books?id=kd4VEZv8oo0C&pg=PA105>) from the original on 11 August 2016.
33. Oppenheimer, J. R.; Volkoff, G. M. (1939). "On Massive Neutron Cores". *Physical Review*. **55** (4): 374–381. Bibcode:1939PhRv...55..374O (<https://ui.adsabs.harvard.edu/abs/1939PhRv...55..374O>). doi:10.1103/PhysRev.55.374 (<https://doi.org/10.1103%2FPhysRev.55.374>).
34. Bombaci, I. (1996). "The Maximum Mass of a Neutron Star". *Astronomy and Astrophysics*. **305**: 871–877. Bibcode:1996A&A...305..871B (<https://ui.adsabs.harvard.edu/abs/1996A&A...305..871B>).

35. Cho, A. (16 February 2018). "A weight limit emerges for neutron stars". *Science* (News). **359** (6377): 724–725. Bibcode:2018Sci...359..724C (<https://ui.adsabs.harvard.edu/abs/2018Sci...359..724C>). doi:10.1126/science.359.6377.724 (<https://doi.org/10.1126%2Fscience.359.6377.724>). PMID 29449468 (<https://pubmed.ncbi.nlm.nih.gov/29449468>).
36. Margalit, B.; Metzger, B. D. (1 December 2017). "Constraining the Maximum Mass of Neutron Stars from Multi-messenger Observations of GW170817" (<https://doi.org/10.3847%2F2041-8213%2Faa991c>). *The Astrophysical Journal Letters* (Article). **850** (2): L19. arXiv:1710.05938 (<https://arxiv.org/abs/1710.05938>). Bibcode:2017ApJ...850L..19M (<https://ui.adsabs.harvard.edu/abs/2017ApJ...850L..19M>). doi:10.3847/2041-8213/aa991c (<https://doi.org/10.3847%2F2041-8213%2Faa991c>). S2CID 119342447 (<https://api.semanticscholar.org/CorpusID:119342447>).
37. Shibata M, Fujibayashi S, Hotokezaka K, Kiuchi K, Kyutoku K, Sekiguchi Y, Tanaka M (22 December 2017). "Modeling GW170817 based on numerical relativity and its implications". *Physical Review D* (Article). **96** (12) 123012. arXiv:1710.07579 (<https://arxiv.org/abs/1710.07579>). Bibcode:2017PhRvD..96i3012S (<https://ui.adsabs.harvard.edu/abs/2017PhRvD..96i3012S>). doi:10.1103/PhysRevD.96.123012 (<https://doi.org/10.1103%2FPhysRevD.96.123012>). S2CID 119206732 (<https://api.semanticscholar.org/CorpusID:119206732>).
38. Ruiz M, Shapiro SL, Tsokaros A (11 January 2018). "GW170817, general relativistic magnetohydrodynamic simulations, and the neutron star maximum mass" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6036631>). *Physical Review D* (Rapid communication). **97** (2) 021501. arXiv:1711.00473 (<https://arxiv.org/abs/1711.00473>). Bibcode:2018PhRvD..97b1501R (<https://ui.adsabs.harvard.edu/abs/2018PhRvD..97b1501R>). doi:10.1103/PhysRevD.97.021501 (<https://doi.org/10.1103%2FPhysRevD.97.021501>). PMC 6036631 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6036631>). PMID 30003183 (<https://pubmed.ncbi.nlm.nih.gov/30003183>).
39. Rezzolla L, Most ER, Weih LR (9 January 2018). "Using Gravitational-wave Observations and Quasi-universal Relations to Constrain the Maximum Mass of Neutron Stars" (<https://doi.org/10.3847%2F2041-8213%2Faaa401>). *The Astrophysical Journal Letters* (Article). **852** (2): L25. arXiv:1711.00314 (<https://arxiv.org/abs/1711.00314>). Bibcode:2018ApJ...852L..25R (<https://ui.adsabs.harvard.edu/abs/2018ApJ...852L..25R>). doi:10.3847/2041-8213/aaa401 (<https://doi.org/10.3847%2F2041-8213%2Faaa401>). S2CID 119359694 (<https://api.semanticscholar.org/CorpusID:119359694>).
40. Ruffini, R.; Wheeler, J. A. (1971). "Introducing the black hole" ([https://web.archive.org/web/20110725133758/http://authors.library.caltech.edu/14972/1/Ruffini2009p1645Phys\\_Today.pdf](https://web.archive.org/web/20110725133758/http://authors.library.caltech.edu/14972/1/Ruffini2009p1645Phys_Today.pdf)) (PDF). *Physics Today* (Feature article). **24** (1): 30–41. Bibcode:1971PhT....24a..30R (<https://ui.adsabs.harvard.edu/abs/1971PhT....24a..30R>). doi:10.1063/1.3022513 (<https://doi.org/10.1063%2F1.3022513>). Archived from the original ([http://authors.library.caltech.edu/14972/1/Ruffini2009p1645Phys\\_Today.pdf](http://authors.library.caltech.edu/14972/1/Ruffini2009p1645Phys_Today.pdf)) (PDF) on 25 July 2011. Retrieved 5 December 2009.
41. Bernstein, Jeremy (2007). "The Reluctant Father of Black Holes" (<https://www.scientificamerican.com/article/the-reluctant-father-of-black-holes-2007-04/>). *Scientific American*. Vol. 17. pp. 4–11. doi:10.1038/scientificamerican0407-4sp (<https://doi.org/10.1038%2Fscientificamerican0407-4sp>). Retrieved 3 August 2023.
42. Oppenheimer, J.R.; Snyder, H. (1939). "On Continued Gravitational Contraction" (<https://doi.org/10.1103%2FPhysRev.56.455>). *Physical Review* (Highlighted article). **56** (5): 455–459. Bibcode:1939PhRv...56..455O (<https://ui.adsabs.harvard.edu/abs/1939PhRv...56..455O>). doi:10.1103/PhysRev.56.455 (<https://doi.org/10.1103%2FPhysRev.56.455>).
43. Finkelstein, D. (1958). "Past-Future Asymmetry of the Gravitational Field of a Point Particle". *Physical Review* (Article). **110** (4): 965–967. Bibcode:1958PhRv..110..965F (<https://ui.adsabs.harvard.edu/abs/1958PhRv..110..965F>). doi:10.1103/PhysRev.110.965 (<https://doi.org/10.1103%2FPhysRev.110.965>).
44. Bartusiak, Marcia (2015). *Black Hole: How an Idea Abandoned by Newtonians, Hated by Einstein, and Gambled On by Hawking Became Loved*. New Haven, CT: Yale University Press. ISBN 978-0-300-21363-8.

45. Luminet, J.-P. (May 1979). "Image of a spherical black hole with thin accretion disk" (<https://ui.adsabs.harvard.edu/abs/1979A&A....75..228L/abstract>). *Astronomy and Astrophysics*. **75**: 228–235. Bibcode:1979A&A....75..228L (<https://ui.adsabs.harvard.edu/abs/1979A&A....75..228L>). ISSN 0004-6361 (<https://search.worldcat.org/issn/0004-6361>).
46. French National Centre for Scientific Research (10 April 2019). "First ever image of a black hole: a CNRS researcher had simulated it as early as 1979" (<https://www.cnrs.fr/en/press/first-ever-image-black-hole-cnrs-researcher-had-simulated-it-early-1979>). CNRS. Retrieved 18 June 2025.
47. Thorne K (2003). "5. Warping spacetime". In Shellard ES, Gibbons GW, Rankin SJ (eds.). *The future of theoretical physics and cosmology: celebrating Stephen Hawking's 60th birthday* (<https://books.google.com/books?id=yLy4b61rfPwC>). Cambridge University Press. p. 74. ISBN 0-521-82081-2.
48. Thorne, Kip (1995). *Black Holes and Time Warps: Einstein's Outrageous Legacy* (Commonwealth Fund Book Program). Commonwealth Fund Book Program Series (1 ed.). Erscheinungsort nicht ermittelbar: W. W. Norton & Company, Incorporated. ISBN 978-0-393-31276-8.
49. Kerr, R. P. (2009). "Discovering the Kerr and Kerr-Schild metrics". In Wiltshire, D. L.; Visser, M.; Scott, S. M. (eds.). *The Kerr Spacetime*. Cambridge University Press. arXiv:0706.1109 (<https://arxiv.org/abs/0706.1109>). Bibcode:2007arXiv0706.1109K (<https://ui.adsabs.harvard.edu/abs/2007arXiv0706.1109K>). ISBN 978-0-521-88512-6.
50. Hooper, Dan (27 December 2019). "The Discovery of Black Holes: From Theory to Actuality" (<https://web.archive.org/web/20220925103539/https://www.wondriumdaily.com/the-discovery-of-black-holes-from-theory-to-actuality/>). *Wondrium Daily*. Virginia, United States: The Teaching Company. Archived from the original (<https://www.wondriumdaily.com/the-discovery-of-black-holes-from-theory-to-actuality/>) on 25 September 2022. Retrieved 29 June 2022.
51. Newman ET, et al. (1965). "Metric of a Rotating, Charged Mass". *Journal of Mathematical Physics* (Research article). **6** (6): 918. Bibcode:1965JMP....6..918N (<https://ui.adsabs.harvard.edu/abs/1965JMP....6..918N>). doi:10.1063/1.1704351 (<https://doi.org/10.1063%2F1.1704351>).
52. Israel, W. (1967). "Event Horizons in Static Vacuum Space-Times". *Physical Review* (Article). **164** (5): 1776. Bibcode:1967PhRv..164.1776I (<https://ui.adsabs.harvard.edu/abs/1967PhRv..164.1776I>). doi:10.1103/PhysRev.164.1776 (<https://doi.org/10.1103%2FPhysRev.164.1776>).
53. Carter, B. (1971). "Axisymmetric Black Hole Has Only Two Degrees of Freedom". Elementary Particles and Fields. *Physical Review Letters* (Letter). **26** (6): 331. Bibcode:1971PhRvL..26..331C (<https://ui.adsabs.harvard.edu/abs/1971PhRvL..26..331C>). doi:10.1103/PhysRevLett.26.331 (<https://doi.org/10.1103%2FPhysRevLett.26.331>).
54. Carter, B. (1977). "The vacuum black hole uniqueness theorem and its conceivable generalisations". *Proceedings of the 1st Marcel Grossmann meeting on general relativity*. pp. 243–254.
55. Chruściel PT, Costa JL, Heusler M (2012). "Stationary Black Holes: Uniqueness and Beyond" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5255892>). *Living Reviews in Relativity* (Article). **15** (7) 7. arXiv:1205.6112 (<https://arxiv.org/abs/1205.6112>). Bibcode:2012LRR....15....7C (<https://ui.adsabs.harvard.edu/abs/2012LRR....15....7C>). doi:10.12942/lrr-2012-7 (<https://doi.org/10.12942%2Flrr-2012-7>). PMC 5255892 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5255892>). PMID 28179837 (<https://pubmed.ncbi.nlm.nih.gov/28179837>).
56. Penrose, Roger (1996). "Chandrasekhar, black holes, and singularities". *Journal of Astrophysics and Astronomy* (Article). **17** (3–4): 213–231. Bibcode:1996JApA...17..213P (<https://ui.adsabs.harvard.edu/abs/1996JApA...17..213P>). doi:10.1007/BF02702305 (<https://doi.org/10.1007%2FBF02702305>).



57. Penrose, R. (1965). "Gravitational Collapse and Space-Time Singularities" (<http://pdfs.semanticscholar.org/faad/1f4358fddf70df2e00c0a290b7e4501c27de.pdf>) (PDF). *Physical Review Letters*. **14** (3): 57. Bibcode:1965PhRvL..14...57P (<https://ui.adsabs.harvard.edu/abs/1965PhRvL..14...57P>). doi:10.1103/PhysRevLett.14.57 (<https://doi.org/10.1103%2FPhysRevLett.14.57>). S2CID 116755736 (<https://api.semanticscholar.org/CorpusID:116755736>). Archived (<https://web.archive.org/web/20201011221750/http://pdfs.semanticscholar.org/faad/1f4358fddf70df2e00c0a290b7e4501c27de.pdf>) (PDF) from the original on 11 October 2020.
58. Ford, L. H. (2003). "The Classical Singularity Theorems and Their Quantum Loopholes". *International Journal of Theoretical Physics*. **42** (6): 1219–1227. arXiv:gr-qc/0301045 (<https://arxiv.org/abs/gr-qc/0301045>). Bibcode:2003gr.qc.....1045F (<https://ui.adsabs.harvard.edu/abs/2003gr.qc.....1045F>). doi:10.1023/A:1025754515197 (<https://doi.org/10.1023%2FA%3A1025754515197>). S2CID 14404560 (<https://api.semanticscholar.org/CorpusID:14404560>).
59. Hawking, Stephen (18 October 1966). "The Occurrence of Singularities in Cosmology" (<http://www.jstor.org/stable/pdf/2415489.pdf>) (PDF). *Proceedings of the Royal Society A*. **294** (1439): 511–521. Bibcode:1966RSPSA.294..511H (<https://ui.adsabs.harvard.edu/abs/1966RSPSA.294..511H>). doi:10.1098/rspa.1966.0221 (<https://doi.org/10.1098%2Frspa.1966.0221>). JSTOR 2415489 (<https://www.jstor.org/stable/2415489>) – via JSTOR.
60. "The Nobel Prize in Physics 2020" (<https://www.nobelprize.org/prizes/physics/2020/summary/>). *NobelPrize.org*. Archived (<https://web.archive.org/web/20210424115309/https://www.nobelprize.org/prizes/physics/2020/summary/>) from the original on 24 April 2021. Retrieved 8 October 2020.
61. Hewish, A.; et al. (1968). "Observation of a Rapidly Pulsating Radio Source". *Nature*. **217** (5130): 709–713. Bibcode:1968Natur.217..709H (<https://ui.adsabs.harvard.edu/abs/1968Natur.217..709H>). doi:10.1038/217709a0 (<https://doi.org/10.1038%2F217709a0>). S2CID 4277613 (<https://api.semanticscholar.org/CorpusID:4277613>).
62. Pilkington, J. D. H.; et al. (1968). "Observations of some further Pulsed Radio Sources". *Nature*. **218** (5137): 126–129. Bibcode:1968Natur.218..126P (<https://ui.adsabs.harvard.edu/abs/1968Natur.218..126P>). doi:10.1038/218126a0 (<https://doi.org/10.1038%2F218126a0>). S2CID 4253103 (<https://api.semanticscholar.org/CorpusID:4253103>).
63. Hewish, A. (1970). "Pulsars". *Annual Review of Astronomy and Astrophysics*. **8** (1): 265–296. Bibcode:1970ARA&A...8..265H (<https://ui.adsabs.harvard.edu/abs/1970ARA&A...8..265H>). doi:10.1146/annurev.aa.08.090170.001405 (<https://doi.org/10.1146%2Fannurev.aa.08.090170.001405>).
64. Boissoneault, Lorraine (28 February 2018). "Fifty Years Ago, a Grad Student's Discovery Changed the Course of Astrophysics" (<https://www.smithsonianmag.com/science-nature/Fifty-years-ago-grad-students-discovery-changed-course-astrophysics-180968288/>). *Smithsonian Magazine*. Retrieved 22 December 2023.
65. Rolston, Bruce (10 November 1997), *The First Black Hole* ([https://web.archive.org/web/20080307181205/http://www.news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](https://web.archive.org/web/20080307181205/http://www.news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)), University of Toronto, archived from the original ([http://news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](http://news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)) on 7 March 2008, retrieved 11 March 2008
66. Shipman, H. L.; Yu, Z; Du, Y.W (1975), "The implausible history of triple star models for Cygnus X-1 Evidence for a black hole", *Astrophysical Letters*, **16** (1): 9–12, Bibcode:1975ApL....16....9S (<https://ui.adsabs.harvard.edu/abs/1975ApL....16....9S>), doi:10.1016/S0304-8853(99)00384-4 (<https://doi.org/10.1016%2FS0304-8853%2899%2900384-4>)

67. Bardeen, J. M.; Carter, B.; Hawking, S. W. (1973). "The four laws of black hole mechanics" (<http://projecteuclid.org/euclid.cmp/1103858973>). *Communications in Mathematical Physics*. **31** (2): 161–170. Bibcode:1973CMaPh..31..161B (<https://ui.adsabs.harvard.edu/abs/1973CMaPh..31..161B>). doi:10.1007/BF01645742 (<https://doi.org/10.1007%2FBF01645742>). MR 0334798 (<https://mathscinet.ams.org/mathscinet-getitem?mr=0334798>). S2CID 54690354 (<https://api.semanticscholar.org/CorpusID:54690354>). Zbl 1125.83309 (<https://zbmath.org/?format=complete&q=an:1125.83309>). Archived (<https://web.archive.org/web/20200516211604/https://projecteuclid.org/euclid.cmp/1103858973>) from the original on 16 May 2020. Retrieved 4 June 2021.
68. Hawking, S. W. (1974). "Black hole explosions?". *Nature*. **248** (5443): 30–31. Bibcode:1974Natur.248...30H (<https://ui.adsabs.harvard.edu/abs/1974Natur.248...30H>). doi:10.1038/248030a0 (<https://doi.org/10.1038%2F248030a0>). S2CID 4290107 (<https://api.semanticscholar.org/CorpusID:4290107>).
69. Remillard, Ronald A.; McClintock, Jeffrey E. (22 September 2006). "X-Ray Properties of Black-Hole Binaries" (<https://www.annualreviews.org/content/journals/10.1146/annurev.astro.44.051905.092532>). *Annual Review of Astronomy and Astrophysics*. **44**: 49–92. arXiv:astro-ph/0606352 (<https://arxiv.org/abs/astro-ph/0606352>). Bibcode:2006ARA&A..44...49R (<https://ui.adsabs.harvard.edu/abs/2006ARA&A..44...49R>). doi:10.1146/annurev.astro.44.051905.092532 (<https://doi.org/10.1146%2Fannurev.astro.44.051905.092532>). ISSN 0066-4146 (<https://search.worldcat.org/issn/0066-4146>).
70. Herbert, Friedman (2002). "From the ionosphere to high energy astronomy – a personal experience". *The Century of Space Science*. Springer. ISBN 0-7923-7196-8.
71. Liu, C. Z.; Li, T. P. (2004). "X-Ray Spectral Variability in Cygnus X-1". *The Astrophysical Journal*. **611** (2): 1084–1090. arXiv:astro-ph/0405246 (<https://arxiv.org/abs/astro-ph/0405246>). Bibcode:2004ApJ...611.1084L (<https://ui.adsabs.harvard.edu/abs/2004ApJ...611.1084L>). doi:10.1086/422209 (<https://doi.org/10.1086%2F422209>). S2CID 208868049 (<https://api.semanticscholar.org/CorpusID:208868049>).
72. Bowyer, S.; Byram, E. T.; Chubb, T. A.; Friedman, H. (1965). "Cosmic X-ray Sources". *Science*. **147** (3656): 394–398. Bibcode:1965Sci...147..394B (<https://ui.adsabs.harvard.edu/abs/1965Sci...147..394B>). doi:10.1126/science.147.3656.394 (<https://doi.org/10.1126%2Fscience.147.3656.394>). PMID 17832788 (<https://pubmed.ncbi.nlm.nih.gov/17832788>). S2CID 206565068 (<https://api.semanticscholar.org/CorpusID:206565068>).
73. Maron, O.; Kijak, J.; Kramer, M.; Wielebinski, R. (2000). "Pulsar spectra of radio emission". *Astronomy and Astrophysics Supplement Series*. **147** (2): 195–203. arXiv:astro-ph/0010233 (<https://arxiv.org/abs/astro-ph/0010233>). Bibcode:2000A&AS..147..195M (<https://ui.adsabs.harvard.edu/abs/2000A&AS..147..195M>). doi:10.1051/aas:2000298 (<https://doi.org/10.1051%2Faas%3A2000298>).
74. Webster, B. Louise; Murdin, Paul (1972). "Cygnus X-1—a Spectroscopic Binary with a Heavy Companion?". *Nature*. **235** (5332): 37–38. Bibcode:1972Natur.235...37W (<https://ui.adsabs.harvard.edu/abs/1972Natur.235...37W>). doi:10.1038/235037a0 (<https://doi.org/10.1038%2F235037a0>). S2CID 4195462 (<https://api.semanticscholar.org/CorpusID:4195462>).
75. Bolton, C. T. (1972). "Identification of Cygnus X-1 with HDE 226868". *Nature*. **235** (5336): 271–273. Bibcode:1972Natur.235..271B (<https://ui.adsabs.harvard.edu/abs/1972Natur.235..271B>). doi:10.1038/235271b0 (<https://doi.org/10.1038%2F235271b0>). S2CID 4222070 (<https://api.semanticscholar.org/CorpusID:4222070>).
76. Rolston, Bruce (10 November 1997). "The First Black Hole" ([https://web.archive.org/web/20080307181205/http://www.news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](https://web.archive.org/web/20080307181205/http://www.news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)). University of Toronto. Archived from the original ([http://news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](http://news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)) on 7 March 2008. Retrieved 11 March 2008.

77. Shipman, H. L.; Yu, Z.; Du, Y. W. (1975). "The implausible history of triple star models for Cygnus X-1 Evidence for a black hole". *Astrophysical Letters*. **16** (1): 9–12. Bibcode:1975ApL....16....9S (<https://ui.adsabs.harvard.edu/abs/1975ApL....16....9S>). doi:10.1016/S0304-8853(99)00384-4 (<https://doi.org/10.1016%2FS0304-8853%2899%2900384-4>).
78. Lynden-Bell, D. (1969). "Galactic Nuclei as Collapsed Old Quasars". *Nature*. **223** (5207): 690–694. Bibcode:1969Natur.223..690L (<https://ui.adsabs.harvard.edu/abs/1969Natur.223..690L>). doi:10.1038/223690a0 (<https://doi.org/10.1038%2F223690a0>).
79. Rees, Martin J. (1984). "Black Hole Models for Active Galactic Nuclei". *Annual Review of Astronomy and Astrophysics*. **22**: 471–506. Bibcode:1984ARA&A..22..471R (<https://ui.adsabs.harvard.edu/abs/1984ARA&A..22..471R>). doi:10.1146/annurev.aa.22.090184.002351 (<https://doi.org/10.1146%2Fannurev.aa.22.090184.002351>).
80. Ferrarese, Laura; Ford, Holland (2005). "Supermassive Black Holes in Galactic Nuclei: Past, Present and Future Research". *Space Science Reviews*. **116** (3–4): 523–624. arXiv:astro-ph/0411247 (<https://arxiv.org/abs/astro-ph/0411247>). Bibcode:2005SSRv..116..523F (<https://ui.adsabs.harvard.edu/abs/2005SSRv..116..523F>). doi:10.1007/s11214-005-3947-6 (<https://doi.org/10.1007%2Fs11214-005-3947-6>).
81. Peterson, Bradley M. (2014). "Measuring the Masses of Supermassive Black Holes". *Space Science Reviews*. **183** (1–4): 253–275. Bibcode:2014SSRv..183..253P (<https://ui.adsabs.harvard.edu/abs/2014SSRv..183..253P>). doi:10.1007/s11214-013-9987-4 (<https://doi.org/10.1007%2Fs11214-013-9987-4>).
82. Merritt, David (1999). "Black holes and galaxy evolution" (<https://archive.org/details/xvthiapmeetingdy0197iapm/page/221>). In Combes, F.; Mamon, G. A.; Charmandaris, V. (eds.). *Dynamics of Galaxies: from the Early Universe to the Present*. Vol. 197. Astronomical Society of the Pacific. pp. 221–232 (<https://archive.org/details/xvthiapmeetingdy0197iapm/page/221>). arXiv:astro-ph/9910546 (<https://arxiv.org/abs/astro-ph/9910546>). Bibcode:2000ASPC..197..221M (<https://ui.adsabs.harvard.edu/abs/2000ASPC..197..221M>). ISBN 978-1-58381-024-8.
83. Tremaine, Scott; Gebhardt, Karl; Bender, Ralf; Bower, Gary; Dressler, Alan; Faber, S. M.; Filippenko, Alexei V.; Green, Richard; Grillmair, Carl; Ho, Luis C.; Kormendy, John; Lauer, Tod R.; Magorrian, John; Pinkney, Jason; Richstone, Douglas (2002). "The Slope of the Black Hole Mass versus Velocity Dispersion Correlation". *The Astrophysical Journal*. **574** (2): 740–753. arXiv:astro-ph/0203468 (<https://arxiv.org/abs/astro-ph/0203468>). Bibcode:2002ApJ...574..740T (<https://ui.adsabs.harvard.edu/abs/2002ApJ...574..740T>). doi:10.1086/341002 (<https://doi.org/10.1086%2F341002>).
84. Ferrarese, Laura; Merritt, David (2000). "A Fundamental Relation between Supermassive Black Holes and Their Host Galaxies". *The Astrophysical Journal*. **539** (1): L9 – L12. arXiv:astro-ph/0006053 (<https://arxiv.org/abs/astro-ph/0006053>). Bibcode:2000ApJ...539L...9F (<https://ui.adsabs.harvard.edu/abs/2000ApJ...539L...9F>). doi:10.1086/312838 (<https://doi.org/10.1086%2F312838>).
85. Nelson, Charles H.; Green, Richard F.; Bower, Gary; Gebhardt, Karl; Weistrop, Donna (2004). "The Relationship Between Black Hole Mass and Velocity Dispersion in Seyfert 1 Galaxies". *The Astrophysical Journal*. **615** (2): 652–661. arXiv:astro-ph/0407383 (<https://arxiv.org/abs/astro-ph/0407383>). Bibcode:2004ApJ...615..652N (<https://ui.adsabs.harvard.edu/abs/2004ApJ...615..652N>). doi:10.1086/424657 (<https://doi.org/10.1086%2F424657>).
86. Genzel, R.; Eckart, A.; Ott, T.; Eisenhauer, F. (1997). "On the nature of the dark mass in the centre of the Milky Way" (<https://doi.org/10.1093%2Fmnras%2F291.1.219>). *Monthly Notices of the Royal Astronomical Society*. **291**: 219–234. doi:10.1093/mnras/291.1.219 (<https://doi.org/10.1093%2Fmnras%2F291.1.219>).

87. Ghez, A. M.; Klein, B. L.; Morris, M.; Becklin, E. E. (1998). "High Proper-Motion Stars in the Vicinity of Sagittarius A\*: Evidence for a Supermassive Black Hole at the Center of Our Galaxy". *The Astrophysical Journal*. **509** (2): 678–686. arXiv:astro-ph/9807210 (<https://arxiv.org/abs/astro-ph/9807210>). Bibcode:1998ApJ...509..678G (<https://ui.adsabs.harvard.edu/abs/1998ApJ...509..678G>). doi:10.1086/306528 (<https://doi.org/10.1086%2F306528>).
88. "Oxford Mathematician Roger Penrose jointly wins the Nobel Prize in Physics" (<https://www.ox.ac.uk/news/2020-10-06-oxford-mathematician-roger-penrose-jointly-wins-nobel-prize-physics>). *University of Oxford*. 6 October 2020. Retrieved 20 October 2025.
89. Poffenberger, Leah. "2020 Nobel Prize in Physics" (<https://www.aps.org/archives/publications/apsnews/202011/nobel-physics.cfm>). *American Physical Society*. Retrieved 20 October 2025.
90. Abbott, B.P.; et al. (2016). "Observation of Gravitational Waves from a Binary Black Hole Merger". *Phys. Rev. Lett.* **116** (6) 061102. arXiv:1602.03837 (<https://arxiv.org/abs/1602.03837>). Bibcode:2016PhRvL.116f1102A (<https://ui.adsabs.harvard.edu/abs/2016PhRvL.116f1102A>). doi:10.1103/PhysRevLett.116.061102 (<https://doi.org/10.1103%2FPhysRevLett.116.061102>). PMID 26918975 (<https://pubmed.ncbi.nlm.nih.gov/26918975>). S2CID 124959784 (<https://api.semanticscholar.org/CorpusID:124959784>).
91. The LIGO Scientific Collaboration and The Virgo Collaboration (2016). "An improved analysis of GW150914 using a fully spin-precessing waveform model". *Physical Review X*. **6** (4) 041014. arXiv:1606.01210 (<https://arxiv.org/abs/1606.01210>). Bibcode:2016PhRvX...6d1014A (<https://ui.adsabs.harvard.edu/abs/2016PhRvX...6d1014A>). doi:10.1103/PhysRevX.6.041014 (<https://doi.org/10.1103%2FPhysRevX.6.041014>). S2CID 18217435 (<https://api.semanticscholar.org/CorpusID:18217435>).
92. Abbott, Benjamin P.; et al. (LIGO Scientific Collaboration and Virgo Collaboration) (2016). "Observation of Gravitational Waves from a Binary Black Hole Merger". *Phys. Rev. Lett.* **116** (6) 061102. arXiv:1602.03837 (<https://arxiv.org/abs/1602.03837>). Bibcode:2016PhRvL.116f1102A (<https://ui.adsabs.harvard.edu/abs/2016PhRvL.116f1102A>). doi:10.1103/PhysRevLett.116.061102 (<https://doi.org/10.1103%2FPhysRevLett.116.061102>). PMID 26918975 (<https://pubmed.ncbi.nlm.nih.gov/26918975>). S2CID 124959784 (<https://api.semanticscholar.org/CorpusID:124959784>).
  - "Observation of Gravitational Waves from a Binary Black Hole Merger" ([https://www.ligo.caltech.edu/system/media\\_files/binaries/301/original/detection-science-summary.pdf](https://www.ligo.caltech.edu/system/media_files/binaries/301/original/detection-science-summary.pdf)) (PDF). *LIGO Scientific Collaboration*.
93. "Facts" (<https://web.archive.org/web/20170704030142/https://www.ligo.caltech.edu/page/facts>). *LIGO*. Archived from the original (<https://www.ligo.caltech.edu/page/facts>) on 4 July 2017. Retrieved 24 August 2017. "This is equivalent to measuring the distance from Earth to the nearest star to an accuracy smaller than the width of a human hair!" (that is, to *Proxima Centauri* at  $4.0208 \times 10^{13}$  km).
94. "The Nobel Prize in Physics 2017" ([https://www.nobelprize.org/nobel\\_prizes/physics/laureates/2017/press.html](https://www.nobelprize.org/nobel_prizes/physics/laureates/2017/press.html)). Nobel Foundation.
95. Burtnyk, Kimberly (20 March 2025). "LIGO-Virgo-KAGRA Announce the 200th Gravitational Wave Detection of O4!" (<https://www.ligo.caltech.edu/news/ligo20250320>). *LIGO Caltech*. Retrieved 22 October 2025.
96. Event Horizon Telescope, The (2019). "First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole" (<https://doi.org/10.3847%2F2041-8213%2Fab0ec7>). *The Astrophysical Journal*. **875** (1): L1. arXiv:1906.11238 (<https://arxiv.org/abs/1906.11238>). Bibcode:2019ApJ...875L...1E (<https://ui.adsabs.harvard.edu/abs/2019ApJ...875L...1E>). doi:10.3847/2041-8213/ab0ec7 (<https://doi.org/10.3847%2F2041-8213%2Fab0ec7>). S2CID 145906806 (<https://api.semanticscholar.org/CorpusID:145906806>).

97. Bouman, Katherine L.; Johnson, Michael D.; Zoran, Daniel; Fish, Vincent L.; Doeleman, Sheperd S.; Freeman, William T. (2016). "Computational Imaging for VLBI Image Reconstruction". *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*. pp. 913–922. arXiv:1512.01413 (<https://arxiv.org/abs/1512.01413>). doi:10.1109/CVPR.2016.105 (<https://doi.org/10.1109%2FCVPR.2016.105>). hdl:1721.1/103077 (<https://hdl.handle.net/1721.1%2F103077>). ISBN 978-1-4673-8851-1. S2CID 9085016 (<https://api.semanticscholar.org/CorpusID:9085016>).
98. Gardiner, Aidan (12 April 2018). "When a Black Hole Finally Reveals Itself, It Helps to Have Our Very Own Cosmic Reporter – Astronomers announced Wednesday that they had captured the first image of a black hole. The Times's Dennis Overbye answers readers' questions" (<https://ghostarchive.org/archive/20220101/https://www.nytimes.com/2019/04/12/reader-center/black-holes-dennis-overbye.html>). *The New York Times*. Archived from the original (<https://www.nytimes.com/2019/04/12/reader-center/black-holes-dennis-overbye.html>) on 1 January 2022. Retrieved 15 April 2019.
99. "How many telescopes are currently in the EHT network?" (<https://eventhorizontelescope.org/faq/how-many-telescopes-are-currently-network>). *Event Horizon Telescope*. Retrieved 22 October 2025.
100. Lutz, Ota (8 October 2024). "How Scientists Captured the First Image of a Black Hole" (<https://www.jpl.nasa.gov/edu/resources/teachable-moment/how-scientists-captured-the-first-image-of-a-black-hole/>). *NASA.gov*. Retrieved 22 October 2025.
101. Chown, Marcus (12 April 2019). "The M87 image will change our understanding of black holes, but why was the photo so hard to capture?" (<https://www.sciencefocus.com/space/m87-why-the-first-black-hole-photo-was-so-hard-to-capture>). *BBC Science Focus Magazine*. Retrieved 22 October 2025.
102. "Astronomers Reveal First Image of the Black Hole at the Heart of Our Galaxy" (<https://eventhorizontelescope.org/blog/astronomers-reveal-first-image-black-hole-heart-our-galaxy>). *Event Horizon Telescope*. 12 May 2022.
103. Chakrabarti, Sukanya; Simon, Joshua D.; Craig, Peter A.; Reggiani, Henrique; Brandt, Timothy D.; Guhathakurta, Puragra; Dalba, Paul A.; Kirby, Evan N.; Chang, Philip; Hey, Daniel R.; Savino, Alessandro; Geha, Marla; Thompson, Ian B. (1 July 2023). "A Noninteracting Galactic Black Hole Candidate in a Binary System with a Main-sequence Star" (<https://doi.org/10.3847%2F1538-3881%2Faccf21>). *The Astronomical Journal*. **166** (1): 6. arXiv:2210.05003 (<https://arxiv.org/abs/2210.05003>). Bibcode:2023AJ....166....6C (<https://ui.adsabs.harvard.edu/abs/2023AJ....166....6C>). doi:10.3847/1538-3881/accf21 (<https://doi.org/10.3847%2F1538-3881%2Faccf21>). ISSN 0004-6256 (<https://search.worldcat.org/issn/0004-6256>).
104. El-Badry, Kareem; Rix, Hans-Walter; Quataert, Eliot; Howard, Andrew W.; Isaacson, Howard; Fuller, Jim; Hawkins, Keith; Breivik, Katelyn; Wong, Kaze W. K.; Rodriguez, Antonio C.; Conroy, Charlie; Shahaf, Sahar; Mazeh, Tsevi; Arenou, Frédéric; Burdge, Kevin B.; Bashi, Dolev; Faigler, Simchon; Weisz, Daniel R.; Seeburger, Rhys; Monter, Silvia Almada; Wojno, Jennifer (2023). "A Sun-like star orbiting a black hole" (<https://doi.org/10.1093%2Fmnras%2Fstac3140>). *Monthly Notices of the Royal Astronomical Society*. **518** (1): 1057–1085. arXiv:2209.06833 (<https://arxiv.org/abs/2209.06833>). Bibcode:2023MNRAS.518.1057E (<https://ui.adsabs.harvard.edu/abs/2023MNRAS.518.1057E>). doi:10.1093/mnras/stac3140 (<https://doi.org/10.1093%2Fmnras%2Fstac3140>).
105. El-Badry, Kareem; Rix, Hans-Walter; et al. (2 November 2022). "A Sun-like star orbiting a black hole" (<https://doi.org/10.1093%2Fmnras%2Fstac3140>). *Monthly Notices of the Royal Astronomical Society*. **518** (1): 1057–1085. arXiv:2209.06833 (<https://arxiv.org/abs/2209.06833>). Bibcode:2023MNRAS.518.1057E (<https://ui.adsabs.harvard.edu/abs/2023MNRAS.518.1057E>). doi:10.1093/mnras/stac3140 (<https://doi.org/10.1093%2Fmnras%2Fstac3140>).
106. "Astronomers Discover Closest Black Hole to Earth" (<https://noirlab.edu/public/news/noirlab2227/>). *noirlab.edu*. NOIRLab. 4 November 2022. Retrieved 4 November 2022.

107. El-Badry, Kareem; Rix, Hans-Walter; Cendes, Yvette; Rodriguez, Antonio C.; Conroy, Charlie; Quataert, Eliot; Hawkins, Keith; Zari, Eleonora; Hobson, Melissa; Breivik, Katelyn; Rau, Arne; Berger, Edo; Shahaf, Sahar; Seeburger, Rhys; Burdge, Kevin B. (1 February 2023). "A red giant orbiting a black hole" (<https://doi.org/10.1093%2Fmnras%2Fstad799>). *Monthly Notices of the Royal Astronomical Society*. **521** (3): 4323–4348. arXiv:2302.07880 (<https://arxiv.org/abs/2302.07880>). Bibcode:2023MNRAS.521.4323E (<https://ui.adsabs.harvard.edu/abs/2023MNRAS.521.4323E>). doi:10.1093/mnras/stad799 (<https://doi.org/10.1093%2Fmnras%2Fstad799>).
108. Panuzzo, P.; Mazeh, T.; Arenou, F.; Holl, B.; Caffau, E. (2024). "Discovery of a dormant 33 solar-mass black hole in pre-release Gaia astrometry" (<https://doi.org/10.1051%2F0004-6361%2F202449763>). *The Astrophysical Journal*. **686**: L2. arXiv:2404.10486 (<https://arxiv.org/abs/2404.10486>). Bibcode:2024A&A...686L...2G (<https://ui.adsabs.harvard.edu/abs/2024A&A...686L...2G>). doi:10.1051/0004-6361/202449763 (<https://doi.org/10.1051%2F0004-6361%2F202449763>).
109. "Pioneering Physicist John Wheeler Dies at 96" (<https://www.scientificamerican.com/article/pioneering-physicist-john-wheeler-dies/>). *Scientific American*. Archived (<https://web.archive.org/web/20161128050759/https://www.scientificamerican.com/article/pioneering-physicist-john-wheeler-dies/>) from the original on 28 November 2016. Retrieved 27 November 2016.
110. Overbye, Dennis (14 April 2008). "John A. Wheeler, Physicist Who Coined the Term 'Black Hole,' Is Dead at 96" (<https://www.nytimes.com/2008/04/14/science/14wheeler.html>). *The New York Times*. Archived (<https://web.archive.org/web/20161122210005/http://www.nytimes.com/2008/04/14/science/14wheeler.html>) from the original on 22 November 2016. Retrieved 27 November 2016.
111. Frolov, Valeri P.; Zelnikov, Andrei (1 December 2011). *Introduction to Black Hole Physics* (1st ed.). Oxford University Press. p. 1. ISBN 978-0-19-969229-3.
112. Booth, Ivan (2005). "Black-hole boundaries". *Canadian Journal of Physics*. **83** (11): 1073–1099. arXiv:gr-qc/0508107 (<https://arxiv.org/abs/gr-qc/0508107>). Bibcode:2005CaJPh..83.1073B (<https://ui.adsabs.harvard.edu/abs/2005CaJPh..83.1073B>). doi:10.1139/p05-063 (<https://doi.org/10.1139%2Fp05-063>).
113. Curiel, Erik (2019). "The many definitions of a black hole". *Nature Astronomy*. **3**: 27–34. arXiv:1808.01507 (<https://arxiv.org/abs/1808.01507>). Bibcode:2019NatAs...3...27C (<https://ui.adsabs.harvard.edu/abs/2019NatAs...3...27C>). doi:10.1038/s41550-018-0602-1 (<https://doi.org/10.1038%2Fs41550-018-0602-1>).
114. Hamilton, Andrew. "A Black Hole is a Waterfall of Space" (<https://jila.colorado.edu/~ajsh/insidebh/waterfall.html>). *Inside Black Holes*. Archived (<https://web.archive.org/web/20250820063124/https://jila.colorado.edu/%7Eajsh/insidebh/waterfall.html>) from the original on 20 August 2025. Retrieved 24 October 2025.
115. Susskind, Leonard (2008). *The black hole war: my battle with Stephen Hawking to make the world safe for quantum mechanics* (1st ed.). New York: Little, Brown. ISBN 978-0-316-01640-7. OCLC 181603165 (<https://search.worldcat.org/oclc/181603165>).
116. **Cite error: The named reference `waterfall` was invoked but never defined (see the help page).**
117. Xu, Hao; Ong, Yen Chin; Yung, Man-Hong (2020). "Cosmic censorship and the evolution of  $d$ -dimensional charged evaporating black holes". *Physical Review D*. **101** (6) 064015. arXiv:1911.11990 (<https://arxiv.org/abs/1911.11990>). Bibcode:2020PhRvD.101f4015X (<https://ui.adsabs.harvard.edu/abs/2020PhRvD.101f4015X>). doi:10.1103/PhysRevD.101.064015 (<https://doi.org/10.1103%2FPhysRevD.101.064015>).



118. Reynolds, Christopher S. (January 2019). "Observing black holes spin" (<https://www.nature.com/articles/s41550-018-0665-z>). *Nature Astronomy*. **3** (1): 41–47. arXiv:1903.11704 (<https://arxiv.org/abs/1903.11704>). Bibcode:2019NatAs...3...41R (<https://ui.adsabs.harvard.edu/abs/2019NatAs...3...41R>). doi:10.1038/s41550-018-0665-z (<https://doi.org/10.1038/s41550-018-0665-z>). ISSN 2397-3366 (<https://search.worldcat.org/issn/2397-3366>). S2CID 85543351 (<https://api.semanticscholar.org/CorpusID:85543351>). Archived (<https://web.archive.org/web/20201118092121/https://www.nature.com/articles/s41550-018-0665-z>) from the original on 18 November 2020. Retrieved 21 August 2020.
119. Steiner, James F.; McClintock, Jeffrey E.; Remillard, Ronald A.; Gou, Lijun; Yamada, Shin'ya; Narayan, Ramesh (2010). "The Constant Inner-Disk Radius of LMC X-3: A Basis for Measuring Black Hole Spin". *The Astrophysical Journal*. **718** (2): L117 – L121. arXiv:1006.5729 (<https://arxiv.org/abs/1006.5729>). Bibcode:2010ApJ...718L.117S (<https://ui.adsabs.harvard.edu/abs/2010ApJ...718L.117S>). doi:10.1088/2041-8205/718/2/L117 (<https://doi.org/10.1088/2041-8205/718/2/L117>).
120. Chatziioannou, Katerina; Lovelace, Geoffrey; Boyle, Michael; Giesler, Matthew; Hemberger, Daniel A.; Katebi, Reza; Kidder, Lawrence E.; Pfeiffer, Harald P.; Scheel, Mark A.; Szilágyi, Béla (2018). "Measuring the properties of nearly extremal black holes with gravitational waves". *Physical Review D*. **98** (4) 044028. arXiv:1804.03704 (<https://arxiv.org/abs/1804.03704>). Bibcode:2018PhRvD..98d4028C (<https://ui.adsabs.harvard.edu/abs/2018PhRvD..98d4028C>). doi:10.1103/PhysRevD.98.044028 (<https://doi.org/10.1103/PhysRevD.98.044028>).
121. Poisson, Eric (1996). "Measuring black-hole parameters and testing general relativity using gravitational-wave data from space-based interferometers". *Physical Review D*. **54** (10): 5939–5953. arXiv:gr-qc/9606024 (<https://arxiv.org/abs/gr-qc/9606024>). Bibcode:1996PhRvD..54.5939P (<https://ui.adsabs.harvard.edu/abs/1996PhRvD..54.5939P>). doi:10.1103/PhysRevD.54.5939 (<https://doi.org/10.1103/PhysRevD.54.5939>). PMID 10020601 (<https://pubmed.ncbi.nlm.nih.gov/10020601>).
122. Echeverria, Fernando (1989). "Gravitational-wave measurements of the mass and angular momentum of a black hole". *Physical Review D*. **40** (10): 3194–3203. Bibcode:1989PhRvD..40.3194E (<https://ui.adsabs.harvard.edu/abs/1989PhRvD..40.3194E>). doi:10.1103/PhysRevD.40.3194 (<https://doi.org/10.1103/PhysRevD.40.3194>). PMID 10011687 (<https://pubmed.ncbi.nlm.nih.gov/10011687>).
123. Anderson, Warren G. (1996). "The Black Hole Information Loss Problem" ([https://web.archive.org/web/2009012223839/http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/info\\_loss.html](https://web.archive.org/web/2009012223839/http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/info_loss.html)). *Usenet Physics FAQ*. Archived from the original ([http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/info\\_loss.html](http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/info_loss.html)) on 22 January 2009. Retrieved 24 March 2009.
124. Preskill, J. (21 October 1994). *Black holes and information: A crisis in quantum physics* ([http://web.archive.org/web/20080518054438/http://www.theory.caltech.edu/~preskill/talks/black\\_holes.pdf](http://web.archive.org/web/20080518054438/http://www.theory.caltech.edu/~preskill/talks/black_holes.pdf)) (PDF). Caltech Theory Seminar. Archived from the original (<http://www.theory.caltech.edu/~preskill/talks/blackholes.pdf>) (PDF) on 18 May 2008. Retrieved 17 May 2009.
125. Danielsson, Ulf H.; Schiffer, Marcelo (1993). "Quantum mechanics, common sense, and the black hole information paradox". *Physical Review D*. **48** (10): 4779–4784. arXiv:gr-qc/9305012 (<https://arxiv.org/abs/gr-qc/9305012>). Bibcode:1993PhRvD..48.4779D (<https://ui.adsabs.harvard.edu/abs/1993PhRvD..48.4779D>). doi:10.1103/PhysRevD.48.4779 (<https://doi.org/10.1103/PhysRevD.48.4779>). PMID 10016132 (<https://pubmed.ncbi.nlm.nih.gov/10016132>).
126. Hawking & Ellis 1973, Appendix B
127. Seeds, Michael A.; Backman, Dana E. (2007). *Perspectives on Astronomy* (<https://books.google.com/books?id=CXom04KGIL8C&pg=PA167>). Cengage Learning. p. 167. ISBN 978-0-495-11352-2. Archived (<https://web.archive.org/web/20160810211808/https://books.google.com/books?id=CXom04KGIL8C&pg=PA167>) from the original on 10 August 2016.

128. Shapiro, S. L.; Teukolsky, S. A. (1983). *Black holes, white dwarfs, and neutron stars: the physics of compact objects*. John Wiley and Sons. p. 357. ISBN 978-0-471-87316-7.
129. Wald, R. M. (1997). "Gravitational Collapse and Cosmic Censorship". In Iyer, B. R.; Bhawal, B. (eds.). *Black Holes, Gravitational Radiation and the Universe*. Dordrecht: Springer. pp. 69–86. arXiv:gr-qc/9710068 (<https://arxiv.org/abs/gr-qc/9710068>). doi:10.1007/978-94-017-0934-7 (<https://doi.org/10.1007%2F978-94-017-0934-7>). ISBN 978-94-017-0934-7.
130. Berger, B. K. (2002). "Numerical Approaches to Spacetime Singularities" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5256073>). *Living Reviews in Relativity*. **5** (1) 1: 2002–1. arXiv:gr-qc/0201056 (<https://arxiv.org/abs/gr-qc/0201056>). Bibcode:2002LRR.....5....1B (<http://ui.adsabs.harvard.edu/abs/2002LRR.....5....1B>). doi:10.12942/lrr-2002-1 (<https://doi.org/10.12942%2Flrr-2002-1>). PMC 5256073 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5256073>). PMID 28179859 (<https://pubmed.ncbi.nlm.nih.gov/28179859>).
131. Joshi, Pankaj S. (1 February 2009). "Naked Singularities". *Scientific American*. Vol. 300, no. 2. pp. 36–43. JSTOR 26001219 (<https://www.jstor.org/stable/26001219>).
132. Reynolds, Christopher S. (2013). "The spin of supermassive black holes". *Classical and Quantum Gravity*. **30** (24). arXiv:1307.3246 (<https://arxiv.org/abs/1307.3246>). Bibcode:2013CQGra..30x4004R (<https://ui.adsabs.harvard.edu/abs/2013CQGra..30x4004R>). doi:10.1088/0264-9381/30/24/244004 (<https://doi.org/10.1088%2F0264-9381%2F30%2F24%2F244004>).
133. Ghosh, Rajes; Mishra, Akash K.; Sarkar, Sudipta (2021). "Overcharging extremal black holes". *Physical Review D*. **104** (10) 104043. arXiv:2106.10667 (<https://arxiv.org/abs/2106.10667>). Bibcode:2021PhRvD.104j4043G (<https://ui.adsabs.harvard.edu/abs/2021PhRvD.104j4043G>). doi:10.1103/PhysRevD.104.104043 (<https://doi.org/10.1103%2FPhysRevD.104.104043>).
134. Zajaček, Michal; Tursunov, Arman; Eckart, Andreas; Britzen, Silke (2018). "On the charge of the Galactic centre black hole" (<https://doi.org/10.1093%2Fmnras%2Fsty2182>). *Monthly Notices of the Royal Astronomical Society*. **480** (4): 4408–4423. arXiv:1808.07327 (<https://arxiv.org/abs/1808.07327>). doi:10.1093/mnras/sty2182 (<https://doi.org/10.1093%2Fmnras%2Fsty2182>).
135. Gong, Yi; Cao, Zhoujian; Gao, He; Zhang, Bing (2019). "On neutralization of charged black holes" (<https://doi.org/10.1093%2Fmnras%2Fstz1904>). *Monthly Notices of the Royal Astronomical Society*. **488** (2): 2722–2731. arXiv:1907.05239 (<https://arxiv.org/abs/1907.05239>). doi:10.1093/mnras/stz1904 (<https://doi.org/10.1093%2Fmnras%2Fstz1904>).
136. Zakharov, A. F.; De Paolis, F.; Ingrosso, G.; Nucita, A. A. (2005). "Direct measurements of black hole charge with future astrometrical missions". *Astronomy & Astrophysics*. **442** (3): 795–799. arXiv:astro-ph/0505286 (<https://arxiv.org/abs/astro-ph/0505286>). Bibcode:2005A&A...442..795Z (<https://ui.adsabs.harvard.edu/abs/2005A&A...442..795Z>). doi:10.1051/0004-6361:20053432 (<https://doi.org/10.1051%2F0004-6361%3A20053432>).
137. Turimov, Bobur; Boboqambarova, Madina; Ahmedov, Bobomurat; Stuchlík, Zdeněk (2022). "Distinguishable feature of electric and magnetic charged black hole". *The European Physical Journal Plus*. **137** (2) 222. doi:10.1140/epjp/s13360-022-02390-7 (<https://doi.org/10.1140%2Fepjp%2Fs13360-022-02390-7>).
138. "Black Hole Basics" (<https://science.nasa.gov/universe/black-holes/>). NASA.gov. 13 March 2024. Retrieved 25 October 2025.
139. Daly, Ruth A. (2019). "Black Hole Spin and Accretion Disk Magnetic Field Strength Estimates for More Than 750 Active Galactic Nuclei and Multiple Galactic Black Holes" (<https://doi.org/10.3847%2F1538-4357%2Fab35e6>). *The Astrophysical Journal*. **886** (1): 37. arXiv:1905.11319 (<https://arxiv.org/abs/1905.11319>). Bibcode:2019ApJ...886...37D (<https://ui.adsabs.harvard.edu/abs/2019ApJ...886...37D>). doi:10.3847/1538-4357/ab35e6 (<https://doi.org/10.3847%2F1538-4357%2Fab35e6>).

140. Eilon, Ehud; Ori, Amos (2016). "Numerical study of the gravitational shock wave inside a spherical charged black hole". *Physical Review D*. **94** (10) 104060. arXiv:1610.04355 (<https://arxiv.org/abs/1610.04355>). Bibcode:2016PhRvD..94j4060E (<https://ui.adsabs.harvard.edu/abs/2016PhRvD..94j4060E>). doi:10.1103/PhysRevD.94.104060 (<https://doi.org/10.1103%2FPhysRevD.94.104060>).
141. Daly, Ruth A.; Donahue, Megan; o'Dea, Christopher P.; Sebastian, Biny; Haggard, Daryl; Lu, Anan (2024). "New black hole spin values for Sagittarius A\* obtained with the outflow method" (<https://doi.org/10.1093%2Fmnras%2Fstad3228>). *Monthly Notices of the Royal Astronomical Society*. **527**: 428–436. doi:10.1093/mnras/stad3228 (<https://doi.org/10.1093%2Fmnras%2Fstad3228>).
142. Tamburini, Fabrizio; Thidé, Bo; Della Valle, Massimo (2020). "Measurement of the spin of the M87 black hole from its observed twisted light" (<https://doi.org/10.1093%2Fmnrasl%2Fslz176>). *Monthly Notices of the Royal Astronomical Society: Letters*. **492**: L22 – L27. arXiv:1904.07923 (<https://arxiv.org/abs/1904.07923>). doi:10.1093/mnras/slz176 (<https://doi.org/10.1093%2Fmnrasl%2Fslz176>).
143. Bambi, Cosimo; Freese, Katherine; Vagnozzi, Sunny; Visinelli, Luca (2019). "Testing the rotational nature of the supermassive object M87\* from the circularity and size of its first image". *Physical Review D*. **100** (4) 044057. arXiv:1904.12983 (<https://arxiv.org/abs/1904.12983>). Bibcode:2019PhRvD.100d4057B (<https://ui.adsabs.harvard.edu/abs/2019PhRvD.100d4057B>). doi:10.1103/PhysRevD.100.044057 (<https://doi.org/10.1103%2FPhysRevD.100.044057>).
144. Abbott, B. P.; et al. (LIGO Scientific Collaboration and Virgo Collaboration) (1 June 2017). "GW170104: Observation of a 50-Solar-Mass Binary Black Hole Coalescence at Redshift 0.2". *Physical Review Letters*. **118** (22) 221101. arXiv:1706.01812 (<https://arxiv.org/abs/1706.01812>). Bibcode:2017PhRvL.118v1101A (<https://ui.adsabs.harvard.edu/abs/2017PhRvL.118v1101A>). doi:10.1103/PhysRevLett.118.221101 (<https://doi.org/10.1103%2FPhysRevLett.118.221101>). PMID 28621973 (<https://pubmed.ncbi.nlm.nih.gov/28621973>). S2CID 206291714 (<https://api.semanticscholar.org/CorpusID:206291714>).
145. Abbott, B. P.; et al. (LIGO Scientific Collaboration & Virgo Collaboration) (16 October 2017). "GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral". *Physical Review Letters*. **119** (16) 161101. arXiv:1710.05832 (<https://arxiv.org/abs/1710.05832>). Bibcode:2017PhRvL.119p1101A (<https://ui.adsabs.harvard.edu/abs/2017PhRvL.119p1101A>). doi:10.1103/PhysRevLett.119.161101 (<https://doi.org/10.1103%2FPhysRevLett.119.161101>). PMID 29099225 (<https://pubmed.ncbi.nlm.nih.gov/29099225>). S2CID 217163611 (<https://api.semanticscholar.org/CorpusID:217163611>).
146. Cromartie, H. T.; Fonseca, E.; Ransom, S. M.; Demorest, P. B.; Arzoumanian, Z.; Blumer, H.; Brook, P. R.; Decesar, M. E.; Dolch, T.; Ellis, J. A.; Ferdman, R. D.; Ferrara, E. C.; Garver-Daniels, N.; Gentile, P. A.; Jones, M. L.; Lam, M. T.; Lorimer, D. R.; Lynch, R. S.; McLaughlin, M. A.; Ng, C.; Nice, D. J.; Pennucci, T. T.; Spiewak, R.; Stairs, I. H.; Stovall, K.; Swiggum, J. K.; Zhu, W. W. (2019). "Relativistic Shapiro delay measurements of an extremely massive millisecond pulsar". *Nature Astronomy*. **4**: 72–76. arXiv:1904.06759 (<https://arxiv.org/abs/1904.06759>). doi:10.1038/s41550-019-0880-2 (<https://doi.org/10.1038%2Fs41550-019-0880-2>).
147. Drischler, Christian; Han, Sophia; Lattimer, James M.; Prakash, Madappa; Reddy, Sanjay; Zhao, Tianqi (2021). "Limiting masses and radii of neutron stars and their implications". *Physical Review C*. **103** (4) 045808. arXiv:2009.06441 (<https://arxiv.org/abs/2009.06441>). Bibcode:2021PhRvC.103d5808D (<https://ui.adsabs.harvard.edu/abs/2021PhRvC.103d5808D>). doi:10.1103/PhysRevC.103.045808 (<https://doi.org/10.1103%2FPhysRevC.103.045808>).

148. Farr, Will M.; Sravan, Niharika; Cantrell, Andrew; Kreidberg, Laura; Bailyn, Charles D.; Mandel, Ilya; Kalogera, Vicky (2011). "The Mass Distribution of Stellar-Mass Black Holes". *The Astrophysical Journal*. **741** (2): 103. arXiv:1011.1459 (<https://arxiv.org/abs/1011.1459>). Bibcode:2011ApJ...741..103F (<https://ui.adsabs.harvard.edu/abs/2011ApJ...741..103F>). doi:10.1088/0004-637X/741/2/103 (<https://doi.org/10.1088%2F0004-637X%2F741%2F2%2F103>).
149. Carr, Bernard; Kohri, Kazunori; Sendouda, Yuuiti; Yokoyama, Jun'ichi (2021). "Constraints on primordial black holes". *Reports on Progress in Physics*. **84** (11). arXiv:2002.12778 (<https://arxiv.org/abs/2002.12778>). Bibcode:2021RPPh...84k6902C (<https://ui.adsabs.harvard.edu/abs/2021RPPh...84k6902C>). doi:10.1088/1361-6633/ac1e31 (<https://doi.org/10.1088%2F1361-6633%2Fac1e31>). PMID 34874316 (<https://pubmed.ncbi.nlm.nih.gov/34874316>).
150. Nakama, Tomohiro; Yokoyama, Jun'ichi (2019). "Micro black holes formed in the early Universe and their cosmological implications". *Physical Review D*. **99** (6) 061303. arXiv:1811.05049 (<https://arxiv.org/abs/1811.05049>). Bibcode:2019PhRvD..99f1303N (<https://ui.adsabs.harvard.edu/abs/2019PhRvD..99f1303N>). doi:10.1103/PhysRevD.99.061303 (<https://doi.org/10.1103%2FPhysRevD.99.061303>).
151. Scardigli, Fabio (2000). "Gravity coupling from micro-black holes". *Nuclear Physics B - Proceedings Supplements*. **88** (1–3): 291–294. arXiv:hep-th/9907150 (<https://arxiv.org/abs/hep-th/9907150>). Bibcode:2000NuPhS..88..291S (<https://ui.adsabs.harvard.edu/abs/2000NuPhS..88..291S>). doi:10.1016/S0920-5632(00)00788-X (<https://doi.org/10.1016%2FS0920-5632%2800%2900788-X>).
152. Belczynski, Krzysztof; Bulik, Tomasz; Fryer, Chris L.; Ruiters, Ashley; Valsecchi, Francesca; Vink, Jorick S.; Hurley, Jarrod R. (2010). "On the Maximum Mass of Stellar Black Holes". *The Astrophysical Journal*. **714** (2): 1217–1226. arXiv:0904.2784 (<https://arxiv.org/abs/0904.2784>). Bibcode:2010ApJ...714.1217B (<https://ui.adsabs.harvard.edu/abs/2010ApJ...714.1217B>). doi:10.1088/0004-637X/714/2/1217 (<https://doi.org/10.1088%2F0004-637X%2F714%2F2%2F1217>).
153. Fryer, Chris L.; Kalogera, Vassiliki (2001). "Theoretical Black Hole Mass Distributions". *The Astrophysical Journal*. **554** (1): 548–560. arXiv:astro-ph/9911312 (<https://arxiv.org/abs/astro-ph/9911312>). Bibcode:2001ApJ...554..548F (<https://ui.adsabs.harvard.edu/abs/2001ApJ...554..548F>). doi:10.1086/321359 (<https://doi.org/10.1086%2F321359>).
154. Fryer, Chris L. (1999). "Mass Limits for Black Hole Formation". *The Astrophysical Journal*. **522** (1): 413–418. arXiv:astro-ph/9902315 (<https://arxiv.org/abs/astro-ph/9902315>). Bibcode:1999ApJ...522..413F (<https://ui.adsabs.harvard.edu/abs/1999ApJ...522..413F>). doi:10.1086/307647 (<https://doi.org/10.1086%2F307647>).
155. Vink, Jorick S.; Higgins, Erin R.; Sander, Andreas A C.; Sabhahit, Gautham N. (2021). "Maximum black hole mass across cosmic time" (<https://doi.org/10.1093%2Fmnras%2Fstab842>). *Monthly Notices of the Royal Astronomical Society*. **504**: 146–154. arXiv:2010.11730 (<https://arxiv.org/abs/2010.11730>). doi:10.1093/mnras/stab842 (<https://doi.org/10.1093%2Fmnras%2Fstab842>).
156. Coleman Miller, M.; Colbert, E. J. M. (2004). "Intermediate-Mass Black Holes". *International Journal of Modern Physics D*. **13** (1): 1–64. arXiv:astro-ph/0308402 (<https://arxiv.org/abs/astro-ph/0308402>). Bibcode:2004IJMPD..13....1M (<https://ui.adsabs.harvard.edu/abs/2004IJMPD..13....1M>). doi:10.1142/S0218271804004426 (<https://doi.org/10.1142%2FS0218271804004426>).
157. Lupi, A.; Haardt, F.; Dotti, M.; Fiacconi, D.; Mayer, L.; Madau, P. (2016). "Growing massive black holes through supercritical accretion of stellar-mass seeds" (<https://doi.org/10.1093%2Fmnras%2Fstv2877>). *Monthly Notices of the Royal Astronomical Society*. **456** (3): 2993–3003. arXiv:1512.02651 (<https://arxiv.org/abs/1512.02651>). doi:10.1093/mnras/stv2877 (<https://doi.org/10.1093%2Fmnras%2Fstv2877>).

158. Dunn, R. J. H.; Fender, R. P.; Körding, E. G.; Belloni, T.; Cabanac, C. (2010). "A global spectral study of black hole X-ray binaries" (<https://doi.org/10.1111%2Fj.1365-2966.2010.16114.x>). *Monthly Notices of the Royal Astronomical Society*. **403** (1): 61–82. arXiv:0912.0142 (<https://arxiv.org/abs/0912.0142>). Bibcode:2010MNRAS.403...61D (<https://ui.adsabs.harvard.edu/abs/2010MNRAS.403...61D>). doi:10.1111/j.1365-2966.2010.16114.x (<https://doi.org/10.1111%2Fj.1365-2966.2010.16114.x>).
159. Lewin, Walter H. G.; Jan, van Paradijs; van den Heuvel, Edward P.J., eds. (28 January 1997). *X-ray Binaries*. Cambridge University Press. p. 1. ISBN 978-0-521-59934-4.
160. Shao, Yong; Li, Xiang-Dong (2020). "Population Synthesis of Black Hole X-Ray Binaries" (<https://doi.org/10.3847%2F1538-4357%2Faba118>). *The Astrophysical Journal*. **898** (2): 143. arXiv:2006.15961 (<https://arxiv.org/abs/2006.15961>). Bibcode:2020ApJ...898..143S (<https://ui.adsabs.harvard.edu/abs/2020ApJ...898..143S>). doi:10.3847/1538-4357/aba118 (<https://doi.org/10.3847%2F1538-4357%2Faba118>).
161. Broekgaarden, Floor S.; Berger, Edo (2021). "Formation of the First Two Black Hole–Neutron Star Mergers (GW200115 and GW200105) from Isolated Binary Evolution" (<https://doi.org/10.3847%2F2041-8213%2Fac2832>). *The Astrophysical Journal Letters*. **920** (1): L13. arXiv:2108.05763 (<https://arxiv.org/abs/2108.05763>). Bibcode:2021ApJ...920L..13B (<https://ui.adsabs.harvard.edu/abs/2021ApJ...920L..13B>). doi:10.3847/2041-8213/ac2832 (<https://doi.org/10.3847%2F2041-8213%2Fac2832>).
162. Chattopadhyay, Debatri; Stevenson, Simon; Broekgaarden, Floor; Antonini, Fabio; Belczynski, Krzysztof (2022). "Modelling the formation of the first two neutron star–black hole mergers, GW200105 and GW200115: Metallicity, chirp masses, and merger remnant spins" (<https://doi.org/10.1093%2Fmnras%2Fstac1283>). *Monthly Notices of the Royal Astronomical Society*. **513** (4): 5780–5789. doi:10.1093/mnras/stac1283 (<https://doi.org/10.1093%2Fmnras%2Fstac1283>).
163. Ziosi, B. M.; Mapelli, M.; Branchesi, M.; Tormen, G. (2014). "Dynamics of stellar black holes in young star clusters with different metallicities - II. Black hole-black hole binaries" (<https://doi.org/10.1093%2Fmnras%2Fstu824>). *Monthly Notices of the Royal Astronomical Society*. **441** (4): 3703–3717. arXiv:1404.7147 (<https://arxiv.org/abs/1404.7147>). doi:10.1093/mnras/stu824 (<https://doi.org/10.1093%2Fmnras%2Fstu824>).
164. "Sources and Types of Gravitational Waves" (<https://www.ligo.caltech.edu/page/gw-sources>). *LIGO Caltech*. Retrieved 26 October 2025.
165. "Black Hole Types" (<https://science.nasa.gov/universe/black-holes/types>). *NASA.gov*. 21 October 2020. Retrieved 26 October 2025.
166. Caputo, Daniel P.; De Vries, Nathan; Patruno, Alessandro; Portegies Zwart, Simon (2017). "On Estimating the Total Number of Intermediate Mass Black Holes" (<https://doi.org/10.1093%2Fmnras%2Fstw3336>). *Monthly Notices of the Royal Astronomical Society* stw3336. doi:10.1093/mnras/stw3336 (<https://doi.org/10.1093%2Fmnras%2Fstw3336>).
167. Coleman Miller, M.; Hamilton, Douglas P. (2002). "Production of intermediate-mass black holes in globular clusters" (<https://doi.org/10.1046%2Fj.1365-8711.2002.05112.x>). *Monthly Notices of the Royal Astronomical Society*. **330** (1): 232–240. arXiv:astro-ph/0106188 (<https://arxiv.org/abs/astro-ph/0106188>). Bibcode:2002MNRAS.330..232C (<https://ui.adsabs.harvard.edu/abs/2002MNRAS.330..232C>). doi:10.1046/j.1365-8711.2002.05112.x (<https://doi.org/10.1046%2Fj.1365-8711.2002.05112.x>).
168. Rizzuto, Francesco Paolo; Naab, Thorsten; Spurzem, Rainer; Giersz, Mirek; Ostriker, J. P.; Stone, N. C.; Wang, Long; Berczik, Peter; Rampp, M. (2021). "Intermediate mass black hole formation in compact young massive star clusters" (<https://doi.org/10.1093%2Fmnras%2Fstaa3634>). *Monthly Notices of the Royal Astronomical Society*. **501** (4): 5257–5273. arXiv:2008.09571 (<https://arxiv.org/abs/2008.09571>). doi:10.1093/mnras/staa3634 (<https://doi.org/10.1093%2Fmnras%2Fstaa3634>).

169. Fujii, Michiko S.; Wang, Long; Tanikawa, Ataru; Hirai, Yutaka; Saitoh, Takayuki R. (2024). "Simulations predict intermediate-mass black hole formation in globular clusters". *Science*. **384** (6703): 1488–1492. arXiv:2406.06772 (<https://arxiv.org/abs/2406.06772>). Bibcode:2024Sci...384.1488F (<https://ui.adsabs.harvard.edu/abs/2024Sci...384.1488F>). doi:10.1126/science.adi4211 (<https://doi.org/10.1126%2Fscience.adi4211>). PMID 38815090 (<https://pubmed.ncbi.nlm.nih.gov/38815090>).
170. Barai, Paramita; De Gouveia Dal Pino, Elisabete M. (2019). "Intermediate-mass black hole growth and feedback in dwarf galaxies at high redshifts" (<https://doi.org/10.1093%2Fmnras%2Fstz1616>). *Monthly Notices of the Royal Astronomical Society*. **487** (4): 5549–5563. arXiv:1807.04768 (<https://arxiv.org/abs/1807.04768>). doi:10.1093/mnras/stz1616 (<https://doi.org/10.1093%2Fmnras%2Fstz1616>).
171. Beckmann, R. S.; Dubois, Y.; Volonteri, M.; Dong-Páez, C. A.; Trebitsch, M.; Devriendt, J.; Kaviraj, S.; Kimm, T.; Peirani, S. (2023). "Population statistics of intermediate-mass black holes in dwarf galaxies using the newhorizon simulation" (<https://doi.org/10.1093%2Fmnras%2Fstad1544>). *Monthly Notices of the Royal Astronomical Society*. **523** (4): 5610–5623. arXiv:2211.13301 (<https://arxiv.org/abs/2211.13301>). doi:10.1093/mnras/stad1544 (<https://doi.org/10.1093%2Fmnras%2Fstad1544>).
172. Ruiz-Rocha, Krystal; Yelikar, Anjali B.; Lange, Jacob; Gabella, William; Weller, Robert A.; o'Shaughnessy, Richard; Holley-Bockelmann, Kelly; Jani, Karan (2025). "Properties of "Lite" Intermediate-mass Black Hole Candidates in LIGO-Virgo's Third Observing Run" (<https://doi.org/10.3847%2F2041-8213%2Fadc5f8>). *The Astrophysical Journal Letters*. **985** (2): L37. arXiv:2502.17681 (<https://arxiv.org/abs/2502.17681>). Bibcode:2025ApJ...985L..37R (<https://ui.adsabs.harvard.edu/abs/2025ApJ...985L..37R>). doi:10.3847/2041-8213/adc5f8 (<https://doi.org/10.3847%2F2041-8213%2Fadc5f8>).
173. Abbott, R.; et al. (2022). "Search for intermediate-mass black hole binaries in the third observing run of Advanced LIGO and Advanced Virgo". *Astronomy & Astrophysics*. **659**: A84. arXiv:2105.15120 (<https://arxiv.org/abs/2105.15120>). Bibcode:2022A&A...659A..84A (<https://ui.adsabs.harvard.edu/abs/2022A&A...659A..84A>). doi:10.1051/0004-6361/202141452 (<https://doi.org/10.1051%2F0004-6361%2F202141452>).
174. "Black Holes" (<https://www.cfa.harvard.edu/research/topic/black-holes>). *Center for Astrophysics*. Retrieved 26 October 2025.
175. Mezcuá, Mar (2021). "Black Holes". *Encyclopedia of Astrobiology*. pp. 1–8. arXiv:2110.08629 (<https://arxiv.org/abs/2110.08629>). doi:10.1007/978-3-642-27833-4\_5510-1 ([https://doi.org/10.1007%2F978-3-642-27833-4\\_5510-1](https://doi.org/10.1007%2F978-3-642-27833-4_5510-1)). ISBN 978-3-642-27833-4.
176. Broderick, Avery E.; Loeb, Abraham; Narayan, Ramesh (2009). "The Event Horizon of Sagittarius A". *The Astrophysical Journal*. **701** (2): 1357–1366. arXiv:0903.1105 (<https://arxiv.org/abs/0903.1105>). Bibcode:2009ApJ...701.1357B (<https://ui.adsabs.harvard.edu/abs/2009ApJ...701.1357B>). doi:10.1088/0004-637X/701/2/1357 (<https://doi.org/10.1088%2F0004-637X%2F701%2F2%2F1357>).
177. Yuan, Feng; Quataert, Eliot; Narayan, Ramesh (2004). "On the Nature of the Variable Infrared Emission from Sagittarius A". *The Astrophysical Journal*. **606** (2): 894–899. arXiv:astro-ph/0401429 (<https://arxiv.org/abs/astro-ph/0401429>). Bibcode:2004ApJ...606..894Y (<https://ui.adsabs.harvard.edu/abs/2004ApJ...606..894Y>). doi:10.1086/383117 (<https://doi.org/10.1086%2F383117>).
178. Volonteri, Marta (2010). "Formation of supermassive black holes". *The Astronomy and Astrophysics Review*. **18** (3): 279–315. arXiv:1003.4404 (<https://arxiv.org/abs/1003.4404>). Bibcode:2010A&ARv..18..279V (<https://ui.adsabs.harvard.edu/abs/2010A&ARv..18..279V>). doi:10.1007/s00159-010-0029-x (<https://doi.org/10.1007%2Fs00159-010-0029-x>).



179. Choi, Jun-Hwan; Shlosman, Isaac; Begelman, Mitchell C. (2013). "Supermassive Black Hole Formation at High Redshifts Via Direct Collapse: Physical Processes in the Early Stage". *The Astrophysical Journal*. **774** (2): 149. arXiv:1304.1369 (<https://arxiv.org/abs/1304.1369>). Bibcode:2013ApJ...774..149C (<https://ui.adsabs.harvard.edu/abs/2013ApJ...774..149C>). doi:10.1088/0004-637X/774/2/149 (<https://doi.org/10.1088%2F0004-637X%2F774%2F2%2F149>).
180. Natarajan, Priyamvada; Treister, Ezequiel (2009). "Is there an upper limit to black hole masses?" (<https://doi.org/10.1111%2Fj.1365-2966.2008.13864.x>). *Monthly Notices of the Royal Astronomical Society*. **393** (3): 838–845. arXiv:0808.2813 (<https://arxiv.org/abs/0808.2813>). Bibcode:2009MNRAS.393..838N (<https://ui.adsabs.harvard.edu/abs/2009MNRAS.393..838N>). doi:10.1111/j.1365-2966.2008.13864.x (<https://doi.org/10.1111%2Fj.1365-2966.2008.13864.x>).
181. Dullo, Bililign T.; Gil De Paz, Armando; Knapen, Johan H. (2021). "Ultramassive Black Holes in the Most Massive Galaxies:  $M_{\text{BH}}-\sigma$  versus  $M_{\text{BH}}-R_{\text{b}}$ " (<https://doi.org/10.3847%2F1538-4357%2Fabceae>). *The Astrophysical Journal*. **908** (2): 134. arXiv:2012.04471 (<https://arxiv.org/abs/2012.04471>). Bibcode:2021ApJ...908..134D (<https://ui.adsabs.harvard.edu/abs/2021ApJ...908..134D>). doi:10.3847/1538-4357/abceae (<https://doi.org/10.3847%2F1538-4357%2Fabceae>).
182. King, Andrew (February 2016). "How big can a black hole grow?" (<https://doi.org/10.1093%2Fmnrasl%2Fslv186>). *Monthly Notices of the Royal Astronomical Society: Letters*. **456** (1): L109 – L112. arXiv:1511.08502 (<https://arxiv.org/abs/1511.08502>). Bibcode:2016MNRAS.456L.109K (<https://ui.adsabs.harvard.edu/abs/2016MNRAS.456L.109K>). doi:10.1093/mnrasl/slv186 (<https://doi.org/10.1093%2Fmnrasl%2Fslv186>). S2CID 40147275 (<https://api.semanticscholar.org/CorpusID:40147275>).
183. Troser, Jaime (5 May 2014). "Is There a Limit to How Large Black Holes Can Become?" (<https://futurism.com/is-there-a-size-limit-to-how-large-a-black-hole-can-become>). *futurism.com*. Retrieved 27 November 2018.
184. Clery, Daniel (21 December 2015). "Limit to how big black holes can grow is astonishing" (<https://www.science.org/content/article/limit-how-big-black-holes-can-grow-astonishing>). *sciencemag.org*. Retrieved 27 November 2018.
185. "Black holes could grow as large as 50 billion suns before their food crumbles into stars, research shows" (<https://web.archive.org/web/20211025005826/https://www2.le.ac.uk/offices/press/press-releases/2015/december/black-holes-could-grow-as-large-as-50-billion-suns-before-their-food-crumbles-into-stars-research-shows>). University of Leicester. Archived from the original (<https://www2.le.ac.uk/offices/press/press-releases/2015/december/black-holes-could-grow-as-large-as-50-billion-suns-before-their-food-crumbles-into-stars-research-shows>) on 25 October 2021. Retrieved 27 November 2018.
186. Wald 1984, pp. 124–125
187. Saa, Alberto; Santarelli, Raphael (18 July 2011). "Destroying a near-extremal Kerr–Newman black hole". *Physical Review D*. **84** (2) 027501. arXiv:1105.3950 (<https://arxiv.org/abs/1105.3950>). Bibcode:2011PhRvD..84b7501S (<https://ui.adsabs.harvard.edu/abs/2011PhRvD..84b7501S>). doi:10.1103/PhysRevD.84.027501 (<https://doi.org/10.1103%2FPhysRevD.84.027501>). S2CID 118487989 (<https://api.semanticscholar.org/CorpusID:118487989>).
188. Visser, Matt (2007). "The Kerr spacetime: A brief introduction". page 35, Fig. 3. arXiv:0706.0622 (<https://arxiv.org/abs/0706.0622>) [gr-qc (<https://arxiv.org/archive/gr-qc>)].
189. Reynolds, Christopher S. (2019). "Observing black holes spin". *Nature Astronomy*. **3**: 41–47. arXiv:1903.11704 (<https://arxiv.org/abs/1903.11704>). Bibcode:2019NatAs...3...41R (<https://ui.adsabs.harvard.edu/abs/2019NatAs...3...41R>). doi:10.1038/s41550-018-0665-z (<https://doi.org/10.1038%2Fs41550-018-0665-z>).
190. Carroll 2004, Ch. 6.6
191. Carroll 2004, Ch. 6.7

192. Lee, Hyun Kyu; Wijers, R.A.M.J.; Brown, G.E. (2000). "The Blandford–Znajek process as a central engine for a gamma-ray burst". *Physics Reports*. **325** (3): 83–114. arXiv:astro-ph/9906213 (<https://arxiv.org/abs/astro-ph/9906213>). Bibcode:2000PhR...325...83L (<https://ui.adsabs.harvard.edu/abs/2000PhR...325...83L>). doi:10.1016/S0370-1573(99)00084-8 (<https://doi.org/10.1016%2FS0370-1573%2899%2900084-8>).
193. "Black Hole Anatomy" (<https://web.archive.org/web/20250424044622/https://science.nasa.gov/universe/black-holes/anatomy/>). *NASA Science*. 2 August 2022. Archived from the original (<https://science.nasa.gov/universe/black-holes/anatomy/>) on 24 April 2025. Retrieved 13 October 2025.
194. Cunha, Pedro; Eiró, Nelson; Herdeiro, Carlos; Lemos, José (16 March 2020). "Lensing and shadow of a black hole surrounded by a heavy accretion disk" (<https://iopscience.iop.org/article/10.1088/1475-7516/2020/03/035>). *Journey of Cosmology and Astroparticle Physics*. **2020** (3): 035. arXiv:1912.08833 (<https://arxiv.org/abs/1912.08833>). Bibcode:2020JCAP...03..035C (<https://ui.adsabs.harvard.edu/abs/2020JCAP...03..035C>). doi:10.1088/1475-7516/2020/03/035 (<https://doi.org/10.1088%2F1475-7516%2F2020%2F03%2F035>) – via IOPscience.
195. "Researchers clarify dynamics of black hole rotational energy" (<https://phys.org/news/2018-02-dynamics-black-hole-rotational-energy.html>). Archived (<https://web.archive.org/web/20180917105330/https://phys.org/news/2018-02-dynamics-black-hole-rotational-energy.html>) from the original on 17 September 2018. Retrieved 17 September 2018.
196. McClintock, J. E.; Remillard, R. A. (2006). "Black Hole Binaries". In Lewin, W.; van der Klis, M. (eds.). *Compact Stellar X-ray Sources*. p. 157. arXiv:astro-ph/0306213 (<https://arxiv.org/abs/astro-ph/0306213>). Bibcode:2006csxs.book..157M (<https://ui.adsabs.harvard.edu/abs/2006csxs.book..157M>). ISBN 978-0-521-82659-4. section 4.1.5.
197. Abramowicz, Marek A.; Fragile, P. Chris (2013). "Foundations of Black Hole Accretion Disk Theory" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5256006>). *Living Reviews in Relativity*. **16** (1) 1. arXiv:1104.5499 (<https://arxiv.org/abs/1104.5499>). Bibcode:2013LRR....16....1A (<https://ui.adsabs.harvard.edu/abs/2013LRR....16....1A>). doi:10.12942/lrr-2013-1 (<https://doi.org/10.12942%2Flrr-2013-1>). PMC 5256006 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5256006>). PMID 28179840 (<https://pubmed.ncbi.nlm.nih.gov/28179840>).
198. Blaes, Omer (2014). "General Overview of Black Hole Accretion Theory". *Space Science Reviews*. **183** (1–4): 21–41. arXiv:1304.4879 (<https://arxiv.org/abs/1304.4879>). Bibcode:2014SSRv..183...21B (<https://ui.adsabs.harvard.edu/abs/2014SSRv..183...21B>). doi:10.1007/s11214-013-9985-6 (<https://doi.org/10.1007%2Fs11214-013-9985-6>).
199. Page, Don N.; Thorne, Kip S. (1974). "Disk-Accretion onto a Black Hole. Time-Averaged Structure of Accretion Disk". *The Astrophysical Journal*. **191**: 499. Bibcode:1974ApJ...191..499P (<https://ui.adsabs.harvard.edu/abs/1974ApJ...191..499P>). doi:10.1086/152990 (<https://doi.org/10.1086%2F152990>).
200. Lasota, Jean-Pierre (2016). "Black Hole Accretion Discs". *Astrophysics of Black Holes*. Astrophysics and Space Science Library. Vol. 440. pp. 1–60. arXiv:1505.02172 (<https://arxiv.org/abs/1505.02172>). doi:10.1007/978-3-662-52859-4\_1 ([https://doi.org/10.1007%2F978-3-662-52859-4\\_1](https://doi.org/10.1007%2F978-3-662-52859-4_1)). ISBN 978-3-662-52857-0.
201. Beloborodov, A. M. (1998). "Super-Eddington accretion discs around Kerr black holes" (<https://doi.org/10.1046%2Fj.1365-8711.1998.01530.x>). *Monthly Notices of the Royal Astronomical Society*. **297** (3): 739–746. arXiv:astro-ph/9802129 (<https://arxiv.org/abs/astro-ph/9802129>). Bibcode:1998MNRAS.297..739B (<https://ui.adsabs.harvard.edu/abs/1998MNRAS.297..739B>). doi:10.1046/j.1365-8711.1998.01530.x (<https://doi.org/10.1046%2Fj.1365-8711.1998.01530.x>).

202. Bisnovatyi-Kogan, Gennady (2019). "Accretion into Black Hole, and Formation of Magnetically Arrested Accretion Disks" (<https://doi.org/10.3390%2Funiverse5060146>). *Universe*. **5** (6): 146. arXiv:1905.13731 (<https://arxiv.org/abs/1905.13731>). Bibcode:2019Univ....5..146B (<https://ui.adsabs.harvard.edu/abs/2019Univ....5..146B>). doi:10.3390/universe5060146 (<https://doi.org/10.3390%2Funiverse5060146>).
203. Zakharov, A. F.; Repin, S. V. (2002). "Model radiation spectrum for an accretion disk near a rotating black hole". *Astronomy Reports*. **46** (5): 360–365. Bibcode:2002ARep...46..360Z (<https://ui.adsabs.harvard.edu/abs/2002ARep...46..360Z>). doi:10.1134/1.1479423 (<https://doi.org/10.1134%2F1.1479423>).
204. Wang, Zi-Liang (2025). "Exploring the role of accretion disk geometry in shaping black hole shadows". *Physical Review D*. **112** (6) 064052. arXiv:2506.21148 (<https://arxiv.org/abs/2506.21148>). Bibcode:2025PhRvD.112f4052W (<https://ui.adsabs.harvard.edu/abs/2025PhRvD.112f4052W>). doi:10.1103/fhqj-wgcm (<https://doi.org/10.1103%2Ffhqj-wgcm>).
205. Gimeno-Soler, Sergio; Font, José A. (2017). "Magnetised Polish doughnuts revisited". *Astronomy & Astrophysics*. **607**: A68. arXiv:1707.03867 (<https://arxiv.org/abs/1707.03867>). Bibcode:2017A&A...607A..68G (<https://ui.adsabs.harvard.edu/abs/2017A&A...607A..68G>). doi:10.1051/0004-6361/201730935 (<https://doi.org/10.1051%2F0004-6361%2F201730935>).
206. Qian, Lei; Abramowicz, M. A.; Fragile, P. C.; Horák, J.; Machida, M.; Straub, O. (2009). "The Polish doughnuts revisited". *Astronomy & Astrophysics*. **498** (2): 471–477. doi:10.1051/0004-6361/200811518 (<https://doi.org/10.1051%2F0004-6361%2F200811518>).
207. Abramowicz, M.A. (2005). "Super-Eddington Black Hole Accretion". *Growing Black Holes: Accretion in a Cosmological Context*. ESO Astrophysics Symposia. pp. 257–273. doi:10.1007/11403913\_49 ([https://doi.org/10.1007%2F11403913\\_49](https://doi.org/10.1007%2F11403913_49)). ISBN 978-3-540-25275-7.
208. Kishimoto, Makoto; Antonucci, Robert; Blaes, Omer; Lawrence, Andy; Boisson, Catherine; Albrecht, Marcus; Leipski, Christian (2008). "The characteristic blue spectra of accretion disks in quasars as uncovered in the infrared". *Nature*. **454** (7203): 492–494. arXiv:0807.3703 (<https://arxiv.org/abs/0807.3703>). Bibcode:2008Natur.454..492K (<https://ui.adsabs.harvard.edu/abs/2008Natur.454..492K>). doi:10.1038/nature07114 (<https://doi.org/10.1038%2Fnature07114>). PMID 18650919 (<https://pubmed.ncbi.nlm.nih.gov/18650919>).
209. Fukue, Jun; Yokoyama, Takushi (1988). "Color Photographs of an Accretion Disk around a Black Hole". *Publications of the Astronomical Society of Japan*. **40**: 15–24. doi:10.1093/pasj/40.1.15 (<https://doi.org/10.1093%2Fpasj%2F40.1.15>).
210. Bonning, E. W.; Cheng, L.; Shields, G. A.; Salvander, S.; Gebhardt, K. (2007). "Accretion Disk Temperatures and Continuum Colors in QSOs". *The Astrophysical Journal*. **659** (1): 211–217. arXiv:astro-ph/0611263 (<https://arxiv.org/abs/astro-ph/0611263>). Bibcode:2007ApJ...659..211B (<https://ui.adsabs.harvard.edu/abs/2007ApJ...659..211B>). doi:10.1086/510712 (<https://doi.org/10.1086%2F510712>).
211. Thorne, Kip (7 November 2014). *The Science of Interstellar*. W. W. Norton & Company. ISBN 978-0-393-35137-8.
212. James, Oliver; Tunzelmann, Eugénie von; Franklin, Paul; Thorne, Kip S. (2015). "Gravitational lensing by spinning black holes in astrophysics, and in the movie *Interstellar*". *Classical and Quantum Gravity*. **32** (6). arXiv:1502.03808 (<https://arxiv.org/abs/1502.03808>). Bibcode:2015CQGra..32f5001J (<https://ui.adsabs.harvard.edu/abs/2015CQGra..32f5001J>). doi:10.1088/0264-9381/32/6/065001 (<https://doi.org/10.1088%2F0264-9381%2F32%2F6%2F065001>).
213. Guo, Sen; Huang, Yu-Xiang; Cui, Yu-Hao; Han, Yan; Jiang, Qing-Quan; Liang, En-Wei; Lin, Kai (2023). "Unveiling the unconventional optical signatures of regular black holes within accretion disk". *The European Physical Journal C*. **83** (11) 1059. arXiv:2310.20523 (<https://arxiv.org/abs/2310.20523>). Bibcode:2023EPJC...83.1059G (<https://ui.adsabs.harvard.edu/abs/2023EPJC...83.1059G>). doi:10.1140/epjc/s10052-023-12208-0 (<https://doi.org/10.1140%2Fepjc%2Fs10052-023-12208-0>).

214. "NASA scientists identify smallest known black hole" ([https://web.archive.org/web/20081227195554/http://www.eurekalert.org/pub\\_releases/2008-04/nsfc-nsi040108.php](https://web.archive.org/web/20081227195554/http://www.eurekalert.org/pub_releases/2008-04/nsfc-nsi040108.php)) (Press release). Goddard Space Flight Center. 1 April 2008. Archived from the original ([http://www.eurekalert.org/pub\\_releases/2008-04/nsfc-nsi040108.php](http://www.eurekalert.org/pub_releases/2008-04/nsfc-nsi040108.php)) on 27 December 2008. Retrieved 14 March 2009.
215. Ingram, Adam R.; Motta, Sara E. (2019). "A review of quasi-periodic oscillations from black hole X-ray binaries: Observation and theory". *New Astronomy Reviews*. **85** 101524. arXiv:2001.08758 (<https://arxiv.org/abs/2001.08758>). Bibcode:2019NewAR..8501524I (<https://ui.adsabs.harvard.edu/abs/2019NewAR..8501524I>). doi:10.1016/j.newar.2020.101524 (<https://doi.org/10.1016%2Fj.newar.2020.101524>).
216. Wagoner, Robert V.; Silbergleit, Alexander S.; Ortega-Rodríguez, Manuel (2001). "Stable Quasi-periodic Oscillations and Black Hole Properties from Diskoseismology". *The Astrophysical Journal*. **559** (1): L25 – L28. arXiv:astro-ph/0107168 (<https://arxiv.org/abs/astro-ph/0107168>). Bibcode:2001ApJ...559L..25W (<https://ui.adsabs.harvard.edu/abs/2001ApJ...559L..25W>). doi:10.1086/323655 (<https://doi.org/10.1086%2F323655>).
217. Mirabel, I. F.; Rodríguez, L. F. (1999). "Sources of Relativistic Jets in the Galaxy". *Annual Review of Astronomy and Astrophysics*. **37**: 409–443. arXiv:astro-ph/9902062 (<https://arxiv.org/abs/astro-ph/9902062>). Bibcode:1999ARA&A..37..409M (<https://ui.adsabs.harvard.edu/abs/1999ARA&A..37..409M>). doi:10.1146/annurev.astro.37.1.409 (<https://doi.org/10.1146%2Fannurev.astro.37.1.409>).
218. "Relativistic Jets" (<https://nustar.caltech.edu/page/relativistic-jets>). *NuSTAR*. Retrieved 9 November 2025.
219. Bagchi, Joydeep; Vivek, M.; Vikram, Vinu; Hota, Ananda; Biju, K. G.; Sirothia, S. K.; Srianand, Raghunathan; Gopal-Krishna; Jacob, Joe (2014). "Megaparsec Relativistic Jets Launched from an Accreting Supermassive Black Hole in an Extreme Spiral Galaxy". *The Astrophysical Journal*. **788** (2): 174. arXiv:1404.6889 (<https://arxiv.org/abs/1404.6889>). Bibcode:2014ApJ...788..174B (<https://ui.adsabs.harvard.edu/abs/2014ApJ...788..174B>). doi:10.1088/0004-637X/788/2/174 (<https://doi.org/10.1088%2F0004-637X%2F788%2F2%2F174>).
220. Nemmen, R. S.; Georganopoulos, M.; Guiriec, S.; Meyer, E. T.; Gehrels, N.; Sambruna, R. M. (2012). "A Universal Scaling for the Energetics of Relativistic Jets from Black Hole Systems". *Science*. **338** (6113): 1445–1448. arXiv:1212.3343 (<https://arxiv.org/abs/1212.3343>). Bibcode:2012Sci...338.1445N (<https://ui.adsabs.harvard.edu/abs/2012Sci...338.1445N>). doi:10.1126/science.1227416 (<https://doi.org/10.1126%2Fscience.1227416>). PMID 23239730 (<https://pubmed.ncbi.nlm.nih.gov/23239730/>).
221. Blandford, Roger; Meier, David; Readhead, Anthony (2019). "Relativistic Jets from Active Galactic Nuclei". *Annual Review of Astronomy and Astrophysics*. **57**: 467–509. arXiv:1812.06025 (<https://arxiv.org/abs/1812.06025>). Bibcode:2019ARA&A..57..467B (<https://ui.adsabs.harvard.edu/abs/2019ARA&A..57..467B>). doi:10.1146/annurev-astro-081817-051948 (<https://doi.org/10.1146%2Fannurev-astro-081817-051948>).
222. Chen 陈, Yongyun 永云; Gu 顾, Qiusheng 秋生; Fan 樊, Junhui 军辉; Zhou 周, Hongyan 宏岩; Yuan 袁, Yefei 业飞; Gu 顾, Weimin 为民; Wu 吴, Qinwen 庆文; Xiong 熊, Dingrong 定荣; Guo 郭, Xiaotong 晓通; Ding 丁, Nan 楠.; Yu 俞, Xiaoling 效龄 (2021). "The Powers of Relativistic Jets Depend on the Spin of Accreting Supermassive Black Holes" (<https://doi.org/10.3847%2F1538-4357%2Fabf4ff>). *The Astrophysical Journal*. **913** (2): 93. arXiv:2104.04242 (<https://arxiv.org/abs/2104.04242>). Bibcode:2021ApJ...913...93C (<https://ui.adsabs.harvard.edu/abs/2021ApJ...913...93C>). doi:10.3847/1538-4357/abf4ff (<https://doi.org/10.3847%2F1538-4357%2Fabf4ff>).

223. Ghisellini, G.; Haardt, F.; Ceca, R. Della; Volonteri, M.; Sbarrato, T. (2013). "The role of relativistic jets in the heaviest and most active supermassive black holes at high redshift" (<https://doi.org/10.1093%2Fmnras%2Fstt637>). *Monthly Notices of the Royal Astronomical Society*. **432** (4): 2818–2823. doi:10.1093/mnras/stt637 (<https://doi.org/10.1093%2Fmnras%2Fstt637>).
224. Ghisellini, G.; Tavecchio, F.; Maraschi, L.; Celotti, A.; Sbarrato, T. (2014). "The power of relativistic jets is larger than the luminosity of their accretion disks". *Nature*. **515** (7527): 376–378. arXiv:1411.5368 (<https://arxiv.org/abs/1411.5368>). Bibcode:2014Natur.515..376G (<https://ui.adsabs.harvard.edu/abs/2014Natur.515..376G>). doi:10.1038/nature13856 (<https://doi.org/10.1038%2Fnature13856>). PMID 25409827 (<https://pubmed.ncbi.nlm.nih.gov/25409827>).
225. Meier, David L. (2003). "The theory and simulation of relativistic jet formation: Towards a unified model for micro- and macroquasars". *New Astronomy Reviews*. **47** (6–7): 667–672. arXiv:astro-ph/0312048 (<https://arxiv.org/abs/astro-ph/0312048>). Bibcode:2003NewAR..47..667M (<https://ui.adsabs.harvard.edu/abs/2003NewAR..47..667M>). doi:10.1016/S1387-6473(03)00120-9 (<https://doi.org/10.1016%2FS1387-6473%2803%2900120-9>).
226. Blandford, R. D.; Znajek, R. L. (1977). "Electromagnetic extraction of energy from Kerr black holes" (<https://doi.org/10.1093%2Fmnras%2F179.3.433>). *Monthly Notices of the Royal Astronomical Society*. **179** (3): 433. arXiv:astro-ph/0506302 (<https://arxiv.org/abs/astro-ph/0506302>). Bibcode:1977MNRAS.179..433B (<https://ui.adsabs.harvard.edu/abs/1977MNRAS.179..433B>). doi:10.1093/mnras/179.3.433 (<https://doi.org/10.1093%2Fmnras%2F179.3.433>).
227. Penrose, R. (1969). "Gravitational Collapse: The Role of General Relativity". *Rivista del Nuovo Cimento*. **1**: 252–276. Bibcode:1969NCimR...1..252P (<https://ui.adsabs.harvard.edu/abs/1969NCimR...1..252P>).
228. Narayan, Ramesh; McClintock, Jeffrey E.; Tchekhovskoy, Alexander (2014). "Energy Extraction from Spinning Black Holes Via Relativistic Jets". *General Relativity, Cosmology and Astrophysics*. pp. 523–535. arXiv:1303.3004 (<https://arxiv.org/abs/1303.3004>). doi:10.1007/978-3-319-06349-2\_25 ([https://doi.org/10.1007%2F978-3-319-06349-2\\_25](https://doi.org/10.1007%2F978-3-319-06349-2_25)). ISBN 978-3-319-06348-5.
229. McClintock, Jeffrey E.; Narayan, Ramesh; Davis, Shane W.; Gou, Lijun; Kulkarni, Akshay; Orosz, Jerome A.; Penna, Robert F.; Remillard, Ronald A.; Steiner, James F. (2011). "Measuring the spins of accreting black holes". *Classical and Quantum Gravity*. **28** (11). arXiv:1101.0811 (<https://arxiv.org/abs/1101.0811>). Bibcode:2011CQGra..28k4009M (<https://ui.adsabs.harvard.edu/abs/2011CQGra..28k4009M>). doi:10.1088/0264-9381/28/11/114009 (<https://doi.org/10.1088%2F0264-9381%2F28%2F11%2F114009>).
230. Misner, Thorne & Wheeler 1973, Box 25.6
231. Jefremov, Paul I.; Tsupko, Oleg Yu.; Bisnovatyi-Kogan, Gennady S. (2015). "Innermost stable circular orbits of spinning test particles in Schwarzschild and Kerr space-times". *Physical Review D*. **91** (12) 124030. arXiv:1503.07060 (<https://arxiv.org/abs/1503.07060>). Bibcode:2015PhRvD..91l4030J (<https://ui.adsabs.harvard.edu/abs/2015PhRvD..91l4030J>). doi:10.1103/PhysRevD.91.124030 (<https://doi.org/10.1103%2FPhysRevD.91.124030>).
232. Bardeen, James M.; Press, William H.; Teukolsky, Saul A. (1 December 1972). "Rotating Black Holes: Locally Nonrotating Frames, Energy Extraction, and Scalar Synchrotron Radiation". *The Astrophysical Journal*. **178**: 347–370. Bibcode:1972ApJ...178..347B (<https://ui.adsabs.harvard.edu/abs/1972ApJ...178..347B>). doi:10.1086/151796 (<https://doi.org/10.1086%2F151796>).
233. Zhang, Yu-Peng; Wei, Shao-Wen; Guo, Wen-Di; Sui, Tao-Tao; Liu, Yu-Xiao (2018). "Innermost stable circular orbit of spinning particle in charged spinning black hole background". *Physical Review D*. **97** (8) 084056. arXiv:1711.09361 (<https://arxiv.org/abs/1711.09361>). Bibcode:2018PhRvD..97h4056Z (<https://ui.adsabs.harvard.edu/abs/2018PhRvD..97h4056Z>). doi:10.1103/PhysRevD.97.084056 (<https://doi.org/10.1103%2FPhysRevD.97.084056>).

234. Tsupko, O. Yu.; Bisnovatyi-Kogan, G. S.; Jefremov, P. I. (2016). "Parameters of innermost stable circular orbits of spinning test particles: Numerical and analytical calculations". *Gravitation and Cosmology*. **22** (2): 138–147. arXiv:1605.04189 (<https://arxiv.org/abs/1605.04189>). Bibcode:2016GrCo...22..138T (<https://ui.adsabs.harvard.edu/abs/2016GrCo...22..138T>). doi:10.1134/S0202289316020158 (<https://doi.org/10.1134%2FS0202289316020158>).
235. Jefremov, Paul I.; Tsupko, Oleg Yu.; Bisnovatyi-Kogan, Gennady S. (2017). "Spin-induced changes in the parameters of ISCO in Kerr spacetime". *The Fourteenth Marcel Grossmann Meeting*. pp. 3715–3721. doi:10.1142/9789813226609\_0486 ([https://doi.org/10.1142%2F9789813226609\\_0486](https://doi.org/10.1142%2F9789813226609_0486)). ISBN 978-981-322-659-3.
236. "First proof of black hole 'plunging regions' " (<https://www.physics.ox.ac.uk/news/first-proof-black-hole-plunging-regions>). Department of Physics. University of Oxford. 16 May 2024.
237. Mummery, Andrew; Ingram, Adam; Davis, Shane; Fabian, Andrew (June 2024). "Continuum emission from within the plunging region of black hole discs" (<https://doi.org/10.1093%2Fmnras%2Fstae1160>). *Monthly Notices of the Royal Astronomical Society*. **531** (1): 366–386. arXiv:2405.09175 (<https://arxiv.org/abs/2405.09175>). doi:10.1093/mnras/stae1160 (<https://doi.org/10.1093%2Fmnras%2Fstae1160>).
238. Machida, Mami; Matsumoto, Ryoji (2003). "Global Three-dimensional Magnetohydrodynamic Simulations of Black Hole Accretion Disks: X-Ray Flares in the Plunging Region". *The Astrophysical Journal*. **585** (1): 429–442. arXiv:astro-ph/0211240 (<https://arxiv.org/abs/astro-ph/0211240>). Bibcode:2003ApJ...585..429M (<https://ui.adsabs.harvard.edu/abs/2003ApJ...585..429M>). doi:10.1086/346070 (<https://doi.org/10.1086%2F346070>).
239. Prisco, Jacopo (17 May 2024). "Study proves black holes have a 'plunging region,' just as Einstein predicted" (<https://www.cnn.com/2024/05/17/world/black-holes-einstein-plunging-region-scn>). *CNN*.
240. Cramer, Claes R. (1997). "Using the Uncharged Kerr Black Hole as a Gravitational Mirror". *General Relativity and Gravitation*. **29** (4): 445–454. arXiv:gr-qc/9510053 (<https://arxiv.org/abs/gr-qc/9510053>). Bibcode:1997GReGr..29..445C (<https://ui.adsabs.harvard.edu/abs/1997GReGr..29..445C>). doi:10.1023/A:1018878515046 (<https://doi.org/10.1023%2FA%3A1018878515046>). S2CID 9517046 (<https://api.semanticscholar.org/CorpusID:9517046>).
241. Lü, H.; Lyu, Hong-Da (2020). "Schwarzschild black holes have the largest size". *Physical Review D*. **101** (4) 044059. arXiv:1911.02019 (<https://arxiv.org/abs/1911.02019>). Bibcode:2020PhRvD.101d4059L (<https://ui.adsabs.harvard.edu/abs/2020PhRvD.101d4059L>). doi:10.1103/PhysRevD.101.044059 (<https://doi.org/10.1103%2FPhysRevD.101.044059>).
242. Qiao, Chen-Kai (2022). "Curvatures, photon spheres, and black hole shadows". *Physical Review D*. **106** (8) 084060. arXiv:2208.01771 (<https://arxiv.org/abs/2208.01771>). Bibcode:2022PhRvD.106h4060Q (<https://ui.adsabs.harvard.edu/abs/2022PhRvD.106h4060Q>). doi:10.1103/PhysRevD.106.084060 (<https://doi.org/10.1103%2FPhysRevD.106.084060>).
243. Ma, Liang; Lü, H. (2020). "Bounds on photon spheres and shadows of charged black holes in Einstein-Gauss-Bonnet-Maxwell gravity". *Physics Letters B*. **807** 135535. arXiv:1912.05569 (<https://arxiv.org/abs/1912.05569>). Bibcode:2020PhLB..80735535M (<https://ui.adsabs.harvard.edu/abs/2020PhLB..80735535M>). doi:10.1016/j.physletb.2020.135535 (<https://doi.org/10.1016%2Fj.physletb.2020.135535>).
244. Nitta, Daisuke; Chiba, Takeshi; Sugiyama, Naoshi (September 2011). "Shadows of colliding black holes". *Physical Review D*. **84** (6) 063008. arXiv:1106.2425 (<https://arxiv.org/abs/1106.2425>). Bibcode:2011PhRvD..84f3008N (<https://ui.adsabs.harvard.edu/abs/2011PhRvD..84f3008N>). doi:10.1103/PhysRevD.84.063008 (<https://doi.org/10.1103%2FPhysRevD.84.063008>). S2CID 119264596 (<https://api.semanticscholar.org/CorpusID:119264596>).
245. Teo, Edward (2003). "Spherical Photon Orbits Around a Kerr Black Hole" (<http://scholarbank.nus.edu.sg/handle/10635/97985>). *General Relativity and Gravitation*. **35** (11): 1909–1926. Bibcode:2003GReGr..35.1909T (<https://ui.adsabs.harvard.edu/abs/2003GReGr..35.1909T>). doi:10.1023/A:1026286607562 (<https://doi.org/10.1023%2FA%3A1026286607562>).



246. Heydarzade, Yaghoub; Vertogradov, Vitalii (2024). "Dynamical photon spheres in charged black holes and naked singularities". *The European Physical Journal C*. **84** (6) 582. arXiv:2311.08930 (<https://arxiv.org/abs/2311.08930>). Bibcode:2024EPJC...84..582H (<https://ui.adsabs.harvard.edu/abs/2024EPJC...84..582H>). doi:10.1140/epjc/s10052-024-12945-w (<https://doi.org/10.1140%2Fepjc%2Fs10052-024-12945-w>).
247. Chen, Ying-Xuan; Huang, Jia-Hui; Jiang, Haoxiang (2023). "Radii of spherical photon orbits around Kerr-Newman black holes". *Physical Review D*. **107** (4) 044066. arXiv:2210.08509 (<https://arxiv.org/abs/2210.08509>). Bibcode:2023PhRvD.107d4066C (<https://ui.adsabs.harvard.edu/abs/2023PhRvD.107d4066C>). doi:10.1103/PhysRevD.107.044066 (<https://doi.org/10.1103%2FPhysRevD.107.044066>).
248. Misner, Thorne & Wheeler 1973, p. 848
249. Davies, Paul (1992). *The New Physics* (<https://books.google.com/books?id=akb2FpZSGnMC>) (illustrated ed.). Cambridge University Press. p. 26. ISBN 978-0-521-43831-5. Archived (<https://web.archive.org/web/20210817161727/https://books.google.com/books?id=akb2FpZSGnMC>) from the original on 17 August 2021. Retrieved 25 September 2020. Extract of page 26 (<https://books.google.com/books?id=akb2FpZSGnMC&pg=PA26>) Archived (<https://web.archive.org/web/20210815222341/https://books.google.com/books?id=akb2FpZSGnMC&pg=PA26>) 15 August 2021 at the Wayback Machine
250. Fleisch, Daniel; Kregenow, Julia (2013). *A Student's Guide to the Mathematics of Astronomy* (<https://books.google.com/books?id=x4gaBQAAQBAJ>) (illustrated ed.). Cambridge University Press. p. 168. ISBN 978-1-107-03494-5. Archived (<https://web.archive.org/web/20210817045139/https://books.google.com/books?id=x4gaBQAAQBAJ>) from the original on 17 August 2021. Retrieved 25 September 2020. Extract of page 168 (<https://books.google.com/books?id=x4gaBQAAQBAJ&pg=PA168>) Archived (<https://web.archive.org/web/20210817113029/https://books.google.be/books?id=x4gaBQAAQBAJ&pg=PA168>) 17 August 2021 at the Wayback Machine
251. Wheeler 2007, p. 179
252. Carroll 2004, Ch. 5.4 and 7.3
253. "Singularities and Black Holes > Lightcones and Causal Structure" (<https://plato.stanford.edu/entries/spacetime-singularities/lightcone.html>). *plato.stanford.edu*. Stanford Encyclopedia of Philosophy. Archived (<https://web.archive.org/web/20190517122738/https://plato.stanford.edu/entries/spacetime-singularities/lightcone.html>) from the original on 17 May 2019. Retrieved 11 March 2018.
254. Carroll 2004, p. 217
255. Carroll 2004, p. 218
256. "Inside a black hole" ([https://web.archive.org/web/20090423053437/http://nrumiano.free.fr/Estars/int\\_bh.html](https://web.archive.org/web/20090423053437/http://nrumiano.free.fr/Estars/int_bh.html)). *Knowing the universe and its secrets*. Archived from the original ([http://nrumiano.free.fr/Estars/int\\_bh.html](http://nrumiano.free.fr/Estars/int_bh.html)) on 23 April 2009. Retrieved 26 March 2009.
257. "What happens to you if you fall into a black hole" ([http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/fall\\_in.html](http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/fall_in.html)). *math.ucr.edu*. John Baez. Archived ([https://web.archive.org/web/20190213124648/http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/fall\\_in.html](https://web.archive.org/web/20190213124648/http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/fall_in.html)) from the original on 13 February 2019. Retrieved 11 March 2018.
258. Carroll 2004, p. 222
259. "Watch: Three Ways an Astronaut Could Fall Into a Black Hole" (<https://web.archive.org/web/20190415101947/https://news.nationalgeographic.com/news/2014/01/140130-black-holes-stephen-hawking-work-space-astronomy/>). 1 February 2014. Archived from the original (<https://news.nationalgeographic.com/news/2014/01/140130-black-holes-stephen-hawking-work-space-astronomy/>) on 15 April 2019. Retrieved 13 March 2018.
260. Smarr, L. (1973). "Surface Geometry of Charged Rotating Black Holes". *Physical Review D*. **7** (2): 289–295. Bibcode:1973PhRvD...7..289S (<https://ui.adsabs.harvard.edu/abs/1973PhRvD...7..289S>). doi:10.1103/PhysRevD.7.289 (<https://doi.org/10.1103%2FPhysRevD.7.289>).

261. Visser, M. (22 January 2009). "The Kerr spacetime: A brief introduction". In Wiltshire, D.L.; Visser, M.; Scott, S.M. (eds.). *Horizon geometry for Kerr black holes with synchronized hair* ([https://books.google.com/books?id=wymJBq\\_80Q0C](https://books.google.com/books?id=wymJBq_80Q0C)). Vol. 97. Cambridge University Press. arXiv:0706.0622 (<https://arxiv.org/abs/0706.0622>). Bibcode:2018PhRvD..97l4012D (<https://ui.adsabs.harvard.edu/abs/2018PhRvD..97l4012D>). doi:10.1103/PhysRevD.97.124012 (<https://doi.org/10.1103%2FPhysRevD.97.124012>). ISBN 978-0-521-88512-6. Archived ([https://web.archive.org/web/20200520134643/https://books.google.com/books?id=wymJBq\\_80Q0C](https://web.archive.org/web/20200520134643/https://books.google.com/books?id=wymJBq_80Q0C)) from the original on 20 May 2020. Retrieved 12 January 2020. `{{cite book}}: |journal= ignored (help)`
262. Delgado, J.F. M.; Herdeiro, C.A. R.; Radu, E. (2018). "Horizon geometry for Kerr black holes with synchronized hair". *Physical Review D*. **97** (12) 124012. arXiv:1804.04910 (<https://arxiv.org/abs/1804.04910>). Bibcode:2018PhRvD..97l4012D (<https://ui.adsabs.harvard.edu/abs/2018PhRvD..97l4012D>). doi:10.1103/PhysRevD.97.124012 (<https://doi.org/10.1103%2FPhysRevD.97.124012>). hdl:10773/24121 (<https://hdl.handle.net/10773%2F24121>). S2CID 55732213 (<https://api.semanticscholar.org/CorpusID:55732213>).
263. Hod, Shahar; Piran, Tsvi (1998). "The Inner Structure of Black Holes". *General Relativity and Gravitation*. **30** (11): 1555–1562. arXiv:gr-qc/9902008 (<https://arxiv.org/abs/gr-qc/9902008>). Bibcode:1998GReGr..30.1555H (<https://ui.adsabs.harvard.edu/abs/1998GReGr..30.1555H>). doi:10.1023/A:1026654519980 (<https://doi.org/10.1023%2FA%3A1026654519980>).
264. Poisson, Eric; Israel, Werner (1990). "Internal structure of black holes". *Physical Review D*. **41** (6): 1796–1809. Bibcode:1990PhRvD..41.1796P (<https://ui.adsabs.harvard.edu/abs/1990PhRvD..41.1796P>). doi:10.1103/PhysRevD.41.1796 (<https://doi.org/10.1103%2FPhysRevD.41.1796>). PMID 10012548 (<https://pubmed.ncbi.nlm.nih.gov/10012548>).
265. Scheel, M. A.; Thorne, K. S. (2014). "Geometrodynamics: The nonlinear dynamics of curved spacetime". *Physics-Uspekhi*. **57** (4): 342–351. arXiv:1706.09078 (<https://arxiv.org/abs/1706.09078>). Bibcode:2014PhyU...57..342S (<https://ui.adsabs.harvard.edu/abs/2014PhyU...57..342S>). doi:10.3367/UFNe.0184.201404b.0367 (<https://doi.org/10.3367%2FUFNe.0184.201404b.0367>).
266. Marolf, Donald; Ori, Amos (2012). "Outgoing gravitational shock wave at the inner horizon: The late-time limit of black hole interiors". *Physical Review D*. **86** (12) 124026. arXiv:1109.5139 (<https://arxiv.org/abs/1109.5139>). Bibcode:2012PhRvD..86l4026M (<https://ui.adsabs.harvard.edu/abs/2012PhRvD..86l4026M>). doi:10.1103/PhysRevD.86.124026 (<https://doi.org/10.1103%2FPhysRevD.86.124026>).
267. Ori, Amos (1991). "Inner structure of a charged black hole: An exact mass-inflation solution". *Physical Review Letters*. **67** (7): 789–792. Bibcode:1991PhRvL..67..789O (<https://ui.adsabs.harvard.edu/abs/1991PhRvL..67..789O>). doi:10.1103/PhysRevLett.67.789 (<https://doi.org/10.1103%2FPhysRevLett.67.789>). PMID 10044989 (<https://pubmed.ncbi.nlm.nih.gov/10044989>).
268. Burko, Lior M. (1997). "Structure of the Black Hole's Cauchy-Horizon Singularity". *Physical Review Letters*. **79** (25): 4958–4961. arXiv:gr-qc/9710112 (<https://arxiv.org/abs/gr-qc/9710112>). Bibcode:1997PhRvL..79.4958B (<https://ui.adsabs.harvard.edu/abs/1997PhRvL..79.4958B>). doi:10.1103/PhysRevLett.79.4958 (<https://doi.org/10.1103%2FPhysRevLett.79.4958>).
269. Burko, Lior M.; Khanna, Gaurav; Zenginoğlu, Anıl (2016). "Cauchy-horizon singularity inside perturbed Kerr black holes". *Physical Review D*. **93** (4) 041501. arXiv:1601.05120 (<https://arxiv.org/abs/1601.05120>). Bibcode:2016PhRvD..93d1501B (<https://ui.adsabs.harvard.edu/abs/2016PhRvD..93d1501B>). doi:10.1103/PhysRevD.93.041501 (<https://doi.org/10.1103%2FPhysRevD.93.041501>).
270. Hamilton, Andrew J. S. (2017). "Mass inflation followed by Belinskii-Khalatnikov-Lifshitz collapse inside accreting, rotating black holes". *Physical Review D*. **96** (8) 084041. arXiv:1703.01921 (<https://arxiv.org/abs/1703.01921>). Bibcode:2017PhRvD..96h4041H (<https://ui.adsabs.harvard.edu/abs/2017PhRvD..96h4041H>). doi:10.1103/PhysRevD.96.084041 (<https://doi.org/10.1103%2FPhysRevD.96.084041>).

271. Barceló, Carlos; Boyanov, Valentin; Carballo-Rubio, Raúl; Garay, Luis J. (2022). "Classical mass inflation versus semiclassical inner horizon inflation". *Physical Review D*. **106** (12) 124006. arXiv:2203.13539 (<https://arxiv.org/abs/2203.13539>). Bibcode:2022PhRvD.106l4006B (<https://ui.adsabs.harvard.edu/abs/2022PhRvD.106l4006B>). doi:10.1103/PhysRevD.106.124006 (<https://doi.org/10.1103%2FPhysRevD.106.124006>).
272. Crowther, Karen; De Haro, Sebastian (29 September 2022). "Four Attitudes Towards Singularities in the Search for a Theory of Quantum Gravity". In Vassallo, Antonio (ed.). *The Foundations of Spacetime Physics* (<https://philarchive.org/archive/CROFAT-5>). New York. arXiv:2208.05946 (<https://arxiv.org/abs/2208.05946>). doi:10.4324/9781003219019 (<https://doi.org/10.4324%2F9781003219019>). ISBN 978-1-003-21901-9.
273. Alesci, Emanuele; Bahrami, Sina; Pranzetti, Daniele (2019). "Quantum gravity predictions for black hole interior geometry". *Physics Letters B*. **797** 134908. arXiv:1904.12412 (<https://arxiv.org/abs/1904.12412>). Bibcode:2019PhLB..79734908A (<https://ui.adsabs.harvard.edu/abs/2019PhLB..79734908A>). doi:10.1016/j.physletb.2019.134908 (<https://doi.org/10.1016%2Fj.physletb.2019.134908>).
274. Koshelev, Alexey S.; Tokareva, Anna (2025). "Nonperturbative quantum gravity denounces singular black holes" (<https://doi.org/10.1103%2FPhysRevD.111.086026>). *Physical Review D*. **111** (8) 086026. Bibcode:2025PhRvD.111h6026K (<https://ui.adsabs.harvard.edu/abs/2025PhRvD.111h6026K>). doi:10.1103/PhysRevD.111.086026 (<https://doi.org/10.1103%2FPhysRevD.111.086026>).
275. Olmedo, Javier; Saini, Sahil; Singh, Parampreet (2017). "From black holes to white holes: A quantum gravitational, symmetric bounce". *Classical and Quantum Gravity*. **34** (22). arXiv:1707.07333 (<https://arxiv.org/abs/1707.07333>). Bibcode:2017CQGra..34v5011O (<https://ui.adsabs.harvard.edu/abs/2017CQGra..34v5011O>). doi:10.1088/1361-6382/aa8da8 (<https://doi.org/10.1088%2F1361-6382%2Faa8da8>).
276. Misner, Charles W. (1969). "Absolute Zero of Time". *Physical Review*. **186** (5): 1328–1333. Bibcode:1969PhRv..186.1328M (<https://ui.adsabs.harvard.edu/abs/1969PhRv..186.1328M>). doi:10.1103/PhysRev.186.1328 (<https://doi.org/10.1103%2FPhysRev.186.1328>).
277. Doran, Rosa; Lobo, Francisco S. N.; Crawford, Paulo (2008). "Interior of a Schwarzschild Black Hole Revisited". *Foundations of Physics*. **38** (2): 160–187. arXiv:gr-qc/0609042 (<https://arxiv.org/abs/gr-qc/0609042>). Bibcode:2008FoPh...38..160D (<https://ui.adsabs.harvard.edu/abs/2008FoPh...38..160D>). doi:10.1007/s10701-007-9197-6 (<https://doi.org/10.1007%2Fs10701-007-9197-6>).
278. Carroll 2004, p. 205
279. Hawking, S. W.; Penrose, R. (1970). "The singularities of gravitational collapse and cosmology". *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*. **314** (1519): 529–548. Bibcode:1970RSPSA.314..529H (<https://ui.adsabs.harvard.edu/abs/1970RSPSA.314..529H>). doi:10.1098/rspa.1970.0021 (<https://doi.org/10.1098%2Frspa.1970.0021>).
280. Carroll 2004, pp. 264–265
281. Carroll 2004, p. 252
282. "Sizes of Black Holes? How Big is a Black Hole?" (<https://www.skyandtelescope.com/astronomy-resources/how-big-is-a-black-hole/>). *Sky & Telescope*. 22 July 2014. Archived (<https://web.archive.org/web/20190403035741/https://www.skyandtelescope.com/astronomy-resources/how-big-is-a-black-hole/>) from the original on 3 April 2019. Retrieved 9 October 2018.

283. Lewis, G. F.; Kwan, J. (2007). "No Way Back: Maximizing Survival Time Below the Schwarzschild Event Horizon" (<https://www.cambridge.org/core/journals/publications-of-the-astronomical-society-of-australia/article/no-way-back-maximizing-survival-time-below-the-schwarzschild-event-horizon/2A1CCF5CB13E7BEFA6441B3038C635A3>). *Publications of the Astronomical Society of Australia*. **24** (2): 46–52. arXiv:0705.1029 (<https://arxiv.org/abs/0705.1029>). Bibcode:2007PASA...24...46L (<https://ui.adsabs.harvard.edu/abs/2007PASA...24...46L>). doi:10.1071/AS07012 (<https://doi.org/10.1071%2FAS07012>). S2CID 17261076 (<https://api.semanticscholar.org/CorpusID:17261076>).
284. Toporensky, Alexei; Popov, Sergei (2023). "How to Delay Death and Look Further into the Future if You Fall into a Black Hole" (<https://link.springer.com/article/10.1007/s12045-023-1602-8>). *Resonance*. **28** (5): 737–749. doi:10.1007/s12045-023-1602-8 (<https://doi.org/10.1007%2Fs12045-023-1602-8>).
285. Wheeler 2007, p. 182
286. Belinskii, V.A.; Lifshitz, E.M.; Khalatnikov, I.M.; Agyei, A.K. (1992). "The oscillatory mode of approach to a singularity in homogeneous cosmological models with rotating axes". *Perspectives in Theoretical Physics*. pp. 677–689. doi:10.1016/B978-0-08-036364-6.50048-X (<https://doi.org/10.1016%2FB978-0-08-036364-6.50048-X>). ISBN 978-0-08-036364-6.
287. Garfinkle, David (2007). "Of singularities and breadmaking" ([https://www.einstein-online.info/en/spotlight/singularities\\_bkl/](https://www.einstein-online.info/en/spotlight/singularities_bkl/)). *Einstein Online*. Archived ([https://web.archive.org/web/20250709022836/https://www.einstein-online.info/en/spotlight/singularities\\_bkl/](https://web.archive.org/web/20250709022836/https://www.einstein-online.info/en/spotlight/singularities_bkl/)) from the original on 9 July 2025. Retrieved 14 October 2025.
288. Carroll 2004, pp. 257–259 and 265–266
289. Droz, S.; Israel, W.; Morsink, S. M. (1996). "Black holes: the inside story". *Physics World*. **9** (1): 34–37. Bibcode:1996PhyW....9...34D (<https://ui.adsabs.harvard.edu/abs/1996PhyW....9...34D>). doi:10.1088/2058-7058/9/1/26 (<https://doi.org/10.1088%2F2058-7058%2F9%2F1%2F26>).
290. Carroll 2004, p. 266
291. Thorne, Kip S. (1993). *Closed Timelike Curves* (<https://www.its.caltech.edu/~kip/scripts/ClosedTimelikeCurves-II121.pdf>) (PDF). General relativity and gravitation.
292. Lan, Chen; Yang, Hao; Guo, Yang; Miao, Yan-Gang (2023). "Regular Black Holes: A Short Topic Review". *International Journal of Theoretical Physics*. **62** (9) 202. arXiv:2303.11696 (<https://arxiv.org/abs/2303.11696>). Bibcode:2023IJTP...62..202L (<https://ui.adsabs.harvard.edu/abs/2023IJTP...62..202L>). doi:10.1007/s10773-023-05454-1 (<https://doi.org/10.1007%2Fs10773-023-05454-1>).
293. Olmo, Gonzalo; Rubiera-Garcia, Diego (2015). "Nonsingular Black Holes in  $f(R)$  Theories" (<https://doi.org/10.3390%2Funiverse1020173>). *Universe*. **1** (2): 173–185. arXiv:1509.02430 (<https://arxiv.org/abs/1509.02430>). Bibcode:2015Univ....1..173O (<https://ui.adsabs.harvard.edu/abs/2015Univ....1..173O>). doi:10.3390/universe1020173 (<https://doi.org/10.3390%2Funiverse1020173>).
294. Mathur, Samir D. (2005). "The fuzzball proposal for black holes: an elementary review". *Fortschritte der Physik*. **53** (7–8): 793. arXiv:hep-th/0502050 (<https://arxiv.org/abs/hep-th/0502050>). Bibcode:2005ForPh..53..793M (<https://ui.adsabs.harvard.edu/abs/2005ForPh..53..793M>). doi:10.1002/prop.200410203 (<https://doi.org/10.1002%2Fprop.200410203>). S2CID 15083147 (<https://api.semanticscholar.org/CorpusID:15083147>).
295. Avery, Steven G.; Chowdhury, Borun D.; Puhm, Andrea (2013). "Unitarity and fuzzball complementarity: "Alice fuzzes but may not even know it!" ". *Journal of High Energy Physics* (9) 12. arXiv:1210.6996 (<https://arxiv.org/abs/1210.6996>). Bibcode:2013JHEP...09..012A (<https://ui.adsabs.harvard.edu/abs/2013JHEP...09..012A>). doi:10.1007/JHEP09(2013)012 (<https://doi.org/10.1007%2FJHEP09%282013%29012>).

296. Bojowald, Martin (2020). "Black-Hole Models in Loop Quantum Gravity" (<https://doi.org/10.3390%2Funiverse6080125>). *Universe*. **6** (8): 125. arXiv:2009.13565 (<https://arxiv.org/abs/2009.13565>). Bibcode:2020Univ....6..125B (<https://ui.adsabs.harvard.edu/abs/2020Univ....6..125B>). doi:10.3390/universe6080125 (<https://doi.org/10.3390%2Funiverse6080125>).
297. Woosley, S. E.; Heger, A.; Weaver, T. A. (7 November 2002). "The evolution and explosion of massive stars" (<https://link.aps.org/doi/10.1103/RevModPhys.74.1015>). *Reviews of Modern Physics*. **74** (4): 1015–1071. Bibcode:2002RvMP...74.1015W (<https://ui.adsabs.harvard.edu/abs/2002RvMP...74.1015W>). doi:10.1103/RevModPhys.74.1015 (<https://doi.org/10.1103%2FRevModPhys.74.1015>). ISSN 0034-6861 (<https://search.worldcat.org/issn/0034-6861>).
298. Zappa, Francesco; Bernuzzi, Sebastiano; Pannarale, Francesco; Mapelli, Michela; Giacobbo, Nicola (25 July 2019). "Black-Hole Remnants from Black-Hole–Neutron-Star Mergers" (<https://link.aps.org/doi/10.1103/PhysRevLett.123.041102>). *Physical Review Letters*. **123** (4) 041102. arXiv:1903.11622 (<https://arxiv.org/abs/1903.11622>). Bibcode:2019PhRvL.123d1102Z (<https://ui.adsabs.harvard.edu/abs/2019PhRvL.123d1102Z>). doi:10.1103/PhysRevLett.123.041102 (<https://doi.org/10.1103%2FPhysRevLett.123.041102>). ISSN 0031-9007 (<https://search.worldcat.org/issn/0031-9007>). PMID 31491270 (<http://pubmed.ncbi.nlm.nih.gov/31491270>).
299. Inayoshi, Kohei; Visbal, Eli; Haiman, Zoltán (18 August 2020). "The Assembly of the First Massive Black Holes" (<https://www.annualreviews.org/content/journals/10.1146/annurev-astro-120419-014455>). *Annual Review of Astronomy and Astrophysics*. **58**: 27–97. arXiv:1911.05791 (<https://arxiv.org/abs/1911.05791>). Bibcode:2020ARA&A..58...27I (<https://ui.adsabs.harvard.edu/abs/2020ARA&A..58...27I>). doi:10.1146/annurev-astro-120419-014455 (<https://doi.org/10.1146%2Fannurev-astro-120419-014455>). ISSN 0066-4146 (<http://search.worldcat.org/issn/0066-4146>).
300. "Ripped Apart by a Black Hole" (<https://web.archive.org/web/20130721014626/http://www.eso.org/public/news/eso1332/>). ESO (Press release). 17 July 2013. Archived from the original (<http://www.eso.org/public/news/eso1332/>) on 21 July 2013. Retrieved 19 July 2013.
301. Janka, H.; Langanke, K.; Marek, A.; Martinezpinedo, G.; Muller, B. (2007). "Theory of core-collapse supernovae". *Physics Reports*. **442** (1–6): 38–74. arXiv:astro-ph/0612072 (<https://arxiv.org/abs/astro-ph/0612072>). Bibcode:2007PhR...442...38J (<https://ui.adsabs.harvard.edu/abs/2007PhR...442...38J>). doi:10.1016/j.physrep.2007.02.002 (<https://doi.org/10.1016%2Fj.physrep.2007.02.002>).
302. Fryer, Chris L.; Holz, Daniel E.; Hughes, Scott A. (2002). "Gravitational Wave Emission from Core Collapse of Massive Stars". *The Astrophysical Journal*. **565** (1): 430–446. arXiv:astro-ph/0106113 (<https://arxiv.org/abs/astro-ph/0106113>). Bibcode:2002ApJ...565..430F (<https://ui.adsabs.harvard.edu/abs/2002ApJ...565..430F>). doi:10.1086/324034 (<https://doi.org/10.1086%2F324034>).
303. Bennett, Jeffrey (2025). "Degeneracy pressure in stars and stellar corpses". *The Physics Teacher*. **63** (3): 212–213. Bibcode:2025PhTea..63c.212B (<https://ui.adsabs.harvard.edu/abs/2025PhTea..63c.212B>). doi:10.1119/5.0260882 (<https://doi.org/10.1119%2F5.0260882>).
304. Carroll 2004, Section 5.8
305. Rees, M. J.; Volonteri, M. (2007). "Massive black holes: Formation and evolution". In Karas, V.; Matt, G. (eds.). *Black Holes from Stars to Galaxies – Across the Range of Masses*. Proceedings of the International Astronomical Union. pp. 51–58. arXiv:astro-ph/0701512 (<https://arxiv.org/abs/astro-ph/0701512>). Bibcode:2007IAUS..238...51R (<https://ui.adsabs.harvard.edu/abs/2007IAUS..238...51R>). doi:10.1017/S1743921307004681 (<https://doi.org/10.1017%2FS1743921307004681>). ISBN 978-0-521-86347-6. S2CID 14844338 (<https://api.semanticscholar.org/CorpusID:14844338>).



306. Pacucci, F.; Ferrara, A.; Grazian, A.; Fiore, F.; Giallongo, E. (2016). "First Identification of Direct Collapse Black Hole Candidates in the Early Universe in CANDELS/GOODS-S" (<https://doi.org/10.1093%2Fmnras%2Fstw725>). *Mon. Not. R. Astron. Soc.* **459** (2): 1432. arXiv:1603.08522 (<https://arxiv.org/abs/1603.08522>). Bibcode:2016MNRAS.459.1432P (<https://ui.adsabs.harvard.edu/abs/2016MNRAS.459.1432P>). doi:10.1093/mnras/stw725 (<https://doi.org/10.1093%2Fmnras%2Fstw725>). S2CID 118578313 (<https://api.semanticscholar.org/CorpusID:118578313>).
307. Bañados, Eduardo; Venemans, Bram P.; Mazzucchelli, Chiara; Farina, Emanuele P.; Walter, Fabian; Wang, Feige; Decarli, Roberto; Stern, Daniel; Fan, Xiaohui; Davies, Frederick B.; Hennawi, Joseph F. (1 January 2018). "An 800-million-solar-mass black hole in a significantly neutral Universe at a redshift of 7.5". *Nature*. **553** (7689): 473–476. arXiv:1712.01860 (<https://arxiv.org/abs/1712.01860>). Bibcode:2018Natur.553..473B (<https://ui.adsabs.harvard.edu/abs/2018Natur.553..473B>). doi:10.1038/nature25180 (<https://doi.org/10.1038%2Fnature25180>). PMID 29211709 (<https://pubmed.ncbi.nlm.nih.gov/29211709>). S2CID 205263326 (<https://api.semanticscholar.org/CorpusID:205263326>).
308. Boylan-Kolchin, Michael; Weisz, Daniel R. (2021). "Uncertain times: The redshift–time relation from cosmology and stars" (<https://doi.org/10.1093%2Fmnras%2Fstab1521>). *Monthly Notices of the Royal Astronomical Society*. **505** (2): 2764–2783. doi:10.1093/mnras/stab1521 (<https://doi.org/10.1093%2Fmnras%2Fstab1521>).
309. Penrose, R. (2002). "Gravitational Collapse: The Role of General Relativity" ([https://web.archive.org/web/20130526224126/http://www.imamu.edu.sa/Scientific\\_selections/abstracts/Physics/Gravitational%20Collapse%20The%20Role%20of%20General.pdf](https://web.archive.org/web/20130526224126/http://www.imamu.edu.sa/Scientific_selections/abstracts/Physics/Gravitational%20Collapse%20The%20Role%20of%20General.pdf)) (PDF). *General Relativity and Gravitation*. **34** (7): 1141. Bibcode:2002GReGr..34.1141P (<https://ui.adsabs.harvard.edu/abs/2002GReGr..34.1141P>). doi:10.1023/A:1016578408204 (<https://doi.org/10.1023%2FA%3A1016578408204>). S2CID 117459073 (<https://api.semanticscholar.org/CorpusID:117459073>). Archived from the original ([http://www.imamu.edu.sa/Scientific\\_selections/abstracts/Physics/Gravitational%20Collapse%20The%20Role%20of%20General.pdf](http://www.imamu.edu.sa/Scientific_selections/abstracts/Physics/Gravitational%20Collapse%20The%20Role%20of%20General.pdf)) (PDF) on 26 May 2013.
310. Yoo, Chul-Moon (2022). "The Basics of Primordial Black Hole Formation and Abundance Estimation" (<https://doi.org/10.3390%2Fgalaxies10060112>). *Galaxies*. **10** (6): 112. arXiv:2211.13512 (<https://arxiv.org/abs/2211.13512>). Bibcode:2022Galax..10..112Y (<https://ui.adsabs.harvard.edu/abs/2022Galax..10..112Y>). doi:10.3390/galaxies10060112 (<https://doi.org/10.3390%2Fgalaxies10060112>).
311. Balzer, Ashley (7 May 2024). "Primordial Black Holes" (<https://svs.gsfc.nasa.gov/14524/>). NASA SVS. Archived (<https://web.archive.org/web/20250827012141/https://svs.gsfc.nasa.gov/14524/>) from the original on 27 August 2025. Retrieved 23 November 2025.
312. Carr, B. J. (2005). "Primordial Black Holes: Do They Exist and Are They Useful?". In Suzuki, H.; Yokoyama, J.; Suto, Y.; Sato, K. (eds.). *First identification of direct collapse black hole candidates in the early Universe in CANDELS/GOODS-S*. Vol. 459. Universal Academy Press. pp. astro-ph/0511743. arXiv:astro-ph/0511743 (<https://arxiv.org/abs/astro-ph/0511743>). Bibcode:2005astro.ph.11743C (<https://ui.adsabs.harvard.edu/abs/2005astro.ph.11743C>). doi:10.1093/mnras/stw725 (<https://doi.org/10.1093%2Fmnras%2Fstw725>). ISBN 978-4-946443-94-7. {{cite book}}: |journal= ignored (help)
313. Carr, Bernard; Kohri, Kazunori; Sendouda, Yuuiti; Yokoyama, Jun'ichi (2021). "Constraints on primordial black holes". *Reports on Progress in Physics*. **84** (11). arXiv:2002.12778 (<https://arxiv.org/abs/2002.12778>). Bibcode:2021RPPh...84k6902C (<https://ui.adsabs.harvard.edu/abs/2021RPPh...84k6902C>). doi:10.1088/1361-6633/ac1e31 (<https://doi.org/10.1088%2F1361-6633%2Fac1e31>). PMID 34874316 (<https://pubmed.ncbi.nlm.nih.gov/34874316>).
314. Philip Gibbs. "Is the Big Bang a black hole?" (<http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/universe.html>). John Baez. Archived (<https://web.archive.org/web/20181231021714/http://math.ucr.edu/home/baez/physics/Relativity/BlackHoles/universe.html>) from the original on 31 December 2018. Retrieved 16 March 2018.



315. Sutter, Paul (21 June 2023). "Why didn't the infant universe collapse into a black hole?" (<http://www.space.com/why-infant-universe-not-collapse-black-hole>). *Space.com*. Archived (<http://web.archive.org/web/20250325152015/https://www.space.com/why-infant-universe-not-collapse-black-hole>) from the original on 25 March 2025. Retrieved 24 November 2025.
316. Musser, George (22 September 2003). "According to the big bang theory, all the matter in the universe erupted from a singularity. Why didn't all this matter--cheek by jowl as it was--immediately collapse into a black hole?" (<https://www.scientificamerican.com/article/according-to-the-big-bang/>). *Scientific American*. Archived (<https://web.archive.org/web/20250426033551/https://www.scientificamerican.com/article/according-to-the-big-bang/>) from the original on 26 April 2025. Retrieved 24 November 2025.
317. Kaloper, Nemanja; Terning, John (2007). "How black holes form in high energy collisions". *General Relativity and Gravitation*. **39** (10): 1525–1532. arXiv:0705.0408 (<https://arxiv.org/abs/0705.0408>). Bibcode:2007GReGr..39.1525K (<https://ui.adsabs.harvard.edu/abs/2007GReGr..39.1525K>). doi:10.1007/s10714-007-0468-5 (<https://doi.org/10.1007%2Fs10714-007-0468-5>).
318. Giddings, S. B.; Thomas, S. (2002). "High energy colliders as black hole factories: The end of short distance physics". *Physical Review D*. **65** (5) 056010. arXiv:hep-ph/0106219 (<http://arxiv.org/abs/hep-ph/0106219>). Bibcode:2002PhRvD..65e6010G (<https://ui.adsabs.harvard.edu/abs/2002PhRvD..65e6010G>). doi:10.1103/PhysRevD.65.056010 (<https://doi.org/10.1103%2FPhysRevD.65.056010>). S2CID 1203487 (<https://api.semanticscholar.org/CorpusID:1203487>).
319. LHC Safety Assessment Group (2008). "Review of the Safety of LHC Collisions" (<http://lsag.web.cern.ch/lsag/LSAG-Report.pdf>) (PDF). *Journal of Physics G: Nuclear Physics*. **35** (11) 115004. arXiv:0806.3414 (<https://arxiv.org/abs/0806.3414>). Bibcode:2008JPhG...35k5004E (<https://ui.adsabs.harvard.edu/abs/2008JPhG...35k5004E>). doi:10.1088/0954-3899/35/11/115004 (<https://doi.org/10.1088%2F0954-3899%2F35%2F11%2F115004>). S2CID 53370175 (<https://api.semanticscholar.org/CorpusID:53370175>). Archived (<https://web.archive.org/web/20100414160742/http://lsag.web.cern.ch/lsag/LSAG-Report.pdf>) (PDF) from the original on 14 April 2010.
320. Cavaglià, M. (2010). "Particle accelerators as black hole factories?" ([https://web.archive.org/web/20130508085759/http://www.einstein-online.info/spotlights/accelerators\\_bh](https://web.archive.org/web/20130508085759/http://www.einstein-online.info/spotlights/accelerators_bh)). *Einstein-Online*. **4**: 1010. Archived from the original ([http://www.einstein-online.info/spotlights/accelerators\\_bh/](http://www.einstein-online.info/spotlights/accelerators_bh/)) on 8 May 2013. Retrieved 8 May 2013.
321. Vesperini, E.; McMillan, S. L. W.; d'Ercole, A.; et al. (2010). "Intermediate-Mass Black Holes in Early Globular Clusters". *The Astrophysical Journal Letters*. **713** (1): L41 – L44. arXiv:1003.3470 (<https://arxiv.org/abs/1003.3470>). Bibcode:2010ApJ...713L..41V (<https://ui.adsabs.harvard.edu/abs/2010ApJ...713L..41V>). doi:10.1088/2041-8205/713/1/L41 (<https://doi.org/10.1088%2F2041-8205%2F713%2F1%2FL41>). S2CID 119120429 (<https://api.semanticscholar.org/CorpusID:119120429>).
322. Zwart, S. F. P.; Baumgardt, H.; Hut, P.; et al. (2004). "Formation of massive black holes through runaway collisions in dense young star clusters". *Nature*. **428** (6984): 724–726. arXiv:astro-ph/0402622 (<https://arxiv.org/abs/astro-ph/0402622>). Bibcode:2004Natur.428..724P (<https://ui.adsabs.harvard.edu/abs/2004Natur.428..724P>). doi:10.1038/nature02448 (<https://doi.org/10.1038%2Fnature02448>). PMID 15085124 (<http://pubmed.ncbi.nlm.nih.gov/15085124>). S2CID 4408378 (<https://api.semanticscholar.org/CorpusID:4408378>).
323. O'Leary, R. M.; Rasio, F. A.; Fregeau, J. M.; et al. (2006). "Binary Mergers and Growth of Black Holes in Dense Star Clusters". *The Astrophysical Journal*. **637** (2): 937–951. arXiv:astro-ph/0508224 (<https://arxiv.org/abs/astro-ph/0508224>). Bibcode:2006ApJ...637..937O (<https://ui.adsabs.harvard.edu/abs/2006ApJ...637..937O>). doi:10.1086/498446 (<https://doi.org/10.1086%2F498446>). S2CID 1509957 (<https://api.semanticscholar.org/CorpusID:1509957>).

324. Brightman, M.; Bachetti, M.; Earnshaw, H. P.; Fürst, F.; García, J.; Grefenstette, B.; Heida, M.; Kara, E.; Madsen, K. K.; Middleton, M. J.; Stern, D.; Tombesi, F.; Walton, D. J. (2019). "Breaking the limit: Super-Eddington accretion onto black holes and neutron stars". *Bulletin of the American Astronomical Society*. **51** (3): 352. arXiv:1903.06844 (<https://arxiv.org/abs/1903.06844>). Bibcode:2019BAAS...51c.352B (<https://ui.adsabs.harvard.edu/abs/2019BAAS...51c.352B>).
325. Regan, John A.; Downes, Turlough P.; Volonteri, Marta; Beckmann, Ricarda; Lupi, Alessandro; Trebitsch, Maxime; Dubois, Yohan (2019). "Super-Eddington accretion and feedback from the first massive seed black holes" (<https://doi.org/10.1093%2Fmnras%2Fstz1045>). *Monthly Notices of the Royal Astronomical Society*. **486** (3): 3892–3906. doi:10.1093/mnras/stz1045 (<https://doi.org/10.1093%2Fmnras%2Fstz1045>).
326. Marziani, Paola; Garnica Luna, Karla; Floris, Alberto; Del Olmo, Ascensión; Deconto-Machado, Alice; Buendia-Rios, Tania M.; Negrete, C. Alenka; Dultzin, Deborah (2025). "Super-Eddington Accretion in Quasars" (<https://doi.org/10.3390%2Funiverse11020069>). *Universe*. **11** (2): 69. arXiv:2502.14713 (<https://arxiv.org/abs/2502.14713>). Bibcode:2025Univ...11...69M (<https://ui.adsabs.harvard.edu/abs/2025Univ...11...69M>). doi:10.3390/universe11020069 (<https://doi.org/10.3390%2Funiverse11020069>).
327. Ryu, Taeho; Perna, Rosalba; Haiman, Zoltán; Ostriker, Jeremiah P.; Stone, Nicholas C. (2018). "Interactions between multiple supermassive black holes in galactic nuclei: A solution to the final parsec problem" (<https://doi.org/10.1093%2Fmnras%2Fstx2524>). *Monthly Notices of the Royal Astronomical Society*. **473** (3): 3410–3433. doi:10.1093/mnras/stx2524 (<https://doi.org/10.1093%2Fmnras%2Fstx2524>).
328. Vasiliev, Eugene; Antonini, Fabio; Merritt, David (2014). "The Final-Parsec Problem in Nonspherical Galaxies Revisited". *The Astrophysical Journal*. **785** (2): 163. arXiv:1311.1167 (<https://arxiv.org/abs/1311.1167>). Bibcode:2014ApJ...785..163V (<https://ui.adsabs.harvard.edu/abs/2014ApJ...785..163V>). doi:10.1088/0004-637X/785/2/163 (<https://doi.org/10.1088%2F0004-637X%2F785%2F2%2F163>).
329. "What powers a black hole's mighty jets?" (<https://www.science.org/content/article/what-powers-black-holes-mighty-jets>). *Science* | AAAS. 19 November 2014. Archived (<https://web.archive.org/web/20190505171217/https://www.sciencemag.org/news/2014/11/what-powers-black-holes-mighty-jets>) from the original on 5 May 2019. Retrieved 19 March 2018.
330. Celotti, A.; Miller, J. C.; Sciama, D. W. (1999). "Astrophysical evidence for the existence of black holes" (<https://web.archive.org/web/20180727052939/https://cds.cern.ch/record/411555/files/9912186.pdf>) (PDF). *Classical and Quantum Gravity*. **16** (12A): A3 – A21. arXiv:astro-ph/9912186 (<https://arxiv.org/abs/astro-ph/9912186>). Bibcode:1999CQGra..16A...3C (<https://ui.adsabs.harvard.edu/abs/1999CQGra..16A...3C>). doi:10.1088/0264-9381/16/12A/301 (<https://doi.org/10.1088%2F0264-9381%2F16%2F12A%2F301>). S2CID 17677758 (<https://api.semanticscholar.org/CorpusID:17677758>). Archived from the original (<https://cds.cern.ch/record/411555/files/9912186.pdf>) (PDF) on 27 July 2018.
331. Winter, L. M.; Mushotzky, R. F.; Reynolds, C. S. (2006). "XMM-Newton Archival Study of the Ultraluminous X-Ray Population in Nearby Galaxies". *The Astrophysical Journal*. **649** (2): 730–752. arXiv:astro-ph/0512480 (<https://arxiv.org/abs/astro-ph/0512480>). Bibcode:2006ApJ...649..730W (<https://ui.adsabs.harvard.edu/abs/2006ApJ...649..730W>). doi:10.1086/506579 (<https://doi.org/10.1086%2F506579>). S2CID 118445260 (<https://api.semanticscholar.org/CorpusID:118445260>).
332. Evans, Charles R.; Kochanek, Christopher S. (1989). "The tidal disruption of a star by a massive black hole". *The Astrophysical Journal*. **346**: L13. Bibcode:1989ApJ...346L..13E (<https://ui.adsabs.harvard.edu/abs/1989ApJ...346L..13E>). doi:10.1086/185567 (<https://doi.org/10.1086%2F185567>).

333. Komossa, S. (2015). "Tidal disruption of stars by supermassive black holes: Status of observations". *Journal of High Energy Astrophysics*. **7**: 148–157. arXiv:1505.01093 (<https://arxiv.org/abs/1505.01093>). Bibcode:2015JHEAp...7..148K (<https://ui.adsabs.harvard.edu/abs/2015JHEAp...7..148K>). doi:10.1016/j.jheap.2015.04.006 (<https://doi.org/10.1016%2Fj.jheap.2015.04.006>).
334. Hayasaki, Kimitake; Stone, Nicholas; Loeb, Abraham (2016). "Circularization of tidally disrupted stars around spinning supermassive black holes" (<https://doi.org/10.1093%2Fmnras%2Fstw1387>). *Monthly Notices of the Royal Astronomical Society*. **461** (4): 3760–3780. doi:10.1093/mnras/stw1387 (<https://doi.org/10.1093%2Fmnras%2Fstw1387>).
335. Page, D. N. (2005). "Hawking radiation and black hole thermodynamics". *New Journal of Physics*. **7** (1): 203. arXiv:hep-th/0409024 (<https://arxiv.org/abs/hep-th/0409024>). Bibcode:2005NJPh....7..203P (<https://ui.adsabs.harvard.edu/abs/2005NJPh....7..203P>). doi:10.1088/1367-2630/7/1/203 (<https://doi.org/10.1088%2F1367-2630%2F7%2F1%2F203>). S2CID 119047329 (<https://api.semanticscholar.org/CorpusID:119047329>).
336. Smerlak, Matteo; Singh, Suprit (2013). "New perspectives on Hawking radiation". *Physical Review D*. **88** (10) 104023. arXiv:1304.2858 (<https://arxiv.org/abs/1304.2858>). Bibcode:2013PhRvD..88j4023S (<https://ui.adsabs.harvard.edu/abs/2013PhRvD..88j4023S>). doi:10.1103/PhysRevD.88.104023 (<https://doi.org/10.1103%2FPhysRevD.88.104023>).
337. Carroll 2004, Ch. 9.6
338. Ruiz, O.; Molina, U.; Vitoria, P. (2019). "Thermodynamic analysis of Kerr-Newman black holes". *Journal of Physics: Conference Series*. **1219** (1) 012016. Bibcode:2019JPhCS1219a2016R (<https://ui.adsabs.harvard.edu/abs/2019JPhCS1219a2016R>). doi:10.1088/1742-6596/1219/1/012016 (<https://doi.org/10.1088%2F1742-6596%2F1219%2F1%2F012016>). "From this, an expression is established for the Hawking temperature of a Kerr-Newman black hole as a function of its mass  $M$ , angular momentum  $J$  and charge  $Q$ . As the black hole loses mass, its temperature increases inversely proportional."
339. Siegel, Ethan (2017). "Ask Ethan: Do Black Holes Grow Faster Than They Evaporate?" (<https://www.forbes.com/sites/startswithabang/2017/08/19/ask-ethan-do-black-holes-grow-faster-than-they-evaporate/>). *Forbes* ("Starts With A Bang" blog). Archived (<https://web.archive.org/web/20181122031830/https://www.forbes.com/sites/startswithabang/2017/08/19/ask-ethan-do-black-holes-grow-faster-than-they-evaporate/>) from the original on 22 November 2018. Retrieved 17 March 2018.
340. Sivaram, C. (2001). "Black hole Hawking radiation may never be observed!". *General Relativity and Gravitation*. **33** (2): 175–181. Bibcode:2001GReGr..33..175S (<https://ui.adsabs.harvard.edu/abs/2001GReGr..33..175S>). doi:10.1023/A:1002753400430 (<https://doi.org/10.1023%2FA%3A1002753400430>). S2CID 118913634 (<https://api.semanticscholar.org/CorpusID:118913634>).
341. "Evaporating black holes?" ([https://web.archive.org/web/20110722055345/http://www.einstein-online.info/elementary/quantum/evaporating\\_bh/?set\\_language=en](https://web.archive.org/web/20110722055345/http://www.einstein-online.info/elementary/quantum/evaporating_bh/?set_language=en)). *Einstein online*. Max Planck Institute for Gravitational Physics. 2010. Archived from the original ([http://www.einstein-online.info/elementary/quantum/evaporating\\_bh/?set\\_language=en](http://www.einstein-online.info/elementary/quantum/evaporating_bh/?set_language=en)) on 22 July 2011. Retrieved 12 December 2010.
342. Giddings, S. B.; Mangano, M. L. (2008). "Astrophysical implications of hypothetical stable TeV-scale black holes". *Physical Review D*. **78** (3) 035009. arXiv:0806.3381 (<https://arxiv.org/abs/0806.3381>). Bibcode:2008PhRvD..78c5009G (<https://ui.adsabs.harvard.edu/abs/2008PhRvD..78c5009G>). doi:10.1103/PhysRevD.78.035009 (<https://doi.org/10.1103%2FPhysRevD.78.035009>). S2CID 17240525 (<https://api.semanticscholar.org/CorpusID:17240525>).
343. Peskin, M. E. (2008). "The end of the world at the Large Hadron Collider?" (<https://doi.org/10.1103%2FPhysics.1.14>). *Physics*. **1** 14. Bibcode:2008PhyOJ...1...14P (<https://ui.adsabs.harvard.edu/abs/2008PhyOJ...1...14P>). doi:10.1103/Physics.1.14 (<https://doi.org/10.1103%2FPhysics.1.14>).

344. Fichtel, C. E.; Bertsch, D. L.; Dingus, B. L.; et al. (1994). "Search of the energetic gamma-ray experiment telescope (EGRET) data for high-energy gamma-ray microsecond bursts". *Astrophysical Journal*. **434** (2): 557–559. Bibcode:1994ApJ...434..557F (<https://ui.adsabs.harvard.edu/abs/1994ApJ...434..557F>). doi:10.1086/174758 (<https://doi.org/10.1086%2F174758>).
345. Carr, Bernard; Kohri, Kazunori; Sendouda, Yuuiti; Yokoyama, Jun'ichi (2021). "Constraints on primordial black holes". *Reports on Progress in Physics*. **84** (11). arXiv:2002.12778 (<https://arxiv.org/abs/2002.12778>). Bibcode:2021RPPh...84k6902C (<https://ui.adsabs.harvard.edu/abs/2021RPPh...84k6902C>). doi:10.1088/1361-6633/ac1e31 (<https://doi.org/10.1088%2F1361-6633%2Fac1e31>). PMID 34874316 (<https://pubmed.ncbi.nlm.nih.gov/34874316>).
346. Naeye, R. "Testing Fundamental Physics" ([https://www.nasa.gov/mission\\_pages/GLAST/science/testing\\_fundamental\\_physics.html](https://www.nasa.gov/mission_pages/GLAST/science/testing_fundamental_physics.html)). NASA. Archived ([https://web.archive.org/web/20080831045232/http://www.nasa.gov/mission\\_pages/GLAST/science/testing\\_fundamental\\_physics.html](https://web.archive.org/web/20080831045232/http://www.nasa.gov/mission_pages/GLAST/science/testing_fundamental_physics.html)) from the original on 31 August 2008. Retrieved 16 September 2008.
347. Ackermann, M.; et al. (2018). "Search for Gamma-Ray Emission from Local Primordial Black Holes with the Fermi Large Area Telescope" (<https://doi.org/10.3847%2F1538-4357%2Faaac7b>). *The Astrophysical Journal*. **857** (1): 49. Bibcode:2018ApJ...857...49A (<https://ui.adsabs.harvard.edu/abs/2018ApJ...857...49A>). doi:10.3847/1538-4357/aaac7b (<https://doi.org/10.3847%2F1538-4357%2Faaac7b>).
348. Frautschi, S. (1982). "Entropy in an Expanding Universe". *Science*. **217** (4560): 593–599. Bibcode:1982Sci...217..593F (<https://ui.adsabs.harvard.edu/abs/1982Sci...217..593F>). doi:10.1126/science.217.4560.593 (<https://doi.org/10.1126%2Fscience.217.4560.593>). PMID 17817517 (<https://pubmed.ncbi.nlm.nih.gov/17817517>). S2CID 27717447 (<https://api.semanticscholar.org/CorpusID:27717447>). See page 596: table 1 and section "black hole decay" and previous sentence on that page.
349. Page, Don N. (1976). "Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole". *Physical Review D*. **13** (2): 198–206. Bibcode:1976PhRvD..13..198P (<https://ui.adsabs.harvard.edu/abs/1976PhRvD..13..198P>). doi:10.1103/PhysRevD.13.198 (<https://doi.org/10.1103%2FPhysRevD.13.198>).. See in particular equation (27).
350. Hiscock, William A. (1981). "Models of evaporating black holes. I". *Physical Review D*. **23** (12): 2813–2822. Bibcode:1981PhRvD..23.2813H (<https://ui.adsabs.harvard.edu/abs/1981PhRvD..23.2813H>). doi:10.1103/PhysRevD.23.2813 (<https://doi.org/10.1103%2FPhysRevD.23.2813>).
351. Stephens, C. R.; Hooft, G 't; Whiting, B. F. (1994). "Black hole evaporation without information loss". *Classical and Quantum Gravity*. **11** (3): 621–647. arXiv:gr-qc/9310006 (<https://arxiv.org/abs/gr-qc/9310006>). Bibcode:1994CQGra..11..621S (<https://ui.adsabs.harvard.edu/abs/1994CQGra..11..621S>). doi:10.1088/0264-9381/11/3/014 (<https://doi.org/10.1088%2F0264-9381%2F11%2F3%2F014>).
352. Saida, Hiromi (2007). "Black hole evaporation in a heat bath as a nonequilibrium process and its final fate". *Classical and Quantum Gravity*. **24** (3): 691–722. arXiv:gr-qc/0701030 (<https://arxiv.org/abs/gr-qc/0701030>). Bibcode:2007CQGra..24..691S (<https://ui.adsabs.harvard.edu/abs/2007CQGra..24..691S>). doi:10.1088/0264-9381/24/3/012 (<https://doi.org/10.1088%2F0264-9381%2F24%2F3%2F012>).
353. Elbert, Oliver D.; Bullock, James S.; Kaplinghat, Manoj (1 January 2018). "Counting black holes: The cosmic stellar remnant population and implications for LIGO" (<https://academic.oup.com/mnras/article/473/1/1186/4060726>). *Monthly Notices of the Royal Astronomical Society*. **473** (1): 1186–1194. arXiv:1703.02551 (<https://arxiv.org/abs/1703.02551>). doi:10.1093/mnras/stx1959 (<https://doi.org/10.1093%2Fmnras%2Fstx1959>). ISSN 0035-8711 (<https://search.worldcat.org/issn/0035-8711>).

354. "April 2017 Observations" (<https://eventhorizontelescope.org/galleries/2017-observations>). *Event Horizon Telescope*. Archived (<https://web.archive.org/web/20190410223148/https://eventhorizontelescope.org/galleries/2017-observations>) from the original on 10 April 2019. Retrieved 11 April 2019.
355. Overbye, Dennis (24 January 2024). "That Famous Black Hole Gets a Second Look - Repeated studies of the supermassive black hole in the galaxy Messier 87 confirm that it continues to act as Einstein's theory predicted it would" (<https://www.nytimes.com/2024/01/24/science/space/black-holes-photography-m87.html>). *The New York Times*. Archived (<https://archive.today/20240124200523/https://www.nytimes.com/2024/01/24/science/space/black-holes-photography-m87.html>) from the original on 24 January 2024. Retrieved 25 January 2024.
356. Overbye, Dennis (10 April 2019). "Darkness Visible, Finally: Astronomers Capture First Ever Image of a Black Hole" (<https://www.nytimes.com/2019/04/10/science/black-hole-picture.html>). *The New York Times*. Archived (<https://web.archive.org/web/20190521020231/https://www.nytimes.com/2019/04/10/science/black-hole-picture.html>) from the original on 21 May 2019. Retrieved 11 April 2019.
357. *Black Holes: The Edge of All We Know* (Motion picture). 2020. 99 minutes in.
358. "Astronomers Reveal the First Picture of a Black Hole" (<https://www.nytimes.com/video/science/space/100000006453594/first-image-black-hole.html>). *The New York Times* (video). AP. 10 April 2019. Archived (<https://web.archive.org/web/20190522052307/https://www.nytimes.com/video/science/space/100000006453594/first-image-black-hole.html>) from the original on 22 May 2019. Retrieved 11 April 2019.
359. Doeleman, Shep (4 April 2016). "The Event Horizon Telescope: Imaging and Time-Resolving a Black Hole" ([https://www.youtube.com/watch?v=\\_yGSgUUYDQ8&t=46m50s](https://www.youtube.com/watch?v=_yGSgUUYDQ8&t=46m50s)). *Physics @ Berkeley*. Event occurs at 46:50. Archived ([https://web.archive.org/web/20161201115303/https://www.youtube.com/watch?v=\\_yGSgUUYDQ8&t=46m50s](https://web.archive.org/web/20161201115303/https://www.youtube.com/watch?v=_yGSgUUYDQ8&t=46m50s)) from the original on 1 December 2016. Retrieved 8 July 2016.
360. Grossman, Lisa; Conover, Emily (10 April 2019). "The first picture of a black hole opens a new era of astrophysics" (<https://www.sciencenews.org/article/black-hole-first-picture-event-horizon-telescope>). *Science News*. Archived (<https://web.archive.org/web/20190427192956/https://www.sciencenews.org/article/black-hole-first-picture-event-horizon-telescope>) from the original on 27 April 2019. Retrieved 11 April 2019.
361. Event Horizon Telescope Collaboration (2021). "First M87 Event Horizon Telescope Results. VII. Polarization of the Ring" (<https://doi.org/10.3847/2F2041-8213/2Fabe71d>). *The Astrophysical Journal*. **910** (1): L12. arXiv:2105.01169 (<https://arxiv.org/abs/2105.01169>). Bibcode:2021ApJ...910L..12E (<https://ui.adsabs.harvard.edu/abs/2021ApJ...910L..12E>). doi:10.3847/2041-8213/abe71d (<https://doi.org/10.3847/2F2041-8213/2Fabe71d>). S2CID 233851995 (<https://api.semanticscholar.org/CorpusID:233851995>).
362. Overbye, Dennis (26 April 2023). "A Fresh View of an Increasingly Familiar Black Hole - Radio astronomers have captured a wide-angle image of one of the most violent locales in the cosmos" (<https://www.nytimes.com/2023/04/26/science/astronomy-black-hole-m87.html>). *The New York Times*. Archived (<https://archive.today/20230426212932/https://www.nytimes.com/2023/04/26/science/astronomy-black-hole-m87.html>) from the original on 26 April 2023. Retrieved 26 April 2023.
363. Lu, Ru-Sen; et al. (26 April 2023). "A ring-like accretion structure in M87 connecting its black hole and jet" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10132962>). *Nature*. **616** (7958): 686–690. arXiv:2304.13252 (<https://arxiv.org/abs/2304.13252>). Bibcode:2023Natur.616..686L (<https://ui.adsabs.harvard.edu/abs/2023Natur.616..686L>). doi:10.1038/s41586-023-05843-w (<https://doi.org/10.1038/2Fs41586-023-05843-w>). PMC 10132962 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10132962>). PMID 37100940 (<https://pubmed.ncbi.nlm.nih.gov/37100940>).

364. Johnson, M. D.; Fish, V. L.; Doeleman, S. S.; Marrone, D. P.; Plambeck, R. L.; Wardle, J. F. C.; Akiyama, K.; Asada, K.; Beaudoin, C. (4 December 2015). "Resolved magnetic-field structure and variability near the event horizon of Sagittarius A\*". *Science*. **350** (6265): 1242–1245. arXiv:1512.01220 (<https://arxiv.org/abs/1512.01220>). Bibcode:2015Sci...350.1242J (<https://ui.adsabs.harvard.edu/abs/2015Sci...350.1242J>). doi:10.1126/science.aac7087 (<https://doi.org/10.1126%2Fscience.aac7087>). PMID 26785487 (<https://pubmed.ncbi.nlm.nih.gov/26785487>). S2CID 21730194 (<https://api.semanticscholar.org/CorpusID:21730194>).
365. "Event Horizon Telescope Reveals Magnetic Fields at Milky Way's Central Black Hole" (<http://www.cfa.harvard.edu/news/2015-28>). *cfa.harvard.edu*. 3 December 2015. Archived (<http://web.archive.org/web/20151231211408/https://www.cfa.harvard.edu/news/2015-28>) from the original on 31 December 2015. Retrieved 12 January 2016.
366. Event Horizon Telescope Collaboration; Akiyama, Kazunori; Alberdi, Antxon; Alef, Walter; Algaba, Juan Carlos; Anantua, Richard; Asada, Keiichi; Azulay, Rebecca; Bach, Uwe; Bacsko, Anne-Kathrin; Ball, David; Baloković, Mislav; Barrett, John; Bauböck, Michi; Benson, Bradford A. (1 May 2022). "First Sagittarius A\* Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole in the Center of the Milky Way" (<https://doi.org/10.3847%2F2041-8213%2Fac6674>). *The Astrophysical Journal Letters*. **930** (2): L12. arXiv:2311.08680 (<https://arxiv.org/abs/2311.08680>). Bibcode:2022ApJ...930L..12E (<https://ui.adsabs.harvard.edu/abs/2022ApJ...930L..12E>). doi:10.3847/2041-8213/ac6674 ([https://doi.org/10.3847/2041-8213%2Fac6674](https://doi.org/10.3847%2F2041-8213%2Fac6674)). ISSN 2041-8205 (<https://search.worldcat.org/issn/2041-8205>). "The ring usually lies near the gravitationally lensed photon orbits that define the boundary of what we hereafter refer to as the black hole "shadow.""
367. "Astronomers Reveal First Image of the Black Hole at the Heart of Our Galaxy" (<https://eventhorizontelescope.org/blog/astronomers-reveal-first-image-black-hole-heart-our-galaxy>). *eventhorizontelescope.org*. 12 May 2022. Retrieved 22 June 2022.
368. C. Bower, Geoffrey (May 2022). "Focus on First Sgr A\* Results from the Event Horizon Telescope" ([https://iopscience.iop.org/journal/2041-8205/page/Focus\\_on\\_First\\_Sgr\\_A\\_Results](https://iopscience.iop.org/journal/2041-8205/page/Focus_on_First_Sgr_A_Results)). *The Astrophysical Journal Letters* (2041–8205).
369. Overbye, Dennis (11 February 2016). "Physicists Detect Gravitational Waves, Proving Einstein Right" (<https://www.nytimes.com/2016/02/12/science/ligo-gravitational-waves-black-holes-einstein.html>). *The New York Times*. Archived (<https://web.archive.org/web/20160211165128/http://www.nytimes.com/2016/02/12/science/ligo-gravitational-waves-black-holes-einstein.html>) from the original on 11 February 2016. Retrieved 11 February 2016.
370. Abbott, Benjamin P.; et al. (LIGO Scientific Collaboration & Virgo Collaboration) (11 February 2016). "Properties of the binary black hole merger GW150914". *Physical Review Letters*. **116** (24) 241102. arXiv:1602.03840 (<https://arxiv.org/abs/1602.03840>). Bibcode:2016PhRvL.116x1102A (<https://ui.adsabs.harvard.edu/abs/2016PhRvL.116x1102A>). doi:10.1103/PhysRevLett.116.241102 (<https://doi.org/10.1103%2FPhysRevLett.116.241102>). PMID 27367378 (<https://pubmed.ncbi.nlm.nih.gov/27367378>). S2CID 217406416 (<https://api.semanticscholar.org/CorpusID:217406416>).
371. Cardoso, V.; Franzin, E.; Pani, P. (2016). "Is the gravitational-wave ringdown a probe of the event horizon?". *Physical Review Letters*. **116** (17) 171101. arXiv:1602.07309 (<https://arxiv.org/abs/1602.07309>). Bibcode:2016PhRvL.116q1101C (<https://ui.adsabs.harvard.edu/abs/2016PhRvL.116q1101C>). doi:10.1103/PhysRevLett.116.171101 (<https://doi.org/10.1103%2FPhysRevLett.116.171101>). PMID 27176511 (<https://pubmed.ncbi.nlm.nih.gov/27176511>). S2CID 206273829 (<https://api.semanticscholar.org/CorpusID:206273829>).



372. Abbott, B. P.; et al. (LIGO Scientific Collaboration & Virgo Collaboration) (2016). "Astrophysical Implications of the Binary Black Hole Merger GW150914" (<https://dcc.ligo.org/P1500262/public>). *Astrophys. J. Lett.* **818** (2): L22. arXiv:1602.03846 (<https://arxiv.org/abs/1602.03846>). Bibcode:2016ApJ...818L..22A (<https://ui.adsabs.harvard.edu/abs/2016ApJ...818L..22A>). doi:10.3847/2041-8205/818/2/L22 (<https://doi.org/10.3847%2F2041-8205%2F818%2F2%2FL22>). hdl:1826/11732 (<https://hdl.handle.net/1826%2F11732>). S2CID 209315965 (<https://api.semanticscholar.org/CorpusID:209315965>). Archived (<https://web.archive.org/web/20160316053938/https://dcc.ligo.org/P1500262/public>) from the original on 16 March 2016.
373. "Detection of gravitational waves" (<https://www.ligo.org/detections.php>). LIGO. Archived (<https://web.archive.org/web/20200520134427/https://www.ligo.org/detections.php>) from the original on 20 May 2020. Retrieved 9 April 2018.
374. Gillessen, S.; Eisenhauer, F.; Trippe, S.; et al. (2009). "Monitoring Stellar Orbits around the Massive Black Hole in the Galactic Center". *The Astrophysical Journal*. **692** (2): 1075–1109. arXiv:0810.4674 (<https://arxiv.org/abs/0810.4674>). Bibcode:2009ApJ...692.1075G (<https://ui.adsabs.harvard.edu/abs/2009ApJ...692.1075G>). doi:10.1088/0004-637X/692/2/1075 (<https://doi.org/10.1088%2F0004-637X%2F692%2F2%2F1075>). S2CID 1431308 (<https://api.semanticscholar.org/CorpusID:1431308>).
375. Ghez, A. M.; Klein, B. L.; Morris, M.; et al. (1998). "High Proper-Motion Stars in the Vicinity of Sagittarius A\*: Evidence for a Supermassive Black Hole at the Center of Our Galaxy". *The Astrophysical Journal*. **509** (2): 678–686. arXiv:astro-ph/9807210 (<https://arxiv.org/abs/astro-ph/9807210>). Bibcode:1998ApJ...509..678G (<https://ui.adsabs.harvard.edu/abs/1998ApJ...509..678G>). doi:10.1086/306528 (<https://doi.org/10.1086%2F306528>). S2CID 18243528 (<https://api.semanticscholar.org/CorpusID:18243528>).
376. Broderick, Avery; Loeb, Abraham; Narayan, Ramesh (August 2009). "The Event Horizon of Sagittarius A\*". *The Astrophysical Journal*. **701** (2): 1357–1366. arXiv:0903.1105 (<https://arxiv.org/abs/0903.1105>). Bibcode:2009ApJ...701.1357B (<https://ui.adsabs.harvard.edu/abs/2009ApJ...701.1357B>). doi:10.1088/0004-637X/701/2/1357 (<https://doi.org/10.1088%2F0004-637X%2F701%2F2%2F1357>). S2CID 12991878 (<https://api.semanticscholar.org/CorpusID:12991878>).
377. "Astronomers Reveal First Image of the Black Hole at the Heart of Our Galaxy" (<https://eventhorizontelescope.org/blog/astronomers-reveal-first-image-black-hole-heart-our-galaxy>). *Event Horizon Telescope*. 12 May 2022. Archived (<https://web.archive.org/web/20250926081853/https://eventhorizontelescope.org/blog/astronomers-reveal-first-image-black-hole-heart-our-galaxy>) from the original on 26 September 2025. Retrieved 2 December 2025.
378. "Press release: The Nobel Prize in Physics 2020" (<https://www.nobelprize.org/prizes/physics/2020/press-release/>). Nobel Foundation. Retrieved 6 October 2020.
379. Overbye, Dennis; Taylor, Derrick Bryson (6 October 2020). "Nobel Prize in Physics Awarded to 3 Scientists for Work on Black Holes – The prize was awarded half to Roger Penrose for showing how black holes could form and half to Reinhard Genzel and Andrea Ghez for discovering a supermassive object at the Milky Way's center" (<https://www.nytimes.com/2020/10/06/science/nobel-prize-physics.html>). *The New York Times*. Retrieved 6 October 2020.
380. Schatz, H.; Rehm, K.E. (2006). "X-ray binaries". *Nuclear Physics A*. **777**: 601–622. arXiv:astro-ph/0607624 (<https://arxiv.org/abs/astro-ph/0607624>). Bibcode:2006NuPhA.777..601S (<https://ui.adsabs.harvard.edu/abs/2006NuPhA.777..601S>). doi:10.1016/j.nuclphysa.2005.05.200 (<https://doi.org/10.1016%2Fj.nuclphysa.2005.05.200>).
381. Quirrenbach, Andreas; Frink, Sabine; Tomsick, John (1 December 2004). "Masses and Luminosities of X-ray Binaries" (<https://ntrs.nasa.gov/api/citations/20050186757/downloads/20050186757.pdf>) (PDF). *SIM PlanetQuest: Science with the Space Interferometry Mission*. National Aeronautics and Space Administration. Bibcode:2002swsi.conf...33Q (<https://ui.adsabs.harvard.edu/abs/2002swsi.conf...33Q>).

382. Cho, Adrian (2018). "A weight limit emerges for neutron stars". *Science*. **359** (6377): 724–725. Bibcode:2018Sci...359..724C (<https://ui.adsabs.harvard.edu/abs/2018Sci...359..724C>). doi:10.1126/science.359.6377.724 (<https://doi.org/10.1126%2Fscience.359.6377.724>). PMID 29449468 (<https://pubmed.ncbi.nlm.nih.gov/29449468>).
383. Rolston, B. (10 November 1997). "The First Black Hole" ([https://web.archive.org/web/20080502230214/http://news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](https://web.archive.org/web/20080502230214/http://news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)). *The bulletin*. University of Toronto. Archived from the original ([http://news.utoronto.ca/bin/bulletin/nov10\\_97/art4.htm](http://news.utoronto.ca/bin/bulletin/nov10_97/art4.htm)) on 2 May 2008. Retrieved 11 March 2008.
384. Orosz, Jerome A.; McClintock, Jeffrey E.; Aufdenberg, Jason P.; Remillard, Ronald A.; Reid, Mark J.; Narayan, Ramesh; Gou, Lijun (9 November 2011). "The Mass of the Black Hole in Cygnus X-1". *The Astrophysical Journal*. **742** (2): 84. arXiv:1106.3689 (<https://arxiv.org/abs/1106.3689>). Bibcode:2011ApJ...742...84O (<https://ui.adsabs.harvard.edu/abs/2011ApJ...742...84O>). doi:10.1088/0004-637x/742/2/84 (<https://doi.org/10.1088%2F0004-637x%2F742%2F2%2F84>). ISSN 0004-637X (<https://search.worldcat.org/issn/0004-637X>).
385. Corral-Santana, J. M.; Casares, J.; Muñoz-Darias, T.; Bauer, F. E.; Martínez-Pais, I. G.; Russell, D. M. (1 March 2016). "BlackCAT: A catalogue of stellar-mass black holes in X-ray transients" (<https://www.aanda.org/articles/aa/abs/2016/03/aa27130-15/aa27130-15.html>). *Astronomy & Astrophysics*. **587**: A61. arXiv:1510.08869 (<https://arxiv.org/abs/1510.08869>). Bibcode:2016A&A...587A..61C (<https://ui.adsabs.harvard.edu/abs/2016A&A...587A..61C>). doi:10.1051/0004-6361/201527130 (<https://doi.org/10.1051%2F0004-6361%2F201527130>). ISSN 0004-6361 (<https://search.worldcat.org/issn/0004-6361>).
386. Chou, Felicia; Anderson, Janet; Watzke, Megan (5 January 2015). "RELEASE 15-001 – NASA's Chandra Detects Record-Breaking Outburst from Milky Way's Black Hole" (<https://www.nasa.gov/press/2015/january/nasa-s-chandra-detects-record-breaking-outburst-from-milky-way-s-black-hole/>). NASA. Archived (<https://web.archive.org/web/20150106100532/http://www.nasa.gov/press/2015/january/nasa-s-chandra-detects-record-breaking-outburst-from-milky-way-s-black-hole/>) from the original on 6 January 2015. Retrieved 6 January 2015.
387. Krolik, J. H. (1999). *Active Galactic Nuclei* (<https://books.google.com/books?id=oRK8otMiWlgC&q=Active+Galactic+Nuclei>). Princeton University Press. Ch. 1.2. ISBN 978-0-691-01151-6. Archived (<https://web.archive.org/web/20210814220336/https://books.google.com/books?id=oRK8otMiWlgC&q=Active+Galactic+Nuclei>) from the original on 14 August 2021. Retrieved 16 October 2020.
388. Sparke, L. S.; Gallagher, J. S. (2000). *Galaxies in the Universe: An Introduction* (<https://books.google.com/books?id=N8Hngab5liQC&q=Galaxies+in+the+Universe:+An+Introduction>). Cambridge University Press. Ch. 9.1. ISBN 978-0-521-59740-1. Archived (<https://web.archive.org/web/20220322141933/https://books.google.com/books?id=N8Hngab5liQC&q=Galaxies+in+the+Universe%3A+An+Introduction>) from the original on 22 March 2022. Retrieved 16 October 2020.
389. Marconi, A.; Risaliti, G.; Gilli, R.; Hunt, L. K.; Maiolino, R.; Salvati, M. (2004). "Local supermassive black holes, relics of active galactic nuclei and the X-ray background" (<https://doi.org/10.1111%2Fj.1365-2966.2004.07765.x>). *Monthly Notices of the Royal Astronomical Society*. **351** (1): 169–185. arXiv:astro-ph/0311619 (<https://arxiv.org/abs/astro-ph/0311619>). Bibcode:2004MNRAS.351..169M (<https://ui.adsabs.harvard.edu/abs/2004MNRAS.351..169M>). doi:10.1111/j.1365-2966.2004.07765.x (<https://doi.org/10.1111%2Fj.1365-2966.2004.07765.x>).
390. Kormendy, J.; Richstone, D. (1995). "Inward Bound – The Search For Supermassive Black Holes In Galactic Nuclei". *Annual Review of Astronomy and Astrophysics*. **33** (1): 581–624. Bibcode:1995ARA&A...33..581K (<https://ui.adsabs.harvard.edu/abs/1995ARA&A...33..581K>). doi:10.1146/annurev.aa.33.090195.003053 (<https://doi.org/10.1146%2Fannurev.aa.33.090195.003053>).

391. Melia, Fulvio; Falcke, Heino (2001). "The Supermassive Black Hole at the Galactic Center". *Annual Review of Astronomy and Astrophysics*. **39**: 309–352. arXiv:astro-ph/0106162 (<https://arxiv.org/abs/astro-ph/0106162>). Bibcode:2001ARA&A..39..309M (<https://ui.adsabs.harvard.edu/abs/2001ARA&A..39..309M>). doi:10.1146/annurev.astro.39.1.309 (<https://doi.org/10.1146%2Fannurev.astro.39.1.309>).
392. King, A. (2003). "Black Holes, Galaxy Formation, and the MBH- $\sigma$  Relation". *The Astrophysical Journal Letters*. **596** (1): 27–29. arXiv:astro-ph/0308342 (<https://arxiv.org/abs/astro-ph/0308342>). Bibcode:2003ApJ...596L..27K (<https://ui.adsabs.harvard.edu/abs/2003ApJ...596L..27K>). doi:10.1086/379143 (<https://doi.org/10.1086%2F379143>). S2CID 9507887 (<https://api.semanticscholar.org/CorpusID:9507887>).
393. Ferrarese, L.; Merritt, D. (2000). "A Fundamental Relation Between Supermassive Black Holes and their Host Galaxies". *The Astrophysical Journal Letters*. **539** (1): 9–12. arXiv:astro-ph/0006053 (<https://arxiv.org/abs/astro-ph/0006053>). Bibcode:2000ApJ...539L...9F (<https://ui.adsabs.harvard.edu/abs/2000ApJ...539L...9F>). doi:10.1086/312838 (<https://doi.org/10.1086%2F312838>). S2CID 6508110 (<https://api.semanticscholar.org/CorpusID:6508110>).
394. Bennett, D. P.; Becker, A. C.; Quinn, J. L.; Tomaney, A. B.; Alcock, C.; Allsman, R. A.; Alves, D. R.; Axelrod, T. S.; Calitz, J. J.; Cook, K. H.; Drake, A. J.; Fragile, P. C.; Freeman, K. C.; Geha, M.; Griest, K. (1 November 2002). "Gravitational Microlensing Events Due to Stellar-Mass Black Holes" (<https://ui.adsabs.harvard.edu/abs/2002ApJ...579..639B/abstract>). *The Astrophysical Journal*. **579** (2): 639–659. arXiv:astro-ph/0109467 (<https://arxiv.org/abs/astro-ph/0109467>). Bibcode:2002ApJ...579..639B (<https://ui.adsabs.harvard.edu/abs/2002ApJ...579..639B>). doi:10.1086/342225 (<https://doi.org/10.1086%2F342225>). ISSN 0004-637X (<http://search.worldcat.org/issn/0004-637X>).
395. Mao, Shude; Smith, Martin C.; Woźniak, P.; Udalski, A.; Szymański, M.; Kubiak, M.; Pietrzyński, G.; Soszyński, I.; Żebruń, K. (1 January 2002). "Optical Gravitational Lensing Experiment OGLE-1999-BUL-32: the longest ever microlensing event - evidence for a stellar mass black hole?" (<https://doi.org/10.1046%2Fj.1365-8711.2002.04986.x>). *Monthly Notices of the Royal Astronomical Society*. **329** (2): 349–354. arXiv:astro-ph/0108312 (<https://arxiv.org/abs/astro-ph/0108312>). Bibcode:2002MNRAS.329..349M (<https://ui.adsabs.harvard.edu/abs/2002MNRAS.329..349M>). doi:10.1046/j.1365-8711.2002.04986.x (<https://doi.org/10.1046%2Fj.1365-8711.2002.04986.x>). ISSN 0035-8711 (<https://search.worldcat.org/issn/0035-8711>).
396. Sahu, K. C. (2022). "An Isolated Stellar-Mass Black Hole Detected Through Astrometric Microlensing" (<https://doi.org/10.3847%2F1538-4357%2Fac739e>). *Astrophysical Journal*. **933** (1): 83. arXiv:2201.13296 (<https://arxiv.org/abs/2201.13296>). Bibcode:2022ApJ...933...83S (<https://ui.adsabs.harvard.edu/abs/2022ApJ...933...83S>). doi:10.3847/1538-4357/ac739e (<https://doi.org/10.3847%2F1538-4357%2Fac739e>). S2CID 246430448 (<https://api.semanticscholar.org/CorpusID:246430448>).
397. Bozza, V. (2010). "Gravitational Lensing by Black Holes". *General Relativity and Gravitation*. **42** (9): 2269–2300. arXiv:0911.2187 (<https://arxiv.org/abs/0911.2187>). Bibcode:2010GReGr..42.2269B (<https://ui.adsabs.harvard.edu/abs/2010GReGr..42.2269B>). doi:10.1007/s10714-010-0988-2 (<https://doi.org/10.1007%2Fs10714-010-0988-2>). S2CID 118635353 (<https://api.semanticscholar.org/CorpusID:118635353>).
398. Kovacs, Z.; Cheng, K. S.; Harko, T. (2009). "Can stellar mass black holes be quark stars?" (<https://doi.org/10.1111%2Fj.1365-2966.2009.15571.x>). *Monthly Notices of the Royal Astronomical Society*. **400** (3): 1632–1642. arXiv:0908.2672 (<https://arxiv.org/abs/0908.2672>). Bibcode:2009MNRAS.400.1632K (<https://ui.adsabs.harvard.edu/abs/2009MNRAS.400.1632K>). doi:10.1111/j.1365-2966.2009.15571.x (<https://doi.org/10.1111%2Fj.1365-2966.2009.15571.x>). S2CID 18263809 (<https://api.semanticscholar.org/CorpusID:18263809>).
399. Kusenko, A. (2006). "Properties and signatures of supersymmetric Q-balls". arXiv:hep-ph/0612159 (<https://arxiv.org/abs/hep-ph/0612159>).

400. Hansson, J.; Sandin, F. (2005). "Preon stars: a new class of cosmic compact objects". *Physics Letters B*. **616** (1–2): 1–7. arXiv:astro-ph/0410417 (<https://arxiv.org/abs/astro-ph/0410417>). Bibcode:2005PhLB..616....1H (<https://ui.adsabs.harvard.edu/abs/2005PhLB..616....1H>). doi:10.1016/j.physletb.2005.04.034 (<https://doi.org/10.1016%2Fj.physletb.2005.04.034>). S2CID 119063004 (<https://api.semanticscholar.org/CorpusID:119063004>).
401. Murk, Sebastian (2023). "Nomen non est omen: Why it is too soon to identify ultra-compact objects as black holes". *International Journal of Modern Physics D*. **32** (14) 2342012: 2342012–2342235. arXiv:2210.03750 (<https://arxiv.org/abs/2210.03750>). Bibcode:2023IJMPD..3242012M (<https://ui.adsabs.harvard.edu/abs/2023IJMPD..3242012M>). doi:10.1142/S0218271823420129 (<https://doi.org/10.1142%2FS0218271823420129>). S2CID 252781040 (<https://api.semanticscholar.org/CorpusID:252781040>).
402. Mazur, Pawel O.; Mottola, Emil (2004). "Gravitational vacuum condensate stars" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC470711>). *Proceedings of the National Academy of Sciences of the United States of America*. **101** (26): 9545–9550. arXiv:gr-qc/0407075 (<https://arxiv.org/abs/gr-qc/0407075>). Bibcode:2004PNAS..101.9545M (<https://ui.adsabs.harvard.edu/abs/2004PNAS..101.9545M>). doi:10.1073/pnas.0402717101 (<https://doi.org/10.1073%2Fpnas.0402717101>). PMC 470711 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC470711>). PMID 15210982 (<https://pubmed.ncbi.nlm.nih.gov/15210982>). S2CID 2607263 (<https://api.semanticscholar.org/CorpusID:2607263>).
403. Choi, Charles Q. (2018). "Black Hole Pretenders Could Really Be Bizarre Quantum Stars" (<https://www.scientificamerican.com/article/black-hole-pretenders-could-really-be-bizarre-quantum-stars/>). *Scientific American*. Archived (<https://web.archive.org/web/20190617164631/https://www.scientificamerican.com/article/black-hole-pretenders-could-really-be-bizarre-quantum-stars/>) from the original on 17 June 2019. Retrieved 17 March 2018.
404. McRae, Mike (20 February 2024). "Bubble-Like 'Stars Within Stars' Could Explain Black Hole Weirdness" (<https://www.sciencealert.com/bubble-like-stars-within-stars-could-explain-black-hole-weirdness>). *ScienceAlert*. Archived (<https://archive.today/20240220150012/https://www.sciencealert.com/bubble-like-stars-within-stars-could-explain-black-hole-weirdness>) from the original on 20 February 2024. Retrieved 20 February 2024.
405. Ball, Philip (31 March 2005). "Black holes 'do not exist' ". *Nature*. doi:10.1038/news050328-8 (<https://doi.org/10.1038%2Fnews050328-8>).
406. Hawking, S. W. (1971). "Gravitational Radiation from Colliding Black Holes". *Physical Review Letters*. **26** (21): 1344–1346. Bibcode:1971PhRvL..26.1344H (<https://ui.adsabs.harvard.edu/abs/1971PhRvL..26.1344H>). doi:10.1103/PhysRevLett.26.1344 (<https://doi.org/10.1103%2FPhysRevLett.26.1344>).
407. Wald, R. M. (2001). "The Thermodynamics of Black Holes" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5253844>). *Living Reviews in Relativity*. **4** (1) 6. arXiv:gr-qc/9912119 (<https://arxiv.org/abs/gr-qc/9912119>). Bibcode:2001LRR.....4....6W (<https://ui.adsabs.harvard.edu/abs/2001LRR.....4....6W>). doi:10.12942/lrr-2001-6 (<https://doi.org/10.12942%2Flrr-2001-6>). PMC 5253844 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5253844>). PMID 28163633 (<https://pubmed.ncbi.nlm.nih.gov/28163633>).
408. 't Hooft, G. (2001). "The Holographic Principle". In Zichichi, A. (ed.). *Basics and Highlights in Fundamental Physics*. Subnuclear series. Vol. 37. World Scientific Publishing. pp. 72–100. arXiv:hep-th/0003004 (<https://arxiv.org/abs/hep-th/0003004>). Bibcode:2001bhfp.conf...72T (<https://ui.adsabs.harvard.edu/abs/2001bhfp.conf...72T>). doi:10.1142/9789812811585\_0005 ([https://doi.org/10.1142%2F9789812811585\\_0005](https://doi.org/10.1142%2F9789812811585_0005)). ISBN 978-981-02-4536-8. S2CID 119383028 (<https://api.semanticscholar.org/CorpusID:119383028>).
409. Kiefer, C. (12 January 2006). "Quantum gravity: general introduction and recent developments" (<https://onlinelibrary.wiley.com/doi/10.1002/andp.200651801-210>). *Annalen der Physik*. **518** (1–2): 129–148. doi:10.1002/andp.200651801-210 (<https://doi.org/10.1002%2Fandp.200651801-210>). ISSN 0003-3804 (<https://search.worldcat.org/issn/0003-3804>).

410. Strominger, A.; Vafa, C. (1996). "Microscopic origin of the Bekenstein-Hawking entropy". *Physics Letters B*. **379** (1–4): 99–104. arXiv:hep-th/9601029 (<https://arxiv.org/abs/hep-th/9601029>). Bibcode:1996PhLB..379...99S (<https://ui.adsabs.harvard.edu/abs/1996PhLB..379...99S>). doi:10.1016/0370-2693(96)00345-0 (<https://doi.org/10.1016%2F0370-2693%2896%2900345-0>). S2CID 1041890 (<https://api.semanticscholar.org/CorpusID:1041890>).
411. Carlip, S. (2009). "Black Hole Thermodynamics and Statistical Mechanics". *Physics of Black Holes*. Lecture Notes in Physics. Vol. 769. Berlin: Springer. pp. 89–123. arXiv:0807.4520 (<https://arxiv.org/abs/0807.4520>). Bibcode:2009LNP...769...89C (<https://ui.adsabs.harvard.edu/abs/2009LNP...769...89C>). doi:10.1007/978-3-540-88460-6\_3 ([https://doi.org/10.1007%2F978-3-540-88460-6\\_3](https://doi.org/10.1007%2F978-3-540-88460-6_3)). ISBN 978-3-540-88459-0. S2CID 15877702 (<https://api.semanticscholar.org/CorpusID:15877702>).
412. Hawking, S. W. "Does God Play Dice?" (<https://web.archive.org/web/20120111012413/http://www.hawking.org.uk/does-god-play-dice.html>). *www.hawking.org.uk*. Archived from the original (<http://www.hawking.org.uk/does-god-play-dice.html>) on 11 January 2012. Retrieved 14 March 2009.
413. Giddings, S. B. (1995). "The black hole information paradox". *Particles, Strings and Cosmology*. Johns Hopkins Workshop on Current Problems in Particle Theory 19 and the PASCOS Interdisciplinary Symposium 5. arXiv:hep-th/9508151 (<https://arxiv.org/abs/hep-th/9508151>). Bibcode:1995hep.th....8151G (<https://ui.adsabs.harvard.edu/abs/1995hep.th....8151G>).
414. Unruh, William G.; Wald, Robert M. (2017). "Information loss". *Reports on Progress in Physics*. **80** (9): 092002. arXiv:1703.02140 (<https://arxiv.org/abs/1703.02140>). Bibcode:2017RPPh...80i2002U (<https://ui.adsabs.harvard.edu/abs/2017RPPh...80i2002U>). doi:10.1088/1361-6633/aa778e (<https://doi.org/10.1088%2F1361-6633%2Faa778e>). PMID 28585922 (<https://pubmed.ncbi.nlm.nih.gov/28585922>). S2CID 39957660 (<https://api.semanticscholar.org/CorpusID:39957660>).
415. Mathur, S. D. (2011). *The information paradox: conflicts and resolutions*. XXV International Symposium on Lepton Photon Interactions at High Energies. *Pramana*. Vol. 79, no. 5. pp. 1059–1073. arXiv:1201.2079 (<https://arxiv.org/abs/1201.2079>). Bibcode:2012Prama..79.1059M (<https://ui.adsabs.harvard.edu/abs/2012Prama..79.1059M>). doi:10.1007/s12043-012-0417-z (<https://doi.org/10.1007%2Fs12043-012-0417-z>).
416. Page, Don N. (1993). "Information in black hole radiation". *Phys. Rev. Lett.* **71** (23): 3743–3746. arXiv:hep-th/9306083 (<https://arxiv.org/abs/hep-th/9306083>). Bibcode:1993PhRvL..71.3743P (<https://ui.adsabs.harvard.edu/abs/1993PhRvL..71.3743P>). CiteSeerX 10.1.1.267.174 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.267.174>). doi:10.1103/PhysRevLett.71.3743 (<https://doi.org/10.1103%2FPhysRevLett.71.3743>). PMID 10055062 (<https://pubmed.ncbi.nlm.nih.gov/10055062>). S2CID 9363821 (<https://api.semanticscholar.org/CorpusID:9363821>).
417. Page, Don N. (1993). "Average entropy of a subsystem". *Phys. Rev. Lett.* **71** (9): 1291–1294. arXiv:gr-qc/9305007 (<https://arxiv.org/abs/gr-qc/9305007>). Bibcode:1993PhRvL..71.1291P (<https://ui.adsabs.harvard.edu/abs/1993PhRvL..71.1291P>). CiteSeerX 10.1.1.339.7694 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.339.7694>). doi:10.1103/PhysRevLett.71.1291 (<https://doi.org/10.1103%2FPhysRevLett.71.1291>). PMID 10055503 (<https://pubmed.ncbi.nlm.nih.gov/10055503>). S2CID 17058654 (<https://api.semanticscholar.org/CorpusID:17058654>).
418. Merali, Zeeya (3 April 2013). "Astrophysics: Fire in the hole!" (<https://doi.org/10.1038%2F496020a>). *Nature*. **496** (7443): 20–23. Bibcode:2013Natur.496...20M (<https://ui.adsabs.harvard.edu/abs/2013Natur.496...20M>). doi:10.1038/496020a (<https://doi.org/10.1038%2F496020a>). PMID 23552926 (<https://pubmed.ncbi.nlm.nih.gov/23552926>).

419. Amheiri, Ahmed; Marolf, Donald; Polchinski, Joseph; Sully, James (2013). "Black holes: Complementarity or Firewalls?". *Journal of High Energy Physics*. **2013** (2): 62. arXiv:1207.3123 (<https://arxiv.org/abs/1207.3123>). Bibcode:2013JHEP...02..062A (<https://ui.adsabs.harvard.edu/abs/2013JHEP...02..062A>). doi:10.1007/JHEP02(2013)062 ([https://doi.org/10.1007/JHEP02\(2013\)062](https://doi.org/10.1007/JHEP02(2013)062)). S2CID 55581818 (<https://api.semanticscholar.org/CorpusID:55581818>).

## Sources

---

- Carroll, Sean M. (2004). *Spacetime and Geometry*. Addison Wesley. ISBN 978-0-8053-8732-2., the lecture notes on which the book was based are available for free from Sean Carroll's website (<https://www.preposterousuniverse.com/spacetimeandgeometry/>) Archived (<https://web.archive.org/web/20170323013522/http://www.preposterousuniverse.com/spacetimeandgeometry/>) 23 March 2017 at the Wayback Machine
- Hawking, S. W.; Ellis, G. F. R. (1973). *Large Scale Structure of space time* ([https://books.google.com/books?id=QagG\\_KI7LI8C](https://books.google.com/books?id=QagG_KI7LI8C)). Cambridge University Press. ISBN 978-0-521-09906-6. Archived ([https://web.archive.org/web/20200721064932/https://books.google.com/books?id=QagG\\_KI7LI8C](https://web.archive.org/web/20200721064932/https://books.google.com/books?id=QagG_KI7LI8C)) from the original on 21 July 2020. Retrieved 16 May 2020.
- Misner, Charles; Thorne, Kip S.; Wheeler, John (1973). *Gravitation*. W. H. Freeman and Company. ISBN 978-0-7167-0344-0.
- Thorne, Kip S. (1994). *Black Holes and Time Warps*. Norton, W. W. & Company, Inc. ISBN 978-0-393-31276-8.
- Wald, Robert M. (1984). *General Relativity* (<https://books.google.com/books?id=9S-hzg6-moYC>). University of Chicago Press. ISBN 978-0-226-87033-5. Archived (<https://web.archive.org/web/20160811035125/https://books.google.com/books?id=9S-hzg6-moYC>) from the original on 11 August 2016. Retrieved 23 February 2016.
- Wheeler, J. Craig (2007). *Cosmic Catastrophes* (2nd ed.). Cambridge University Press. ISBN 978-0-521-85714-7.

## Further reading

---

### Popular reading

- Begelman, Mitchell C.; Rees, Martin J. (2021). *Gravity's fatal attraction: black holes in the universe* (<https://books.google.com/books?id=ZPwAEAAAQBAJ>) (3rd ed.). Cambridge, United Kingdom ; New York, NY: Cambridge University Press. ISBN 978-1-108-87112-9.
- Ferguson, Kitty (1991). *Black Holes in Space-Time*. Watts Franklin. ISBN 978-0-531-12524-3.
- Hawking, Stephen (1988). *A Brief History of Time*. Bantam Books, Inc. ISBN 978-0-553-38016-3.
- Hawking, Stephen; Penrose, Roger (1996). *The Nature of Space and Time* (<https://books.google.com/books?id=LstaQTXP65cC>). Princeton University Press. ISBN 978-0-691-03791-2. Archived (<https://web.archive.org/web/20211018031528/https://books.google.com/books?id=LstaQTXP65cC>) from the original on 18 October 2021. Retrieved 16 May 2020.
- Levin, Janna (2020). *Black hole survival guide* (<https://books.google.com/books?id=9rITEAAQBAJ>). New York: Alfred A. Knopf. ISBN 978-0-525-65822-1. Archived ([https://web.archive.org/web/20220322141933/https://www.google.com/books/edition/Black\\_Hole\\_Survival\\_Guide/9rITEAAQBAJ?hl=en&gbpv=1&dq=janna+levin&printsec=frontcover](https://web.archive.org/web/20220322141933/https://www.google.com/books/edition/Black_Hole_Survival_Guide/9rITEAAQBAJ?hl=en&gbpv=1&dq=janna+levin&printsec=frontcover)) from the original on 22 March 2022. Retrieved 6 November 2021.



- Melia, Fulvio (2003). *The Black Hole at the Center of Our Galaxy* (<https://archive.org/details/blackholeatcente0000meli>). Princeton U Press. ISBN 978-0-691-09505-9.
- Melia, Fulvio (2003). *The Edge of Infinity. Supermassive Black Holes in the Universe* (<http://archive.org/details/edgeofinfinitysu00meli>). Cambridge U Press. ISBN 978-0-521-81405-8.
- Pickover, Clifford (1998). *Black Holes: A Traveler's Guide*. Wiley, John & Sons, Inc. ISBN 978-0-471-19704-1.
- Susskind, Leonard (2008). *The black hole war: my battle with Stephen Hawking to make the world safe for quantum mechanics* (1st ed.). New York: Little, Brown. ISBN 978-0-316-01640-7. OCLC 181603165 (<https://search.worldcat.org/oclc/181603165>).

## University textbooks and monographs

- Carter, B. (1973). "Black hole equilibrium states". In DeWitt-Morette, Cécile; DeWitt, Bryce S. (eds.). *Black holes*. New York: Gordon and Breach. ISBN 978-0-677-15610-1.
- Chandrasekhar, Subrahmanyan (1999). *Mathematical Theory of Black Holes*. Oxford University Press. ISBN 978-0-19-850370-5.
- Frolov, Valeri P.; Novikov, Igor D. (1998). *Black Hole Physics. Fundamental Theories of Physics*. Vol. 96. doi:10.1007/978-94-011-5139-9 (<https://doi.org/10.1007%2F978-94-011-5139-9>). ISBN 978-0-7923-5146-7.
- Frolov, Valeri P.; Zelnikov, Andrei (2011). *Introduction to Black Hole Physics* ([https://books.google.com/books?id=r\\_I5AK9DdXsC&pg=PA34](https://books.google.com/books?id=r_I5AK9DdXsC&pg=PA34)). Oxford: Oxford University Press. ISBN 978-0-19-969229-3. Zbl 1234.83001 (<https://zbmath.org/?format=complete&q=an:1234.83001>). Archived ([https://web.archive.org/web/20220322142008/https://books.google.com/books?id=r\\_I5AK9DdXsC&pg=PA34](https://web.archive.org/web/20220322142008/https://books.google.com/books?id=r_I5AK9DdXsC&pg=PA34)) from the original on 22 March 2022. Retrieved 2 January 2022.
- Melia, Fulvio (2007). *The Galactic Supermassive Black Hole*. Princeton U Press. ISBN 978-0-691-13129-0.
- Taylor, Edwin F.; Wheeler, John Archibald (2000). *Exploring Black Holes*. Addison Wesley Longman. ISBN 978-0-201-38423-9.
- Wald, Robert M. (1992). *Space, Time, and Gravity: The Theory of the Big Bang and Black Holes*. University of Chicago Press. ISBN 978-0-226-87029-8.
- Price, Richard; Creighton, Teviet (2008). "Black holes" (<https://doi.org/10.4249%2Fscholarpedia.4277>). *Scholarpedia*. **3** (1): 4277. Bibcode:2008SchpJ...3.4277C (<https://ui.adsabs.harvard.edu/abs/2008SchpJ...3.4277C>). doi:10.4249/scholarpedia.4277 (<https://doi.org/10.4249%2Fscholarpedia.4277>).

## Review papers

- Hughes, Scott A. (2005). "Trust but verify: The case for astrophysical black holes". arXiv:hep-ph/0511217 (<https://arxiv.org/abs/hep-ph/0511217>). Lecture notes from 2005 SLAC Summer Institute.
- Gallo, Elena; Marolf, Donald (2009). "Resource Letter BH-2: Black Holes". *American Journal of Physics*. **77** (4): 294–307. arXiv:0806.2316 (<https://arxiv.org/abs/0806.2316>). Bibcode:2009AmJPh..77..294G (<https://ui.adsabs.harvard.edu/abs/2009AmJPh..77..294G>). doi:10.1119/1.3056569 (<https://doi.org/10.1119%2F1.3056569>). S2CID 118494056 (<https://api.semanticscholar.org/CorpusID:118494056>).
- Cardoso, Vitor; Pani, Paolo (2019). "Testing the nature of dark compact objects: a status report". *Living Reviews in Relativity*. **22** (1): 4. arXiv:1904.05363 (<https://arxiv.org/abs/1904.05363>). Bibcode:2019LRR....22....4C (<https://ui.adsabs.harvard.edu/abs/2019LRR....22....4>).

C). doi:10.1007/s41114-019-0020-4 (<https://doi.org/10.1007%2Fs41114-019-0020-4>). S2CID 256465740 (<https://api.semanticscholar.org/CorpusID:256465740>).

- Mann, Robert B.; Murk, Sebastian; Terno, Daniel R. (2022). "Black holes and their horizons in semiclassical and modified theories of gravity". *International Journal of Modern Physics D*. **31** (9): 2230015–2230276. arXiv:2112.06515 (<https://arxiv.org/abs/2112.06515>). Bibcode:2022IJMPD..3130015M (<https://ui.adsabs.harvard.edu/abs/2022IJMPD..3130015M>). doi:10.1142/S0218271822300154 (<https://doi.org/10.1142%2FS0218271822300154>). S2CID 245123647 (<https://api.semanticscholar.org/CorpusID:245123647>).

## External links

---

- Black Holes (<https://www.bbc.co.uk/programmes/p00547f4>) on *In Our Time* at the BBC
- *Stanford Encyclopedia of Philosophy*: "Singularities and Black Holes (<https://plato.stanford.edu/entries/spacetime-singularities/>)" by Erik Curiel and Peter Bokulich.
- Black Holes: Gravity's Relentless Pull (<https://hubblesite.org/contents/articles/black-holes.html>) – Interactive multimedia Web site about the physics and astronomy of black holes from the Space Telescope Science Institute (HubbleSite)
- ESA's Black Hole Visualization (<https://www.esa.int/gsp/ACT/phy/Projects/Blackholes/WebGL.html>) Archived (<https://web.archive.org/web/20190503070935/https://www.esa.int/gsp/ACT/phy/Projects/Blackholes/WebGL.html>) 3 May 2019 at the Wayback Machine
- Frequently Asked Questions (FAQs) on Black Holes ([https://apod.nasa.gov/htmltest/gifcity/bh\\_pub\\_faq.html](https://apod.nasa.gov/htmltest/gifcity/bh_pub_faq.html))
- Schwarzschild Geometry (<https://web.archive.org/web/19980118051503/http://casa.colorado.edu/~ajsh/schwp.html>)
- Black holes - basic (NYT; April 2021) (<https://www.nytimes.com/2021/04/22/science/black-holes-astrophysics-names.html>)

## Videos

- 16-year-long study tracks stars orbiting Sagittarius A\* (<https://www.eso.org/public/videos/eso0846b/>)
- Movie of Black Hole Candidate from Max Planck Institute (<https://web.archive.org/web/20040925044354/http://www.mpe.mpg.de/ir/GC/index.php>)
- Cowen, Ron (20 April 2015). "3D simulations of colliding black holes hailed as most realistic yet" (<https://doi.org/10.1038%2Fnature.2015.17360>). *Nature*. doi:10.1038/nature.2015.17360 (<https://doi.org/10.1038%2Fnature.2015.17360>).
- Computer visualisation of the signal detected by LIGO (<https://www.bbc.com/news/science-environment-35524440>)
- Two Black Holes Merge into One (based upon the signal GW150914) ([https://www.youtube.com/watch?v=I\\_88S8DWbcU](https://www.youtube.com/watch?v=I_88S8DWbcU))

---

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Black\\_hole&oldid=1325611753](https://en.wikipedia.org/w/index.php?title=Black_hole&oldid=1325611753)"