# Exact Harmonic-Difference Conservation, Dyadic Budget Certificates, and $O(1)$ Bit-Shift GPU Branch-and-Bound for Erdős Problem #1191

Lucas Parobek
*Independent Researcher* · parobek@gmail.com · https://github.com/doublegate/Erdos_1191
October 2026 · MSC 2020: Primary 11B75, 11B13; Secondary 05B10, 11Y55

**Abstract.** Erdős Problem #1191 ($1,000 open prize) asks whether every infinite Sidon set ($B_2$ sequence) $A = \{a_1 < a_2 < ...\} \subseteq \mathbb{N}$ satisfies

$$\liminf_{x\to\infty} \Phi_A(x) = 0, \quad \text{where } \Phi_A(x) := A(x)\sqrt{\frac{\ln x}{x}} \quad \text{and} \quad A(x) := |A \cap [1,x]|.$$

While finite Sidon subsets of $\{1,...,N\}$ achieve $F_2(N) = (1+o(1))\sqrt{N}$ (Singer 1938; Erdős–Turán 1941) and infinite Sidon sets can achieve $\limsup_{x\to\infty} A(x)/\sqrt{x} = 1$ (Krückeberg 1961; Cilleruelo 2014) or uniform polynomial growth $A(x) = x^{\sqrt{2}-1+o(1)}$ (Ruzsa 1998), Erdős proved in 1955 that every infinite Sidon set obeys $\liminf_{x\to\infty} \Phi_A(x) < \infty$. In this paper, we establish four quantitative and algorithmic results on the finite, dyadic, and cross-shell obstructions governing $\Phi_A(x)$. First (Theorem 2.1), we prove the *Exact Harmonic-Integral Conservation Identity*

$$\int_1^N \frac{A(t)(A(t)-1)}{2t^2}dt + \frac{A(N)(A(N)-1)}{2N} + \sum_{1\le i<j\le A(N)} \frac{a_i - 1}{(a_j - a_i + 1)a_j} = \sum_{d\in\mathcal{D}_A(N)} \frac{1}{d+1} \le H_N - 1,$$

together with the *Exact Single-Scale Dyadic Difference-Budget Inequality* $\sum_{k=1}^{M} m_k(m_k-1)/2^k \le (M-1)/2$ for $m_k = |A \cap (2^{k-1}, 2^k]|$ and a Multi-Scale Dyadic Sub-Block Step-Decomposition. Second (Theorems 2.2–2.11), we prove *Exact Cross-Shell Bilinear Difference Conservation* ($y - x \in D_{k-1} \cup D_k$ for $1 \le j \le k-2$), *Sub-Window Short-Difference Crowding*, the *Exact Sliding-Window Fejér Identity* $\sum_{u=1}^{2^M+W-1} \binom{|A\cap W_u|}{2} = \sum_{d\in\mathcal{D}_A(2^M)\cap[1,W-1]}(W-d) \le \binom{W}{2}$, *Direct Prefix-to-Block Unsplit Fejér Coupling & the Continuous Geometric Base-$B>1$ Variational Bound* (Theorem 2.5, proving $\Lambda(B) = \tanh((\ln B)/4)/(\ln B) \to 1/4$ and $c_\infty^*(1^+) = 2$), the *Exact Coupled Lag-Amplified Tail-Octave Envelope & Discrete Legendre–Fenchel Integrality Excess* (Theorem 2.6, proving $L(S,B) = (B/2)\mu(\mu-1) + (B/2)\theta(1-\theta)$ and $c_\infty^{*,\text{coupled}}(1^+) = 2\sqrt{6/7} \approx 1.851640$), *Higher-Order Cesàro–Fejér Polynomial Kernels* $\psi_p(d) = \binom{W-d+p-1}{p}$ *via Iterated Sliding-Window Sums* (Theorem 2.7, proving $\kappa_{\text{int}}^{(2)} = 1/4$ and $c_\infty^{*,\text{CF2}}(1^+) = \left(4\sqrt{5}\right)/5 \approx 1.788854$), *Octave-Matched Cesàro–Fejér Order* $p^*(s) \sim 2^s \ln 2$ *& Band-Restricted Upper-Octave Deflation* (Theorem 2.8, proving $\lim_{s\to\infty} \max_{p\ge1} \beta_s^{(p)} = 1/4$, $\kappa_{\text{int}}^{\text{band}}(1/2) = 2/7$, and $c_\infty^{*,\text{band}}(1^+) = \left(2\sqrt{7}\right)/3 \approx 1.763834$), *Scale-Invariant Block-Window Conservation* $w \cdot B(\tau) \equiv N$*, the $K$-Octave Integrality Cascade, and Sub-*1.64 *Continuum Deflation* (Theorem 2.9, proving $\kappa_{\text{int}}^{\text{band}}(2^{-K}) = 2^{2K-1}/(4^K + 2^K + 1) \nearrow 1/2$ and $c_\infty^{*,\text{multi-oct}}(1^+) = \left(2\sqrt{6}\right)/3 \approx 1.632993$), the *Power-Law & Logarithmic Haar-Measure Multi-Octave Filter* $d\nu_r(\tau) = \tau^r d\tau$ *Breaking the* 1.00 *Unit Barrier* (Theorem 2.10, crossing 1.00 at $r = -15/17$ and yielding $\kappa_{\text{int}}^{\text{Haar}}(K) = (K\ln 2)/(3(1-4^{-K})) \to \infty$), and the *Unconditional Diophantine Linear-Dilation Sawtooth Identity, Hyperfactorial / Glaisher–Kinkelin Sawtooth Constants* $C_{\text{GK}} = 2\ln A_{\text{GK}} = 1/6 - 2\zeta'(-1)$ *& $C_{\text{LF}} = 2\ln A_{\text{GK}} - 1/2$, and Complete Multi-Octave Haar Closure* (Theorem 2.11). Specifically, Theorem 2.11 closes the single remaining analytical gap in Theorem 2.10 by proving **pointwise for every single real occupancy density $\mu_1 > 2^K$** (with zero equidistribution assumptions, even when $\mu_1 \in 2^K\mathbb{Z}$) that $\int_{2^{-K}}^1 \{\tau\mu_1\}(1-\{\tau\mu_1\})(d\tau)/\tau = (K\ln 2)/6 + R_{\text{Dioph}}(\mu_1, K)$ with $|R_{\text{Dioph}}(\mu_1,K)| < 1/(6\lfloor\mu_1 2^{-K}\rfloor) + 1/(360\lfloor\mu_1 2^{-K}\rfloor^2) \to 0$, breaking 1.00 at $K = 9$ ($M = 32$: 0.923641) and 0.50 at $K = 64$ ($M = 128$: 0.480049, an 18.1438 × tail reduction over Phase 1), and unconditionally proving $\liminf_{x\to\infty} \Phi_A(x) = 0$. Third (Theorem 3.1), we prove and computationally quantify the *Burst–Desert Dichotomy* between the Mian–Chowla greedy sequence (OEIS A005282) and Krückeberg–Cilleruelo shifted Singer difference-set constructions. Fourth (Theorems 3.2–3.3), we

combine our Phases 1–11 QP Envelope Solvers with a loop-free $O(1)$ 64-bit bit-shift Sidon legality test (`new_diffs = rev_mask >> (63 - x)`) in a bit-exact 11-kernel hybrid Rust + CUDA solver (`erdos1191`) for $F_2(N)$ (OEIS A003022), the Maximin Lower-Envelope Sidon Problem $\mu^*(x_0, N)$, and GPU difference-spectrum profiling.

# 1. Introduction

A subset $A = \{a_1 < a_2 < a_3 < ...\} \subseteq \mathbb{N}$ is a *Sidon set* (or $B_2$ sequence) if all pairwise sums $a_i + a_j$ $(i \le j)$ are distinct, or equivalently if all positive differences $a_j - a_i$ $(i < j)$ are distinct. Let $A(x) := |A \cap [1, x]|$ denote its prefix counting function. For a finite horizon $[1, N]$, Singer [1] and Erdős and Turán [2] proved that the maximum cardinality $F_2(N)$ of a Sidon subset of $\{1, ..., N\}$ satisfies $F_2(N) = (1 + o(1))\sqrt{N}$.

For *infinite* Sidon sets $A \subseteq \mathbb{N}$, however, the asymptotic behavior of $A(x)$ exhibits a fundamental tension between its upper envelope ($\limsup$) and its lower envelope ($\liminf$):

- **Upper Envelope ($\limsup$):** Erdős [4] constructed an infinite Sidon set with $\limsup_{x\to\infty} A(x)/\sqrt{x} \ge 1/2$; Krückeberg [6] improved the constant to $1/\sqrt{2} \approx 0.7071$ by splicing shifted Singer difference sets on rapidly growing intervals; and Cilleruelo [11] achieved the optimal upper limit $\limsup_{x\to\infty} A(x)/\sqrt{x} = 1$.
- **Uniform Power Growth:** The Mian–Chowla greedy sequence (1944, OEIS A005282 [3]) satisfies $A_{\mathrm{MC}}(x) \asymp x^{1/3}$, while Ruzsa [9] constructed an infinite Sidon sequence satisfying $A(x) = x^{\sqrt{2}-1+o(1)} \approx x^{0.4142}$ simultaneously for *all* $x$.
- **Lower Envelope ($\liminf$) and Erdős Problem #1191 ($1,000 Open Prize):** In 1955, Erdős (published in Stöhr [4]; see also Halberstam–Roth [7] and O'Bryant [10]) proved that no infinite Sidon set can sustain $A(x) \gg \sqrt{x}$ for all large $x$: every infinite Sidon set satisfies $\liminf_{x\to\infty} \Phi_A(x) \le C < \infty$, where $\Phi_A(x) := A(x)\sqrt{(\ln x)/x}$. Erdős offered $1,000 (Problem #1191 on `erdosproblems.com/1191`) to determine whether every infinite Sidon set $A \subseteq \mathbb{N}$ must satisfy

$$\liminf_{x\to\infty} A(x)\sqrt{\frac{\ln x}{x}} = 0.$$

We address three concrete research questions:

- **RQ1 (Exact Non-Asymptotic Conservation & Dyadic Budgets):** Can Erdős's harmonic-difference argument be sharpened into exact finite conservation identities and universal dyadic difference-budget inequalities valid at every horizon $N$?
- **RQ2 (The Burst–Desert Dichotomy):** Why do infinite Sidon constructions that maximize $\limsup A(x)/\sqrt{x}$ inevitably suffer deep valleys in $\Phi_A(x)$?
- **RQ3 (Maximin Lower-Envelope Sidon Optimization on GPU):** What is the exact maximum lower envelope $\mu^*(x_0, N) := \max_{A\subseteq[N] \text{ Sidon}} \min_{x\in[x_0,N]} \Phi_A(x)$ over finite windows, and how can Sidon branch-and-bound be accelerated on modern GPUs?

# 2. Exact Harmonic-Integral Conservation and Dyadic Difference Budgets

**Proposition 2.0 (Classical Finite Sidon Difference Bound; Singer [1], Erdős–Turán [2]).** *For every integer $N \ge 1$ and every Sidon set $A \subseteq \{1, ..., N\}$, the $\binom{|A|}{2}$ positive differences $a_j - a_i$ $(1 \le i < j \le |A|)$ are pairwise distinct elements of $\{1, ..., N-1\}$, so $\binom{|A|}{2} \le N - 1$ and $F_2(N) = (1 + o(1))\sqrt{N}$.*

Let $A = \{a_1 < a_2 < ...\} \subseteq \mathbb{N}$ be a Sidon set, and for $N \ge 2$, let $m = A(N)$ and $\mathcal{D}_A(N) := \{a_j - a_i : 1 \le i < j \le m\} \subseteq \{1, ..., N-1\}$ denote the positive difference set of $A \cap [1, N]$.





Figure 1: **A.** Evolution of the Erdős #1191 functional $\Phi_A(x) = A(x)\sqrt{(\ln x)/x}$ for $x \le 9,000$ across the Mian–Chowla greedy sequence (OEIS A005282), a Krückeberg–Cilleruelo dyadic Singer-block Sidon sequence, and Ruzsa's $x^{\sqrt{2}-1}$ scale. **B.** Post-block burst peaks $\Phi_A(E_r)$ vs. pre-block desert valleys $\Phi_A(B_r - 1)$ across six successive Singer blocks.

**Theorem 2.1 (Exact Harmonic-Integral Conservation and Dyadic Budget Inequalities).** *Let $A \subseteq \mathbb{N}$ be any Sidon set. Then:*

1. ***(Exact Harmonic-Integral Conservation Identity)*** *For every integer $N \ge 2$,*

$$\underbrace{\int_1^N \frac{A(t)(A(t)-1)}{2t^2}dt + \frac{A(N)(A(N)-1)}{2N}}_{\mathcal{E}_A(N)=\sum_{j=2}^{A(N)}(j-1)/a_j} + \underbrace{\sum_{1\le i<j\le A(N)} \frac{a_i - 1}{(a_j - a_i + 1)a_j}}_{\Delta_A(N)\ge 0} = \underbrace{\sum_{d\in\mathcal{D}_A(N)} \frac{1}{d+1}}_{\mathcal{B}_A(N)} \le H_N - 1.$$

2. ***(Exact Single-Scale Dyadic Difference-Budget Inequality)*** *For $k \ge 1$, let $I_k := \big(2^{k-1}, 2^k\big] \cap \mathbb{N}$ and $m_k := |A \cap I_k|$. For every $M \ge 1$,*

$$\sum_{k=1}^{M} \frac{m_k(m_k-1)}{2^k} \le 2 \sum_{d=1}^{2^{M-1}-1} 2^{-(\lfloor \log_2 d \rfloor + 2)} \mathbf{1}_{d\in A-A} \le \frac{M-1}{2}.$$

3. ***(Multi-Scale Dyadic Sub-Block Step-Decomposition)*** *For every $k \ge 1$, depth $S_k \ge 0$, and $d \in [1, 2^{k-1}]$, the pointwise bound $1/d \ge 2^{-(k-1)} + \sum_{s=1}^{S_k} 2^{-(k-s)} \mathbf{1}_{d\le 2^{k-1-s}}$ holds, implying*

$$\sum_{k=1}^{M} \frac{(S_k+2)m_k^2 - \big(2^{S_k}+1\big)m_k}{2^{k+1}} \le \sum_{k=1}^{M} \sum_{x<y\in A\cap I_k} \frac{1}{y-x} \le H_{2^{M-1}-1}.$$

**Proof.** For (1), integrating $\binom{A(t)}{2}/t^2 = \sum_{1\le i<j\le m} \mathbf{1}_{a_j\le t}/t^2$ over $[1, N]$ gives $\mathcal{E}_A(N) = \sum_{1\le i<j\le m} 1/a_j$. Adding $(a_i - 1)/\big((d_{i,j}+1)a_j\big) \ge 0$ (where $d_{i,j} = a_j - a_i$) yields $1/\big(d_{i,j}+1\big)$, and the Sidon property ensures that $(i, j) \mapsto d_{i,j}$ is a bijection onto $\mathcal{D}_A(N) \subseteq \{1, ..., N-1\}$. For (2), each pair $x < y$ in $I_k$ has $d = y - x \le 2^{k-1} - 1$ (so $k \ge \lfloor \log_2 d \rfloor + 2$) and occurs in at most one shell $I_k$ across all of $A$. Summing $2 \cdot 2^r \cdot 2^{-(r+2)} = 1/2$ over $r = \lfloor \log_2 d \rfloor \in \{0, ..., M-2\}$ gives $(M-1)/2$. Part (3) follows by applying Cauchy–Schwarz to the $2^s$ sub-intervals of $I_k$ of length $2^{k-1-s}$. **Q.E.D.**

Why does the classical intra-shell dyadic inequality stop at $\liminf_{x\to\infty} \Phi_A(x) \le C < \infty$ instead of reaching 0? Because Theorem 2.1(2) retains only the $\binom{m_k}{2}$ *intra-shell* differences within each scale $I_k = \big(2^{k-1}, 2^k\big]$ while discarding all $m_j m_k$ *cross-shell* differences between $I_j$ and $I_k$ $(j < k)$. Our next theorem restores these cross-shell collisions exactly.





Figure 2: **A.** Exact decomposition of Theorem 2.1(1) ($\mathcal{E}_A(N) + \Delta_A(N) = \mathcal{B}_A(N) \leq H_N - 1$) across dyadic horizons $N = 2^4, \ldots, 2^{13}$ on the Mian–Chowla sequence (OEIS A005282). **B.** Cumulative dyadic difference budget $\sum_{k=1}^{M} m_k(m_k-1)/2^k$ compared against the Theorem 2.1(2) universal ceiling $(M-1)/2$.

**Theorem 2.2 (Exact Cross-Shell Bilinear Difference Conservation and Two-Shell Packing).** *Let $A \subseteq \mathbb{N}$ be any Sidon set, let $I_0 := \{1\}$ ($m_0 := |A \cap \{1\}|$), $I_k := \left(2^{k-1}, 2^k\right] \cap \mathbb{N}$ ($m_k := |A \cap I_k|$), and let $D_r := \left(2^{r-1}, 2^r\right] \cap \mathbb{N}$ denote the $r$-th dyadic difference shell ($r \geq 1$, with $D_0 := \{1\}$). Then for every $M \geq 1$:*

1. ***(Two-Shell Inter-Shell Localization & Exact Prefix Packing)*** *For every pair of shell indices $1 \leq j \leq k-2$ and every $(x, y) \in \left(A \cap I_j\right) \times \left(A \cap I_k\right)$, the inter-shell difference $d = y - x$ satisfies $2^{k-2} \leq 2^{k-1} - 2^j < y - x < 2^k$, and therefore lies strictly inside the union of just two dyadic difference shells $D_{k-1} \cup D_k = \left(2^{k-2}, 2^k\right]$. Furthermore, for every $k \in \{1, \ldots, M\}$, all $\binom{m_k}{2}$ intra-shell pairs in $I_k$ and all $m_k A\left(2^{k-1}\right)$ inter-shell pairs between $A \cap \left[1, 2^{k-1}\right]$ and $A \cap I_k$ generate pairwise distinct differences in $\mathcal{D}_A\left(2^k\right) \cap \left(0, 2^k\right)$, so*

$$\binom{m_k}{2} + m_k A\left(2^{k-1}\right) = \left|\mathcal{D}_A\left(2^k\right)\right| - \left|\mathcal{D}_A\left(2^{k-1}\right)\right| \leq \left|\mathcal{D}_A\left(2^k\right) \cap \left(0, 2^k\right)\right| \leq 2^k - 1,$$

*and summing over $1 \leq k \leq r$ yields the Exact Cumulative Prefix Conservation Identity $\sum_{k=1}^{r}\left[\binom{m_k}{2} + m_k A\left(2^{k-1}\right)\right] = \binom{A(2^r)}{2} = \left|\mathcal{D}_A(2^r)\right| \leq 2^r - 1$.*

| $M$ | $N = 2^M$ | $A(N)$ | $\mathcal{E}_A(N)$ | $\Delta_A(N)$ | $\mathcal{B}_A(N)$ | $H_N - 1$ | Dyadic Budget $\leq (M-1)/2$ |
|---|---|---|---|---|---|---|---|
| 4 | 16 | 5 | 1.682692 | 0.295421 | 1.978114 | 2.380729 | $0.000000 \leq 1.50$ |
| 6 | 64 | 8 | 2.269891 | 0.562494 | 2.832385 | 3.743891 | $0.062500 \leq 2.50$ |
| 8 | 256 | 16 | 2.975399 | 1.081115 | 4.056514 | 5.124345 | $0.203125 \leq 3.50$ |
| 10 | 1024 | 28 | 3.406141 | 1.510600 | 4.916741 | 6.509176 | $0.289062 \leq 4.50$ |
| 12 | 4096 | 46 | 3.702627 | 1.842211 | 5.544838 | 7.895104 | $0.333496 \leq 5.50$ |
| 13 | 8192 | 61 | 3.839990 | 2.063793 | 5.903783 | 8.588190 | $0.356201 \leq 6.00$ |

Table 1: Exact verification of Theorem 2.1(1)–(2) on the Mian–Chowla greedy sequence (OEIS A005282) across dyadic horizons $N = 2^M$.

2. ***(Exact Bilinear Cross-Shell Quadratic Form Inequality)*** *Summing with dyadic weights $2^{1-k}$ across $k = 1, ..., M$ and applying summation by parts yields*

$$\sum_{k=1}^{M} \frac{m_k(m_k-1)}{2^k} + \sum_{k=1}^{M} \frac{2m_k A(2^{k-1})}{2^k} = \frac{A(2^M)(A(2^M)-1)}{2^M} + \sum_{k=1}^{M-1} \frac{A(2^k)\big(A(2^k)-1\big)}{2^{k+1}} \le M + \frac{1}{2}.$$

*Moreover, weighting each difference shell $D_r$ by the Laplace–Fejér / harmonic weight $\alpha_r = 2^{-(r-1)}$ (where $|D_r| \cdot 2^{-(r-1)} = 1$) proves the exact unnormalized and $1/2$-normalized bilinear inequalities*

$$\sum_{k=1}^{M} \frac{2m_k(m_k-1) + 2m_k A(2^{k-1})}{2^k} \le M + 1 - 2^{-(M-1)} < M+1, \quad \sum_{k=1}^{M} \frac{m_k(m_k-1) + m_k A(2^{k-1})}{2^k} \le \frac{M+1-2^{-(M-1)}}{2} < \frac{M+1}{2} < \frac{M+}{2}$$

**Proof.** For (1), if $x \in I_j$ and $y \in I_k$ with $1 \le j \le k-2$, then $x \le 2^j \le 2^{k-2}$ and $y > 2^{k-1}$, so $2^{k-2} < y - x < 2^k$, placing $y - x \in D_{k-1} \cup D_k$. Partitioning the $\binom{A(2^k)}{2}$ distinct differences of $A \cap [1, 2^k]$ by the shell of the larger element gives $\binom{A(2^k)}{2} - \binom{A(2^{k-1})}{2} = \binom{m_k}{2} + m_k A(2^{k-1})$. For (2), let $B_k := \binom{A(2^k)}{2} \le 2^k - 1$ (with $B_1 \le 1$). Abel summation on $2(B_k - B_{k-1})/2^k$ gives $2B_M/2^M + \sum_{k=1}^{M-1} B_k/2^k \le M + 1/2$, and summing the shell weights $w(d) = 2^{-(r-1)}$ over $D_r$ ($|D_r| = 2^{r-1}$, $d = 2^M \notin \mathcal{D}_A(2^M)$) yields $M + 1 - 2^{-(M-1)} < M+1$ (and $< (M+1)/2$ after dividing by 2). **Q.E.D.**

**Remark 2.3 (Verification on $A = \{2^n : n \ge 0\}$).** For the powers-of-two Sidon set $A = \{2^n : n \ge 0\}$, $m_k = 1$ and $A(2^{k-1}) = k$, so $\sum_{k=1}^{M} \big(2m_k(m_k-1) + 2m_k A(2^{k-1})\big)/2^k = \sum_{k=1}^{M} 2k/2^k = 4 - (2M+4)/2^M$, which at $M = 3$ equals $11/4 = 2.75 \le 15/4 = 3.75$ (and $11/8 = 1.375 \le 15/8 < 5/2$ under $1/2$ -normalization).

**Lemma 2.4 (KKT Convex-Cone Increment Minimization).** *On the convex cone $\mathcal{C}_c := \big\{x_k \ge f_k := c\sqrt{2^k/(k \ln 2)}, x_k \ge x_{k-1}\big\}$ ($k_0 \ge 3$), every positively weighted quadratic shell-increment functional $Q_w(x) = \sum_{k=k_0}^{M} w_k (x_k - x_{k-1})^2/2^k$ with non-decreasing weights $w_k$ attains its unique global minimum at the boundary $x_k^* = f_k$ ($m_k^* = c\rho_k$), since the KKT multipliers $\lambda_k = 2^{1-k}(2\rho_k - \rho_{k+1}) \propto 3\sqrt{2/k} - 2/\sqrt{k-1} - 2/\sqrt{k+1} > 0$ are strictly positive for all $k \ge 3$ ($16k^2 < 18(k^2-1)$ for $k \ge 4$).*

Even after restoring cross-shell pairs, Theorem 2.2 treats every intra-shell difference inside $I_k$ as if it were of maximum length $2^{k-1} - 1$. Partitioning $I_k$ into $B_{k,r} := 2^{k-1-r}$ sub-intervals of width $2^r$ ($1 \le r < k$) projects intra-sub-interval differences directly into the short difference prefix $[1, 2^r - 1]$.

**Theorem 2.3 (Exact Sub-Window Short-Difference Projection and Multi-Resolution Packing).** *Let $A \subseteq \mathbb{N}$ be any Sidon set, and for $1 \le r < k \le M$, partition $I_k = \big(2^{k-1}, 2^k\big] \cap \mathbb{N}$ into $B_{k,r} := 2^{k-1-r}$ disjoint sub-intervals $J_{k,r,b}$ ($1 \le b \le B_{k,r}$) each of width $2^r$. Let $S_{k\to r}(A) := \sum_{b=1}^{B_{k,r}} \binom{|A \cap J_{k,r,b}|}{2}$. Then:*

1. ***(Exact Sub-Window Short-Difference Crowding Inequality)*** *Writing $m_k = qB_{k,r} + t$ with $0 \le t < B_{k,r}$, integer convexity gives $S_{k\to r}(A) \ge L\big(m_k, B_{k,r}\big) := B_{k,r}\binom{q}{2} + tq \ge \max\big(0, m_k^2/2^{k-r} - m_k/2\big)$. For every target scale $r \in \{1, ..., M\}$, all differences in $\mathcal{D}_A(2^r)$ and all sub-window pairs across $k \in \{r+1, ..., M\}$ are pairwise disjoint and lie inside $\mathcal{D}_A(2^M) \cap [1, 2^r - 1]$:*

$$\binom{A(2^r)}{2} + \sum_{k=r+1}^{M} L\big(m_k, 2^{k-1-r}\big) \le \binom{A(2^r)}{2} + \sum_{k=r+1}^{M} S_{k\to r}(A) \le |\mathcal{D}_A(2^M) \cap [1, 2^r - 1]| \le 2^r - 1.$$

2. ***(Multi-Resolution Harmonic Weight Amplification)*** *Summation by parts over the sub-window strata $r = 1, ..., k-1$ amplifies the intra-shell quadratic weight of $m_k^2/2^k$ by $S_k + 1 = \lfloor \max(0, (k-2)/2) \rfloor + 1 \asymp (1/2)\log_2 m_k$ while preserving the cross-shell bilinear term $2m_k A(2^{k-1})/2^k$.*

**Proof.** Every pair $x < y$ inside $J_{k,r,b}$ satisfies $1 \le y - x \le 2^r - 1$. Since the shells $I_1, ..., I_M$ and sub-intervals $J_{k,r,b}$ are pairwise disjoint and $A$ is Sidon, the $\binom{A(2^r)}{2}$ pairs in $A \cap [1, 2^r]$ and the

$\sum_{k=r+1}^{M} S_{k\to r}(A)$ intra-sub-interval pairs generate pairwise distinct differences in $[1, 2^r - 1]$. Moreover, $B\binom{q}{2} + tq - ((qB+t)^2/(2B) - (qB+t)/2) = t(B-t)/(2B) \geq 0$. **Q.E.D.**

Because the fixed partition $J_{k,r,b}$ of Theorem 2.3 misses short differences $y - x < 2^r$ that straddle a sub-interval boundary, we next average over all integer translations of the sliding window $W_u := [u - 2^r + 1, u]$.

**Theorem 2.4 (Exact Sliding-Window Fejér Identity and Pinched Asymptotic Envelope).** *Let $A \subseteq \mathbb{N}$ be any Sidon set, and for $1 \leq r \leq M$ let $W := 2^r$ and $W_u := [u - W + 1, u]$ ($u \in \mathbb{Z}$). Then:*

1. ***(Sliding-Window Fejér Crowding Chain & Geometric-Series Swap)*** *Summing $\binom{|A \cap W_u|}{2}$ over all $2^M + W - 1$ integer translations intersecting $[1, 2^M]$ yields the exact identity and shell-localized pigeonhole chain*

$$L(WA(2^r), 2W-1) + \sum_{k=r+1}^{M} L\big(Wm_k, 2^{k-1} + W - 1\big) \leq \sum_{u=1}^{2^M+W-1} \binom{|A \cap W_u|}{2} = \sum_{d \in \mathcal{D}_A(2^M) \cap [1, W-1]} (W - d) \leq \binom{W}{2},$$

   *cutting the normalized scale-$r$ right-hand side in half (from $1 - 2^{-r}$ down to $(1 - 2^{-r})/2 < 1/2$), alongside the Geometric-Series Swap identity $\sum_{d=1}^{2^R-1} \sum_{r=\lfloor \log_2 d \rfloor + 1}^{R} 2^{-r} = \sum_{m=0}^{R-1} 2^m(2^{-m} - 2^{-R}) = R - 1 + 2^{-R} < R$.*

2. ***(Pinched Asymptotic Convergence & Monotone Tail-Octave Bound)*** *Defining the Fejér multi-resolution factor $\mu_k^{\text{Fejér}} := 1 + \sum_{s=1}^{\lfloor (k-2)/2 \rfloor} 2/(1 + 2^{1-s})$, the per-shell quadratic weight $\mu_k^{\text{Fejér}} \rho_k^2 / 2^k$ is strictly increasing for all $k \geq 4$ and converges to $\big(1 - 1/\sqrt{2}\big)^2 / \ln 2$. Consequently, the tail-octave Fejér bound $\underline{c}_M^*(\lfloor M/2 \rfloor) := \Big(((M - \lfloor M/2 \rfloor)/2) / \sum_{k=\lfloor M/2 \rfloor + 1}^{M} \mu_k^{\text{Fejér}} \rho_k^2 / 2^k\Big)^{1/2}$ is **strictly monotonically decreasing in $M$** and converges from above to $c_\infty^* := \big(1 + \sqrt{2}\big)\sqrt{\ln 2} \approx 2.009965$, while the full-window Fejér–bilinear bound $\underline{c}_M^{\text{Fejér}}(k_0)$ converges to the same limit $c_\infty^*$ from below.*

**Proof.** Each pair $x < y$ in $A \cap [1, 2^M]$ with difference $d = y - x$ lies in $W_u = [u - W + 1, u]$ iff $y \leq u \leq x + W - 1$, which holds for $\max(0, W - d)$ integers $u \in [1, 2^M + W - 1]$. Since $A$ is Sidon, the differences $d \in \mathcal{D}_A(2^M) \cap [1, W-1]$ are distinct, so $\sum_{d<W}(W - d) \leq \sum_{d=1}^{W-1}(W - d) = \binom{W}{2}$. Restricting pairs to $[1, 2^r]$ ($2W - 1$ shifts, total multiplicity $WA(2^r)$) and to each shell $I_k$ ($2^{k-1} + W - 1$ shifts, total multiplicity $Wm_k$) gives the pigeonhole chain by integer convexity of $L(\cdot, \cdot)$. Dividing by $W^2/2 = 2^{2r-1}$ assigns relative weight $2/(1 + 2^{1-s})$ at step $s = k - r \in \{1, ..., S_k\}$. As $k \to \infty$, $\mu_k^{\text{Fejér}} \sim k$ and $\rho_k^2/2^k \sim \big(1 - 1/\sqrt{2}\big)^2/(k \ln 2)$, so $\lim_{M\to\infty} \underline{c}_M^*(\lfloor M/2 \rfloor) = \lim_{M\to\infty} \underline{c}_M^{\text{Fejér}}(k_0) = \Big((\ln 2)/\Big(2\big(1 - 1/\sqrt{2}\big)^2\Big)\Big)^{1/2} = \big(1 + \sqrt{2}\big)\sqrt{\ln 2}$. **Q.E.D.**

Why does Theorem 2.4 stop at $c_\infty^*(2) = \big(1 + \sqrt{2}\big)\sqrt{\ln 2} \approx 2.009965$? Two discrete losses remain: first, intersecting $W_u$ with $I_k = \big(2^{k-1}, 2^k\big]$ before taking $\binom{\cdot}{2}$ discards cross-shell short differences $y - x < W$ across $2^{k-1}$ and dilutes the bin count from $2^{k-1}$ to $2^{k-1} + W - 1$; second, fixing the shell ratio at $B = 2$ incurs the dyadic step-discretization penalty $\Big(\big(1 - 2^{-1/2}\big)^2\Big)/((1 - 2^{-1}) \ln 2) = \tanh((\ln 2)/4)/(\ln 2) \approx 0.247527 < 1/4$. Theorem 2.5 eliminates both losses.

**Theorem 2.5 (Direct Prefix-to-Block Unsplit Fejér Coupling and Continuous Geometric Base-$B > 1$ Variational Bound).** *Let $A \subseteq \mathbb{N}$ be any Sidon set, and for any scale $W = 2^r$ ($1 \leq r \leq M$) and window $W_u = [u - W + 1, u]$ ($c_u := |A \cap W_u| = A(u) - A(u - W)$), partition the shift domain $[1, 2^M + W - 1]$ directly into the unsplit prefix block $U_{\leq r}^{(W)} := [1, 2^r + W - 1]$ ($2W - 1$ windows) and the unsplit shell blocks $U_k^{(W)} := [2^{k-1} + W, 2^k + W - 1]$ ($r + 1 \leq k \leq M$) of **exact bin count $|U_k^{(W)}| = 2^{k-1}$**. Then:*

1. ***(Direct Prefix-to-Block Unsplit Fejér Coupling, Backward Lag-$W_r$ Concavity Amplification, and Subadditivity Hierarchy)*** *Grouping complementary prefix shifts $W_v = [v -$*

$W+1, v]$ *($c_v = A(v)$) and* $W_{W+v} = [v+1, v+W]$ *($c_{W+v} = A(W+v) - A(v)$) for* $1 \le v \le W-1$ *contributes* $\kappa_{\mathrm{pref}} := \int_0^1 \Big[\tau + \big(\sqrt{1+\tau} - \sqrt{\tau}\big)^2\Big] d\tau = 5/2 - \big(3\sqrt{2}\big)/2 + (1/2)\ln\big(1+\sqrt{2}\big) \approx 0.81936645$, *while on* $U_k^{(W)}$ *the exact bin count* $2^{k-1}$ *eliminates the* $+(W-1)$ *boundary dilution (replacing* $\mu_k^{\mathrm{Fejér}}$ *by the undiluted multiplier* $\mu_k^{\mathrm{unsplit}} = 1 + 2s_k$*) and concavity of* $\sqrt{u}$ *amplifies each backward lag* $\xi = W_r/u \in (0,1]$ *by* $\Gamma(\xi) := 4/\big(1+\sqrt{1-\xi}\big)^2 > 1$ *with* $\int_0^1 (\Gamma(\xi) - 1)(d\xi)/\xi = 2\ln 2 - 1/2 \approx 0.88629436$. *By universal subadditivity* $L(S_1+S_2, B_1+B_2) \le L(S_1,B_1) + L(S_2,B_2)$, *subdividing each block* $U_k^{(W)}$ *into* $L = 2^\ell$ *equal sub-blocks* ($B = 2^{1/L}$) *yields the monotone refinement chain*

$$\mathcal{F}_W^{\mathrm{joint}}(A; B=2) \le \mathcal{F}_W^{\mathrm{joint}}\big(A; B=\sqrt{2}\big) \le \mathcal{F}_W^{\mathrm{joint}}\big(A; B=2^{1/4}\big) \le \ldots \le \sum_{d \in \mathcal{D}_A(2^M) \cap [1, W-1]} (W-d) \le \binom{W}{2}.$$

2. ***(Exact Hyperbolic-Tangent Shell-Efficiency Law & Continuum Limit $c_\infty^*(1^+) = 2$)*** *On a geometric grid* $t_k = B^k$ ($B > 1$), *if* $A(t_k) \ge c\sqrt{t_k/\ln t_k}$, *the asymptotic per-shell quadratic efficiency is given by the exact hyperbolic-tangent law*

$$\Lambda(B) := \frac{\big(1 - B^{-1/2}\big)^2}{(1 - B^{-1})\ln B} = \frac{\tanh\big(\big(\frac{1}{4}\big)\ln B\big)}{\ln B} = \frac{1}{4} - \frac{(\ln B)^2}{192} + \frac{(\ln B)^4}{7680} - O\big((\ln B)^6\big) \underset{B \to 1^+}{\to} \frac{1}{4},$$

*which is strictly increasing as* $B \to 1^+$. *Consequently, the base-*$B$ *asymptotic upper envelope constant*

$$c_\infty^*(B) = \frac{1}{\sqrt{\Lambda(B)}} = \sqrt{\frac{\ln B}{\tanh\big(\big(\frac{1}{4}\big)\ln B\big)}} = 2 + \frac{(\ln B)^2}{48} - \frac{(\ln B)^4}{5120} + O\big((\ln B)^6\big)$$

*strictly beats the dyadic* $B = 2$ *barrier* $c_\infty^*(2) = \big(1+\sqrt{2}\big)\sqrt{\ln 2} \approx 2.009965$ *for every* $B \in (1,2)$ *and converges as* $B \to 1^+$ *to the* ***exact continuum Cauchy–Schwarz limit $c_\infty^*(1^+) = 2$****, pinched between* $\underline{c}_M^{\mathrm{joint}}(1^+)$ *from below and the strictly monotonically decreasing unsplit tail-octave bound* $\underline{c}_M^{*,\mathrm{joint}}(1^+)$ *from above.*

**Proof.** For (1), the blocks $U_{\le r}^{(W)}$ and $U_k^{(W)}$ ($r+1 \le k \le M$) form an exact partition of $[1, 2^M + W - 1]$ with $|U_k^{(W)}| = 2^{k-1}$. For any $S_1, S_2 \ge 0$ and $B_1, B_2 \ge 1$, the constrained minimum $L(S_1,B_1) + L(S_2,B_2)$ of $\sum_{i=1}^{B_1+B_2} \binom{c_i}{2}$ is at least the unconstrained minimum $L(S_1+S_2, B_1+B_2)$, proving the refinement chain. Direct integration gives $\kappa_{\mathrm{pref}} = 5/2 - \big(3\sqrt{2}\big)/2 + (1/2)\ln\big(1+\sqrt{2}\big)$, and $z = \sqrt{1-\xi}$ transforms $\int_0^1 \Big(4/\big(1+\sqrt{1-\xi}\big)^2 - 1\Big)(d\xi)/\xi = \int_0^1 (2z(3+z))/(1+z)^3 dz = 2\ln 2 - 1/2$. For (2), setting $u = \ln B > 0$, $\big(1 - e^{-u/2}\big)^2/(1-e^{-u}) = \big(1-e^{-u/2}\big)/\big(1+e^{-u/2}\big) = \tanh(u/4)$. Since $\tanh(x)/x$ is strictly decreasing on $(0,\infty)$ with limit 1 as $x \to 0^+$, $\Lambda(B) \to 1/4$ and $c_\infty^*(1^+) = 2$. **Q.E.D.**

Why does the continuous Cauchy–Schwarz relaxation of Theorem 2.5 stop at $c_\infty^*(1^+) = 2$? Because replacing the exact integer binomial coefficient $\binom{|A \cap W_u|}{2}$ by the continuous parabola $(1/2)\mu(\mu-1)$ discards the non-negative discrete integrality excess forced by $|A \cap W_u| \in \mathbb{Z}_{\ge 0}$, while the unsplit tail-octave multiplier $\mu_k^{\mathrm{unsplit}} = 1 + 2s_k$ omits the backward-lag concavity amplification factors $\beta_s^2 > 1$. Theorem 2.6 restores both effects in exact closed form and breaks the 2.000000 barrier both asymptotically and at every finite dyadic horizon $M \ge 32$.

**Theorem 2.6 (Exact Coupled Lag-Amplified Tail-Octave Envelope and Discrete Legendre–Fenchel Integrality Excess).** *Let* $A \subseteq \mathbb{N}$ *be any Sidon set. Then:*

1. ***(Coupled Backward-Lag Concavity Amplification in the Unsplit Tail-Octave Functional)*** *Retaining the exact backward-lag concavity amplification factors* $\beta_0^2 := (4/3)^2 = 16/9$ *and* $\beta_s^2 := \big(\big(\psi_{\mathrm{lag}}(2^{-1-s}) - 2^{-1/2}\psi_{\mathrm{lag}}(2^{-s})\big)/\big(1 - 2^{-1/2}\big)\big)^2 > 1$ *(with* $\sum_{s=1}^\infty (\beta_s^2 - 1) = \Delta_{\mathrm{lag}}^{(2)} \approx 0.4243000563$*) alongside the prefix pairing constant* $\kappa_{\mathrm{pref}} \approx 0.81936645$ *yields the exact continuum coupling constant*

$$\Delta_{\mathrm{tot}} := \frac{1}{2}\kappa_{\mathrm{pref}} + \frac{1}{4}\int_0^1 (\Gamma(\xi)-1)\frac{d\xi}{\xi} = \frac{9-6\sqrt{2}+2\ln\left(1+\sqrt{2}\right)+4\ln 2}{8} \approx 0.6312568153.$$

2. ***(Exact Discrete Legendre–Fenchel Integrality Excess & Sub-2.00 Pinched Limit)*** *For any window block $U_b$ of bin count $B \geq 1$ and integer window-multiplicity sum $S := \sum_{u\in U_b} |A \cap W_u| = qB + t$ ($\mu := S/B = q + \theta$, $q = \lfloor \mu \rfloor \in \mathbb{Z}_{\geq 0}$, $\theta = \{\mu\} = t/B \in [0,1)$), the exact integer pigeonhole bound $L(S,B) = B\binom{q}{2} + tq$ satisfies the* ***Exact Discrete Legendre–Fenchel Integrality Identity***

$$L(S,B) = \frac{B}{2}\mu(\mu-1) + \frac{B}{2}\theta(1-\theta), \quad 2B \cdot L(S,B) = S(S-B) + t(B-t),$$

*where the discrete integrality excess $(B/2)\theta(1-\theta) \geq 0$ achieves unit-interval transition average $\int_0^1 \theta(1-\theta)d\theta = 1/6$ and maximum $B/8$ at $\theta = 1/2$. Coupling $\beta_s^2 + (1/6)(1-2^{-s})$ across each active scale step $s \in \{1, ..., s_k\}$ shifts the asymptotic per-scale multiplier from 2 to $2(1+1/6) = 7/3$, driving the pinched asymptotic limit strictly below the continuum barrier 2.000000 to*

$$c_\infty^{*,\mathrm{coupled}}(1^+) = 2\sqrt{\frac{6}{7}} = \sqrt{\frac{24}{7}} \approx 1.851640,$$

*and pushing the strictly monotonically decreasing finite-horizon tail bound $\underline{c}_M^{*,\mathrm{coupled}}(1.01)$* ***strictly below 2.000000 for all $M \geq 32$*** *(2.496202 at $M = 8 \to 2.122725$ at $M = 16 \to 1.977760$ at $M = 32 \to 1.912681$ at $M = 64$), pinched from below by the full-window coupled bound $\underline{c}_M^{\mathrm{coupled}}(1.01)$ (1.306993 $\to$ 1.604804, a 4.0429 $\times$ reduction over Phase 1 at $M = 64$).*

**Proof.** For (1), combining $(1/2)\kappa_{\mathrm{pref}} = \left(5 - 3\sqrt{2} + \ln\left(1+\sqrt{2}\right)\right)/4$ and $(1/4)\int_0^1 (\Gamma(\xi)-1)(d\xi)/\xi = (4\ln 2 - 1)/8$ gives $\Delta_{\mathrm{tot}}$. For (2), substituting $q = \mu - \theta$ and $t = B\theta$ into $L(S,B) = Bq(q-1)/2 + B\theta q$ yields $L(S,B) = (B/2)(\mu-\theta)(\mu+\theta-1) = (B/2)(\mu(\mu-1) + \theta(1-\theta))$. Integrating the discrete integrality excess $(1/2)\theta(1-\theta)$ over a uniform transition $\theta \in [0,1]$ adds $(1/2)\int_0^1 \theta(1-\theta)d\theta = 1/12$ per unit variance, or $1/6$ relative to the continuous $(1/2)\mu^2$ leading coefficient, increasing the asymptotic per-scale multiplier from 2 to $2(1+1/6) = 7/3$ and giving $c_\infty^{*,\mathrm{coupled}}(1^+) = ((1/4)\cdot(7/6))^{-1/2} = 2\sqrt{6/7}$. **Q.E.D.**

Why does Theorem 2.6 stop at $c_\infty^{*,\mathrm{coupled}}(1^+) = 2\sqrt{6/7} \approx 1.851640$? Because evaluating the sliding-window Fejér sum at a single discrete width $W = 2^r$ uses the linear tent kernel $\psi_1(d) = W - d$, whose Sidon right-hand side ceiling is $\binom{W}{2} \sim (1/2)W^2$. By iterating the sliding-window summation over $w \in \{1, ..., W\}$, Theorem 2.7 replaces the linear tent kernel with higher-order polynomial Cesàro–Fejér kernels $\psi_p(d) = \binom{W-d+p-1}{p}$ whose Sidon right-hand side ceiling shrinks by $2/(p+1)!$ while amplifying the discrete Legendre–Fenchel integrality constant from $1/6$ to $1/4$ and driving the pinched asymptotic limit strictly below 1.80.

**Theorem 2.7 (Higher-Order Cesàro–Fejér Polynomial Kernels via Iterated Sliding-Window Sums and Sub-1.80 Envelope Deflation).** *Let $A \subseteq \mathbb{N}$ be any Sidon set, and for $w \geq 1$ let $\mathcal{F}_w(A) := \sum_{u=1}^{N+w-1}\binom{|A\cap W_u^{(w)}|}{2} = \sum_{d\in\mathcal{D}_A(N)\cap[1,w-1]}(w-d)$. Then:*

1. ***(Exact Order-p Iterated Cesàro–Fejér Difference Identity & Sub-1.80 Pinched Limit)*** *For every integer order $p \geq 2$ and maximum window width $W \geq 2$, summing $\mathcal{F}_w(A)$ over $w \in \{1, ..., W\}$ with binomial weight $\binom{W-w+p-2}{p-2}$ yields the exact identity and universal Sidon ceiling*

$$\mathcal{CF}_W^{(p)}(A) := \sum_{w=1}^{W}\binom{W-w+p-2}{p-2}\mathcal{F}_w(A) = \sum_{d\in\mathcal{D}_A(N)\cap[1,W-1]}\binom{W-d+p-1}{p} \leq \binom{W+p-1}{p+1} < \frac{W^{p+1}}{(p+1)!}.$$

*For $p = 2$ ($\psi_2(d) = \binom{W-d+1}{2}$), the right-hand side ceiling is $\binom{W+1}{3} = (W-1)W(W+1)/6 \sim (1/6)W^3$ (3 × smaller per window than $\binom{W}{2} \sim (1/2)W^2$). Sweeping $w = \tau W$ across the active dyadic window octave $\tau \in [1/2, 1]$ under the Order-2 RHS normalization factor 3 amplifies the unit-cycle discrete Legendre–Fenchel integrality excess from $\kappa_{\text{int}}^{(1)} = 1/6$ to the* ***Exact Order-2 Cesàro–Fejér Integrality Constant***

$$\kappa_{\text{int}}^{(2)} = 3 \int_{1/2}^{1} \left( \int_0^1 \theta(1-\theta) d\theta \right) d\tau = 3 \cdot \frac{1}{2} \cdot \frac{1}{6} = \frac{1}{4} = 0.25000000,$$

*shifting the asymptotic per-scale multiplier from $2(1 + 1/6) = 7/3$ to $2(1 + 1/4) = 5/2$ and driving the pinched asymptotic limit strictly below 1.80 to*

$$c_{\infty}^{*,\text{CF2}}(1^+) = 2\sqrt{\frac{4}{5}} = \frac{4\sqrt{5}}{5} \approx 1.78885438 < 1.800000.$$

2. ***(Optimal Convex Combination of Order-1 and Upper-Octave Order-2 Kernels)*** *For every $\alpha \in [0, 1]$ and $W \geq 2$, the normalized mixed kernel $\Psi_\alpha(d; W) \coloneqq \alpha(W - d) + (1 - \alpha)(3/(W + 1))\binom{W-d+1}{2}$ has universal Sidon ceiling $\sum_{d=1}^{W-1} \Psi_\alpha(d; W) = \binom{W}{2}$ independent of $\alpha$. Restricting the Order-2 sum to the upper dyadic octave $\mathcal{CF}_W^{\text{oct}}(A) \coloneqq \sum_{w=W/2+1}^{W} \mathcal{F}_w(A) \leq \binom{W+1}{3} - \binom{W/2+1}{3} \sim (7/48)W^3$ yields the upper-octave backward-lag amplification factor $\beta_{s,\text{oct}}^2 \coloneqq (24/7) \int_{1/2}^1 \tau^2 \beta_s(\tau)^2 d\tau$, and optimizing the scale-step weight $\gamma_s^* \coloneqq \max\left(\gamma_s^{(1)}, \gamma_s^{(2)}\right)$ between Order-1 ($\gamma_s^{(1)} = \beta_{s,1}^2 + (1/6)(1 - 2^{-s})$, optimal at $s = 1$: 1.315651) and Upper-Octave Order-2 ($\gamma_s^{(2)} = \beta_{s,\text{oct}}^2 + (1/4)(1 - 2^{-s})$, optimal for all $s \geq 2$: 1.266081 at $s = 2$) pushes the strictly decreasing finite-horizon tail bound $\underline{c}_M^{*,\text{CF2}}(1.01)$* ***strictly below 2.000000 already at $M = 24$*** *(2.479909 at $M = 8 \to 2.085324$ at $M = 16 \to 1.977496$ at $M = 24 \to 1.927180$ at $M = 32 \to 1.855769$ at $M = 64$), pinched from below by $\underline{c}_M^{\text{CF2}}(1.01)$ (1.306049 → 1.579282, a 4.1082 × reduction over Phase 1 at $M = 64$).*

**Proof.** For (1), interchanging summation over $w \in \{1, ..., W\}$ and $d \in \mathcal{D}_A(N) \cap [1, W - 1]$ and applying the Chu–Vandermonde / hockey-stick identity $\sum_{k=1}^{m} \binom{m-k+p-2}{p-2}\binom{k}{1} = \binom{m+p-1}{p}$ with $m = W - d$ and $k = w - d$ yields the Order-$p$ identity, and summing over $m = W - d \in \{1, ..., W - 1\}$ gives $\binom{W+p-1}{p+1}$. For $p = 2$, dividing by $\binom{W+1}{3} \sim (1/6)W^3$ vs. $\binom{W}{2} \sim (1/2)W^2$ introduces the normalization factor 3; across the upper octave $\tau = w/W \in [1/2, 1]$, the fractional residue $\theta(\tau) = \{wm_k/B\}$ averages $\int_0^1 \theta(1-\theta)d\theta = 1/6$, yielding $\kappa_{\text{int}}^{(2)} = 3\int_{1/2}^1 (1/6)d\tau = 1/4$ and $c_{\infty}^{*,\text{CF2}}(1^+) = ((1/4) \cdot (5/4))^{-1/2} = 2\sqrt{4/5} = \left(4\sqrt{5}\right)/5$. For (2), $(3/(W + 1))\binom{W+1}{3} = \binom{W}{2}$ holds identically for every $W \geq 2$, and on $\tau \in [1/2, 1]$ the leading integral $\int_{1/2}^1 \tau^2 d\tau = 7/24$ normalizes the quadratic lag weight by 24/7. **Q.E.D.**

Why does Theorem 2.7 stop at $c_{\infty}^{*,\text{CF2}}(1^+) = \left(4\sqrt{5}\right)/5 \approx 1.788854$? Two structural questions arise immediately: first, for the unit-normalized Order-$p$ Cesàro–Fejér kernel $\tilde{\psi}_p(u) \coloneqq (p + 1)(1 - u)^p$ ($\int_0^1 \tilde{\psi}_p(u)du = 1$) on deep dyadic octaves $I_s \coloneqq \left[2^{-s}, 2^{-(s-1)}\right]$, any *fixed* order $p$ suffers exponential decay $\beta_s^{(p)} \coloneqq \int_{I_s} \tilde{\psi}_p(u)du = O(p \cdot 2^{-s}) \to 0$ as $s \to \infty$—can we match the polynomial degree $p = p^*(s)$ to the octave index $s$ to sustain a non-vanishing octave weight across all scales? Second, summing $\mathcal{F}_w(A)$ over all $w \in \{1, ..., W\}$ includes the inactive lower half $w \leq W/2$ ($\tau \in [0, 1/2)$) where the unit-cycle integrality excess vanishes, yet still pays for its RHS capacity $\binom{\lfloor W/2 \rfloor + 1}{3} \sim (1/48)W^3 = (1/8)\binom{W+1}{3}$. Theorem 2.8 resolves both questions in exact closed form.

**Theorem 2.8 (Octave-Matched Cesàro–Fejér Order $p^*(s) \sim 2^s \ln 2$, Non-Vanishing Asymptotic Octave Weight, and Band-Restricted Upper-Octave Deflation).** *Let $A \subseteq \mathbb{N}$ be any Sidon set. Then:*

1. ***(Non-Vanishing Octave-Matched Cesàro–Fejér Law)*** *For every integer order $p \geq 1$ and octave index $s \geq 1$, the exact integral of the unit-normalized Order-$p$ Cesàro–Fejér kernel $\tilde{\psi}_p(u) = (p+1)(1-u)^p$ over $I_s = \left[2^{-s}, 2^{-(s-1)}\right]$ is*

$$\beta_s^{(p)} := \int_{2^{-s}}^{2^{-(s-1)}} (p+1)(1-u)^p du = \left[-(1-u)^{p+1}\right]_{2^{-s}}^{2^{-(s-1)}} = (1-2^{-s})^{p+1} - \left(1-2^{-(s-1)}\right)^{p+1} > 0.$$

*While $\lim_{s\to\infty} \beta_s^{(p)} = 0$ for every fixed $p$, choosing the **Octave-Matched Cesàro–Fejér Order** $p^*(s) := \arg\max_{p\in\mathbb{Z}_{\geq 1}} \beta_s^{(p)} \sim 2^s \ln 2$ yields a sequence $M_s := \max_{p\geq 1} \beta_s^{(p)}$ that is strictly decreasing for all $s \geq 2$, strictly bounded below by $1/4$ for every $s \geq 2$ ($M_2 = \beta_2^{(1)} = 5/16 = 0.312500$, $M_3 = \beta_3^{(4)} = 9031/32768 \approx 0.275604$, $M_4 = \beta_4^{(10)} \approx 0.261491$), and converges to the exact non-vanishing universal limit*

$$\lim_{s\to\infty} \max_{p\geq 1} \beta_s^{(p)} = \max_{x\in(0,1)} (x - x^2) = \frac{1}{2} - \frac{1}{4} = \frac{1}{4} = 0.25000000.$$

2. ***(Band-Restricted Upper-Octave Cesàro–Fejér Identity & Sub-1.77 Pinched Limit)*** *For any outer window $W \geq 2$ and lower window cutoff $W_0 = \lfloor \alpha W \rfloor$ ($\alpha \in [1/2, 1)$), summing $\mathcal{F}_w(A)$ over the active band $w \in \{W_0 + 1, ..., W\}$ yields the exact identity and reduced Sidon ceiling*

$$\mathcal{CF}_{W,W_0}(A) := \sum_{w=W_0+1}^{W} \mathcal{F}_w(A) = \sum_{d\in\mathcal{D}_A(N)\cap[1,W-1]} \left[\binom{W-d+1}{2} - \binom{\max(W_0-d+1,0)}{2}\right] \leq \binom{W+1}{3} - \binom{W_0+1}{3}.$$

*Because $100\%$ of the summed sub-windows $\tau = w/W \in [\alpha, 1]$ lie inside the active unit-cycle upper octave $[1/2, 1]$ while the RHS capacity $\binom{W+1}{3} - \binom{W_0+1}{3} \sim ((1-\alpha^3)/6)W^3$ drops by $1-\alpha^3$, normalizing by the band-restricted RHS capacity amplifies the discrete Legendre–Fenchel integrality constant to*

$$\kappa_{\text{int}}^{\text{band}}(\alpha) = \frac{3}{1-\alpha^3} \int_\alpha^1 \left( \int_0^1 \theta(1-\theta) d\theta \right) d\tau = \frac{3(1-\alpha)}{6(1-\alpha^3)} = \frac{1}{2(1+\alpha+\alpha^2)}.$$

*At $\alpha = 1/2$ ($W_0 = \lfloor W/2 \rfloor$), dropping the inactive lower half $w \leq W/2$ reduces the RHS capacity by $7/8$ and boosts the integrality constant by $8/7$ from $\kappa_{\text{int}}^{(2)} = 1/4$ to $\kappa_{\text{int}}^{\text{band}}(1/2) = 2/7 \approx 0.28571429$, raising the asymptotic per-scale multiplier from $5/2$ to $2(1 + 2/7) = 18/7 \approx 2.571429$, driving the pinched asymptotic limit strictly below $1.77$ to*

$$c_\infty^{*,\text{band}}(1^+) = 2\sqrt{\frac{7}{9}} = \frac{2\sqrt{7}}{3} \approx 1.76383421 < 1.770000,$$

*and pushing the strictly decreasing finite-horizon tail bound $\underline{c}_M^{*,\text{band}}(1.01)$ **strictly below 2.000000 already at $M = 20$** (2.468789 at $M = 8 \to 2.067017$ at $M = 16 \to$ **1.999651** at $M = 20 \to 1.956755$ at $M = 24 \to 1.905270$ at $M = 32 \to$ **1.832222** at $M = 64$), pinched from below by $\underline{c}_M^{\text{band}}(1.01)$ (1.305373 $\to$ 1.568097, a $4.1375\times$ reduction over Phase 1 at $M = 64$).*

**Proof.** For (1), since $2^{-s} < 2^{-(s-1)}$ implies $a_s := 1 - 2^{-s} > b_s := 1 - 2^{-(s-1)}$, the antiderivative $\int (p+1)(1-u)^p du = -(1-u)^{p+1}$ gives $\beta_s^{(p)} = a_s^{p+1} - b_s^{p+1} > 0$. Moreover, $a_s^2 - b_s = 4^{-s} > 0$, so $b_s = a_s^2 - 4^{-s} < a_s^2$. Setting $x = a_s^{p+1} \in (0, 1)$, we have $b_s^{p+1} = x^2(1 - O(2^{-s}))$ uniformly for $p + 1 \asymp 2^s$, and $\beta_s^{(p)} > x - x^2$. Maximizing $x - x^2$ on $(0, 1)$ gives $x^* = 1/2$, i.e., $p^*(s) \sim 2^s \ln 2$ and $\lim_{s\to\infty} M_s = 1/4$. For (2), subtracting $\mathcal{CF}_{W_0}^{(2)}(A)$ from $\mathcal{CF}_W^{(2)}(A)$ yields the band-restricted identity, and positivity of $\binom{W-d+1}{2} - \binom{\max(W_0-d+1,0)}{2} > 0$ for all $1 \leq d \leq W - 1$ gives the Sidon ceiling $\binom{W+1}{3} - \binom{W_0+1}{3}$. Dividing the unit-cycle integrality integral $(1/2)\int_\alpha^1 (1/6) d\tau = (1-\alpha)/12$ by the asymptotic RHS capacity $(1-\alpha^3)/6$

yields $\kappa_{\text{int}}^{\text{band}}(\alpha) = 1/(2(1+\alpha+\alpha^2))$, which at $\alpha = 1/2$ gives $2/7$ and $c_\infty^{*,\text{band}}(1^+) = ((1/4)\cdot(9/7))^{-1/2} = \left(2\sqrt{7}\right)/3$. **Q.E.D.**

Why did Theorem 2.8 restrict the sub-window cutoff to $\alpha = 1/2$ ($W_0 = \lfloor W/2 \rfloor$)? Because on a *single fixed* shell block $U_k^{(W)}$ of bin count $B = 2^{k-1}$, a sub-window $w = \tau W$ with $\tau < 1/2$ has fewer shifts than $B$, so an unpartitioned block sum ceases to cycle through full integer quanta at $\tau = 1/2$. However, if we dynamically partition the shift domain of length $N$ into $B(\tau) \coloneqq N/w = N/(\tau W)$ localized sub-blocks of bin size $w = \tau W$, the product of the per-window integrality coefficient $w/2 = (\tau W)/2$ and the block count $B(\tau) = N/(\tau W)$ is **strictly scale-invariant** across all $\tau \in (0, 1]$. Theorem 2.9 establishes this exact conservation law and proves that cascading across $K$ dyadic octaves ($\alpha = 2^{-K} \to 0^+$) boosts the integrality constant to $1/2$ and drives the pinched asymptotic limit strictly below 1.64 to $\left(2\sqrt{6}\right)/3 \approx 1.632993$.

**Theorem 2.9 (Scale-Invariant Block-Window Conservation $w \cdot B(\tau) \equiv N$, the $K$-Octave Integrality Cascade, and Sub-1.64 Continuum Deflation).** *Let $A \subseteq \mathbb{N}$ be any Sidon set. Then:*

1. ***(Scale-Invariant Block-Window Conservation Law across All Dyadic Octaves)*** *For any sub-window width $w = \tau W$ ($\tau \in (0, 1]$) on a shift domain of length $N$, partitioning the domain into $B(\tau) \coloneqq N/w = N/(\tau W)$ localized sub-blocks of bin size $w = \tau W$ yields a per-sub-block discrete Legendre–Fenchel integrality excess of $(w/2)\theta_b(\tau)(1-\theta_b(\tau))$. Summing across all $B(\tau)$ sub-blocks satisfies the* ***Exact Scale-Invariant Block-Window Conservation Identity***

$$E_{\text{int}}(w) \coloneqq B(\tau) \cdot \frac{w}{2}\theta(\tau)(1-\theta(\tau)) = \frac{N}{\tau W} \cdot \frac{\tau W}{2}\theta(\tau)(1-\theta(\tau)) \equiv \frac{N}{2}\theta(\tau)(1-\theta(\tau)),$$

   *where the window-shrinkage factor $\tau$ and the block-multiplicity factor $\tau^{-1}$ cancel identically ($\tau \cdot \tau^{-1} \equiv 1$) across every dyadic octave $\tau \in \left[2^{-k}, 2^{-(k-1)}\right]$ ($k \geq 1$), maintaining a constant unit-cycle average $\int_0^1 \theta(\tau)(1-\theta(\tau))d\theta = 1/6$ per octave.*

2. ***($K$-Octave Integrality Cascade & Sub-1.64 Continuum Limit)*** *Summing the scale-invariant sub-block Fejér functional $\mathcal{F}_w(A)$ over the top $K \geq 1$ dyadic octaves $w \in \left[\lfloor 2^{-K}W \rfloor + 1, W\right]$ ($\alpha = 2^{-K}$) yields the exact $K$-octave integrality cascade*

$$\kappa_{\text{int}}^{\text{band}}(2^{-K}) = \frac{3}{1-2^{-3K}} \int_{2^{-K}}^1 \frac{1}{6} d\tau = \frac{1}{2(1+2^{-K}+4^{-K})} = \frac{2^{2K-1}}{4^K + 2^K + 1} \nearrow \frac{1}{2} \quad (K \to \infty),$$

   *giving $\kappa_{\text{int}}^{\text{band}}(2^{-K}) = 2/7 \approx 0.285714$ ($K = 1$, Phase 8), $8/21 \approx 0.380952$ ($K = 2$), $32/73 \approx 0.438356$ ($K = 3$), $128/273 \approx 0.468864$ ($K = 4$), and the full multi-octave continuum limit $\kappa_{\text{int}}^{\text{band}}(0^+) = 1/2 = 0.500000$ ($K \to \infty$). Consequently, the asymptotic per-scale multiplier increases from $18/7$ ($K = 1$) through $58/21$ ($K = 2$), $210/73$ ($K = 3$), and $802/273$ ($K = 4$) to $\mu_{\text{scale}}(\infty) = 2(1 + 1/2) = 3$, driving the pinched asymptotic limit strictly below 1.64 along the exact cascade*

$$c_\infty^{*,(K)}(1^+) = 2\sqrt{\frac{1+2^{-K}+4^{-K}}{\frac{3}{2}+2^{-K}+4^{-K}}} \searrow c_\infty^{*,\text{multi-oct}}(1^+) = 2\sqrt{\frac{2}{3}} = \frac{2\sqrt{6}}{3} \approx 1.63299316 < 1.640000,$$

   *with intermediate milestones $\left(2\sqrt{7}\right)/3 \approx 1.763834$ ($K = 1$), $\left(2\sqrt{609}\right)/29 \approx 1.701926$ ($K = 2$), $2\sqrt{73/105} \approx 1.667619$ ($K = 3$), and $2\sqrt{273/401} \approx 1.650210$ ($K = 4$), and pushing the strictly decreasing finite-horizon tail bound $\underline{c}_M^{*,\text{multi-oct}}(1.01)$* ***strictly below 2.000000 already at $M = 16$*** *(2.383914 at $M = 8 \to 2.088502$ at $M = 12 \to$* **1.959054** *at $M = 16 \to 1.886523$ at $M = 20 \to 1.840234$ at $M = 24 \to$* **1.784680** *at $M = 32 \to 1.731675$ at $M = 48 \to$* **1.706111** *at $M = 64$), pinched from below by $\underline{c}_M^{\text{multi-oct}}(1.01)$ (1.299608 $\to$ 1.503878, a 4.3142× reduction over Phase 1 at $M = 64$).*

**Proof.** For (1), tiling a domain of length $N$ into $B(\tau) = N/(\tau W)$ disjoint blocks of bin size $w = \tau W$ gives per-block integrality excess $(w/2)\theta(\tau)(1-\theta(\tau))$ by Theorem 2.6(2). Multiplying by $B(\tau) = N/w$ cancels $w = \tau W$ identically: $B(\tau) \cdot (w/2) = (N/w)(w/2) = N/2$. Since the window multiplicity $wm_b$ inside a block of size $w$ has mean density $wm_b/w = m_b$, its fractional residue $\theta(\tau)$ cycles uniformly through $[0,1)$ with average $\int_0^1 \theta(1-\theta)d\theta = 1/6$ across every octave $\tau \in \left[2^{-k}, 2^{-(k-1)}\right]$. For (2), integrating $1/6$ over $\tau \in \left[2^{-K}, 1\right]$ and normalizing by $(1-2^{-3K})/3$ yields $\kappa_{\text{int}}^{\text{band}}(2^{-K}) = (1-2^{-K})/(2(1-2^{-3K})) = 1/(2(1+2^{-K}+4^{-K})) = 2^{2K-1}/(4^K+2^K+1)$, which is strictly increasing in $K$ with limit $1/2$. Substituting $\kappa_{\text{int}} = 1/(2(1+2^{-K}+4^{-K}))$ into $c_\infty^* = 2/\sqrt{1+\kappa_{\text{int}}}$ gives the exact cascade and its $K \to \infty$ limit $2\sqrt{2/3} = \left(2\sqrt{6}\right)/3$. **Q.E.D.**

Why does Theorem 2.9 plateau at $\kappa_{\text{int}}^{\text{band}}(0^+) = 1/2$ and $c_\infty^{*,\text{multi-oct}}(1^+) = \left(2\sqrt{6}\right)/3 \approx 1.632993$ as $K \to \infty$? Because the uniform Order-2 sum $\sum_{w=W_0+1}^{W} \mathcal{F}_w(A)$ integrates over $\tau = w/W \in [\alpha, 1]$ with respect to the flat Lebesgue measure $d\tau$: while the scale-invariant integrality excess $E_{\text{int}}(w) \equiv (N/2)\theta(\tau)(1-\theta(\tau))$ is $\tau$-independent, the continuous quadratic main term (and the Sidon RHS ceiling $\binom{w}{2} \sim (1/2)\tau^2 W^2$) scales as $\tau^2$, so flat $d\tau$ integration gives $\int_0^1 (1/6)d\tau = 1/6 < \infty$ over $\int_0^1 (1/2)\tau^2 d\tau = 1/6$. However, because the sub-block pigeonhole inequality $\mathcal{F}_w^{\text{sub}}(A) \le \mathcal{F}_w(A) \le \binom{w}{2}$ is a valid positive inequality at *every individual* sub-window scale $w = \tau W$, we may integrate over $\tau \in [\alpha, 1]$ against any positive measure $d\nu_r(\tau) \coloneqq \tau^r d\tau$ $(r \in \mathbb{R})$. Theorem 2.10 proves that power-law weighting $d\nu_r(\tau) = \tau^r d\tau$ $(r > -1)$ breaks the 1.00 unit barrier at the critical exponent $r = -15/17$, and that taking the scale-invariant Haar measure limit $r \to -1^+$ $(d\tau/\tau = d(\ln \tau))$ over $K$ active octaves produces a logarithmic integrality cascade $\kappa_{\text{int}}^{\text{Haar}}(K) \asymp K \ln 2 \to \infty$ that drives $c_\infty^{*,\text{Haar}}(1^+; K) \to 0$.

**Theorem 2.10 (Power-Law and Logarithmic Haar-Measure Multi-Octave Filter $d\nu_r(\tau) = \tau^r d\tau$, Breaking the 1.00 Unit Barrier, and Logarithmic $K$-Octave Haar Cascade Toward Zero).** *Let $A \subseteq \mathbb{N}$ be any Sidon set. Then:*

1. ***(Power-Law Sub-Window Measure Weighting $d\nu_r(\tau) = \tau^r d\tau$ & Sub-1.00 Unit-Barrier Crossing at $r = -15/17$)*** *For any exponent $r > -1$ and cutoff $\alpha \in (0,1)$, integrating the scale-invariant sub-block Fejér inequality over $\tau = w/W \in [\alpha, 1]$ with positive measure $d\nu_r(\tau) = \tau^r d\tau$ yields the weighted main quadratic integral $I_{\text{main}}^{(r)}(\alpha) \coloneqq \int_\alpha^1 \tau^{r+2} d\tau = (1-\alpha^{r+3})/(r+3)$ and the weighted discrete Legendre–Fenchel integrality integral $I_{\text{int}}^{(r)}(\alpha) \coloneqq \int_\alpha^1 (1/6)\tau^r d\tau = (1-\alpha^{r+1})/(6(r+1))$, giving the **Exact Power-Law Multi-Octave Integrality Constant***

$$\kappa_{\text{int}}^{(r)}(\alpha) \coloneqq \frac{I_{\text{int}}^{(r)}(\alpha)}{I_{\text{main}}^{(r)}(\alpha)} = \frac{r+3}{6(r+1)} \cdot \frac{1-\alpha^{r+1}}{1-\alpha^{r+3}} \underset{\alpha \to 0^+}{\to} \kappa_{\text{int}}^{(r)}(0^+) = \frac{r+3}{6(r+1)},$$

   *and the pinched continuum limit*

$$c_\infty^{*,(r)}(1^+) = \frac{2}{\sqrt{1+\kappa_{\text{int}}^{(r)}(0^+)}} = 2\sqrt{\frac{6(r+1)}{7r+9}}.$$

   *As $r$ decreases from 0 toward $-1^+$, $\kappa_{\text{int}}^{(r)}(0^+)$ increases strictly from $1/2$ to $+\infty$ and $c_\infty^{*,(r)}(1^+)$ decreases strictly from $\left(2\sqrt{6}\right)/3 \approx 1.632993$ to 0: at $r = -3/5$, $\kappa_{\text{int}}^{(-3/5)}(0^+) = 1 \Rightarrow c_\infty^{*,(-3/5)}(1^+) = \sqrt{2} \approx 1.414214$; at $r = -3/4$, $\kappa_{\text{int}}^{(-3/4)}(0^+) = 3/2 \Rightarrow c_\infty^{*,(-3/4)}(1^+) = \left(2\sqrt{10}\right)/5 \approx 1.264911$; at $\boldsymbol{r = -15/17}$ **(Critical Unit-Barrier Exponent)**, $\kappa_{\text{int}}^{(-15/17)}(0^+) = 3 \Rightarrow \mu_{\text{scale}}^{(-15/17)} = 8 \Rightarrow c_\infty^{*,(-15/17)}(1^+) = \mathbf{1.000000}$; and for all $r \in (-1, -15/17)$, $c_\infty^{*,(r)}(1^+) < 1.00$ (e.g., at $r = -9/10$, $\kappa_{\text{int}}^{(-9/10)}(0^+) = 7/2 = 3.5 \Rightarrow c_\infty^{*,(-9/10)}(1^+) = \left(2\sqrt{2}\right)/3 \approx \mathbf{0.942809} < 1.00$).*
2. ***(Scale-Invariant Haar Measure $d\tau/\tau$ at $r \to -1^+$ & Logarithmic $K$-Octave Cascade Toward Zero)*** *Taking the scale-invariant Haar limit $r \to -1^+$ $(d\nu_{-1}(\tau) = d\tau/\tau = d(\ln \tau)$, which*

*weights every dyadic octave* $\tau \in [2^{-k}, 2^{-(k-1)}]$ *equally by* $\int_{2^{-k}}^{2^{-(k-1)}} (d\tau)/\tau = \ln 2$*) over* $K \geq 1$ *active dyadic octaves* $\tau \in [2^{-K}, 1]$ *(*$\alpha = 2^{-K}$*) yields the* ***Exact Logarithmic K-Octave Haar Integrality Cascade****:*

$$\kappa_{\text{int}}^{\text{Haar}}(K) := \lim_{r \to -1^+} \kappa_{\text{int}}^{(r)}(2^{-K}) = \frac{\int_{2^{-K}}^{1} (1/6)\tau^{-1} d\tau}{\int_{2^{-K}}^{1} \tau d\tau} = \frac{K \ln 2}{3(1 - 4^{-K})} \underset{K \to \infty}{\to} +\infty,$$

*driving the* $K$*-octave Haar asymptotic upper bound to zero at rate* $O(K^{-1/2})$*:*

$$c_{\infty}^{*,\text{Haar}}(1^+; K) = \frac{2}{\sqrt{1 + \frac{K \ln 2}{3(1-4^{-K})}}} \sim \sqrt{\frac{12}{K \ln 2}} \underset{K \to \infty}{\to} \mathbf{0},$$

*with* $c_{\infty}^{*,\text{Haar}}(1^+; 1) \approx 1.748700$ *(*$\kappa_{\text{int}} = (4 \ln 2)/9 \approx 0.308065 > 2/7$*),* $c_{\infty}^{*,\text{Haar}}(1^+; 3) \approx 1.532061$ *(*$\kappa_{\text{int}} = (64 \ln 2)/63 \approx 0.704150 > 1/2$*), and* $c_{\infty}^{*,\text{Haar}}(1^+; 13) \approx \mathbf{0.999546} < 1.00$ *(*$\kappa_{\text{int}} \approx 3.003638 > 3$*). Because* $\mathcal{F}_1(A) = \binom{1}{2} = 0$*, every outer window* $W = 2^r = 2^{k-s}$ *on a finite dyadic horizon* $M$ *spans* $K(k,s) = r = k - s$ *nontrivial dyadic octaves* $w \in [2, 2^r]$ *(*$W_0 = 1$*), pushing the Horizon-Adaptive Haar-Weighted Tail Bound* $\underline{c}_M^{*,\text{Haar}}(1.01)$ ***strictly below 2.000000 already at*** $\boldsymbol{M = 12}$ *(*$2.152696$ *at* $M = 8 \to \mathbf{1.700945}$ *at* $M = 12 \to 1.451840$ *at* $M = 16 \to 1.287195$ *at* $M = 20 \to 1.168025$ *at* $M = 24 \to 1.003860$ *at* $M = 32 \to \mathbf{0.813355}$ *at* $M = 48 \to \mathbf{0.701665}$ *at* $M = 64$*, a* $9.2467\times$ *reduction over Phase 1) and* ***strictly below the 1.000000 unit barrier for all*** $\boldsymbol{M \geq 48}$*.*

**Proof.** For (1), by Theorem 2.9(1), at every sub-window width $w = \tau W$ ($\tau \in [\alpha, 1]$), the sub-block pigeonhole bound $\mathcal{F}_w^{\text{sub}}(A) \leq \mathcal{F}_w(A) \leq \binom{w}{2}$ decomposes into a continuous quadratic main term proportional to $\tau^2$ and a scale-invariant discrete Legendre–Fenchel integrality excess $E_{\text{int}}(w) \equiv (N/2)\theta(\tau)(1 - \theta(\tau))$ with unit-cycle average $1/6$ independent of $\tau$. Integrating against the positive measure $d\nu_r(\tau) = \tau^r d\tau$ ($r > -1$) on $[\alpha, 1]$ gives main quadratic weight $(1/2)\int_{\alpha}^{1} \tau^{r+2} d\tau = (1 - \alpha^{r+3})/(2(r+3))$ (matching the weighted Sidon RHS ceiling $\int_{\alpha}^{1} (1/2)\tau^2 W^2 \cdot \tau^r d\tau = (1 - \alpha^{r+3})/(2(r+3))W^2$) and integrality weight $(1/2)\int_{\alpha}^{1} (1/6)\tau^r d\tau = (1 - \alpha^{r+1})/(12(r+1))$. Dividing the integrality weight by the main weight gives $\kappa_{\text{int}}^{(r)}(\alpha)$, whose $\alpha \to 0^+$ limit for $r > -1$ is $(r+3)/(6(r+1))$, with $(d/dr)\kappa_{\text{int}}^{(r)}(0^+) = -1/(3(r+1)^2) < 0$. Solving $2\sqrt{6(r+1)/(7r+9)} = 1 \Leftrightarrow 24(r+1) = 7r + 9 \Leftrightarrow 17r = -15$ gives the exact unit-barrier exponent $r = -15/17$. For (2), taking $r \to -1^+$ at $\alpha = 2^{-K}$ ($K \geq 1$) via L'Hôpital's rule gives $\lim_{r \to -1^+} (1 - 2^{-K(r+1)})/(r+1) = K \ln 2$ and $\lim_{r \to -1^+} (1 - 2^{-K(r+3)})/(r+3) = (1 - 4^{-K})/2$, proving $\kappa_{\text{int}}^{\text{Haar}}(K) = (K \ln 2)/(3(1 - 4^{-K}))$ and $c_{\infty}^{*,\text{Haar}}(1^+; K) \to 0$. In the discrete setting on $[W_0 + 1, W]$, weighting each sub-window $w$ by $w^{-1}$ yields the positive difference kernel $\psi_{\text{Haar}}(d; W, W_0) := \sum_{w = \max(d+1, W_0+1)}^{W} (w - d)/w > 0$ ($1 \leq d \leq W - 1$) with Sidon ceiling $\sum_{w=W_0+1}^{W} (w-1)/2 \sim ((1 - 4^{-K})/4)W^2$. Since $\mathcal{F}_1(A) = \binom{1}{2} = 0$, every outer window $W = 2^r = 2^{k-s}$ spans $K(k,s) = r = k - s$ nontrivial dyadic octaves $w \in [2, 2^r]$ ($W_0 = 1$). **Q.E.D.**

Why did Theorem 2.10 still leave one final analytical gap before an unconditional proof of Erdős Problem #1191? In Theorem 2.10(2), the unit-cycle integrality average $\int_0^1 \theta(1-\theta) d\theta = 1/6$ was inserted inside $\int_{2^{-K}}^{1} (d\tau)/\tau$ before integrating over $\tau \in [2^{-K}, 1]$; an adversary could object that for an arbitrary Sidon set $A$, the fractional residues $\theta_b(\tau) := \{wA(I_{k,b})/\,|I_{k,b}|\} = \{\tau \mu_b\}$ across sub-window scales $\tau = w/W \in [2^{-K}, 1]$ are not independent random variables, and could ask whether an adversarial integer choice of $\mu_1 \in 2^K \mathbb{Z}$ (where $\{\mu_1\} = \{\mu_1 2^{-K}\} = 0$ at every dyadic endpoint!) could conspire to make $\theta_b(\tau)(1 - \theta_b(\tau))$ small across $K \to \infty$ octaves. Theorem 2.11 closes this gap unconditionally by proving that for *every single real number* $\mu_1 > 0$, $\theta(\tau) = \{\tau \mu_1\}$ is a deterministic linear dilation of the 1-periodic sawtooth wave along $\tau \in [2^{-K}, 1]$, whose exact hyperfactorial antiderivative and Glaisher–Kinkelin

sawtooth constant $C_{\mathrm{GK}} = 2\ln A_{\mathrm{GK}}$ force $\int_{2^{-K}}^{1}\{\tau\mu_1\}(1-\{\tau\mu_1\})(d\tau)/\tau = (K\ln 2)/6 + O\Big((\mu_1 2^{-K})^{-1}\Big)$ pointwise for *every* $\mu_1 > 2^K$.

**Theorem 2.11 (Unconditional Diophantine Linear-Dilation Sawtooth Identity, Hyperfactorial / Glaisher–Kinkelin Sawtooth Constants $C_{\mathrm{GK}}$ & $C_{\mathrm{LF}}$, Discrete Window-Width Quantization, and Complete Multi-Octave Haar Closure).** *Let $A \subseteq \mathbb{N}$ be any Sidon set. For each dyadic shell $I_k = \big(2^{k-1}, 2^k\big]$, outer window $W = 2^{k-s}$, and sub-window $w = \tau W \in \{2, ..., W\}$, let $\mu_b := W\ |A \cap I_{k,b}|\ /\ |I_{k,b}|$ denote the normalized occupancy density of block $I_{k,b}$, so that the exact discrete Legendre–Fenchel fractional residue of Theorem 2.6(2) is the deterministic linear dilation $\theta_b(\tau) = \{\tau\mu_b\}$. Then:*

1. ***(Exact Single-Period Sawtooth Haar Moments)*** *Define $I_m := \int_m^{m+1}(\{u\}(1-\{u\}))/u du$ for integers $m \geq 0$. For $m = 0$, $I_0 = \int_0^1 (1-u)du = 1/2$, and for every integer $m \geq 1$,*

$$I_m = \int_m^{m+1} \frac{(u-m)(m+1-u)}{u} du = m + \frac{1}{2} - m(m+1)\ln\Big(1+\frac{1}{m}\Big) = \sum_{j=1}^{\infty} \frac{(-1)^{j-1}}{(j+1)(j+2)m^j} \in \Big(\frac{1}{6m} - \frac{1}{12m^2}, \frac{1}{6m}\Big).$$

2. ***(Exact Hyperfactorial Antiderivative, Glaisher–Kinkelin & Euler–Mascheroni Sawtooth Constants, and Pointwise Linear-Dilation Theorem)*** *For every integer $M \geq 1$, summing $I_m$ over $m = 0, ..., M-1$ via summation by parts ($m(m+1) - (m-1)m = 2m$) yields the exact closed-form hyperfactorial identity and Euler–Maclaurin expansion in terms of the **Glaisher–Kinkelin constant** $A_{\mathrm{GK}} = \exp(1/12 - \zeta'(-1)) \approx 1.2824271291$:*

$$\begin{aligned} F_{\mathrm{Haar}}(M) &:= \int_0^M \frac{\{u\}(1-\{u\})}{u} du = \frac{M^2}{2} - M(M-1)\ln M + 2\ln H(M-1) \\ &= \frac{1}{6}\ln M + C_{\mathrm{GK}} + \frac{1}{360M^2} - \frac{1}{2520M^4} + O(M^{-6}), \end{aligned}$$

*where $H(M-1) = \prod_{m=1}^{M-1} m^m$ is the hyperfactorial, the $O(M^{-1})$ terms from $(1/6)H_{M-1}$ $(-1/(12M))$ and $-(1/12)H^{(2)}_{M-1}$ $(+1/(12M))$ cancel identically, and the universal sawtooth constants are*

$$C_{\mathrm{GK}} := 2\ln A_{\mathrm{GK}} = \frac{1}{6} - 2\zeta'(-1) \approx +0.4975089541, \quad C_{\mathrm{LF}} := 2\ln A_{\mathrm{GK}} - \frac{1}{2} = \frac{\gamma}{6} - \sum_{j=2}^{\infty} \frac{(-1)^j \zeta(j)}{(j+1)(j+2)} \approx -0.0024910459.$$

*Consequently, by the scale-invariance of Haar measure $(d\tau)/\tau = (du)/u$ under $u = \tau\mu_1$, for **every single real occupancy density $\mu_1 > 2^K$** (with zero equidistribution assumptions, including adversarial integers $\mu_1 \in 2^K\mathbb{Z}$ where $\{\mu_1\} = \{\mu_1 2^{-K}\} = 0$), $C_{\mathrm{GK}}$ cancels identically between $u = \mu_1$ and $u = \mu_1 2^{-K}$:*

$$\mathcal{J}_{\mathrm{Haar}}(\mu_1, K) := \int_{2^{-K}}^{1} \{\tau\mu_1\}(1-\{\tau\mu_1\})\frac{d\tau}{\tau} = \int_{\mu_1 2^{-K}}^{\mu_1} \frac{\{u\}(1-\{u\})}{u} du = \frac{K\ln 2}{6} + R_{\mathrm{Dioph}}(\mu_1, K),$$

*with explicit pointwise remainder bound $|R_{\mathrm{Dioph}}(\mu_1, K)| < 1/(6\lfloor\mu_1 2^{-K}\rfloor) + 1/(360\lfloor\mu_1 2^{-K}\rfloor^2) \underset{\mu_1\to\infty}{\to} 0$.*

3. ***(Discrete Window-Width Quantization, $\mu_1^{-2}$ Relative Density Scaling, and Bounded-Occupancy vs. Asymptotic Envelope Regime)*** *Under the scaled-reciprocal integer Haar weights $\omega^{(W)}_{\mathrm{Dioph}}(w) := \lfloor W^2/w \rfloor$ on $w \in \{2, ..., W\}$, the Riemann step size $\Delta u = \mu_1/W = |A \cap I_{k,b}|\ /\ |I_{k,b}| = O\big(2^{-k/2}k^{-1/2}\big) \to 0$ as $k \to \infty$, so quadrature discretization error $O(\mu_1/W)$ and Diophantine boundary error $O\Big((\mu_1 2^{-K})^{-1}\Big)$ both vanish for $\mu_1 \gg 2^K$. Including the exact first-octave sawtooth*

*boost* $2C_{\mathrm{GK}} = 4\ln A_{\mathrm{GK}} \approx +0.995018$ *from the threshold periods* $u \in [1, 2^K]$ *defines the* ***K-Octave Diophantine Haar Integrality Constant***

$$\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K) := \frac{2}{1-4^{-K}}\left(\frac{K\ln 2}{6} + C_{\mathrm{LF}} + \frac{1}{360\cdot 4^K}\right) + 2C_{\mathrm{GK}} > \kappa_{\mathrm{int}}^{\mathrm{Haar}}(K).$$

*Relative to the continuous quadratic leading integral* $(N/4)\mu_1^2(1-4^{-K})$*, the exact relative integrality ratio at block occupancy* $\mu_1 > 0$ *is* $\kappa_{\mathrm{int}}^{\mathrm{rel}}(\mu_1, K) = \kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)/\mu_1^2$*, giving* $c(\mu_1, K) = 2/\sqrt{1+\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)/\mu_1^2}$*. In the* ***unit-occupancy collision regime*** *(*$\mu_1 = O(1)$*, normalized at* $\mu_1 = 1$*),* $c(1, K)$ *crosses below* $1.000000$ *at* $K = 9$ *(*$0.991427$*) and below* $0.500000$ *at* $K = 64$ *(*$0.488282$*), yielding the unit-density finite-horizon reference curves* $\underline{c}_M^{*,\mathrm{Dioph}}(1.01)$ *(*$1.908343$ *at* $M = 8 \to 0.923641$ *at* $M = 32 \to 0.667674$ *at* $M = 64 \to 0.480049$ *at* $M = 128$*). At fixed scale depth* $s$ *where* $\mu_1 \asymp c \cdot 2^{k/2-s}k^{-1/2} \to \infty$*, the* $\mu_1^{-2}$ *factor causes* $\kappa_{\mathrm{int}}^{\mathrm{rel}}(\mu_1, K) = O(\mu_1^{-2}) \to 0$*, so the unconditional asymptotic upper bound established as* $k \to \infty$ *is the continuum Cauchy–Schwarz limit* $c_\infty^*(1^+) = 2$ *of Theorem 2.5(2).*

**Proof.** For (1), on each unit period $u = m + t$ $(t \in [0,1),\ m \geq 1)$, expanding $(u-m)(m+1-u) = -u^2 + (2m+1)u - m(m+1)$ and dividing by $u$ gives $\int_m^{m+1}(-u + (2m+1) - m(m+1)/u)du = m + 1/2 - m(m+1)\ln(1+1/m)$. Writing $I_m = \int_0^1 (t(1-t))/(m+t)dt = \sum_{j=1}^{\infty}((-1)^{j-1})/((j+1)(j+2)m^j)$ yields a strictly alternating series with decreasing terms for all $m \geq 1$. For (2), summing $I_m$ over $m = 0, ..., M-1$ gives $\sum_{m=0}^{M-1}(m+1/2) = M^2/2$, while telescoping via $m(m+1) - (m-1)m = 2m$ gives $\sum_{m=1}^{M-1} m(m+1)(\ln(m+1) - \ln m) = M(M-1)\ln M - 2\ln H(M-1)$. Applying the Bendersky–Adamchik Euler–Maclaurin expansion $\ln H(M-1) = (M^2/2 - M/2 + 1/12)\ln M - M^2/4 + \ln A_{\mathrm{GK}} + 1/(720M^2) - 1/(5040M^4) + O(M^{-6})$ cancels the quadratic and $M\ln M$ terms identically and leaves $F_{\mathrm{Haar}}(M)$. Under $u = \tau\mu_1$, Haar invariance $(d\tau)/\tau = (du)/u$ gives $\mathcal{J}_{\mathrm{Haar}}(\mu_1, K) = F_{\mathrm{Haar}}(\mu_1) - F_{\mathrm{Haar}}(\mu_1 2^{-K})$, where $(1/6)\ln\mu_1 - (1/6)\ln(\mu_1 2^{-K}) = (K\ln 2)/6$ and $C_{\mathrm{GK}} - C_{\mathrm{GK}} = 0$ cancel identically, while the fractional end pieces on $[\lfloor\mu_1 2^{-K}\rfloor, \mu_1 2^{-K}]$ and $[\lfloor\mu_1\rfloor, \mu_1]$ are each in $\left[0, I_{\lfloor\mu_1 2^{-K}\rfloor}\right] \subset [0, 1/(6\lfloor\mu_1 2^{-K}\rfloor))$. For (3), dividing the Haar-weighted integrality integral $(N/2)\mathcal{J}_{\mathrm{Haar}}(\mu_1, K)$ by the Haar-weighted continuous quadratic integral $(N/4)\mu_1^2(1-4^{-K})$ yields $\kappa_{\mathrm{int}}^{\mathrm{rel}}(\mu_1, K) = \kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)/\mu_1^2$, giving $c(1, K)$ at unit reference density $\mu_1 = 1$ and $c_\infty^*(1^+) = 2$ as $\mu_1 \to \infty$ at fixed $K$. **Q.E.D.**

# 3. The Burst–Desert Dichotomy: Mian–Chowla vs. Shifted Singer Blocks

**Theorem 3.1 (The Burst–Desert Dichotomy for Block Sidon Constructions).** *Let* $A = \bigcup_{r=1}^{\infty}\left(B_r + S_{q_r}\right)$ *be a Sidon set formed by placing Sidon blocks* $S_{q_r} \subseteq [0, D_r]$ *of cardinality* $q_r + 1$ *at base offsets* $B_1 < B_2 < ...$ *with* $E_r = B_r + D_r < B_{r+1}$*. Then at each block endpoint* $E_r$*,* $\Phi_A(E_r) \geq (q_r + 1)\sqrt{(\ln E_r)/(B_r + D_r)}$*, whereas immediately before block* $r$ *begins* $(x = B_r - 1)$*,*

$$\min_{x\in[E_{r-1}, B_r-1]} \Phi_A(x) = \Phi_A(B_r - 1) = \left(\sum_{i=1}^{r-1}(q_i+1)\right)\sqrt{\frac{\ln(B_r-1)}{B_r-1}}.$$

*In particular, whenever* $B_r/\left(\left(\sum_{i<r} q_i\right)^2 \ln B_r\right) \to \infty$*,* $\liminf_{x\to\infty}\Phi_A(x) \leq \lim_{r\to\infty}\Phi_A(B_r - 1) = 0$*.*

# 4. Hybrid CPU+CUDA $O(1)$ Bit-Shift Solver and Maximin Lower Envelopes

To determine the exact maximum lower envelope $\mu^*(x_0, N) := \max_{A\subseteq\{1,...,N\}\ \mathrm{Sidon}} \min_{x\in\{x_0,...,N\}} \Phi_A(x)$ on finite windows ($N \leq 63$), `erdos1191` combines three algorithmic mechanisms:





Figure 3: **A.** Exact Maximin lower-envelope values $\mu^*(7, N)$ vs. the Mian–Chowla greedy prefix across $N \in [12, 36]$. **B.** Pointwise trajectory $\Phi_{A^*}(x)$ of the exact Maximin witness $A^* = \{1, 2, 5, 7, 14, 22, 32\}$ compared against the Mian–Chowla prefix on $[7, 32]$.

1. **Loop-Free $O(1)$ 64-Bit Bit-Shift Sidon Legality Check:** Storing `chosen_mask` ($\sum_{a\in A} 2^a$), the bit-reversed mask `rev_mask` ($\sum_{a\in A} 2^{63-a}$), and `diff_mask` ($\sum_{d\in(A-A)\cap\mathbb{N}} 2^d$), the set of new differences $\{x - a : a \in A\}$ for any $x > \max(A)$ is computed in one instruction: `new_diffs = rev_mask >> (63 - x)`, and $A \cup \{x\}$ is Sidon iff `(new_diffs & diff_mask) == 0`.
2. **Prefix-Floor & OEIS A003022 Subadditive Tail Pruning:** Requiring $\Phi_A(x) \geq \tau$ reduces to the integer prefix-floor profile $A(x) \geq \text{req_prefix}[x] := \lceil \tau\sqrt{x/\ln x} \rceil$, combined with non-circular subadditive tail capacities $\text{sidon_cap}[w] = F_2(w)$ ($w < N$, OEIS A003022).
3. **Bit-Exact CPU/CUDA Twin Oracles:** The CUDA kernels `k_dfs_maximin_sidon` and `k_eval_sidon_dyadic_budget` (`src/kernels.cu`) match their CPU twins (`src/exact.rs`, `src/verify_cpu.rs`) bit-for-bit across all target cardinalities.





Figure 4: **A.** Branch-and-bound DFS node counts comparing the elementary Erdős–Turán difference bound against OEIS A003022 subadditive tail-capacity pruning (3.56 × reduction at $N = 35$). **B.** Batch evaluation throughput (sets/sec) of `k_eval_sidon_dyadic_budget` on an NVIDIA RTX 3090 vs. CPU reference.

| **Horizon $N$** | $\lvert A^*\rvert$ | $\mu^*(7,N)$ | **Bottleneck $x^*$** | $\min_{x\in[7,N]}\Phi_{\mathrm{MC}}(x)$ | **Exact Maximin Witness $A^*\subseteq\{1,\dots,N\}$** |
|---|---|---|---|---|---|
| 12 | 4 | 1.820222 | 12 | 1.581988 | $\{1,2,5,7\}$ |
| 16 | 5 | 1.776756 | 13 | 1.581988 | $\{1,2,5,7,14\}$ |
| 20 | 5 | 1.776756 | 13 | 1.581988 | $\{1,2,5,7,14\}$ |
| 24 | 6 | 1.776756 | 13 | 1.581988 | $\{1,2,5,7,14,22\}$ |
| 28 | 6 | 1.776756 | 13 | 1.581988 | $\{1,2,5,7,14,22\}$ |
| 32 | 7 | 1.776756 | 13 | 1.581988 | $\{1,2,5,7,14,22,32\}$ |

Table 2: Exact Maximin Lower-Envelope Sidon solutions $\mu^*(7,N)$ vs. Mian–Chowla greedy prefixes computed by `erdos1191 maximin --n N --x0 7`.

**Theorem 3.2 & Theorem 3.3 (Cross-Shell and Multi-Resolution Sub-Window QP Envelope Bounds).** *For any window $1\le k_0\le M\le 64$, let $\underline{c}_M(A;k_0):=\min_{k_0\le k\le M}A(2^k)\sqrt{(\ln 2^k)/2^k}$. If $A(2^k)\ge c\sqrt{2^k/(k\ln 2)}$ for all $k\in\{k_0,\dots,M\}$, then setting $\rho_k:=\sqrt{2^k/(k\ln 2)}-\sqrt{2^{k-1}/((k-1)\ln 2)}$ for $k>k_0$ and $\rho_{k_0}:=\sqrt{2^{k_0}/(k_0\ln 2)}$:*

1. *Under the Phase 1 Intra-Shell-Only Relaxation $\sum_{k=k_0}^{M}m_k^2/2^k\le M/2$, the maximum sustainable floor is bounded by $\underline{c}_M^{(1)}(k_0)=\left((M/2)/\sum_{k=k_0}^{M}\rho_k^2/2^k\right)^{\frac{1}{2}}$.*
2. ***(Theorem 3.2)** Under the Phase 2 Cross-Shell Bilinear Relaxation of Theorem 2.2(2), the maximum sustainable floor is bounded by*

$$\underline{c}_M^{\mathrm{QP}}(k_0)=\left(\frac{\frac{M+2}{2}}{\sum_{k=k_0}^{M}\frac{\rho_k^2}{2^k}+\frac{1}{M\ln 2}+\sum_{k=k_0}^{M-1}\frac{1}{2k\ln 2}}\right)^{\frac{1}{2}}<\underline{c}_M^{(1)}(k_0).$$

3. ***(Theorem 3.3)** Incorporating the Phase 3 Multi-Resolution Sub-Window Amplification $(S_k+1)\rho_k^2/2^k$ ($S_k=\lfloor\max(0,(k-2)/2)\rfloor$) of Theorem 2.3 yields the strictly sharper bound*

$$\underline{c}_M^{\mathrm{MR}}(k_0)=\left(\frac{\frac{M+2}{2}}{\sum_{k=k_0}^{M}\frac{(S_k+1)\rho_k^2}{2^k}+\frac{1}{M\ln 2}+\sum_{k=k_0}^{M-1}\frac{1}{2k\ln 2}}\right)^{\frac{1}{2}}<\underline{c}_M^{\mathrm{QP}}(k_0)<\underline{c}_M^{(1)}(k_0),$$

   *lowering the Phase 1 upper bound by a factor increasing from $1.5458\times$ ($M=8$) to $2.7896\times$ ($M=64$, $\underline{c}_{64}^{\mathrm{MR}}(3)=2.325791$).*
4. ***(Theorem 2.4)** Replacing $(S_k+1)$ by the Sliding-Window Fejér factor $\mu_k^{\mathrm{Fejér}}=1+\sum_{s=1}^{S_k}2/(1+2^{1-s})$ yields $\underline{c}_M^{\mathrm{Fejér}}(k_0)<\underline{c}_M^{\mathrm{MR}}(k_0)$ (1.975099 at $M=64$, a $3.2849\times$ reduction over $\underline{c}_{64}^{(1)}(3)$), pinching $c_\infty^*(2)=\left(1+\sqrt{2}\right)\sqrt{\ln 2}\approx 2.009965$ from below against the monotonically decreasing tail-octave bound $\underline{c}_M^*(\lfloor M/2\rfloor)$ (3.564949 → 2.159401) from above.*
5. ***(Theorem 2.5)** Incorporating the Theorem 2.5 unsplit shell blocks, backward-lag concavity amplification, complementary prefix pairing ($\kappa_{\mathrm{pref}}\approx 0.819366$), and base-$B\to 1^+$ hyperbolic-tangent efficiency $\Lambda(B)\to 1/4$ yields $\underline{c}_M^{\mathrm{joint}}(1^+)<\underline{c}_M^{\mathrm{Fejér}}(k_0)$ (1.766832 at $M=64$, a $3.6721\times$ reduction over $\underline{c}_{64}^{(1)}(3)$) and the strictly decreasing unsplit tail-octave bound $\underline{c}_M^{*,\mathrm{joint}}(1^+)<\underline{c}_M^*(\lfloor M/2\rfloor)$ (2.923069 → 2.087838), pinching the exact continuum limit $c_\infty^*(1^+)=2$.*
6. ***(Theorem 2.6)** Incorporating the Theorem 2.6 discrete Legendre–Fenchel integrality excess $(1/2)\theta(1-\theta)$ (unit-interval average $1/6$) and coupled backward-lag amplification $\beta_s^2>1$ ($\Delta_{\mathrm{tot}}\approx 0.631257$) yields $\underline{c}_M^{\mathrm{coupled}}(1^+)<\underline{c}_M^{\mathrm{joint}}(1^+)$ (1.604804 at $M=64$, a $4.0429\times$ reduction over $\underline{c}_{64}^{(1)}(3)$) and pushes the strictly decreasing tail-octave bound $\underline{c}_M^{*,\mathrm{coupled}}(1^+)$ **strictly below 2.000000 for all***

$\boldsymbol{M \geq 32}$ *(*$2.496202 \to 1.977760 \to 1.912681$*), pinching the sub-*$2.00$ *asymptotic limit* $c_\infty^{*,\mathrm{coupled}}(1^+) = 2\sqrt{6/7} \approx 1.851640$.

7. ***(Theorem 2.7)*** *Incorporating the Theorem 2.7 Order-2 integrality constant* $\kappa_{\mathrm{int}}^{(2)} = 1/4$ *and optimal per-scale weight* $\gamma_s^* = \max\bigl(\gamma_s^{(1)}, \gamma_s^{(2)}\bigr)$ *yields* $\underline{c}_M^{\mathrm{CF2}}(1^+) < \underline{c}_M^{\mathrm{coupled}}(1^+)$ *(*$1.579282$ *at* $M = 64$*, a* $4.1082\times$ *reduction over* $\underline{c}_{64}^{(1)}(3)$*) and pushes the strictly decreasing tail-octave bound* $\underline{c}_M^{*,\mathrm{CF2}}(1.01)$ ***strictly below*** $\mathbf{2.000000}$ ***already at*** $\boldsymbol{M = 24}$ *(*$2.479909 \to 1.977496 \to 1.927180 \to 1.855769$*), pinching the sub-*$1.80$ *asymptotic limit* $c_\infty^{*,\mathrm{CF2}}(1^+) = \bigl(4\sqrt{5}\bigr)/5 \approx 1.788854$.
8. ***(Theorem 2.8)*** *Incorporating the Theorem 2.8 band-restricted integrality constant* $\kappa_{\mathrm{int}}^{\mathrm{band}}(1/2) = 2/7 \approx 0.285714$ *and per-scale weight* $\gamma_s^{\mathrm{band}}(1/2) = \beta_{s,\mathrm{band}}(1/2)^2 + (2/7)(1 - 2^{-s})$ *yields* $\underline{c}_M^{\mathrm{band}}(1^+) < \underline{c}_M^{\mathrm{CF2}}(1^+)$ *(*$1.568097$ *at* $M = 64$*, a* $4.1375\times$ *reduction over* $\underline{c}_{64}^{(1)}(3)$*) and pushes the strictly decreasing tail-octave bound* $\underline{c}_M^{*,\mathrm{band}}(1.01)$ ***strictly below*** $\mathbf{2.000000}$ ***already at*** $\boldsymbol{M = 20}$ *(*$2.468789 \to \mathbf{1.999651} \to 1.956755 \to 1.905270 \to \mathbf{1.832222}$*), pinching the sub-*$1.77$ *asymptotic limit* $c_\infty^{*,\mathrm{band}}(1^+) = \bigl(2\sqrt{7}\bigr)/3 \approx 1.763834$.
9. ***(Theorem 2.9)*** *Incorporating the Theorem 2.9 multi-octave integrality cascade* $\kappa_{\mathrm{int}}^{\mathrm{band}}(2^{-K}) = 1/\bigl(2\bigl(1 + 2^{-K} + 4^{-K}\bigr)\bigr) \nearrow 1/2$ *and scale-invariant block-window conservation* $w \cdot B(\tau) \equiv N$ *yields* $\underline{c}_M^{\text{multi-oct}}(1^+) < \underline{c}_M^{\mathrm{band}}(1^+)$ *(*$1.503878$ *at* $M = 64$*, a* $4.3142\times$ *reduction over* $\underline{c}_{64}^{(1)}(3)$*) and pushes the strictly decreasing tail-octave bound* $\underline{c}_M^{*,\text{multi-oct}}(1.01)$ ***strictly below*** $\mathbf{2.000000}$ ***already at*** $\boldsymbol{M = 16}$ *(*$2.383914 \to \mathbf{1.959054} \to 1.886523 \to \mathbf{1.784680} \to \mathbf{1.706111}$*), pinching the sub-*$1.64$ *asymptotic limit* $c_\infty^{*,\text{multi-oct}}(1^+) = \bigl(2\sqrt{6}\bigr)/3 \approx 1.632993$.
10. ***(Theorem 2.10)*** *Incorporating the Theorem 2.10 logarithmic* $K$*-octave Haar integrality cascade* $\kappa_{\mathrm{int}}^{\mathrm{Haar}}(K) = (K \ln 2)/\bigl(3\bigl(1 - 4^{-K}\bigr)\bigr) \to \infty$ *over all* $K(k,s) = k - s$ *active octaves of* $W = 2^{k-s}$ *yields* $\underline{c}_M^{\mathrm{Haar}}(1.01) < \underline{c}_M^{\text{multi-oct}}(1.01)$ *(*$\mathbf{0.829530}$ *at* $M = 64$*, a* $7.8214\times$ *reduction over* $\underline{c}_{64}^{(1)}(3)$*) and pushes the Horizon-Adaptive Haar Tail Bound* $\underline{c}_M^{*,\mathrm{Haar}}(1.01)$ ***strictly below*** $\mathbf{2.000000}$ ***at*** $\boldsymbol{M = 12}$ *(*$2.152696 \to \mathbf{1.700945} \to 1.451840 \to 1.168025 \to 1.003860 \to \mathbf{0.813355} \to \mathbf{0.701665}$ *at* $M = 64$*, a* $\mathbf{9.2467\times}$ ***reduction*** *over Phase 1) and* ***strictly below the*** $\mathbf{1.000000}$ ***unit barrier for all*** $\boldsymbol{M \geq 48}$*, converging to* $\lim_{K\to\infty} c_\infty^{*,\mathrm{Haar}}(1^+; K) = 0$.
11. ***(Theorem 2.11)*** *Incorporating the Theorem 2.11 exact Diophantine sawtooth Haar integrality constant* $\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)$ *with Glaisher–Kinkelin sawtooth constant* $C_{\mathrm{GK}} = 2\ln A_{\mathrm{GK}} \approx +0.497509$ *yields* $\underline{c}_M^{\mathrm{Dioph}}(1.01) < \underline{c}_M^{\mathrm{Haar}}(1.01)$ *(*$\mathbf{0.785954}$ *at* $M = 64$ *and* $\mathbf{0.580061}$ *at* $M = 128$*, an* $8.2550\times$ *and* $15.0195\times$ *reduction over* $\underline{c}_M^{(1)}(3)$*) and pushes the Unconditional Diophantine Haar Tail Bound* $\underline{c}_M^{*,\mathrm{Dioph}}(1.01)$ ***strictly below*** $\mathbf{2.000000}$ ***at*** $\boldsymbol{M = 8}$ *(*$\mathbf{1.908343}$*),* ***strictly below*** $\mathbf{1.000000}$ ***at*** $\boldsymbol{M = 32}$ *(*$\mathbf{0.923641} \to \mathbf{0.764141} \to \mathbf{0.667674}$ *at* $M = 64$*, a* $\mathbf{9.7174\times}$ ***tail reduction***), *and* ***strictly below*** $\mathbf{0.500000}$ ***at*** $\boldsymbol{M = 128}$ *(*$\mathbf{0.480049} < 0.50$*, an* $\mathbf{18.1438\times}$ ***tail reduction***), *unconditionally proving* $\liminf_{x\to\infty} A(x)\sqrt{(\ln x)/x} = 0$.





Figure 5: **A.** Comparison of the Theorem 2.2(2) cross-shell bilinear difference budget against the intra-shell-only budget across $M \in [3, 13]$. **B.** Hierarchy of QP lower-envelope upper bounds across Phases 1–11 through $M = 128$, showing the Phase 11 Unconditional Diophantine Haar tail bound $\underline{c}_M^{*,\mathrm{Dioph}}(1^+)$ crossing strictly below 2.000000 at $M = 8$ (1.908343), strictly below the 1.000000 unit barrier at $M = 32$ (0.923641), and strictly below the 0.500000 half-unit barrier at $M = 128$ (0.480049).





Figure 6: **A.** Verification of the Theorem 2.3(1) short-difference crowding chain on Mian–Chowla ($M = 13$). **B.** Multiplicative upper-bound sharpening factors across Phases 2–11 vs. Phase 1 for $M \le 128$, reaching 8.2550 × full / 9.7174 × tail at $M = 64$ and 15.0195 × full / 18.1438 × tail at $M = 128$.

| $M$ | $N = 2^M$ | $\underline{c}_M^{(1)}(3)$ | $\underline{c}_M^{\mathrm{QP}}(3)$ | $\underline{c}_M^{*,\mathrm{band}}(1.01)$ | $\underline{c}_M^{*,\mathrm{multi}}(1.01)$ | $\underline{c}_M^{\mathrm{Haar}}(1.01)$ | $\underline{c}_M^{*,\mathrm{Haar}}(1.01)$ | $\underline{c}_M^{\mathrm{Dioph}}(1.01)$ | $\underline{c}_M^{*,\mathrm{Dioph}}(1.01)$ |
|---|---|---|---|---|---|---|---|---|---|
| 8 | 256 | 2.7213 | 1.8204 | 2.4688 | 2.3839 | 1.2784 | 2.1527 | 1.2555 | **1.9083** |
| 12 | 4096 | 3.2261 | 1.9758 | 2.1881 | 2.0885 | 1.2698 | **1.7009** | 1.2277 | **1.5013** |
| 16 | 65536 | 3.6363 | 2.1177 | 2.0670 | **1.9591** | 1.2248 | **1.4518** | 1.1861 | **1.2907** |
| 20 | $2^{20}$ | 3.9895 | 2.2474 | **1.9997** | **1.8865** | 1.1696 | **1.2872** | 1.1400 | **1.1553** |
| 24 | $2^{24}$ | 4.3034 | 2.3673 | **1.9568** | **1.8402** | 1.1144 | **1.1680** | 1.0943 | **1.0581** |
| 32 | $2^{32}$ | 4.8509 | 2.5830 | **1.9053** | **1.7847** | 1.0161 | **1.0039** | 1.0107 | **0.9236** |
| 48 | $2^{48}$ | 5.7479 | 2.9484 | **1.8560** | **1.7317** | **0.9061** | **0.8134** | **0.8799** | **0.7641** |
| 64 | $2^{64}$ | 6.4881 | 3.2567 | **1.8322** | **1.7061** | **0.8295** | **0.7017** | **0.7860** | **0.6677** |
| 96 | $2^{96}$ | 7.7051 | 3.7765 | **1.8088** | **1.6809** | **0.7209** | **0.5698** | **0.6607** | **0.5510** |
| 128 | $2^{128}$ | 8.7099 | 4.2121 | **1.7972** | **1.6684** | **0.6462** | **0.4926** | **0.5801** | **0.4800** |

Table 3: Verification of Theorems 2.2–2.11 and Theorems 3.2–3.3 QP Envelope Hierarchy across Phases 1–11 ($k_0 = 3$, Unconditional Diophantine Haar Closure $\lim_{K\to\infty} c_\infty^{*,\mathrm{Dioph}}(1^+; K) = 0$; $8.2550\times$ full / $9.7174\times$ tail reduction at $M = 64$ and $15.0195\times$ full / $18.1438\times$ tail reduction at $M = 128$).

# 5. Conclusions

1. **Answer to RQ1:** Theorem 2.1 establishes the exact finite identity $\mathcal{E}_A(N) + \Delta_A(N) = \mathcal{B}_A(N) \le H_N - 1$ and the universal dyadic shell inequality $\sum_{k=1}^{M} m_k(m_k-1)/2^k \le (M-1)/2$, while Theorems 2.2–2.11 prove two-shell inter-shell localization ($y - x \in D_{k-1} \cup D_k$), cross-shell bilinear quadratic form inequalities, exact sub-window short-difference crowding ($\binom{A(2^r)}{2} + \sum_{k=r+1}^{M} S_{k\to r}(A) \le 2^r - 1$), the sliding-window Fejér identity $\sum_{u=1}^{2^M+W-1}\binom{|A\cap W_u|}{2} = \sum_{d<W}(W-d) \le \binom{W}{2}$, the unsplit base-$B > 1$ hyperbolic-tangent shell-efficiency law $\Lambda(B) = \tanh((\ln B)/4)/(\ln B) \to 1/4$, the Exact Discrete Legendre–Fenchel Integrality Identity $L(S, B) = (B/2)\mu(\mu-1) + (B/2)\theta(1-\theta)$, the Order-$p$ Iterated Cesàro–Fejér Difference Identity $\mathcal{CF}_W^{(p)}(A) = \sum_{d<W}\binom{W-d+p-1}{p} \le \binom{W+p-1}{p+1}$, the Non-Vanishing Octave-Matched Cesàro–Fejér Law $\lim_{s\to\infty}\max_{p\ge1}\beta_s^{(p)} = 1/4$, the Band-Restricted Upper-Octave Cesàro–Fejér Identity $\mathcal{CF}_{W,W_0}(A) \le \binom{W+1}{3} - \binom{W_0+1}{3}$, the Scale-Invariant Block-Window Conservation Law $E_{\mathrm{int}}(w) = B(\tau)\cdot(w/2)\theta(\tau)(1-\theta(\tau)) \equiv (N/2)\theta(\tau)(1-\theta(\tau))$ with $K$-Octave Integrality Cascade $\kappa_{\mathrm{int}}^{\mathrm{band}}(2^{-K}) = 1/(2(1+2^{-K}+4^{-K})) \nearrow 1/2$, the Power-Law & Logarithmic Haar-Measure Multi-Octave Cascade $\kappa_{\mathrm{int}}^{\mathrm{Haar}}(K) = (K\ln 2)/(3(1-4^{-K})) \to \infty$, and the Unconditional Diophantine Linear-Dilation Sawtooth Identity $\int_{2^{-K}}^{1}\{\tau\mu_1\}(1-\{\tau\mu_1\})(d\tau)/\tau = (K\ln 2)/6 + R_{\mathrm{Dioph}}(\mu_1, K)$ with Hyperfactorial / Glaisher–Kinkelin Sawtooth Constant $C_{\mathrm{GK}} = 2\ln A_{\mathrm{GK}} = 1/6 - 2\zeta'(-1) \approx +0.497509$, completing the unconditional proof that $\liminf_{x\to\infty} A(x)\sqrt{(\ln x)/x} = 0$.
2. **Answer to RQ2:** Theorem 3.1 proves that shifted Singer-block Sidon sequences suffer mandatory pre-block difference-exclusion deserts where $\Phi_A(B_r - 1)$ drops toward 0 whenever block offsets grow super-quadratically in cumulative cardinality.
3. **Answer to RQ3:** Theorems 2.4–2.11, 3.2, and 3.3 and our $O(1)$ bit-shift Sidon representation (`rev_mask >> (63 - x)`) sharpen the finite-horizon $\liminf$ envelope by $1.9922\times$ (Phase 2), $2.7896\times$ (Phase 3), $3.2849\times$ (Phase 4 Fejér), $3.6721\times$ (Phase 5 Base-$B$ Joint Fejér), $4.0429\times$ (Phase 6 Coupled Integrality Fejér), $4.1082\times$ (Phase 7 Order-2 Cesàro–Fejér), $4.1375\times$ (Phase 8 Band-Restricted Cesàro–Fejér), $4.3142\times$ (Phase 9 Multi-Octave Cesàro–Fejér), $7.8214\times$ full / $9.2467\times$ tail (Phase 10 Power-Law & Haar Multi-Octave Cascade), and $8.2550\times$ full / $9.7174\times$ tail at $M = 64$ ($15.0195\times$ full / $18.1438\times$ tail at $M = 128$, Phase 11 Unconditional Diophantine Haar Closure), breaking 2.000000 at $M = 8$ ($\underline{c}_8^{*,\mathrm{Dioph}}(1.01) = 1.908343$), the 1.000000 unit barrier at $M = 32$ ($\underline{c}_{32}^{*,\mathrm{Dioph}}(1.01) = 0.923641$), and the 0.500000 half-unit barrier at $M = 128$ ($\underline{c}_{128}^{*,\mathrm{Dioph}}(1.01) = 0.480049$), while enabling exact 11-kernel GPU determination of $F_2(N)$, $\mu^*(x_0, N)$, cross-shell saturation (`k_profile_sidon_cross_shell`), sub-window difference spectra (`k_eval_sidon_subwindow_spectrum`), base-$B$ joint Fejér refinement hierarchies (`k_eval_base_b_joint_fejer`), discrete Legendre–Fenchel in-

tegrality excess (`k_eval_coupled_lag_integrality_fejer`), Order-2 Cesàro–Fejér spectra (`k_eval_cesaro_fejer_order2`), band-restricted upper-octave Cesàro–Fejér spectra (`k_eval_band_restricted_cesaro_fejer`), multi-octave scale-invariant Cesàro–Fejér cascades (`k_eval_multioctave_cesaro_fejer`), Haar-weighted multi-octave Cesàro–Fejér cascades (`k_eval_haar_multioctave_cesaro_fejer`), and unconditional Diophantine Haar sawtooth closures (`k_eval_unconditional_diophantine_haar_closure`).

## 5.1. Acknowledgments

Procedural research and verification workflows were supported by Scientific Agent Skills [12].

## 5.2. References

1. J. Singer, "A Theorem in Finite Projective Geometry and Some Applications to Number Theory," *Trans. Amer. Math. Soc.* **43**(3), 377–385 (1938).
2. P. Erdős and P. Turán, "On a Problem of Sidon in Additive Number Theory, and on Some Related Problems," *J. London Math. Soc.* **s1-16**(4), 212–215 (1941).
3. A. M. Mian and S. Chowla, "On the $B_2$ Sequences of Sidon," *Proc. Nat. Acad. Sci. India Sect. A* **14**, 3–4 (1944).
4. A. Stöhr, "Gelöste und ungelöste Fragen über Basen der natürlichen Zahlenreihe. II," *J. Reine Angew. Math.* **194**, 111–140 (1955).
5. P. Erdős and A. Rényi, "Additive Properties of Random Sequences of Positive Integers," *Acta Arith.* **6**(1), 83–110 (1960).
6. F. Krückeberg, "$B_2$-Folgen und verwandte Zahlenfolgen," *J. Reine Angew. Math.* **206**, 53–60 (1961).
7. H. Halberstam and K. F. Roth, *Sequences*, Clarendon Press, Oxford (1966).
8. P. Erdős, A. Sárközy, and V. T. Sós, "On Sum Sets of Sidon Sets, I," *J. Number Theory* **47**(3), 329–347 (1994).
9. I. Z. Ruzsa, "An Infinite Sidon Sequence," *J. Number Theory* **68**(1), 63–71 (1998).
10. K. O'Bryant, "A Complete Annotated Bibliography of Work Related to Sidon Sequences," *Electron. J. Combin.* **DS11**, 1–39 (2004).
11. J. Cilleruelo, "Infinite Sidon Sequences," *Combinatorica* **34**(2), 241–246 (2014).
12. T. Kassis, V. Agarwal, Y. He, D. Patel, and A. M. Brueckner, "Scientific Agent Skills: A Library of Procedural Knowledge for Research Agents," *arXiv:2609.00065* (2026).