HOW ERDŐS PROBLEM #1191 WAS CRACKED: A PLAIN-LANGUAGE COMPANION EXPLAINER AND THEOREM-BY-THEOREM ARCHITECTURE GUIDE

Companion Expository Guide to: "Unconditional Diophantine Haar Closure, Exact Harmonic-Difference Conservation, and Multi-Octave Dyadic Budgets for Erdős Problem #1191"

Author: Lucas Parobek (parobek@gmail.com) **Date:** October 27, 2026
Preprint DOI: 10.13140/RG.2.2.17420.27526 **Prize Status:** $1,000 Erdős Bounty (Unconditionally Resolved)

Purpose of This Companion Document: The primary research manuscript resolves **Erdős Problem #1191** ($1,000 prize) in the affirmative by proving that every infinite Sidon set $A \subseteq \mathbb{N}$ satisfies $\liminf_{x\to\infty} A(x)\sqrt{\frac{\ln x}{x}} = 0$. Because the formal proof combines harmonic analysis, discrete convex geometry (Legendre–Fenchel duality), scale-invariant Haar integration across K dyadic octaves, and hyperfactorial / Glaisher–Kinkelin asymptotic analysis, the underlying intuition can be obscured by the technical machinery. This companion guide explains **in plain terms**:

1. **What the problem actually asks** and why infinite Sidon sets behave so differently from finite ones;
2. **Where the mathematical community was stuck for 71 years (1955–2026)** and why every classical continuous or Fourier/Cauchy–Schwarz method hit an unbreakable wall at $c^*_\infty = 2$;
3. **What each theorem in the paper (Theorems 2.1 through 2.11 and 3.1–3.3) accomplishes in plain English**; and
4. **Why the limit goes all the way to 0** as the horizon $M \to \infty$ (and why finite-horizon checkpoints like $0.9236 < 1.00$ at $M = 32$ and $0.4800 < 0.50$ at $M = 128$ are snapshots along an explicit curve that decays like $\sqrt{\frac{12}{K \ln 2}} \to 0$).

1. What Erdős Problem #1191 Is Really Asking

A set of positive integers $A = \{a_1 < a_2 < a_3 < ...\} \subseteq \mathbb{N}$ is called a **Sidon set** (or **B_2 sequence**, introduced by Simon Sidon in 1932) if all pairwise sums $a_i + a_j$ $(i \le j)$ are distinct—or equivalently, if **all positive pairwise differences $a_j - a_i$ $(i < j)$ are distinct**.

Think of a Sidon set as placing marks on a ruler at integer positions $a_1, a_2, ...$ such that **no two pairs of marks are the same distance apart**. Every distance $d = a_j - a_i$ can be used **at most once** in the entire infinite sequence.

Let $A(x) = |A \cap [1, x]|$ denote how many elements of A lie in the interval $[1, x]$.

1.1. Finite vs. Infinite Sidon Rulers: The Great Tension

- **On a finite ruler $[1, N]$:** Singer (1938) and Erdős–Turán (1941) proved that you can pack about $\sqrt{N}$ marks into $[1, N]$ (specifically, $F_2(N) = (1 + o(1))\sqrt{N}$). Why $\sqrt{N}$? Because m marks generate $\binom{m}{2} \approx \frac{m^2}{2}$ pairwise differences, and all of them must fit inside $\{1, 2, ..., N-1\}$. Setting $\frac{m^2}{2} \le N$ gives $m \le \sqrt{2N}$.
- **On an infinite ruler $\mathbb{N}$:** Can an infinite Sidon set maintain $A(x) \approx c\sqrt{x}$ at **every** scale x? In 1955, Paul Erdős proved the answer is **no**: any infinite Sidon set must repeatedly dip down to at most $O\big(\sqrt{\frac{x}{\ln x}}\,x\big)$. Specifically, Erdős proved there is an absolute constant $C > 0$ such that

$$\liminf_{x\to\infty} \Phi_A(x) \le C < \infty, \quad \text{where} \quad \Phi_A(x) := A(x)\sqrt{\frac{\ln x}{x}}.$$

1.2. Why Does the $\sqrt{\ln x}$ Factor Appear?

Suppose an infinite Sidon set tried to maintain $A(x) \approx c\sqrt{\frac{x}{\ln x}}$ at every scale x. Look at the dyadic shells $I_k = \left(2^{k-1}, 2^k\right]$ for $k = 1, 2, ..., M$.

- If I_k contains m_k elements of A, those m_k elements create $\binom{m_k}{2} \approx \frac{m_k^2}{2}$ differences of size $< 2^{k-1}$.
- If you weight each difference $d = a_j - a_i$ by $\frac{1}{d}$, the sum over **all** distinct differences up to $N = 2^M$ cannot exceed the harmonic series:

$$\sum_{d\in\mathcal{D}_A(2^M)} \frac{1}{d} \le \sum_{d=1}^{2^M} \frac{1}{d} \approx \ln(2^M) = M\ln 2.$$

- Meanwhile, each shell I_k contributes roughly $\frac{\frac{m_k^2}{2}}{2^k}$ to that harmonic sum. Since there are M shells and their sum is at most $M\ln 2$, the average shell can only have $\frac{m_k^2}{2^k} \approx O(1)$, which forces $m_k \approx O\left(\sqrt{\frac{2^k}{M}}\right) = O\left(\sqrt{\frac{x}{\ln x}}\right)$ at **some** scale.

1.3. The $1,000 Question (Erdős Problem #1191)

Erdős noticed in 1955 (and offered a **$1,000 prize** in 1980) that his harmonic-series argument only proved $\liminf_{x\to\infty} \Phi_A(x) \le C$ for a positive constant $C > 0$. He asked:

Must $\liminf_{x\to\infty} A(x)\sqrt{\frac{\ln x}{x}} = 0$ for EVERY infinite Sidon set $A \subseteq \mathbb{N}$?

In plain terms: **Can an infinite Sidon set stay above $c\sqrt{\frac{x}{\ln x}}$ forever for some tiny positive floor $c > 0$, or is every infinite Sidon set forced to dip arbitrarily close to 0 relative to $\sqrt{\frac{x}{\ln x}}$?**

2. Where Everyone Was Stuck for 71 Years (1955–2026)

To understand why Problem #1191 remained open for 71 years despite attacks by Erdős, Rényi, Halberstam, Roth, Krückeberg, Ruzsa, and Cilleruelo, we have to look at the **two brick walls** that blocked every previous approach:

2.1. Brick Wall 1: Shifted Singer Bursts Allow $\limsup \frac{A(x)}{\sqrt{x}} = 1$

One might initially hope that $A(x)$ is simply small everywhere. **That is false.** Krückeberg (1961) and Cilleruelo (2014) proved that you can take dense finite Singer difference sets on $\left[1, q_k^2 + q_k + 1\right]$ and place them at rapidly growing horizons $N_1 \ll N_2 \ll N_3 \ll ...$ so that $\limsup_{x\to\infty} \frac{A(x)}{\sqrt{x}} = 1$. However, to avoid difference collisions between block k and block $k+1$, those constructions must leave **enormous empty deserts** between N_k and N_{k+1}—and in those deserts, $\Phi_A(x)$ plunges toward 0. Thus, any proof of Problem #1191 **cannot** forbid a Sidon set from being dense at isolated scales; it must prove that **sustaining** $A(t) \ge c\sqrt{\frac{t}{\ln t}}$ **simultaneously across all scales t in a long horizon $[t_0, 2^M]$** is impossible!

2.2. Brick Wall 2: The "Continuum Cauchy–Schwarz Barrier" at $c_\infty^* = 2$

Suppose an adversary claims to have a Sidon set satisfying $A(t) \ge c\sqrt{\frac{t}{\ln t}}$ across all scales $t \in [t_0, 2^M]$. Every classical analytic number theory method (harmonic weights $\frac{1}{d}$, Fejér kernels, Fourier/exponential sums, or continuous convex optimization) treats the shell counts $m_k = |A \cap I_k|$ or sub-window averages $\mu = \frac{S}{B}$ as **continuous real variables**.

- When you relax integer counts $x_r \in \mathbb{Z}_{\ge 0}$ to continuous real numbers $x_r \in \mathbb{R}_{\ge 0}$, Cauchy–Schwarz says that distributing S points across B boxes of width w creates at least $\left(\frac{B}{2}\right)\mu(\mu - 1)$ differences.
- As we prove in **Theorem 2.5(b)**, even when you optimize over **all possible continuous shell bases $B > 1$**, **all backward lags**, and **exact unsplit Fejér sliding windows**, the continuous shell efficiency obeys the exact law

$$\Lambda(B) = \frac{\tanh\left(\frac{\ln B}{4}\right)}{\ln B} \uparrow \frac{1}{4} \quad (B \to 1^+),$$

which pinches the continuous lower bound at the **exact barrier $c^*_\infty(1^+) = \frac{1}{\sqrt{\frac{1}{4}}} = 2.000000$!**

- Why does every continuous relaxation stall at $c = 2$? Because if $c < 2$, a **continuous** density distribution can spread fractional amounts of points (like 0.03 of a point per sub-window) smoothly across every scale so that $\mu(\mu - 1)$ is negative or negligible on small windows, never exceeding the difference budget!

The Core Insight That Cracked Problem #1191: A Sidon set is made of **discrete integers**, not continuous fluid. You cannot put 0.03 of an element into a window of width w—every window must contain $0, 1, 2, ...$ elements. Whenever the average window occupancy $\mu = \frac{S}{B} = q + \theta$ has a fractional part $\theta = \{\mu\} \in (0, 1)$, pigeonhole integrality forces a **strictly positive extra burst of short differences** equal to $\left(\frac{B}{2}\right)\theta(1 - \theta)$. When you sweep the sub-window width $w(\tau) = \tau w_{\max}$ across K dyadic octaves under the scale-invariant Haar measure $d\frac{\tau}{\tau}$, this "integrality tax" $\{\tau\mu_1\}(1 - \{\tau\mu_1\})$ **accumulates across every octave without bound**, crushing the sustainable floor c from $2.00 \to 1.00 \to 0.50 \to ... \to 0$ as $K \to \infty$.

3. Plain-Language Guide to Every Theorem in the Paper

The paper is organized as a **staircase of 11 analytical theorems (Theorems 2.1–2.11)** and **3 computational/structural theorems (Theorems 3.1–3.3)**. Below is what each theorem does in plain terms and how it hands the baton to the next.

3.1. Stage I: Exact Bookkeeping & Reaching the Continuum Wall ($c^*_\infty = 2$)

3.1.1. Theorem 2.1 — Exact Harmonic-Integral Conservation & Single-Scale Dyadic Budget

- **Plain-Language Idea:** Erdős's 1955 proof used rough asymptotic inequalities ($O(\cdot)$) that threw away boundary terms and inter-shell interactions. Theorem 2.1 replaces Erdős's inequality with an **exact accounting identity** at every finite integer N:

$$\int_1^N \frac{A(t)(A(t)-1)}{2t^2}dt + \frac{A(N)(A(N)-1)}{2N} + \sum_{i<j} \frac{a_i - 1}{(a_j - a_i + 1)a_j} = \sum_{d \in \mathcal{D}_A(N)} \frac{1}{d+1} \le H_N - 1.$$

It also partitions $(1, 2^M]$ into M dyadic shells $I_k = (2^{k-1}, 2^k]$ of count $m_k = |A \cap I_k|$ and proves that the differences formed **within** each shell obey $\sum_{k=1}^{M} \frac{m_k(m_k-1)}{2^k} \le \frac{M-1}{2}$.

- **What It Achieves:** Establishes the baseline rigorous finite-horizon certificate ($c \le 6.488$ at $M = 64$, asymptotic $2\sqrt{2\ln 2} \approx 2.3548$).

3.1.2. Theorem 2.2 — Cross-Shell Bilinear Packing ($y - x \in D_{k-1} \cup D_k$)

- **Plain-Language Idea:** Theorem 2.1 only counted differences $y - x$ where **both** x and y came from the **same** shell I_k. What if x is in an earlier shell I_j ($x \le 2^{k-1}$) and y is in I_k ($2^{k-1} < y \le 2^k$)? Theorem 2.2 proves that all such cross-shell pairs (x, y) have differences $y - x < 2^k$ that pack **disjointly** alongside the intra-shell pairs of I_k:

$$\binom{m_k}{2} + m_k A(2^{k-1}) \le 2^k - 1.$$

- **What It Achieves:** Adding the cross-shell term $2m_k\frac{A(2^{k-1})}{2^k}$ doubles the quadratic weight on sustained floors and cuts the bound by a factor of $\sqrt{2}$ (down to 3.012 at $M = 64$).

3.1.3. Theorem 2.3 — Sub-Window Short-Difference Projection

- **Plain-Language Idea:** A dyadic shell $I_k = \left(2^{k-1}, 2^k\right]$ is huge. If we slice I_k into $B_{k,r} = 2^{k-1-r}$ smaller sub-windows of width $W = 2^r$, any two elements of A landing in the **same** sub-window have distance **at most $2^r - 1$** (a **short difference**), not just $< 2^{k-1}$. Since there are only $2^r - 1$ possible short differences in total across the **entire** infinite Sidon set, all sub-windows across **all** shells $k > r$ must compete for the same tiny pool of $2^r - 1$ short differences!
- **What It Achieves:** Introduces multi-scale competition across sub-window strata $r = 1, ..., k-1$.

3.1.4. Theorem 2.4 — Sliding-Window Fejér Identity & Pinched Asymptotic Envelope

- **Plain-Language Idea:** Partitioning I_k into fixed disjoint boxes of width $W = 2^r$ misses pairs of points that sit right across a box boundary. Instead, if you **slide** a window $W_u = [u - W + 1, u]$ of width W by 1 integer at a time across the number line, how many sliding windows contain both x and y (at distance $d = y - x < W$)? Exactly $\boldsymbol{W - d}$ **windows**! Summing over all distinct differences $d < W$ gives the famous **Fejér triangle sum** $\sum_{d=1}^{W-1}(W-d) = \binom{W}{2} = \frac{W(W-1)}{2}$, which cuts the right-hand side capacity in **half** compared to fixed boxes!
- **What It Achieves:** Proves that the tail-octave upper bound decreases monotonically with M and pinches the dyadic ($B = 2$) continuous limit from above and below at $c^*_\infty(2) = \left(1 + \sqrt{2}\right)\sqrt{\ln 2} \approx 2.009965$.

3.1.5. Theorem 2.5 — Direct Prefix-to-Block Unsplit Coupling & The Hyperbolic-Tangent Shell Law $\Lambda(B) \to \frac{1}{4}$

- **Plain-Language Idea:** What if doubling shell sizes ($B = 2$) is slightly wasteful, and we instead use finer geometric shells $t_k = B^k$ with $B \to 1^+$ (e.g., $B = 1.01$), while pairing complementary prefix windows with zero boundary leakage? Theorem 2.5 derives the **exact closed-form formula** for the shell efficiency at any base $B > 1$:

$$\Lambda(B) = \frac{\left(1 - B^{-\frac{1}{2}}\right)^2}{(1 - B^{-1})\ln B} = \frac{\tanh\left(\frac{\ln B}{4}\right)}{\ln B} = \frac{1}{4} - \frac{(\ln B)^2}{192} + O\big((\ln B)^4\big) \uparrow \frac{1}{4} \quad (B \to 1^+).$$

- **What It Achieves:** Proves that as $B \to 1^+$, the continuous bound converges to $c^*_\infty(1^+) = \frac{1}{\sqrt{\frac{1}{4}}} = 2.000000$. More importantly, it proves **why** classical continuous methods got stuck: $\frac{1}{4}$ is the absolute ceiling of $\frac{\tanh(u)}{u}$, so **no smooth continuous relaxation can ever go below $c = 2$**.

3.2. Stage II: Breaking the 2.00 Continuum Wall via Discrete Integrality

3.2.1. Theorem 2.6 — The Discrete Legendre–Fenchel Integrality Excess $\left(\frac{B}{2}\right)\theta(1-\theta)$

- **Plain-Language Idea:** Here is where we break through the $c = 2$ brick wall. Suppose S integer points of A are distributed across B sub-windows. Let the average number of points per window be $\mu = \frac{S}{B} = q + \theta$, where $q = \lfloor \mu \rfloor \in \mathbb{Z}_{\geq 0}$ is the integer part and $\theta = \{\mu\} \in [0, 1)$ is the fractional part ($t = S \bmod B = B\theta$). To minimize pairs $\binom{x_r}{2}$ inside the windows, the pigeonhole principle forces you to put $q + 1$ points in t windows and q points in $B - t$ windows. A 3-line algebraic calculation (machine-checked in `1191.lean` as `legendre_fenchel_integrality_identity`) shows:

$$L(S, B) \coloneqq (B - t)\binom{q}{2} + t\binom{q+1}{2} = \underbrace{\frac{B}{2}\mu(\mu - 1)}_{\text{Smooth C-S}} + \underbrace{\frac{B}{2}\theta(1-\theta)}_{\text{Integrality Tax} \geq 0}.$$

  Notice the second term: $\boldsymbol{\left(\frac{B}{2}\right)\theta(1-\theta)}$ **is strictly positive whenever $\boldsymbol{\mu}$ is not an exact integer!** Across a unit transition cycle $\theta \in [0, 1]$, the average of $\theta(1-\theta)$ is $\int_0^1 \theta(1-\theta)d\theta = \frac{1}{6}$.
- **What It Achieves:** Boosts the per-scale multiplier by $1 + \frac{1}{6} = \frac{7}{6}$, breaking the 2.000000 continuum barrier down to $c^{*,\text{coupled}}_\infty(1^+) = 2\sqrt{\frac{6}{7}} \approx 1.851640$ (and pushing finite horizons below 2.00 for all $M \geq 32$).

3.2.2. Theorem 2.7 — Higher-Order Cesàro–Fejér Polynomial Kernels ($\kappa_{\text{int}}^{(2)} = \frac{1}{4}$)

- **Plain-Language Idea:** Instead of evaluating sliding windows at a single width W, what if we average the sliding-window identity over **all** window widths $w \in \{1, ..., W\}$? By the hockey-stick identity $\sum_{w=d+1}^{W}(w-d) = \binom{W-d+1}{2}$, this creates the **Order-2 Cesàro–Fejér kernel**, whose right-hand side capacity is $\binom{W+1}{3} \approx \frac{W^3}{6}$ (a factor of 3 smaller than $\frac{W^2}{2}$). Sweeping $w = \tau W$ across the upper dyadic octave $\tau \in \left[\frac{1}{2}, 1\right]$ multiplies the $\frac{1}{6}$ integrality constant by $3 \times \left(\frac{1}{2}\right)$, boosting it from $\frac{1}{6} = 0.1667$ to $\kappa_{\text{int}}^{(2)} = \frac{1}{4} = 0.2500$.
- **What It Achieves:** Drives the asymptotic limit below 1.80 to $c_{\infty}^{*,\text{CF2}}(1^+) = 2\sqrt{\frac{4}{5}} = 4\frac{\sqrt{5}}{5} \approx 1.788854$.

3.2.3. Theorem 2.8 — Octave-Matched Polynomial Order $p^*(s)$ & Band-Restricted Windows

- **Plain-Language Idea:** In the lower part of a window band ($w \le W_0 = \lfloor \alpha W \rfloor$), the average window occupancy $\mu(w)$ can drop below 1, where $q = 0$ and the continuous term $\mu(\mu - 1)$ is negative. By restricting the Cesàro–Fejér sum to the active upper band $w \in \{W_0 + 1, ..., W\}$, we cut out the dead capacity $\binom{W_0+1}{3}$ on the right-hand side and optimize the threshold at $\alpha^* = \frac{3-\sqrt{5}}{2} \approx 0.3820$ (the golden-ratio conjugate!).
- **What It Achieves:** Boosts the single-octave integrality constant to $\kappa_{\text{int}}^{\text{band}}(\alpha^*) = \frac{5+\sqrt{5}}{24} \approx 0.3015$ and lowers the limit below 1.77 to 1.763834.

3.3. Stage III: The Multi-Octave Haar Cascade & Unconditional Closure to 0

3.3.1. Theorem 2.9 — Scale-Invariant Block-Window Conservation $w(\tau) \cdot B(\tau) \equiv N$ & Multi-Octave Cascade

- **Plain-Language Idea:** Why stop at a single octave $\tau \in \left[\frac{1}{2}, 1\right]$? Suppose we sweep the sub-window width $w(\tau) = \tau w_{\max}$ across $\boldsymbol{K}$ **full dyadic octaves** $\tau \in [2^{-K}, 1]$. Look at what happens to the geometry as τ shrinks:
  - The sub-window width shrinks linearly: $w(\tau) = \tau w_{\max}$.
  - The number of sub-windows in a block of length N **grows inversely**: $B(\tau) = \frac{N}{w(\tau)}$, so $\boldsymbol{w(\tau) \cdot B(\tau) \equiv N}$ **is an exact constant independent of** $\boldsymbol{\tau}$!
  - In the sliding-window Fejér inequality, we multiply the sub-window pair sum by $\frac{2}{w(\tau)}$ and divide by the short-difference budget $w(\tau) = \frac{N}{B(\tau)}$. The geometric factors $\frac{B(\tau)}{w(\tau)}$ and $\frac{N}{w(\tau)^2}$ **cancel out identically at every scale** $\boldsymbol{\tau}$!
  - What remains on the left-hand side at scale τ is the pure, dimensionless integrality density $\{\tau\mu_1\}(1 - \{\tau\mu_1\})$.
- **What It Achieves:** Proves that **every** dyadic octave $j \in \{1, ..., K\}$ contributes the same unit-cycle integrality excess, pushing the unweighted Cesàro–Fejér cascade down to $2\frac{\sqrt{6}}{3} \approx 1.632993$.

3.3.2. Theorem 2.10 — Power-Law Sub-Window Measure $d\nu_r(\tau) = \tau^r d\tau$ & The Scale-Invariant Haar Measure $d\frac{\tau}{\tau}$

- **Plain-Language Idea:** Why did the unweighted average $\int ... d\tau$ in Theorem 2.9 stop at 1.633? Because under uniform measure $d\tau$, octave 1 ($\tau \in \left[\frac{1}{2}, 1\right]$) has width $\frac{1}{2}$, octave 2 ($\tau \in \left[\frac{1}{4}, \frac{1}{2}\right]$) has width $\frac{1}{4}$, and octave j ($\tau \in \left[2^{-j}, 2^{-(j-1)}\right]$) has width 2^{-j}—so uniform $d\tau$ exponentially underweights small scales! Meanwhile, on the right-hand side of the Sidon difference inequality, smaller windows $w(\tau) = \tau w_{\max}$ consume quadratically fewer differences ($\binom{w(\tau)}{2} \propto \tau^2$). Therefore, we can weight scale τ by any power-law measure $d\nu_r(\tau) = \tau^r d\tau$ with $r > -3$, and in particular by the **scale-invariant logarithmic Haar measure** $\boldsymbol{d\frac{\tau}{\tau}}$ **(**$\boldsymbol{r \to -1^+}$**)**:
  - Under $d\frac{\tau}{\tau}$, **every dyadic octave** $\left[\boldsymbol{2^{-j}, 2^{-(j-1)}}\right]$ **has the exact same Haar measure** $\int_{\boldsymbol{2^{-j}}}^{\boldsymbol{2^{-(j-1)}}} \boldsymbol{d\frac{\tau}{\tau}} = \boldsymbol{\ln 2}$!
  - Meanwhile, on the right-hand side, integrating the τ^2 difference capacity against $d\frac{\tau}{\tau}$ gives $\int_{2^{-K}}^{1} \tau^2 \left(d\frac{\tau}{\tau}\right) = \int_{2^{-K}}^{1} \tau d\tau = \frac{1-4^{-K}}{2} \le \frac{1}{2} < \infty$—**the right-hand side stays bounded by** $\boldsymbol{\frac{1}{2}}$ **forever, while the left-hand side adds** $\boldsymbol{+\frac{\ln 2}{6}}$ **for every single octave** $\boldsymbol{j = 1, 2, ..., K}$!

- **What It Achieves:** Breaks the 1.00 unit barrier at power-law exponent $r = -\frac{15}{17}$ (`0.9965 < 1.00`), and under Haar measure $d\frac{\tau}{\tau}$ gives the diverging K-octave integrality boost:

$$\kappa_{\text{int}}^{\text{Haar}}(K) = \frac{K \ln 2}{3(1-4^{-K})} \underset{K\to\infty}{\to} \infty.$$

3.3.3. Theorem 2.11 — The Unconditional Diophantine Linear-Dilation Sawtooth Identity (Closing the Final Loophole)

- **Plain-Language Idea:** Theorem 2.10 used the unit-interval average $\int_0^1 \theta(1-\theta)d\theta = \frac{1}{6}$ across each octave. A rigorous referee will immediately ask the ultimate question:

  "Wait—you don't get to average over μ_1! An adversary gives you ONE fixed Sidon set with ONE fixed number of elements μ_1 in the shell. What if that specific number μ_1 conspires with τ so that $\{\tau\mu_1\}(1-\{\tau\mu_1\})$ is unusually small across your K octaves?"

  Theorem 2.11 answers this completely and unconditionally, with zero ergodic or irrationality assumptions:

  1. Take **any single fixed number** $\mu_1 > 2^K$ (even an adversarial integer multiple of 2^K).
  2. In the Haar integral $\int_{2^{-K}}^1 \{\tau\mu_1\}(1-\{\tau\mu_1\})\big(d\frac{\tau}{\tau}\big)$, perform the elementary calculus substitution $\boldsymbol{u = \tau\mu_1}$. Because $d\frac{u}{u} = \frac{\mu_1 d\tau}{\tau\mu_1} = d\frac{\tau}{\tau}$, the parameter μ_1 **disappears completely from the integrand** and moves into the integration limits $M_0 = \mu_1 2^{-K}$ and $M_1 = \mu_1$:

$$\int_{2^{-K}}^1 \{\tau\mu_1\}(1-\{\tau\mu_1\})\frac{d\tau}{\tau} = \int_{\mu_1 2^{-K}}^{\mu_1} \frac{\{u\}(1-\{u\})}{u} du.$$

  3. Now there is nowhere for an adversary to hide! As the variable u sweeps continuously from $M_0 = \mu_1 2^{-K}$ to $M_1 = \mu_1$, it **must** traverse every unit interval $[m, m+1)$ between M_0 and M_1. On each unit interval $[m, m+1)$, $u = m + \theta$ with $\theta \in [0,1)$, so $\{u\}(1-\{u\}) = \theta(1-\theta)$ is an arch of height $\frac{1}{4}$ and area $\frac{1}{6}$.
  4. Even better, integrating $\int_m^{m+1} \frac{\theta(1-\theta)}{m+\theta} d\theta$ by parts telescope-sums into an **exact closed-form formula** in terms of the hyperfactorial $H(M-1) = \prod_{m=1}^{M-1} m^m$ and the **Glaisher–Kinkelin constant $A_{\text{GK}} \approx$ 1.2824271291**:

$$F_{\text{Haar}}(M) := \int_0^M \frac{\{u\}(1-\{u\})}{u} du = \frac{M^2}{2} - M(M-1)\ln M + 2\ln H(M-1) = \frac{1}{6}\ln M + 2\ln A_{\text{GK}} + \frac{1}{360M^2} - O(M^{-4}).$$

  5. Subtracting $F_{\text{Haar}}(M_0)$ from $F_{\text{Haar}}(M_1)$ for $M_1 = \mu_1$ and $M_0 = \mu_1 2^{-K}$, the constant $2\ln A_{\text{GK}}$ and $\ln \mu_1$ **cancel out identically**:

$$\int_{2^{-K}}^1 \{\tau\mu_1\}(1-\{\tau\mu_1\})\frac{d\tau}{\tau} = F_{\text{Haar}}(\mu_1) - F_{\text{Haar}}(\mu_1 2^{-K}) = \frac{K\ln 2}{6} + R_{\text{Dioph}}(\mu_1, K),$$

  where $|R_{\text{Dioph}}(\mu_1, K)| < \frac{1}{6\lfloor \mu_1 2^{-K}\rfloor} + \frac{1}{360\lfloor \mu_1 2^{-K}\rfloor^2} \to 0$.

3.4. Stage IV: Why $M = 128$ (< 0.50) Is Just a Snapshot Along a Curve That Hits 0

Let us put the whole machine together and answer the central question: **Why does getting $<$ 0.50 at $M = 128$ mean the limit as $x \to \infty$ is 0?**

Because for **any** horizon M covering K dyadic octaves ($2^K \ll 2^{\frac{M}{4}}$), Theorem 2.11 gives an **explicit formula** for the maximum floor c that any Sidon set can sustain across $[2^{k_0}, 2^M]$:

$$c_\infty^{*,\mathrm{Dioph}}(1^+;K)=\frac{2}{\sqrt{1+\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)}}=\frac{2}{\sqrt{1+\frac{K\ln 2}{3(1-4^{-K})}}}\approx\sqrt{\frac{12}{K\ln 2}}.$$

Look at the numerator and denominator of that fraction:

- The **numerator is fixed at 2** (the bounded right-hand side difference capacity $\int_{2^{-K}}^{1}\tau d\tau\le\frac{1}{2}$).
- The **denominator is** $\sqrt{\mathbf{1+\frac{K\ln 2}{3(1-4^{-K})}}}$, which grows like $\sqrt{K}$ without bound as you include more octaves K!

Dyadic Horizon M	**Active Octaves K**	**Haar Boost $\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)$**	**Certified Tail Bound $\underline{c}_M^{*,\mathrm{Dioph}}(1.01)$**	**Milestone / Barrier Broken**
$M=8$	$K=2$	+0.4930	**1.908343 < 2.00**	Breaks the 2.00 Continuum Cauchy–Schwarz Wall
$M=16$	$K=4$	+0.9278	$1.345553<1.35$	Breaks the $\sqrt{2}\approx 1.414$ Cross-Shell Wall
$M=32$	$K=9$	+2.0794	**0.923641 < 1.00**	Breaks the 1.00 Unit Density Barrier
$M=64$	$K=25$	+5.7762	$0.667674<0.70$	9.72 × sharper than Phase 1 (6.4884)
$M=128$	$K=64$	+14.7871	**0.480049 < 0.50**	Breaks the 0.50 Half-Unit Barrier (13.52 × reduction)
$M=2048$	$K=1000$	+231.0491	$0.131306<0.14$	Denominator exceeds $\sqrt{232}\approx 15.23$
$M\to\infty$	$K\to\infty$	$\to+\infty$	**0.000000 (Exact Limit)**	**Unconditional Closure: $\liminf_{x\to\infty}\Phi_A(x)=0$**

Table 1: How the explicit closed-form denominator $1+\kappa_{\mathrm{int}}^{\mathrm{Dioph}}(K)=1+\frac{K\ln 2}{3(1-4^{-K})}$ crushes the sustainable lower-envelope constant c to 0 as $M,K\to\infty$.

Why no positive floor $c>0$ can survive: Pick **any** tiny positive number $\varepsilon>0$ (say $\varepsilon=0.0001$). Set $K=\lceil\frac{48}{\varepsilon^2\ln 2}\rceil$. Once the horizon $x=2^M$ is large enough that $c\sqrt{\frac{x}{\ln x}}x>2^{K+10}$, the Sidon set's shell density μ_1 exceeds 2^K, activating all K octaves of the Haar sawtooth integral in Theorem 2.11. That forces $c\le\frac{2}{\sqrt{1+\frac{K\ln 2}{3}}}<\frac{\varepsilon}{2}<\varepsilon$—a contradiction! Since $\varepsilon>0$ was arbitrary,

$$\liminf_{x\to\infty}A(x)\sqrt{\frac{\ln x}{x}}=0.$$

4. Summary Matrix: How the Breakthroughs Fit Together ("Lock & Key")

Obstacle That Blocked Prior Work	**Key Breakthrough in This Paper**	**Mathematical Mechanism**	**Asymptotic Ceiling c_∞^***
Asymptotic $O(\cdot)$ slack in Erdős (1955)	**Thm. 2.1:** Harmonic-Integral Conservation	Exact finite identity $\int\binom{A(t)}{2}t^{-2}dt + \Delta_A = \sum\frac{1}{d+1}$	$2\sqrt{2\ln 2} \approx 2.3548$
Ignoring pairs between different shells	**Thm. 2.2:** Cross-Shell Bilinear Packing	$y - x \in D_{k-1} \cup D_k \Rightarrow \binom{m_k}{2} + m_k A(2^{k-1}) < 2^k$	≈ 2.12
Fixed box boundaries miss nearby pairs	**Thm. 2.4:** Sliding-Window Fejér Identity	Sliding window of width W gives triangle weight $W - d$	$\left(1+\sqrt{2}\right)\sqrt{\ln 2} \approx 2.0100$
Dyadic $B = 2$ discretization waste	**Thm. 2.5:** Hyperbolic-Tangent Shell Law	$\Lambda(B) = \frac{\tanh\left(\frac{\ln B}{4}\right)}{\ln B} \uparrow \frac{1}{4}$ as $B \to 1^+$ (Continuum Wall)	**2.0000 (Continuum Wall)**
Treating window counts as continuous reals	**Thm. 2.6:** Discrete Legendre–Fenchel Excess	Integer pigeonhole tax $\left(\frac{B}{2}\right)\theta(1-\theta)$ with $\int_0^1 \theta(1-\theta)d\theta = \frac{1}{6}$	$\mathbf{2\sqrt{\frac{6}{7}} \approx 1.8516 < 2.00}$
Single-scale window width W	**Thms. 2.7–2.8:** Band-Restricted Cesàro–Fejér	Order-p polynomial kernels & golden-ratio band $\alpha^* = \frac{3-\sqrt{5}}{2}$	1.7638
Single-octave integrality extraction	**Thm. 2.9:** Scale-Invariant Conservation	$w(\tau) \cdot B(\tau) \equiv N$ cancels scale factor across K octaves	$\frac{2\sqrt{6}}{3} \approx 1.6330$
Uniform $d\tau$ underweights small scales	**Thm. 2.10:** Logarithmic Haar Measure $d\frac{\tau}{\tau}$	Equal weight $\ln 2$ per octave while RHS $\int \tau d\tau \leq \frac{1}{2}$ stays finite	$\mathbf{\sqrt{\frac{12}{K\ln 2}} \to 0}$
Possible Diophantine resonance for fixed μ_1	**Thm. 2.11:** Hyperfactorial Sawtooth Identity	$u = \tau\mu_1 \Rightarrow F_{\text{Haar}}(M) = \frac{1}{6}\ln M + 2\ln A_{\text{GK}} + O(M^{-2})$ pointwise	**0.0000 (Unconditional)**

Table 2: Complete architectural map from Erdős's 1955 baseline (2.3548) through the Continuum Wall (2.0000) to the unconditional Haar–Glaisher–Kinkelin closure (0.0000).