

ISO TC 20/SC 14 N 188

Date: 2006-05-17

ISO/CD 22009.8

ISO TC 20/SC 14/WG 4

Secretariat: ANSI

Space Environment (Natural and Artificial) — Model of the Earth's magnetospheric magnetic field

Warning

This document is not an ISO International Standard. It is distributed for review and comment. It is subject to change without notice and may not be referred to as an International Standard.

Recipients of this document are invited to submit, with their comments, notification of any relevant patent rights of which they are aware and to provide supporting documentation.

Document type: International Standard
Document subtype:
Document stage: (30) Committee
Document language: E

Copyright notice

This ISO document is a working draft or committee draft and is copyright-protected by ISO. While the reproduction of working drafts or committee drafts in any form for use by participants in the ISO standards development process is permitted without prior permission from ISO, neither this document nor any extract from it may be reproduced, stored or transmitted in any form for any other purpose without prior written permission from ISO.

Requests for permission to reproduce this document for the purpose of selling it should be addressed as shown below or to ISO's member body in the country of the requester:

[Indicate the full address, telephone number, fax number, telex number, and electronic mail address, as appropriate, of the Copyright Manager of the ISO member body responsible for the secretariat of the TC or SC within the framework of which the working document has been prepared.]

Reproduction for sales purposes may be subject to royalty payments or a licensing agreement.

Violators may be prosecuted.

Contents	Page
1 Scope	1
2 Terms and definitions	1
2.1 Internal magnetic field	1
2.2 External magnetic field	1
2.3 Magnetospheric sources of magnetic field	2
2.4 Geomagnetic dipole tilt angle	2
2.5 Solar-magnetospheric coordinates	2
2.6 Magnetopause stand-off distance	2
3 General concepts and assumptions	2
3.1 Magnetic field induction in the Earth's magnetosphere	2
3.2 Magnetospheric magnetic field standardisation: process based approach	2
4 Model requirements	3
4.1 The magnetospheric magnetic field sources	3
4.2 Parameterization	3
4.3 Magnetospheric dynamics	3
4.4 Model testing and comparison with measurements	3
5 List of criteria	3
Annex A (normative) Paraboloid model of the magnetospheric magnetic field: Calculation of Induction of the Magnetic Field of the Magnetospheric Currents	4
A.1 Paraboloid model of the magnetic field of magnetospheric currents	4
A.1.1 Parameters	4
A.1.2 Submodels	5
A.2 Magnetic field of the magnetopause currents screening the geomagnetic dipole	5
A.3 Magnetic field of the tail current system	6
A.4 Ring current magnetic field	6
A.5 Magnetic field of the magnetopause currents screening the ring current	7
A.6 Magnetic field of field-aligned currents	7
Annex B (normative) Submodels	9
B.1 Submodels: Calculation of the Main Parameters of Magnetospheric Current Systems	9
B.1.1 The tilt angle of geomagnetic dipole	9
B.1.2 The distance from the Earth to the subsolar point on the magnetopause	9
B.1.3 The distance to the earthward edge of the geomagnetic tail current sheet	9
B.1.4 Magnetic flux through the magnetotail lobes	10
B.1.5 The ring current magnetic field at the Earth's centre	10
B.1.6 The total Region 1 field-aligned current	10
Annex C (informative)	12
C.1 Spherical coordinates (1)	12
C.2 Spherical coordinates (2)	12
C.3 Parabolic coordinates	12
C.4 Magnetic Field of Geomagnetic Dipole	13
C.5 Dataset for model testing	13

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 22009 was prepared by Technical Committee ISO/TC 20, *Aircraft and space vehicles*, Subcommittee SC 14, *Space systems and operations*.

This second/third/... edition cancels and replaces the first/second/... edition (ISO 4:2005, ISO 22009:2006, ISO 22009:2006), [clause(s) / subclause(s) / table(s) / figure(s) / annex(es)] of which [has / have] been technically revised.

Introduction

This standard describes the main requirements to the Earth's magnetospheric magnetic field model. The model satisfying the set of requirements is described in the Annex as working example. The model can be used in scientific and engineering applications and is intended to calculate the magnetic induction field generated from a variety of current systems located on the boundaries and within the boundaries of the Earth's magnetosphere under a wide range of environmental conditions, quiet and disturbed, affected by Solar-Terrestrial interactions simulated by Solar activity such as Solar Flares and related phenomena which induce terrestrial magnetic disturbances such as Magnetic Storms.

Space Environment (Natural and Artificial). Model of Earth's Magnetic Field.

1 Scope

Standard describes the main magnetospheric large-scale current systems and the magnetic field in the Earth's magnetosphere and provides the main requirements to the model of the magnetospheric magnetic field. The standard gives in Annex the working example of the model and establishes the parameters of magnetospheric large-scale current systems which are changing in accordance with conditions in the space environment. Standard can be used to develop the new models of magnetospheric magnetic field. Such models are useful in investigate physical processes in the Earth's magnetosphere as well as in calculations, developing, testing and estimating the results of exploitation of spacecrafts and other equipment operating in the space environment.

The main goals of standardisation of the Earth's magnetospheric magnetic field are:

- providing the unambiguous presentation of the magnetic field in the Earth's magnetosphere;
- providing compatibility of results of interpretation and analysis of space experiments;
- providing less labour-consuming character of calculations of the magnetic field of magnetospheric currents in the space at geocentric distances of 1-6.6 Earth's radii (R_E);
- providing the most reliable calculations of all elements of the geomagnetic field in the space environment.

Magnetic field model presented in the Annex of this standard can be used to forecast radiation situation in the space, including the periods of intense magnetic disturbances (magnetic storms) when developing systems of spacecraft magnetic orientation, when forecasting the influence of magnetic disturbances on transcontinental piping and power transmission lines.

2 Terms and definitions

2.1 Internal magnetic field

Internal (main) magnetic field is the magnetic field produced by the sources inside the Earth's core. It can be presented in the form of a series of spherical harmonic functions. The expansion coefficients (IGRF model) undergo very slight changes in time. The International Association of Geomagnetism and Aeronomy (IAGA) is responsible for IGRF model development and modifications and approves its coefficients every 5 years. Internal magnetic field is not the subject of this Standard.

2.2 External magnetic field

External (magnetospheric) magnetic field is the magnetic field produced by magnetospheric sources of magnetic field.

2.3 Magnetospheric sources of magnetic field

Magnetospheric sources of magnetic field are

- Currents flowing over the magnetopause and screening the geomagnetic dipole magnetic field.
- Currents flowing inside the Earth's magnetosphere:
 - tail current, produced by currents across the geomagnetic tail and closure currents on the magnetopause;
 - ring current, including symmetrical ring current, circling around the Earth and carried by trapped particles and partial ring current, produced by currents flowing inside the region of symmetrical ring current, mostly in the pre-midnight sector of equatorial plane, closed by field-aligned and ionospheric currents;
 - Field-aligned currents, produced by currents flowing along the auroral magnetic field lines, closed by currents on the magnetopause and on the ionospheric surface.
- Currents flowing over the magnetopause and screening the ring current and partial ring current magnetic fields.

2.4 Geomagnetic dipole tilt angle

Geomagnetic dipole tilt angle is the angle of inclination of the geomagnetic dipole to the plane orthogonal to the Earth-Sun line.

2.5 Solar-magnetospheric coordinates

Solar-magnetospheric (GSM) coordinates are the Cartesian geocentric coordinates, where X-axis is directed to the Sun, Z-axis lies in the one plane with OX axis and geomagnetic dipole axis and Y-axis supplements the X and Z axes to the right-hand system.

2.6 Magnetopause stand-off distance

Magnetopause stand-off distance is the geocentric distance to the subsolar point on the magnetopause.

3 General concepts and assumptions

3.1 Magnetic field induction in the Earth's magnetosphere

Vector of magnetic field induction \vec{B}_M in the Earth's magnetosphere is calculated by the formula

$$\vec{B}_M = \vec{B}_1 + \vec{B}_2, \text{ nT}, \quad (1)$$

where \vec{B}_1 is the vector of induction of the internal magnetic field, \vec{B}_2 is the vector of induction of the external magnetospheric magnetic field. The magnetic field of the magnetospheric currents (external magnetic field), \vec{B}_2 , is calculated in terms of the quantitative model of the magnetosphere.

3.2 Magnetospheric magnetic field standardisation: process based approach

The magnetospheric magnetic field standard does not specify a single magnetospheric model, theoretical or empirical. In order to encourage continual improvements in magnetospheric modelling, this standard is process-

based standard for determining the magnetospheric magnetic field. Magnetospheric magnetic field model, after its development, may satisfy the requirements in Section 4 and the list of criteria presented in Section 5. The working example of the model is presented in the Annex A and must be reconsidered every 5 years on the base of competitions of the candidate models. The current working example is presented in the Annex A.

4 Model requirements

- The model of the magnetic field of magnetospheric currents (referred to below as "model") presents the vector of induction of magnetospheric currents in solar-magnetospheric coordinates.
- The model describes a regular part of the magnetic field in the region from $1 R_E$ to $6.6 R_E$.
- The model reflects compression of the Earth's magnetosphere in the dayside due to interaction with the solar wind, day-night asymmetry (the field on the nightside is weakened), day and season variations.
- The model takes into account geomagnetic dipole tilt angle, varying in the range from -35° to $+35^\circ$.

4.1 The magnetospheric magnetic field sources

The standardised magnetospheric magnetic field is produced by currents described in Section 2.3.

4.2 Parameterization

Each magnetospheric source of magnetic field depends on parameters which are calculated from empirical data.

4.3 Magnetospheric dynamics

The magnetospheric dynamics is determined to be a sequence of its instant states.

4.4 Model testing and comparison with measurements

The model testing is carrying out with the help of databases which include the spacecraft based and on-ground measurements. The dataset used for the model testing is presented in the Annex C.

5 List of criteria

The compliance criteria for this standard consist from the activities common for any candidate magnetospheric magnetic field model. These criteria specify the compliance process which includes the model documenting, publishing and testing.

- The candidate model is required to include the statement of the modeling approach used (empirical or theoretical model). The empirical models are required to include a clear specification of the input data used to derive the model and where the these data were measured. Theoretical models are required to include a description of the physical principles and approaches that are used as the basis of the model.
- The statement about the candidate model area of application and domain applicability should be included.
- The statement about the rms errors during the model calculations comparison with observational data obtained from measurements should be included.
- The description and implementation of the magnetospheric magnetic field model should be published in an internationally-accessible refereed journals.

Annex A (normative)

Paraboloid model of the magnetospheric magnetic field: Calculation of Induction of the Magnetic Field of the Magnetospheric Currents

The magnetospheric magnetic field calculated by paraboloid model is the solution of the magnetostatic problem inside the paraboloid of revolution.

A.1 Paraboloid model of the magnetic field of magnetospheric currents

Vector of induction of the magnetic field of magnetospheric currents is calculated by formula

$$\vec{B}_2 = \vec{B}_{sd}(\psi, R_1) + \vec{B}_t(\psi, R_1, R_2, \Phi_\infty) + \vec{B}_r(\psi, b_t) + \vec{B}_{sr}(\psi, R_1, b_r) + \vec{B}_{fac}(I_0) . \quad (2)$$

Here

- \vec{B}_{sd} is the magnetic field of currents on the magnetopause screening the dipole field;
- \vec{B}_t is the magnetic field of the magnetospheric tail;
- \vec{B}_r is the magnetic field of the ring current;
- \vec{B}_{sr} is the magnetic field of currents on the magnetopause, screening the ring current field;
- \vec{B}_{fac} is the magnetic field of Region 1 field-aligned currents.

The components of the magnetic field of magnetospheric currents, \vec{B}_{sd} , \vec{B}_t , \vec{B}_r , \vec{B}_{sr} , \vec{B}_{fac} are calculated separately in terms of the paraboloid model of the magnetosphere in the form of series in the Bessel functions or Legendre polynomials.

A.1.1 Parameters

The components of the magnetic field of magnetospheric currents, \vec{B}_{sd} , \vec{B}_t , \vec{B}_r , \vec{B}_{sr} , \vec{B}_{fac} are determined by the values of parameters of the magnetospheric current systems:

- ψ is the geomagnetic dipole tilt angle, degrees;
- \vec{R}_1 is the distance to the subsolar point at the magnetopause, R_E ;
- \vec{R}_2 is the distance to the earthward edge of the magnetospheric tail current sheet, R_E ;
- Φ_∞ is the magnetic flux in the tail lobes, defining the current intensity in the magnetotail, Wb;

- b_r is the intensity of the ring current magnetic field at the Earth's centre, nT;
- I_0 being for total Region 1 field-aligned currents intensity, MA.

A.1.2 Submodels

The instant values of the parameters of the magnetospheric current systems, ψ , \vec{R}_1 , \vec{R}_2 , Φ_∞ , b_r , I_0 are determined using a limited set of empirical data in terms of the so-called submodels (see Annex B)..

A.2 Magnetic field of the magnetopause currents screening the geomagnetic dipole

The, \vec{B}_{sd} , is calculated as

$$\vec{B}_{sd} = -\nabla U_{sd},$$

where the scalar potential U_{sd} of the magnetic field of magnetopause currents is presented in spherical coordinates R, θ, φ (see Annex C.1):

$$U_{sd} = -\frac{M_E}{R_1^2} \sum_{n=1}^{\infty} \left(\frac{R}{R_1} \right)^n \left[d_n^{\parallel} \sin \psi P_n(\cos \theta) + d_n^{\perp} \cos \psi P_n^1(\cos \theta) \right] \quad (3)$$

$$P_n = (2^n n!)^{-1} \cdot (d^n (x^2 - 1)^n / dx^n), \quad P_n^1(x) = \sqrt{1 - x^2} \cdot (dP_n / dx).$$

$M_E = B_0 \cdot R_E^3$ is the magnetic moment of the geomagnetic dipole, B_0 is the magnetic field at the geomagnetic equator of the Earth. The first six dimensionless coefficients d_n^{\parallel} and d_n^{\perp} are listed in Table 1.

Table A.1 — Expansion coefficients for the scalar potential of the magnetic field of magnetopause currents

n	d_n^{\perp}	d_n^{\parallel}
1	0.6497	0.9403
2	0.2165	0.4650
3	0.0434	0.1293
4	-0.0008	-0.0148
5	-0.0049	-0.0160
6	-0.0022	-0.0225

A.3 Magnetic field of the tail current system

The magnetic field of the tail current system \vec{B}_t is calculated from the equation

$$\vec{B}_t = -\nabla U_t + \vec{B}_{t_{in}} \quad (4)$$

Where U_t is determined by the series

$$U_t = b_t R_1 \left\{ \begin{aligned} & \sum_{k,n=1}^{\infty} (b_{nk} + c_{nk} K'_n(\lambda_{nk} \alpha_0) \lambda_{nk}) \cos n\varphi \cdot J_n(\lambda_{nk} \beta) I_n(\lambda_{nk} \alpha) \quad \text{for } \alpha < \alpha_0 \\ & \beta_t \alpha_0 \ln \alpha \cdot \text{sign}\left(\frac{\pi}{2} - |\varphi|\right) + \\ & \sum_{nk} c_{nk} \cos n\varphi \cdot I'_n(\lambda_{nk} \alpha_0) \lambda_{nk} J_n(\lambda_{nk} \beta) K_n(\lambda_{nk} \alpha) \quad \text{for } \alpha \geq \alpha_0 \end{aligned} \right.$$

$$\text{here } c_{nk} = b_{nk} \lambda_{nk} I'_n(\lambda_{nk} \alpha_0), \quad b_{nk} = \frac{2\lambda_{nk} \int_0^1 \int_{-\pi}^{\pi} J_n(\lambda_{nk} \beta) f(\beta, \varphi) \cos n\varphi d\varphi d\beta}{\pi(\lambda_{nk}^2 - n^2) J_n^2(\lambda_{nk}) I'_n(\lambda_{nk} \alpha_0)}.$$

$$f(\beta, \varphi) = \begin{cases} \frac{\alpha_0}{\beta_t} \beta \cos \varphi, & \text{for } \alpha_0 \beta \cos \varphi < \beta_t \\ \text{sign}\left(\frac{\pi}{2} - |\varphi|\right), & \text{for } \alpha_0 \beta \cos \varphi \geq \beta_t \end{cases},$$

where λ_{nk} are zeros of the $J' = 0$ equation, $\alpha_0 = \sqrt{1 - 2R_2/R_1}$ is the parabolic α -coordinate of the inner edge of the tail current sheet, $\beta_t = \frac{d}{R_1}$, d is the half thickness of the current sheet, $b_t = \frac{2\Phi_{\infty}}{\pi R_1^2} \sqrt{R_1/(2R_2 + R_1)}$, is a magnetic field in the tail lobe at the inner edge of the tail current sheet.

The magnetic field inside the current sheet, $B_{t_{in}}$, is calculated from the relations

$$B_{t_{in}\alpha} = b_t \frac{\alpha_0}{\alpha} \frac{\beta}{\beta_t} \frac{\cos \varphi}{\sqrt{\alpha^2 + \beta^2}}, \quad B_{t_{in}\beta} = 0, \quad B_{t_{in}\varphi} = 0.$$

Description of the paraboloid coordinates is presented in the Annex B.3.

A.4 Ring current magnetic field

The ring current magnetic field \vec{B}_r is determined by the expressions

$$\vec{B}_r = \frac{M_R}{M_E} \cdot \begin{cases} \left(\frac{R}{R_{rc}} \right)^5 \cdot \vec{B}_d + 2B_0 \frac{R_E^3}{R_2^3} \left(\frac{R_2^5}{R_{rc}^5} - 1 \right) \vec{e}_z & \text{for } 0 \leq R \leq R_2 \\ \vec{B}_d & \text{for } R \geq R_2 \end{cases} \quad (5)$$

where $R_{rc} = \sqrt{0.5(R^2 + R_2^2)}$, $M_R = 0.5b_r \cdot R_2^3 / (4\sqrt{2} - 1)$ is the magnetic moment of the ring current, \vec{B}_d is the magnetic field of the geomagnetic dipole, \vec{e}_z being a unite vector directed oppositely to the geomagnetic dipole.

Expressions for \vec{B}_d and \vec{e}_z in the solar-magnetospheric coordinates are presented in Annex C.4.

A.5 Magnetic field of the magnetopause currents screening the ring current

The magnetic field of the magnetopause currents screening the ring current \vec{B}_{sr} is calculated from the equation

$$\vec{B}_{sr} = -\nabla U_{sr},$$

where the scalar potential U_{sr} of the magnetic field of magnetospheric currents presented in spherical coordinates R, θ, φ (see Annex C.2) reads

$$U_{sr} = -\frac{M_R}{R_1^2} \sum_1^\infty \left(\frac{R}{R_1} \right)^n \left[d_n^{\parallel} \sin \psi P_n(\cos \theta) + d_n^{\perp} \cos \psi P_n^1(\cos \theta) \right] \quad (6)$$

Coefficients d_n^{\parallel} and d_n^{\perp} are listed in Table 1.

A.6 Magnetic field of field-aligned currents

The magnetic field of field-aligned currents \vec{B}_{fac} is calculated from the equation

$$\vec{B}_{fac} = \text{curl} \vec{A}_{fac}$$

where the vector potential \vec{A}_{fac} of the magnetic field of field-aligned currents is presented in spherical coordinates R, θ, φ with polar axis directed opposite the Earth's dipole (see Annex B.2):

$$\vec{A}_{fac} = \frac{\mu_0 I_0 \sin \varphi}{2(1 + \cos \theta_m)} \begin{cases} \frac{\tan(\theta/2)}{\tan(\theta_m/2)} & \text{for } 0 \leq \theta \leq \theta_m \\ \frac{\sin \theta_m}{\sin \theta} & \text{for } \theta_m \leq \theta \leq \pi - \theta_m \\ \frac{\cot(\theta/2)}{\tan(\theta_m/2)} & \text{for } \pi - \theta_m \leq \theta \leq \pi \end{cases}$$

θ_m - is polar cap radius in radians: $\sin^2 \theta_m = 3.9 \cdot \Phi_\infty [\text{MWb}] / |B_0 [\text{nT}]|$.

A.7 References

- 1) Alexeev, I. I. and Feldstein, Y. I.: Modelling of geomagnetic field during magnetic storms and comparison with observations, *J. Atmos. Sol.-Terr. Phys.*, 63, 331–340, 2001.
- 2) Alexeev I. I., Kalegaev V. V., Belenkaya E. S., Bobrovnikov S. Y., Feldstein Y. I., and Gromova L. I.: Dynamical model of the magnetosphere: case study for 9–12 January 1997, *J. Geophys. Res.*, 106, 25 638–25 693, 2001.
- 3) Alexeev I.I., E.S.Belenkaya, S.Y.Bobrovnikov, V.V.Kalegaev, Modelling of the electromagnetic field in the interplanetary space and in the Earth's magnetosphere, *Space Science Review*, 107, N1/2, 7-26, 2003.
- 4) E. S. Belenkaya, I. I. Alexeev, and C. R. Clauer Jr, Field-aligned current distribution in the transition current system, *J. Geophys. Res.*, 109, A11207, doi:10.1029/2004JA010484, 2004.

Annex B (normative)

Submodels

B.1 Submodels: Calculation of the Main Parameters of Magnetospheric Current Systems

In the paraboloid model of the magnetosphere the values of parameters of the magnetospheric current systems are calculated using submodels. The submodels represent empirical relations or auxiliary models to relate parameters of the magnetospheric current systems to the measured data. While the magnetic field dependence on parameters is fixed, the parameters dependence on empirical data can be changed by model's user. Below are the simple submodels allowing to calculate the model input parameters. These submodels are not the standardisation objects.

B.1.1 The tilt angle of geomagnetic dipole

The tilt angle of geomagnetic dipole, ψ , is calculated by the formula

$$\sin \psi = -\sin \beta \cos \alpha_1 + \cos \beta \sin \alpha_1 \cos \varphi_m, \quad (7)$$

where

- $\alpha_1 = 11.43^\circ$ is the angle between the Earth's axis and the geomagnetic dipole moment,
- β is the Sun's deflection ($\sin \beta = \sin \alpha_2 \cos \varphi_{se}$),
- $\alpha_2 = 23.5^\circ$ is the angle between the Earth's axis and the normal to the ecliptic plane,
- $\varphi_{se} = 0.9856263(172 - I_{day})$ is angle between Earth-Sun line and the projection of the Earth's axis at the ecliptic plane,
- I_{day} is the number of the day in a year,
- $\varphi_m = UT \cdot 15^\circ - 69.76^\circ$ is the angle between the midnight geographic meridian plane and northern magnetic pole meridian plane,
- UT is the universal time in hours.

B.1.2 The distance from the Earth to the subsolar point on the magnetopause

The geocentric distance R_1 to the subsolar point is calculated using solar wind data: solar wind dynamical pressure and IMF B_z component [Shue et al., 1998]:

$$R_1 = \{10.22 + 1.29 \tanh[0.184(B_z + 8.14)]\} (nv^2)^{-\frac{1}{6.6}}.$$

Here B_z is IMF z-component in nT, n, v are solar wind concentration and velocity in cm^{-3} and km/s, respectively.

B.1.3 The distance to the earthward edge of the geomagnetic tail current sheet

The distance to the earthward edge of the geomagnetic tail current sheet, R_2 , is calculated by the formula

$$R_2 = 1 / \cos^2 \varphi_k, \quad (9)$$

where R_2 is expressed in R_E , and φ_k is the latitude of the equatorward boundary of the auroral oval at midnight.

B.1.4 Magnetic flux through the magnetotail lobes

Magnetic flux through the magnetotail lobes, Φ_∞ , is calculated by the formula

$$\Phi_\infty = \Phi_0 + \Phi_s \quad (10)$$

where Φ_0 is the magnetic flux in the magnetotail during quiet periods, Φ_s being the time-dependent magnetic flux in the lobes associated with intensification of the magnetotail current system during disturbances

$$\begin{aligned} \Phi_0 &= 3.7 \cdot 10^8 \text{ Wb} \\ \Phi_s &= -AL \frac{\pi R_1^2}{14} \sqrt{\frac{2R_2}{R_1} + 1}, \end{aligned} \quad (11)$$

where AL is the auroral index of geomagnetic activity.

B.1.5 The ring current magnetic field at the Earth's centre

The ring current intensity is characterised by the value of ring current magnetic field at the Earth's centre, which is calculated by the Dessler-Parker-Scopke relation

$$b_r = -\frac{2}{3} B_0 \frac{\varepsilon_r}{\varepsilon_d}, \quad (12)$$

where ε_r is the total energy of ring current particles, $\varepsilon_d = \frac{1}{3} B_0 M_E$ is the geomagnetic dipole energy.

B.1.6 The total Region 1 field-aligned current

The total Region 1 field-aligned current intensity is calculated by the formula

$$I_0 = 2 \sqrt{\frac{v}{400}} \cdot \left(\frac{5}{n}\right)^{1/8} \cdot \begin{cases} 0.327744 & \text{for } b_z > -1.6nT \\ -1.017 \cdot \frac{b_z}{5} & \text{for } b_z \leq -1.6nT \end{cases} \quad (13)$$

where b_z is IMF north-south component, nT, solar wind velocity v and proton concentration n are the same as in A.1.2.

B.2 References

- 1) Alexeev, I. I. and Feldstein, Y. I.: Modelling of geomagnetic field during magnetic storms and comparison with observations, *J. Atmos. Sol.-Terr. Phys.*, 63, 331–340, 2001.
- 2) Alexeev I. I., Kalegaev V. V., Belenkaya E. S., Bobrovnikov S. Y., Feldstein Y. I., and Gromova L. I.: Dynamical model of the magnetosphere: case study for 9–12 January 1997, *J. Geophys. Res.*, 106, 25 638–25 693, 2001.
- 3) Alexeev I.I., E.S.Belenkaya, S.Y.Bobrovnikov, V.V.Kalegaev, Modelling of the electromagnetic field in the interplanetary space and in the Earth's magnetosphere, *Space Science Review*, 107, N1/2, 7-26, 2003.
- 4) E. S. Belenkaya, I. I. Alexeev, and C. R. Clauer Jr, Field-aligned current distribution in the transition current system, *J. Geophys. Res.*, 109, A11207, doi:10.1029/2004JA010484, 2004.
- 5) Feldstein, Y.I., A.I. Levitin, J.U. Kozyra, B.T. Tsurutani, A. Prigancova, L. Alperovich, W.D. Gonzalez, U. Mall, I.I. Alekseev, L.I. Gromova, and L.A. Dremukhina, Self-consistent modeling of the large-scale distortions in the geomagnetic field during the 24–27 September 1998 major magnetic storm, *J. Geophys. Res.*, 110, A11214, doi:10.1029/2004JA010584, 2005.
- 6) V.V. Kalegaev, N. Yu. Ganushkina, T. I. Pulkkinen, M. V. Kubyshkina, H. J. Singer, C. T. Russell, Relation between the ring current and the tail current during magnetic storms. *Annales Geophysicae*, V. 23, No. 2, p. 523-533, 2005.
- 7) Shue J.-H., Song P., Russel C. T., Steinberg J. T., Chao J. K., Zastenker G., Vaisberg O. L., Kokubun S., Singer H. S., Detman T. R., Kawano H., Magnetopause location under extreme solar wind conditions, *J. Geophys. Res.*, 103, 17691-17700, 1998.

Annex C (informative)

C.1 Spherical coordinates (1)

Spherical coordinates R, θ, φ with the polar axis plotted along the Sun-Earth axis are determined by the expressions

$$\begin{aligned} x / R_1 &= R \cos \theta \\ y / R_1 &= R \sin \theta \sin \varphi \\ z / R_1 &= R \sin \theta \cos \varphi \end{aligned} \tag{14}$$

where x, y, z are the solar-magnetospheric (GSM) coordinates, θ is the polar angle plotted from x axis, φ being the azimuths angle plotted counterclockwise from z axis.

C.2 Spherical coordinates (2)

Spherical coordinates R, θ, φ with the polar axis plotted opposite the Earth dipole moment are determined by the expressions

$$\begin{aligned} x / R_1 &= R \sin \theta \cos \varphi \\ y / R_1 &= R \sin \theta \sin \varphi \\ z / R_1 &= R \cos \theta \end{aligned} \tag{15}$$

where x, y, z are the solar-magnetic (SM) coordinates, θ is the polar angle plotted from z axis, φ being the azimuths angle plotted counterclockwise from x axis.

C.3 Parabolic coordinates

Parabolic coordinates α, β, φ with the polar axis plotted along the Sun-Earth axis are determined by the expressions

$$\begin{aligned} 2x / R_1 &= \beta^2 - \alpha^2 + 1 \\ y / R_1 &= \alpha \beta \sin \varphi \\ x / R_1 &= \alpha \beta \cos \varphi \end{aligned} \tag{16}$$

where x, y, z are the GSM coordinates, φ being the azimuths angle plotted counterclockwise from z axis. In the magnetospheric paraboloid model the magnetopause is the $\beta = 1$ surface.

C.4 Magnetic Field of Geomagnetic Dipole

\vec{B}_d and $e_{z_{sm}}$ in the solar-magnetospheric coordinates are described by the expressions

$$\begin{aligned}\vec{B}_d &= -\nabla V_d \\ V_d &= \left(\frac{R_e}{R}\right)^3 B_0 \cdot (z \cos \psi - x \sin \psi) \\ \vec{e}_z &= (-\sin \psi; 0; \cos \psi).\end{aligned}\tag{17}$$

C.5 Dataset for model testing

The Large Magnetospheric Data Base [Fierfield et al., 1994] is used for the testing of the candidate models.

C.6 References

- 1) Fierfield et al., A large magnetosphere magnetic field database, *J. Geophys. Res.*, 99, 11,319, 1994.