

Acetaminophen (paracetamol): Drug information - UpToDate

Acetaminophen (paracetamol): Drug information - UpToDate

Official reprint from UpToDate®

www.uptodate.com © 2026 UpToDate, Inc. and/or its affiliates. All Rights Reserved.

Acetaminophen (paracetamol): Drug information

2026© UpToDate, Inc. and its affiliates and/or licensors. All Rights Reserved.

Contributor Disclosures

For additional information see "Acetaminophen (paracetamol): Patient drug information" and "Acetaminophen (paracetamol): Pediatric drug information"

For abbreviations, symbols, and age group definitions show table

ALERT: US Boxed Warning

Risk of medication errors and hepatotoxicity (injection):

Take care when prescribing, preparing, and administering acetaminophen injection to avoid dosing errors that could result in accidental overdose and death. In particular, be careful to ensure the following: the dose in milligrams and milliliters is not confused; the dosing is based on weight for patients less than 50 kg; infusion pumps are properly programmed; and the total daily dose of acetaminophen from all sources does not exceed maximum daily limits.

Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed the maximum daily limits, and often involve more than 1 acetaminophen-containing product.

Brand Names: US

- 8 Hour Pain Reliever [OTC];
- 8 HR Arthritis Pain Relief [OTC];
- Acetaminophen 8 Hour [OTC];
- Aminofen [OTC];
- Apra [OTC];
- Arthritis Pain Relief [OTC] [DSC];
- Aurophen Childrens [OTC] [DSC];
- BetaTemp Childrens [OTC];
- Childrens Acetaminophen [OTC] [DSC];
- Childrens APAP [OTC];
- Childrens Non-Aspirin [OTC];
- Childrens Silapap [OTC] [DSC];
- CuraNOL [OTC];
- E4-APAP [OTC];
- ElixSure Fever/Pain [OTC] [DSC];
- FeverAll Adults [OTC] [DSC];

- FeverAll Childrens [OTC];
- FeverAll Infants [OTC];
- FeverAll Junior Strength [OTC];
- FT 8 Hour Pain Relief [OTC];
- FT Arthritis Pain Reliever [OTC] [DSC];
- FT Children's Pain/Fever [OTC];
- FT Pain & Fever Childrens [OTC];
- FT Pain & Fever Infants [OTC];
- FT Pain Relief Adult Extra St [OTC];
- FT Pain Relief Extra Strength [OTC];
- FT Pain Relief [OTC];
- FT Pain Reliever Adults [OTC];
- FT Pain Reliever Children [OTC];
- FT Pain Reliever Ex Str Adult [OTC];
- FT Rapid Release Pain Relief [OTC];
- GoodSense Pain & Fever Child [OTC];
- GoodSense Pain & Fever Infants [OTC];
- GoodSense Pain Relief Extra St [OTC];
- Healthy Mama Shake That Ache [OTC];
- Liquid Pain Relief [OTC];
- M-PAP [OTC];
- Mapap Acetaminophen Extra Str [OTC];
- Mapap Arthritis Pain [OTC] [DSC];
- Mapap Childrens [OTC];
- Mapap [OTC];
- Max Relief Jr Child Pain/Fever [OTC];
- Max Relief Junior [OTC];
- Non-Aspirin Extra Strength [OTC];
- Non-Aspirin [OTC];
- Ofirmev [DSC];
- Pain & Fever Childrens [OTC];
- Pain and Fever Relief Kids [OTC];
- Pain Relief Childrens [OTC];
- Pain Relief Extra Strength [OTC];
- Pain Relief Regular Strength [OTC];
- Pain Relief [OTC];
- Panadol Childrens [OTC];
- Panadol Extra Strength [OTC];
- Panadol Infants [OTC];
- Pharbeto Extra Strength [OTC];
- Pharbeto [OTC];
- Triaminic Fever Reducer [OTC];
- Tylenol 8 Hour Arthritis Pain [OTC];
- Tylenol 8 Hour [OTC];
- Tylenol Childrens Chewables [OTC];
- Tylenol Childrens Pain + Fever [OTC];
- Tylenol Childrens [OTC];
- Tylenol Dissolve Packs [OTC];
- Tylenol Extra Strength [OTC];
- Tylenol for Children + Adults [OTC];
- Tylenol Infants Pain+Fever [OTC];
- Tylenol [OTC]

Pharmacologic Category

- Analgesic, Nonopioid

Dosing: Adult

Pain and/or fever

Pain (mild to moderate) and/or fever (monotherapy or as an adjunct):

Note: When calculating total daily dose, confirm that all sources (eg, prescription, OTCs, combinations) are included. Life-threatening acetaminophen-induced hepatotoxicity has been associated with doses >4 g/day. Although doses up to 4 g/day are generally well tolerated, hepatotoxicity has been reported rarely even with doses below this limit (Ref). Due to risk for hepatotoxicity, some experts recommend a lower maximum dose of 3 g/day in adults with normal liver function, particularly when used for longer durations (eg, >7 days) (Ref). An even lower total daily dose (eg, 2 g/day) or avoidance may be preferred in patients with certain risk factors for hepatotoxicity, such as heavy alcohol use, malnutrition, fasting, low body weight, advanced age, febrile illness, select liver disease, and concomitant use of interacting drugs (Ref).

Oral: 325 to 650 mg every 4 to 6 hours as needed **or** 1 g every 6 hours as needed; maximum dose: 4 g/day (Ref). See "Note" above regarding maximum dose.

OTC labeling (patient-guided therapy): **Note:** Dosage recommendations, including maximum doses, vary among OTC manufacturers.

Immediate release:

Regular strength (325 mg/tablet): 2 tablets (650 mg) every 4 to 6 hours as needed; maximum daily dose: 10 tablets/day (3.25 g/day).

Extra strength (500 mg/tablet): 2 tablets (1 g) every 6 hours as needed; maximum daily dose: 6 tablets/day (3 g/day).

Extended release (650 mg/tablet): 2 tablets (1.3 g) every 8 hours as needed; maximum daily dose: 6 tablets/day (3.9 g/day).

IV:

≥50 kg: 650 mg every 4 hours **or** 1 g every 6 hours; maximum single dose: 1 g/dose; maximum daily dose: 4 g/day.

<50 kg: 12.5 mg/kg every 4 hours **or** 15 mg/kg every 6 hours; maximum single dose: 15 mg/kg/dose (≤750 mg/dose); maximum daily dose: 75 mg/kg/day (≤3.75 g/day). **Note:** Some experts recommend this reduced dosing if used in patients with chronic alcoholism, malnutrition, or dehydration regardless of weight (Ref).

Rectal: 325 to 650 mg every 4 to 6 hours as needed (Ref); maximum daily dose: 3.9 g/day. **Note:** Absorption is irregular; bioavailability may be reduced by ~10% to 20% relative to oral administration (Ref).

Dosing: Kidney Impairment: Adult

The renal dosing recommendations are based upon the best available evidence and clinical expertise. Senior Editorial Team: Bruce Mueller, PharmD, FCCP, FASN, FNKF; Jason A. Roberts, PhD, BPharm (Hons), B App Sc, FSHP, FISAC; Michael Heung, MD, MS.

Altered kidney function: Mild to severe impairment: IV, Oral, Rectal: No dosage adjustment likely to be necessary. The manufacturer's labeling for IV acetaminophen states that longer dosing intervals and a reduced total daily dose may be warranted in patients with severe kidney impairment (CrCl ≤30 mL/minute); however, acetaminophen concentrations and half-life are increased but similar to those in patients with normal renal function (Ref). Glucuronide and sulfate conjugate metabolites accumulate in renal impairment, but the clinical effects are unknown (Ref).

Hemodialysis, intermittent (thrice weekly): IV, Oral, Rectal: Acetaminophen and its conjugates are readily dialyzable (Ref): No dosage adjustment necessary (Ref).

Peritoneal dialysis: IV, Oral, Rectal: Not dialyzed (Ref): No dosage adjustment necessary (Ref).

CRRT: IV, Oral, Rectal: Dialyzed (Ref): No dosage adjustment necessary (Ref).

PIRRT (eg, sustained, low-efficiency diafiltration): IV, Oral, Rectal: No dosage adjustment necessary (Ref).

Dosing: Liver Impairment: Adult

The liver dosing recommendations are based upon the best available evidence and clinical expertise. Senior Editorial Team: Matt Harris, PharmD, MHS, BCPS, FAST, FCCP; Jeong Park, PharmD, MS, BCTXP, FCCP, FAST; Arun Jesudian, MD; Sasan Sakiani, MD.

Note: Risk of acetaminophen-induced liver injury with preexisting liver insufficiency has been observed to occur in an increased dose-duration dependent fashion, especially in the presence of certain risk factors (eg, excessive alcohol intake [>3 drinks/day for males; >2 drinks/day for females], malnutrition, fasted state [eg, ≥ 1 day], low BMI) (Ref).

Liver impairment prior to treatment initiation:

Initial or dose adjustment in patients with preexisting liver cirrhosis:

Child-Turcotte-Pugh class A:

Patients actively consuming alcohol while taking acetaminophen (regardless of amount or frequency): IV, oral, rectal: Use of the lowest effective dose is preferred (eg, 325 to 500 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 2 g per day** from all acetaminophen sources (any duration of treatment) (Ref).

Patients who are not actively consuming alcohol: IV, oral, rectal:

Short-term use (≤ 14 days): No dosage adjustment necessary. **Maximum total daily dose: 4 g per day** from all acetaminophen sources (Ref).

Long-term use (>14 days): Use of the lowest effective dose is preferred (eg, 325 to 500 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 3 g per day** from all acetaminophen sources (Ref).

Child-Turcotte-Pugh class B:

Patients actively consuming alcohol while taking acetaminophen (regardless of amount or frequency): IV, oral, rectal: Use of the lowest effective dose is preferred (eg, 325 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 2 g per day** from all acetaminophen sources (any duration of treatment) (Ref).

Patients who are not actively consuming alcohol: IV, oral, rectal:

Short-term use (≤ 14 days): Use of the lowest effective dose is preferred (eg, 325 to 500 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 3 g per day** from all acetaminophen sources (Ref).

Long-term use (>14 days): Use of the lowest effective dose is preferred (eg, 325 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 2 g per day** from all acetaminophen sources (Ref).

Child-Turcotte-Pugh class C:

All patients (regardless of alcohol consumption):

IV: Use generally not recommended. If use of IV acetaminophen is deemed necessary, intermittent (eg, as needed) dosing instead of scheduled (eg, around-the-clock) dosing is preferred due to lack of data regarding accumulation; use with extreme caution. **Maximum total daily dose: 2 g per day** from all acetaminophen sources (Ref). **Note:** The manufacturer's labeling for the IV formulation states use is contraindicated in severe hepatic impairment or severe active liver disease.

Oral, rectal: Use of the lowest effective dose is preferred (eg, 325 mg every 4 to 6 hours); use with caution. **Maximum total daily dose: 2 g per day** from all acetaminophen sources (any duration of treatment) (Ref).

Dosing: Older Adult

Pain (acute) or fever: Oral, IV: Refer to adult dosing.

Persistent pain (off-label): Adults ≥ 75 years: Oral:

Initial: 325 to 500 mg every 4 hours **or** 500 to 1,000 mg every 6 hours

Maximum: $\leq 4,000$ mg/day. In older adults with hepatic impairment or history of alcohol abuse being treated for persistent pain, do not exceed a maximum of 2,000 to 3,000 mg/day (Ref).

Dosing: Pediatric

(For additional information see "Acetaminophen (paracetamol): Pediatric drug information")

Dosage guidance:

Dosage form information: Oral liquids are available in multiple concentrations (eg, 160 mg per 5 mL, 500 mg per 5 mL, 500 mg per 15 mL); verify appropriate product concentration; dose should be clearly presented as "mg."

Pain or fever

Pain (mild to moderate) or fever:

Note: Consider all sources of acetaminophen (eg, prescription, OTC, combination products) when evaluating a patient's maximum daily dose. To lower the risk for hepatotoxicity, limit daily dose to ≤ 75 mg/kg/**day** (maximum of 5 daily doses), not to exceed 4,000 mg/**day**. While recommended doses are generally considered safe, hepatotoxicity has been reported rarely, even with doses below recommendations (Ref).

Oral:

Weight-directed dosing: Infants, Children, and Adolescents: 10 to 15 mg/kg/dose every 4 to 6 hours as needed; do not exceed 5 doses in 24 hours; maximum daily dose: 75 mg/kg/**day** not to exceed 4,000 mg/**day** (Ref).

Fixed dosing:

Oral suspension, chewable tablets: Infants and Children < 12 years: Consult specific product formulations for appropriate age groups. See table; use of weight to select dose is preferred; if weight is not available, then use age; doses may be repeated every 4 hours; maximum: 5 doses/day.

Acetaminophen Dosing (Oral)

Weight (preferred)^A

Age

Dosage

(mg)

kg

lbs

^AManufacturer's recommendations are based on weight in pounds (OTC labeling); weight in kg listed here is derived from pounds and rounded; kg weight listed also is adjusted to allow for continuous weight ranges in kg. OTC labeling instructs consumer to consult with physician for dosing instructions in infants and children under 2 years of age.

2.7 to 5.3

6 to 11

0 to 3 mo

40

5.4 to 8.1

12 to 17

4 to 11 mo

80

8.2 to 10.8

18 to 23

1 to 2 y

120

10.9 to 16.3

24 to 35

2 to 3 y

160

16.4 to 21.7

36 to 47

4 to 5 y

240

21.8 to 27.2

48 to 59

6 to 8 y

320 to 325

27.3 to 32.6

60 to 71

9 to 10 y

325 to 400

32.7 to 43.2

72 to 95

11 y

480 to 500

Immediate-release solid dosage formulations: **Note:** Actual OTC dosing recommendations may vary by product and/or manufacturer:

Children 6 to 11 years: 325 mg every 4 to 6 hours; maximum daily dose: 1,625 mg/**day**; **Note:** Do not use more than 5 days unless directed by a physician.

Children ≥ 12 years and Adolescents:

Regular strength: 650 mg every 4 to 6 hours; maximum daily dose: 3,250 mg/**day** unless directed by a physician; under physician supervision daily doses $\leq 4,000$ mg may be used.

Extra strength: 1,000 mg every 6 hours; maximum daily dose: 3,000 mg/**day** unless directed by a physician; under physician supervision daily doses $\leq 4,000$ mg may be used.

Extended release: Children ≥ 12 years and Adolescents: 1,300 mg every 8 hours; maximum daily dose: 3,900 mg/**day**.

IV:

Infants and Children <2 years:

Manufacturer's labeling: Fever: 15 mg/kg/dose every 6 hours; maximum daily dose: 60 mg/kg/**day**.

Alternate dosing: Limited data available: Pain and fever: 7.5 to 15 mg/kg/dose every 6 hours; maximum daily dose: 60 mg/kg/**day** (Ref).

Children ≥2 years (Ref):

<50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 750 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 3,750 mg/**day**.

≥50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 1,000 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 4,000 mg/**day**.

Adolescents:

<50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 750 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 3,750 mg/**day**.

≥50 kg: 1,000 mg every 6 hours **or** 650 mg every 4 hours; maximum single dose: 1,000 mg; maximum daily dose: 4,000 mg/**day**.

Rectal:

Weight-directed dosing: Limited data available: Infants and Children <12 years: 10 to 20 mg/kg/dose every 4 to 6 hours as needed; **do not exceed 5 doses in 24 hours** (Ref); maximum daily dose: 75 mg/kg/**day** not to exceed 1,625 mg/**day**.

Fixed dosing:

Infants 6 to 11 months: 80 mg every 6 hours; maximum daily dose: 320 mg/**day**.

Infants and Children 12 to 36 months: 80 mg every 4 to 6 hours; maximum daily dose: 400 mg/**day**.

Children >3 to 6 years: 120 mg every 4 to 6 hours; maximum daily dose: 600 mg/**day**.

Children >6 up to 12 years: 325 mg every 4 to 6 hours; maximum daily dose: 1,625 mg/**day**.

Children ≥12 years and Adolescents: 650 mg every 4 to 6 hours; maximum daily dose: 3,900 mg/**day**.

Pain; peri-/postoperative management; adjunct to opioid therapy

Pain; peri-/postoperative management; adjunct to opioid therapy:

Note: Consider all sources of acetaminophen (eg, prescription, OTC, combination products) when evaluating a patient's maximum daily dose.

IV:

Infants and Children <2 years: Limited data available: 7.5 to 15 mg/kg/dose every 6 hours; maximum daily dose: 60 mg/kg/**day** (Ref).

Children ≥2 years (Ref):

<50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 750 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 3,750 mg/**day**.

≥50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 1,000 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 4,000 mg/**day**.

Adolescents:

<50 kg: 15 mg/kg/dose every 6 hours **or** 12.5 mg/kg/dose every 4 hours; maximum single dose: 15 mg/kg up to 750 mg; maximum daily dose: 75 mg/kg/**day** not to exceed 3,750 mg/**day**.

≥50 kg: 1,000 mg every 6 hours **or** 650 mg every 4 hours; maximum single dose: 1,000 mg; maximum daily dose: 4,000 mg/**day**.

Rectal: Limited data available: Children:

Loading dose: 40 mg/kg for 1 dose, in most trials, the dose was administered postoperatively (Ref); a maximum dose of 1,000 mg was most frequently reported. However, in one trial evaluating 24 older pediatric patients (all patients ≥ 25 kg; mean age: ~ 13 years), the data suggested that a dose of 1,000 mg does not produce therapeutic serum concentrations (target for study: >10 mcg/mL) compared to a 40 mg/kg dose (up to $\sim 2,000$ mg); the resultant $C_{\max}$ was: 7.8 mcg/mL (1,000 mg dose group) vs 15.9 mcg/mL (40 mg/kg dose group). **Note:** Therapeutic serum concentrations for analgesia have not been well-established (Ref).

Maintenance dose: 20 to 25 mg/kg/dose every 6 hours as needed for 2 to 3 days has been suggested if further pain control is needed postoperatively; maximum daily dose: 100 mg/kg/**day** not to exceed 4,000 mg/**day**; therapy longer than 5 days has not been evaluated (Ref).

Note: In the majority of trials, suppositories were not divided due to unequal distribution of drug within suppository; doses were rounded to the nearest mg amount using 1 or 2 suppositories of available product strengths.

Dosing: Kidney Impairment: Pediatric

Altered kidney function:

Infants, Children, and Adolescents:

Oral, rectal: Mild to severe impairment: There are no dosage adjustments provided in the manufacturer's labeling; based on adult pharmacokinetic studies, dosage adjustment may not be necessary for short courses. In adult pharmacokinetic studies, plasma concentrations of acetaminophen did not differ in renal impairment patients when compared to healthy patients for short courses of treatment (ie, 3 days); however, accumulation of the glucuronide and sulfate conjugate metabolites in renal impairment has been described following a single dose of acetaminophen up to repeat dosing for 10 days; the clinical significance of this finding is unknown (Ref).

IV:

Mild to moderate impairment: There are no dosage adjustments provided in the manufacturer's labeling.

Severe impairment ($\text{CrCl} < 30$ mL/minute): The manufacturer's labeling for IV acetaminophen states that longer dosing intervals and a reduced total daily dose may be warranted in patients with severe kidney impairment; use with caution.

Hemodialysis, intermittent: Acetaminophen and its conjugates are readily dialyzable (Ref): No dosage adjustment necessary when used for mild to moderate pain (Ref).

Peritoneal dialysis: Not dialyzed (Ref): No dosage adjustment necessary when used for mild to moderate pain (Ref).

Dosing: Liver Impairment: Pediatric

Use with caution. Limited, low-dose therapy is usually well-tolerated in hepatic disease/cirrhosis; however, cases of hepatotoxicity at daily acetaminophen dosages $< 4,000$ mg/day have been reported. Avoid chronic use in hepatic impairment.

Adverse Reactions (Significant): Considerations

Hepatotoxicity

Acute **hepatotoxicity** may result from intentional or unintentional overdose in adult and pediatric patients. In pediatric patients, unintentional overdose can be a result of accidental ingestion, supratherapeutic dosing, more frequent administration than recommended, and use of multiple acetaminophen-containing products; hepatotoxicity has also been rarely reported with recommended dosages (Ref).

Spontaneous resolution occurs with or without treatment in $\sim 65\%$ of cases; although, some cases may progress to acute liver failure leading to liver transplantation or death (Ref); a mortality rate

of ~0.4% has been reported (Ref). Acetaminophen is one of the most reported products causing drug-induced liver injury (Ref), with ~50% of cases of **acute hepatic failure** in the US attributed to acetaminophen (Ref).

Mechanism: Dose-related; direct toxic effect through formation of toxic metabolite, N-acetyl-p-benzoquinoneimine (NAPQI) that binds to cellular proteins, including mitochondrial proteins. Toxic free radicals, including peroxynitrite, may also cause damage inside the mitochondria (Ref).

Onset: Rapid. Nonspecific symptoms (eg, nausea, vomiting, abdominal pain) may occur within the first 24 hours post-ingestion, followed by resolution of nonspecific symptoms and initial increases in serum AST and ALT within 24 to 72 hours. Marked increases in AST and ALT occur within 72 to 96 hours post-ingestion, along with a return of nonspecific symptoms and additional symptoms (eg, jaundice, encephalopathy, coagulopathy, lactic acidosis) (Ref).

Risk factors:

- **Dose:**

- o Pediatric: Toxicity is likely to occur with single ingestions >150 mg/kg or when the maximum daily acetaminophen dose is >75 mg/kg/day (maximum of 5 daily doses) up to 4,000 mg/day from all sources (Ref).

- o Adult: Toxicity is likely to occur with single ingestions >250 **mg/kg** or >12,000 **mg** over a 24-hour period (Ref). Asymptomatic elevation of ALT may occur following maximal therapeutic doses of acetaminophen (4,000 **mg/day**) for ≥4 days (Ref).

- Multiple acetaminophen-containing products: An unintentional overdose may occur in adult and pediatric patients who take multiple acetaminophen or acetaminophen-containing combination products (Ref).

- Chronic alcohol ingestion: Patients with chronic alcohol use disorder who take therapeutic doses of acetaminophen are NOT at an increased risk of hepatotoxicity (Ref). In contrast, patients with chronic alcohol use disorder who ingest repeated supratherapeutic doses of acetaminophen are at an increased risk for hepatotoxicity (Ref).

- Concomitant medications and herbal products: Although use of products that induce CYP2E1 enzymes (eg, carbamazepine, phenobarbital, phenytoin, isoniazid, rifampin) have been postulated to predispose to acetaminophen hepatotoxicity by enhanced production of NAPQI, there is little evidence, aside from case reports, that drug interactions increase the risk of liver injury (Ref).

- Nutritional status: Malnutrition and fasting may increase the risk (Ref)

- Age: Pediatric patients are less susceptible, whereas elderly patients are at a higher risk (Ref)

- Delay to treatment with N-acetylcysteine (NAC): Most patients with acetaminophen overdose who receive treatment with NAC within 8 hours of ingestion will not develop hepatotoxicity (Ref)

Adverse Reactions

The following adverse drug reactions and incidences are derived from product labeling unless otherwise specified.

Oral, Rectal: Frequency not defined:

Dermatologic: Erythema of skin, skin blister, skin rash

Otic: Hearing loss

IV:

>10%: Gastrointestinal: Nausea (adults: 34%; neonates, infants, children, and adolescents: ≥5%), vomiting (adults: 15%; neonates, infants, children, and adolescents: ≥5%)

1% to 10%:

Cardiovascular: Hypertension, hypotension, peripheral edema (adults)

Dermatologic: Pruritus (neonates, infants, children, and adolescents: $\geq 5\%$)

Endocrine & metabolic: Hypoalbuminemia (neonates, infants, children, and adolescents), hypokalemia, hypomagnesemia (neonates, infants, children, and adolescents), hypophosphatemia (neonates, infants, children, and adolescents)

Gastrointestinal: Constipation (neonates, infants, children, and adolescents: $\geq 5\%$), diarrhea (neonates, infants, children, and adolescents)

Genitourinary: Oliguria (neonates, infants, children, and adolescents)

Hematologic & oncologic: Anemia

Hepatic: Increased serum aspartate aminotransferase (Watkins 2006)

Local: Infusion-site pain, pain at injection site

Nervous system: Agitation (neonates, infants, children, and adolescents), anxiety (adults), fatigue (adults), headache, insomnia (adults: 7%), trismus (adults)

Neuromuscular & skeletal: Muscle spasm ($\geq 1\%$)

Respiratory: Abnormal breath sounds (adults), atelectasis (neonates, infants, children, and adolescents), dyspnea (adults), pleural effusion (neonates, infants, children, and adolescents), pulmonary edema (neonates, infants, children, and adolescents), stridor (adults), wheezing (adults)

Postmarketing (all formulations):

Dermatologic: Acute generalized exanthematous pustulosis (FDA 2016), Stevens-Johnson syndrome (FDA 2016), toxic epidermal necrolysis (Watanabe 2016; FDA 2016)

Hepatic: Acute hepatic failure, hepatotoxicity (Ramachandran 2019, Yoon 2016), increased serum alanine aminotransferase (Watkins 2006)

Hypersensitivity: Anaphylaxis (Ho 2008, Numata 2016), hypersensitivity reaction (Thompson 2019)

Contraindications

Injection: Hypersensitivity to acetaminophen or any component of the formulation; severe hepatic impairment or severe active liver disease

OTC labeling: When used for self-medication, do not use with other drug products containing acetaminophen or if allergic to acetaminophen or any of the inactive ingredients

Warnings/Precautions

Disease-related concerns:

- Bariatric surgery: Presurgical assessment of the indication for use, symptoms, and goals of therapy should be documented to enable postsurgical assessment. Small studies identified changes to pharmacokinetic parameters postoperatively; however, total exposure was similar to normal weight individuals (Chen 2020; Goday Arno 2017; Porat 2019). Monitor for efficacy and toxicity after bariatric surgery and consider switching to an alternative formulation or medication if symptoms worsen.
- Hepatic impairment: Use with caution in patients with hepatic impairment or active liver disease; use of the IV formulation is contraindicated in patients with severe hepatic impairment or severe active liver disease.
- Hypovolemia: Use the IV formulation with caution in patients with severe hypovolemia (eg, due to dehydration or blood loss).

Dosage form specific issues:

- Aspartame: Some products may contain aspartame, which is metabolized to phenylalanine and must be avoided (or used with caution) in patients with phenylketonuria.
- Benzyl alcohol and derivatives: Some dosage forms may contain benzyl alcohol and/or sodium benzoate/benzoic acid; benzoic acid (benzoate) is a metabolite of benzyl alcohol; large amounts

of benzyl alcohol (≥ 99 mg/kg/day) have been associated with a potentially fatal toxicity (“gasping syndrome”) in neonates; the “gasping syndrome” consists of metabolic acidosis, respiratory distress, gasping respirations, CNS dysfunction (including convulsions, intracranial hemorrhage), hypotension and cardiovascular collapse (AAP [“Inactive” 1997]; CDC 1982); some data suggests that benzoate displaces bilirubin from protein binding sites (Ahlfors 2001); avoid or use dosage forms containing benzyl alcohol and/or benzyl alcohol derivative with caution in neonates. See manufacturer’s labeling.

- Polysorbate 80: Some dosage forms may contain polysorbate 80 (also known as Tweens). Hypersensitivity reactions, usually a delayed reaction, have been reported following exposure to pharmaceutical products containing polysorbate 80 in certain individuals (Isaksson 2002; Lucente 2000; Shelley 1995). Thrombocytopenia, ascites, pulmonary deterioration, and renal and hepatic failure have been reported in premature neonates after receiving parenteral products containing polysorbate 80 (Alade 1986; CDC 1984). See manufacturer’s labeling.

- Propylene glycol: Some dosage forms may contain propylene glycol; large amounts are potentially toxic and have been associated hyperosmolality, lactic acidosis, seizures and respiratory depression; use caution (AAP [“Inactive” 1997]; Zar 2007).

Other warnings/precautions:

- Dosage limit: Limit acetaminophen dose from all sources (prescription, OTC, combination products) and all routes of administration (IV, oral, rectal) to <4 g/day (adults).

- Self-medication (OTC use): When used for self-medication, patients should be instructed to contact health care provider if symptoms get worse or new symptoms appear, redness or swelling is present in the painful area, fever lasts >3 days (all ages), or pain (excluding sore throat) lasts longer than: Children ≥ 12 years, Adolescents, and Adults: 10 days; Infants and Children <12 years: 5 days. When treating children with sore throat, if sore throat is severe, persists for >2 days, or is followed by fever, rash, headache, nausea, or vomiting, consult health care provider immediately.

Warnings: Additional Pediatric Considerations

Prophylactic use of acetaminophen to reduce fever and discomfort associated with vaccination is not recommended by the Advisory Committee on Immunization Practices (ACIP). Additionally, the ACIP does not recommend prophylactic acetaminophen to reduce risk of febrile seizure in infants and children with or without a history of febrile seizures. Antipyretics have not been shown to prevent febrile seizures (NCIRD/ACIP 2011). One study reported that routine prophylactic administration of acetaminophen to prevent fever prior to vaccination decreased the immune response of some vaccines; in the trial evaluating 459 infants (including 226 who received acetaminophen), antibody geometric mean concentrations (GMCs) for targeted vaccine immune response markers were lower in significantly more infants in the acetaminophen group compared with control. Before the booster dose, children who received prophylactic acetaminophen had lower antibody GMCs for all vaccine serotypes than children in the control group; this effect persisted after boosting even in the absence of additional acetaminophen doses. The clinical significance of this reduction in immune response has not been established (Prymula 2009). Antipyretics may be used to treat fever or discomfort following vaccination (NCIRD/ACIP 2011).

Some dosage forms may contain propylene glycol; in neonates, large amounts of propylene glycol delivered orally, intravenously (eg, $>3,000$ mg/day), or topically have been associated with potentially fatal toxicities which can include metabolic acidosis, seizures, renal failure, and CNS depression; toxicities have also been reported in children and adults including hyperosmolality, lactic acidosis, seizures, and respiratory depression; use caution (AAP 1997; Shehab 2009).

Dosage Forms: US

Excipient information presented when available (limited, particularly for generics); consult specific product labeling. [DSC] = Discontinued product

Capsule, Oral:

Mapap: 500 mg [aspirin free; contains fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye)]

Tylenol: 325 mg [aspirin free; contains fd&c red #40 (allura red ac dye), soybeans (glycine soja)]

Elixir, Oral:

Apra: 160 mg/5 mL (120 mL, 240 mL, 480 mL, 3840 mL) [contains alcohol, usp]

Pain Relief Childrens: 160 mg/5 mL (118 mL, 237 mL, 473 mL) [alcohol free, aspirin free; contains fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), saccharin sodium, sodium benzoate, sorbitol]

Gel, Oral:

ElixSure Fever/Pain: 160 mg/5 mL (120 mL [DSC]) [alcohol free, aspirin free; contains butylparaben (butyl p-hydroxybenzoate), carbomer 934p, polyethylene glycol (macrogol); bubble-gum flavor]

ElixSure Fever/Pain: 160 mg/5 mL (120 mL [DSC]) [alcohol free, aspirin free; contains butylparaben (butyl p-hydroxybenzoate), carbomer 934p, polyethylene glycol (macrogol); cherry flavor]

ElixSure Fever/Pain: 160 mg/5 mL (120 mL [DSC]) [alcohol free, aspirin free; contains butylparaben (butyl p-hydroxybenzoate), carbomer 934p, polyethylene glycol (macrogol); grape flavor]

Liquid, Oral:

Childrens Silapap: 160 mg/5 mL (118 mL [DSC], 237 mL [DSC], 473 mL [DSC]) [alcohol free, sugar free; contains fd&c red #40 (allura red ac dye), methylparaben (methyl p-hydroxybenzoate), propylene glycol, saccharin sodium, sodium benzoate; cherry flavor]

CuraNOL: 160 mg/5 mL (473 mL) [alcohol free, dye free, gluten free, sugar free; contains methylparaben (methyl p-hydroxybenzoate), propylene glycol, propylparaben (propyl p-hydroxybenzoate); grape flavor]

Ed-APAP: 160 mg/5 mL (236 mL) [alcohol free; contains fd&c red #40 (allura red ac dye), saccharin sodium, sodium benzoate; cherry flavor]

Liquid Pain Relief: 160 mg/5 mL (473 mL) [alcohol free, aspirin free; contains fd&c red #40 (allura red ac dye), methylparaben (methyl p-hydroxybenzoate), polyethylene glycol (macrogol), propylene glycol, propylparaben (propyl p-hydroxybenzoate)]

M-PAP: 160 mg/5 mL (118 mL, 473 mL) [contains fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), propylene glycol, saccharin sodium, sodium benzoate; cherry flavor]

M-PAP: 160 mg/5 mL (120 mL, 473 mL) [alcohol free, aspirin free, sugar free; contains fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), saccharin sodium, sodium benzoate; cherry flavor]

Mapap Acetaminophen Extra Str: 500 mg/15 mL (237 mL) [contains fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), propylene glycol, saccharin sodium, sodium benzoate; cherry flavor]

Max Relief Jr Child Pain/Fever: 160 mg/5 mL (118 mL, 236 mL, 473 mL) [alcohol free, aspirin free, dye free; contains polyethylene glycol (macrogol), saccharin sodium, sodium benzoate; cherry flavor]

Max Relief Jr Child Pain/Fever: 160 mg/5 mL (118 mL, 236 mL, 473 mL) [alcohol free, aspirin free, dye free; contains polyethylene glycol (macrogol), saccharin sodium, sodium benzoate; grape flavor]

Max Relief Junior: 160 mg/5 mL (118 mL, 236 mL, 473 mL) [alcohol free, aspirin free, dye free; contains polyethylene glycol (macrogol), saccharin sodium, sodium benzoate; bubble-gum flavor]

Pain and Fever Relief Kids: 160 mg/5 mL (473 mL) [alcohol free, aspirin free, ibuprofen free, sugar free; contains fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), saccharin sodium, sodium benzoate; cherry flavor]

Pain Relief: 500 mg/15 mL (237 mL) [alcohol free, aspirin free; contains fd&c red #40 (allura red ac dye), methylparaben (methyl p-hydroxybenzoate), polyethylene glycol (macrogol), propylene glycol, propylparaben (propyl p-hydroxybenzoate)]

Generic: 160 mg/5 mL (5 mL [DSC], 118 mL, 236 mL, 473 mL); 500 mg/15 mL (237 mL); 1000 mg/30 mL (237 mL)

Packet, Oral:

Tylenol Childrens Pain + Fever: 160 mg (18 ea, 30 ea) [aspirin free, ibuprofen free; wild berry flavor]

Tylenol Dissolve Packs: 500 mg (12 ea, 32 ea) [berry flavor]

Solution, Intravenous:

Generic: 10 mg/mL (50 mL, 100 mL); 1000 mg/100 mL (100 mL)

Solution, Intravenous [preservative free]:

Ofirmev: 10 mg/mL (100 mL [DSC])

Generic: 10 mg/mL (50 mL, 65 mL, 100 mL)

Solution, Oral:

Generic: 160 mg/5 mL (5 mL, 118 mL, 473 mL); 325 mg/10.15 mL (10.15 mL); 650 mg/20.3 mL (20.3 mL)

Suppository, Rectal:

FeverAll Adults: 650 mg (50 ea [DSC]) [contains polysorbate 80]

FeverAll Childrens: 120 mg (6 ea, 50 ea) [contains polysorbate 80]

FeverAll Infants: 80 mg (1 ea, 6 ea, 50 ea) [contains polysorbate 80]

FeverAll Junior Strength: 325 mg (6 ea, 50 ea) [contains polysorbate 80]

FT Pain Reliever Adults: 650 mg (12 ea)

FT Pain Reliever Children: 120 mg (12 ea)

Generic: 120 mg (1 ea [DSC], 12 ea [DSC], 100 ea [DSC]); 650 mg (12 ea, 50 ea [DSC], 100 ea [DSC])

Suspension, Oral:

Aurophen Childrens: 160 mg/5 mL (118 mL [DSC]) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate]

BetaTemp Childrens: 160 mg/5 mL (118 mL) [contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate]

Childrens Acetaminophen: 160 mg/5 mL (5 mL [DSC]) [alcohol free, aspirin free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), polysorbate 80, propylene glycol, sodium benzoate]

Childrens Acetaminophen: 160 mg/5 mL (5 mL [DSC]) [alcohol free, aspirin free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), polysorbate 80, propylene glycol, sodium benzoate; strawberry flavor]

Childrens Non-Aspirin: 160 mg/5 mL (118 mL)

FT Pain & Fever Childrens: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, gluten free, ibuprofen free, paraben free; contains carrageenan, fd&c blue #1 (brilliant blue), propylene glycol, sodium benzoate; grape flavor]

FT Pain & Fever Childrens: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, gluten free, ibuprofen free, paraben free; contains carrageenan, fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate; bubble-gum flavor]

FT Pain & Fever Childrens: 160 mg/5 mL (473 mL) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c red #40 (allura red ac dye), sodium benzoate; strawberry flavor]

FT Pain & Fever Infants: 160 mg/5 mL (60 mL) [alcohol free, aspirin free, dye free, gluten free, ibuprofen free, paraben free; contains carrageenan, propylene glycol, sodium benzoate; grape flavor]

GoodSense Pain & Fever Child: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, dye free, gluten free, ibuprofen free; contains propylene glycol, propylparaben (propyl p-hydroxybenzoate), sodium benzoate, sorbitol]

GoodSense Pain & Fever Child: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c blue #1 (brilliant blue), propylene glycol, sodium benzoate, sorbitol; grape flavor]

GoodSense Pain & Fever Child: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate, sorbitol]

GoodSense Pain & Fever Child: 160 mg/5 mL (118 mL) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate, sorbitol; cherry flavor]

GoodSense Pain & Fever Infants: 160 mg/5 mL (30 mL) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c blue #1 (brilliant blue), propylene glycol, sodium benzoate, sorbitol]

GoodSense Pain & Fever Infants: 160 mg/5 mL (30 mL) [alcohol free, aspirin free, gluten free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate, sorbitol]

Max Relief Jr Child Pain/Fever: 160 mg/5 mL (60 mL, 118 mL, 236 mL, 473 mL) [contains fd&c blue #1 (brilliant blue); grape flavor]

Max Relief Jr Child Pain/Fever: 160 mg/5 mL (118 mL, 236 mL, 473 mL) [dye free; cherry flavor]

Pain & Fever Childrens: 160 mg/5 mL (118 mL) [aspirin free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate; cherry flavor]

Pain Relief Childrens: 160 mg/5 mL (118 mL) [cherry flavor]

Panadol Childrens: 160 mg/5 mL (118 mL) [aspirin free, ibuprofen free; contains benzoic acid, fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), propylene glycol, saccharin sodium; raspberry flavor]

Panadol Infants: 160 mg/5 mL (54.7 mL) [aspirin free, ibuprofen free; contains benzoic acid, fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), propylene glycol, saccharin sodium; raspberry flavor]

Tylenol Childrens: 160 mg/5 mL (120 mL) [alcohol free; cherry flavor]

Tylenol Childrens: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, ibuprofen free; contains butylparaben (butyl p-hydroxybenzoate), fd&c red #40 (allura red ac dye), propylene glycol, sodium benzoate; strawberry flavor]

Tylenol Childrens: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, ibuprofen free; contains fd&c blue #1 (brilliant blue), sodium benzoate; grape flavor]

Tylenol Childrens: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, ibuprofen free; contains fd&c red #40 (allura red ac dye), sodium benzoate]

Tylenol Childrens: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, ibuprofen free; contains sodium benzoate, sorbitol; grape flavor]

Tylenol Childrens Pain + Fever: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, dye free, ibuprofen free, paraben free]

Tylenol Childrens Pain + Fever: 160 mg/5 mL (120 mL, 240 mL) [alcohol free, aspirin free, dye free, ibuprofen free, paraben free; cherry flavor]

Tylenol Childrens Pain + Fever: 160 mg/5 mL (120 mL) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c red #40 (allura red ac dye), sodium benzoate; bubble-gum flavor]

Tylenol for Children + Adults: 160 mg/5 mL (240 mL) [alcohol free, aspirin free, dye free, ibuprofen free, paraben free; cherry flavor]

Tylenol Infants Pain+Fever: 160 mg/5 mL (60 mL) [alcohol free, aspirin free, dye free, ibuprofen free]

Tylenol Infants Pain+Fever: 160 mg/5 mL (30 mL) [alcohol free, aspirin free, dye free, ibuprofen free; cherry flavor]

Tylenol Infants Pain+Fever: 160 mg/5 mL (60 mL) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c blue #1 (brilliant blue), sodium benzoate]

Tylenol Infants Pain+Fever: 160 mg/5 mL (30 mL) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c blue #1 (brilliant blue), sodium benzoate; grape flavor]

Tylenol Infants Pain+Fever: 160 mg/5 mL (60 mL [DSC]) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c red #40 (allura red ac dye), sodium benzoate]

Tylenol Infants Pain+Fever: 160 mg/5 mL (60 mL) [alcohol free, aspirin free, ibuprofen free, paraben free; contains fd&c red #40 (allura red ac dye), sodium benzoate; cherry flavor]

Generic: 160 mg/5 mL (5 mL, 10.15 mL, 20.3 mL, 59 mL, 118 mL, 236 mL, 473 mL); 650 mg/20.3 mL (20.3 mL)

Syrup, Oral:

Triaminic Fever Reducer: 160 mg/5 mL (59 mL, 118 mL) [alcohol free, aspirin free, ibuprofen free; contains benzoic acid, edetate (edta) disodium, fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol); bubble-gum flavor]

Triaminic Fever Reducer: 160 mg/5 mL (59 mL, 118 mL) [alcohol free, aspirin free, ibuprofen free; contains edetate (edta) disodium, fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), polyethylene glycol (macrogol), sodium benzoate; grape flavor]

Tablet, Oral:

Aminofen: 325 mg [antihistamine free, caffeine free, salt free, sugar free; contains corn starch]

FT Pain Relief: 325 mg [DSC]

FT Pain Relief: 325 mg [scored]

FT Pain Relief Adult Extra St: 500 mg

FT Pain Relief Adult Extra St: 500 mg [contains fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), fd&c yellow #6 (sunset yellow)]

FT Pain Relief Extra Strength: 500 mg

FT Pain Reliever Ex Str Adult: 500 mg [gluten free; contains corn starch]

FT Pain Reliever Ex Str Adult: 500 mg [gluten free; contains corn starch, fd&c red #40(allura red ac)aluminum lake]

FT Rapid Release Pain Relief: 500 mg [gluten free; contains fd&c red #40(allura red ac)aluminum lake, fd&c yellow #6(sunset yellow)alumin lake]

GoodSense Pain Relief Extra St: 500 mg [contains fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), fd&c yellow #6 (sunset yellow)]

GoodSense Pain Relief Extra St: 500 mg [gluten free; contains fd&c red #40(allura red ac)aluminum lake, fd&c yellow #6(sunset yellow)alumin lake]

Healthy Mama Shake That Ache: 500 mg

Non-Aspirin: 325 mg

Non-Aspirin: 325 mg, 500 mg [contains corn starch]

Non-Aspirin Extra Strength: 500 mg

Pain Relief Extra Strength: 500 mg

Pain Relief Extra Strength: 500 mg [contains corn starch]

Pain Relief Extra Strength: 500 mg [aspirin free]

Pain Relief Extra Strength: 500 mg [aspirin free; contains fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye)]

Pain Relief Regular Strength: 325 mg [contains methylparaben (methyl p-hydroxybenzoate), propylparaben (propyl p-hydroxybenzoate)]

Panadol Extra Strength: 500 mg [contains fd&c blue #1 (brill blue) aluminum lake, fd&c red #40 (allura red ac) aluminum lake, quinoline (d&c yellow #10) aluminum lake]

Pharbetol: 325 mg

Pharbetol Extra Strength: 500 mg [aspirin free]

Pharbetol Extra Strength: 500 mg [aspirin free; contains corn starch]

Tylenol: 325 mg

Tylenol: 325 mg [contains corn starch, fd&c red #40 (allura red ac) aluminum lake]

Tylenol Extra Strength: 500 mg [contains butylparaben (butyl p-hydroxybenzoate), edetate (edta) calcium disodium, fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), methylparaben (methyl p-hydroxybenzoate), polysorbate 80, propylparaben (propyl p-hydroxybenzoate), quinoline yellow (d&c yellow #10)]

Tylenol Extra Strength: 500 mg [contains butylparaben (butyl p-hydroxybenzoate), edetate (edta) calcium disodium, fd&c blue #1 (brilliant blue), fd&c red #40 (allura red ac dye), methylparaben (methyl p-hydroxybenzoate), propylparaben (propyl p-hydroxybenzoate), quinoline yellow (d&c yellow #10)]

Tylenol Extra Strength: 500 mg [contains corn starch, fd&c red #40 (allura red ac) aluminum lake]

Tylenol Extra Strength: 500 mg [contains fd&c red #40 (allura red ac dye), fd&c yellow #6 (sunset yellow)]

Tylenol Extra Strength: 500 mg [contains fd&c red #40 (allura red ac) aluminum lake]

Tylenol Extra Strength: 500 mg [contains fd&c red #40 (allura red ac) aluminum lake, fd&c yellow #6 (sunset yellow) aluminum lake]

Generic: 325 mg, 500 mg

Tablet Chewable, Oral:

Childrens APAP: 80 mg [scored; contains aspartame, fd&c yellow #6 (sunset yellow) aluminum lake; fruit flavor]

Childrens Non-Aspirin: 80 mg

FT Children's Pain/Fever: 160 mg [scored; aspirin free, ibuprofen free; bubble-gum flavor]

Mapap Childrens: 80 mg [aspirin free, ibuprofen free; contains fd&c blue #1 (brill blue) aluminum lake; grape flavor]

Mapap Childrens: 160 mg [scored; aspirin free, ibuprofen free; bubble-gum flavor]

Tylenol Childrens Chewables: 160 mg [aspirin free, ibuprofen free]

Tylenol Childrens Chewables: 160 mg [aspirin free, ibuprofen free; contains fd&c blue #1 (brilliant blue)]

Generic: 80 mg, 160 mg [DSC], 325 mg [DSC]

Tablet Extended Release, Oral:

8 Hour Pain Reliever: 650 mg

8 HR Arthritis Pain Relief: 650 mg [aspirin free]

Acetaminophen 8 Hour: 650 mg [aspirin free; contains corn starch]

Arthritis Pain Relief: 650 mg [DSC]

FT 8 Hour Pain Relief: 650 mg

FT Arthritis Pain Reliever: 650 mg [DSC]

Mapap Arthritis Pain: 650 mg [DSC] [gluten free]

Tylenol 8 Hour: 650 mg

Tylenol 8 Hour Arthritis Pain: 650 mg

Generic: 650 mg

Therapy Pack, Oral:

Tylenol Childrens: 160 & 160 MG &MG/5ML (1 ea) [aspirin free, ibuprofen free; contains fd&c blue #1 (brilliant blue), sodium benzoate]

Generic Equivalent Available: US

May be product dependent

Pricing: US

Capsules (Tylenol Oral)

325 mg (per each): \$0.20

Chewable (Mapap Childrens Oral)

80 mg (per each): \$0.08

160 mg (per each): \$0.10

Chewable (Tylenol Childrens Chewables Oral)

160 mg (per each): \$0.29

Liquid (Acetaminophen Oral)

160 mg/5 mL (per mL): \$0.06

Liquid (Mapap Acetaminophen Extra Str Oral)

500 mg/15 mL (per mL): \$0.02

Liquid (Max Relief Jr Child Pain/Fever Oral)

160 mg/5 mL (per mL): \$0.01

Liquid (Max Relief Junior Oral)

160 mg/5 mL (per mL): \$0.01

Pack (Tylenol Childrens Pain + Fever Oral)

160 mg (per each): \$0.38

Pack (Tylenol Dissolve Packs Oral)

500 mg (per each): \$0.36

Solution (Acetaminophen Childrens Oral)

160 mg/5 mL (per mL): \$0.02

Solution (Acetaminophen Intravenous)

10 mg/mL (per mL): \$0.09 - \$0.45

Solution (Acetaminophen Oral)

160 mg/5 mL (per mL): \$0.18 - \$0.26

Suppository (Acetaminophen Rectal)

650 mg (per each): \$0.67

Suppository (FeverAll Childrens Rectal)

120 mg (per each): \$1.12

Suppository (FeverAll Infants Rectal)

80 mg (per each): \$1.12

Suppository (FeverAll Junior Strength Rectal)

325 mg (per each): \$1.22

Suspension (Acetaminophen Oral)

160 mg/5 mL (per mL): \$0.22 - \$0.55

Suspension (Max Relief Jr Child Pain/Fever Oral)

160 mg/5 mL (per mL): \$0.02

Suspension (Tylenol Childrens Oral)

160 mg/5 mL (per mL): \$0.06

Suspension (Tylenol Childrens Pain + Fever Oral)

160 mg/5 mL (per mL): \$0.07

Suspension (Tylenol for Children + Adults Oral)

160 mg/5 mL (per mL): \$0.05

Suspension (Tylenol Infants Pain+Fever Oral)

160 mg/5 mL (per mL): \$0.16

Tablet, controlled release (Acetaminophen ER Oral)

650 mg (per each): \$0.07 - \$0.10

Tablet, controlled release (Tylenol 8 Hour Arthritis Pain Oral)

650 mg (per each): \$0.14

Tablet, controlled release (Tylenol 8 Hour Oral)

650 mg (per each): \$0.18

Tablets (Acetaminophen Oral)

325 mg (per each): \$0.02 - \$0.06

500 mg (per each): \$0.02 - \$0.08

Tablets (Healthy Mama Shake That Ache Oral)

500 mg (per each): \$0.06

Tablets (Pharbetol Extra Strength Oral)

500 mg (per each): \$0.03

Tablets (Pharbetol Oral)

325 mg (per each): \$0.02

Tablets (Tylenol Extra Strength Oral)

500 mg (per each): \$0.12

Tablets (Tylenol Oral)

325 mg (per each): \$0.08

Therapy Pack (Tylenol Childrens Oral)

160 & 160 mg&mg/5 mL (per each): \$12.36

Disclaimer: A representative AWP (Average Wholesale Price) price or price range is provided as reference price only. A range is provided when more than one manufacturer's AWP price is available and uses the low and high price reported by the manufacturers to determine the range. The pricing data should be used for benchmarking purposes only, and as such should not be used alone to set or adjudicate any prices for reimbursement or purchasing functions or considered to be an exact price for a single product and/or manufacturer. Medi-Span expressly disclaims all warranties of any kind or nature, whether express or implied, and assumes no liability with respect to accuracy of price or price range data published in its solutions. In no event shall Medi-Span be liable for special, indirect, incidental, or consequential damages arising from use of price or price range data. Pricing data is updated monthly.

Dosage Forms: Canada

Excipient information presented when available (limited, particularly for generics); consult specific product labeling.

Solution, Intravenous:

Generic: 10 mg/mL (10 mL, 50 mL, 100 mL)

Administration: Adult

Oral: May administer without regard to food; may administer with food to decrease possible GI upset; shake drops and suspension well before use; do not crush or chew ER products.

Bariatric surgery: Acetaminophen is available as an ER formulation. Bariatric surgery may significantly alter the release characteristics in an unknown manner. Acetaminophen is also available in an IR formulation. Acetaminophen liquid formulations may contain nonabsorbable sugars (eg, sorbitol, mannitol, xylitol) that can cause dumping syndrome (Ref). Refer to product labeling. Providers should determine if the condition being treated can be safely monitored or if a switch to an alternative formulation or medication is necessary (Ref).

Injection: For IV infusion only. Administer undiluted over 15 minutes. Attach an administration set in accordance with the manufacturer's recommendations; may vary by product. For doses <1,000 mg, withdraw appropriate dose and place into separate empty, sterile container (eg, glass bottle, plastic IV container, syringe) for administration, to avoid inadvertent administration of the total volume of the commercially available container.

Rectal: Remove wrapper; insert suppository well up into the rectum.

Preparation for Administration: Adult

Injectable solution may be administered without further dilution. For doses <1,000 mg, withdraw appropriate dose and place into separate empty, sterile container (eg, glass bottle, plastic IV container, syringe) for administration, to avoid inadvertent administration of the total volume of the commercially available container.

Administration: Pediatric

Oral: Administer with food to decrease GI upset; shake drops and suspension well before use; do not crush or chew extended-release products.

Parenteral: For IV infusion only. May administer undiluted over 15 minutes per the manufacturer. In neonatal patients, infusion of undiluted and diluted solutions over 15 to 30 minutes has been reported (Ref). Use within 6 hours of opening vial or transferring to another container. Discard any unused portion; single-use vials only.

Rectal: Remove wrapper; insert suppository well up into the rectum.

Preparation for Administration: Pediatric

Parenteral: Injectable solution may be administered directly from the vial without further dilution. Use within 6 hours of opening vial or transferring to another container. Discard any unused portion; single-use vials only.

Doses <1,000 mg (<50 kg): Withdraw appropriate dose from vial and transfer to a separate sterile container (eg, glass bottle, plastic IV container, syringe) for administration. Small volume pediatric doses (up to 600 mg [60 mL]) may be placed in a syringe.

Doses of 1,000 mg (≥50 kg): Insert vented IV set through vial stopper.

Storage/Stability

Injection: Store intact vials and bags at 20°C to 25°C (68°F to 77°F); do not refrigerate or freeze. Some products must be protected from light (eg, B. Braun PAB container); refer to manufacturer's labeling for product-specific details. Nonintact (in use) storage time varies by product (eg, use immediately [B. Braun PAB container] or within 6 hours [Baxter Viaflo container; Ofirmev]; refer to manufacturer's labeling for product-specific details). Discard any unused portion.

Oral formulations: Store at 20°C to 25°C (68°F to 77°F); avoid excessive heat (40°C [104°F]). Avoid high humidity (chewable tablets).

Suppositories: Store at 2°C to 27°C (25°F to 80°F); do not freeze.

Use: Labeled Indications

Fever: Temporary reduction of fever.

Pain:

Injection: Management of mild to moderate pain in patients ≥2 years of age; management of moderate to severe pain when combined with opioid analgesia in patients ≥2 years.

Oral, Rectal: Temporary relief of minor aches, pains, and headache.

Medication Safety Issues

Sound-alike/look-alike issues:

Acephen may be confused with AcipHex

Acetaminophen may be confused with acetaZOLAMIDE

FeverALL may be confused with Fiberall

Triaminic Children's Fever Reducer Pain Reliever may be confused with Triaminic cough and cold products

Tylenol may be confused with atenolol, timolol, Tylenol PM, Tylox

Infusion bottles of ropivacaine and IV acetaminophen look similar. Potentially fatal mix-ups have been reported in which a glass bottle of Naropin was mistaken for Ofirmev in perioperative areas.

High alert medication:

The Institute for Safe Medication Practices (ISMP) includes this medication among its list of drugs (pediatric liquid medications requiring measurement) which have a heightened risk of causing significant patient harm when used in error (ISMP List of High-Alert Medications in Community/Ambulatory Care Settings ©ISMP 2021).

Other safety concerns:

APAP is an error-prone abbreviation (may not be recognized as acetaminophen)

Duplicate therapy issues: This product contains acetaminophen, which may be a component of combination products. Do not exceed the maximum recommended daily dose of acetaminophen.

Infant concentration change: All children's and infant acetaminophen products are available as 160 mg/5 mL. Some remaining infant concentrated solutions of 80 mg/0.8 mL and 100 mg/mL may still be available on pharmacy shelves or in patient homes. Check concentrations closely prior to administering or dispensing and verify concentration available to patients prior to recommending a dose (November 2011).

Injection: Reports of 10-fold overdose errors using the parenteral product have occurred in the U.S. and Europe; calculation of doses in "mg" and subsequent administration of the dose in "mL" using the commercially available concentration of 10 mg/mL contributed to these errors. Expressing doses as mg **and** mL, as well as pharmacy preparation of doses, may decrease error potential (Dart, 2012; ISMP, 2012).

International issues:

Anacin-3 [Taiwan] may be confused with Anacin brand name for lidocaine/prilocaine [Korea]; Anacin 81 brand name for aspirin [Puerto Rico]; Anacin New brand name for acetaminophen/caffeine [India]

Depon [Greece] may be confused with Depen brand name for penicillamine [US]; Depin brand name for nifedipine [India]; Dipen brand name for diltiazem [Greece]

Duorol [Spain] may be confused with Diuril brand name for chlorothiazide [US, Canada]

Paralen [Czech Republic] may be confused with Aralen brand name for chloroquine [US, Mexico]

Procet [Kenya] may be confused with Percocet brand name for oxycodone/acetaminophen [US, Israel, Puerto Rico]

Procet brand name for acetaminophen [Kenya], but also brand name for cetirizine [Bangladesh]

Metabolism/Transport Effects

Substrate of CYP1A2 (Minor), CYP2A6 (Minor), CYP2C9 (Minor), CYP2D6 (Minor), CYP2E1 (Major with inducers), CYP2E1 (Minor with inhibitors), CYP3A4 (Minor), UGT1A1, UGT1A6, UGT1A9, UGT2B15; **Note:** Assignment of Major/Minor substrate status based on clinically relevant drug interaction potential

Drug Interactions

(For additional information: Launch drug interactions program)

Note: Interacting drugs may **not be individually listed below** if they are part of a group interaction (eg, individual drugs within "CYP3A4 Inducers [Strong]" are NOT listed). For a complete list of drug interactions by individual drug name and detailed management recommendations, use the drug interactions program by clicking on the "Launch drug interactions program" link above.

Alcohol (Ethyl): May increase hepatotoxic effects of Acetaminophen. *Risk C: Monitor*

Busulfan: Acetaminophen may increase serum concentrations of Busulfan. *Risk C: Monitor*

CarBAMazepine: May increase metabolism of Acetaminophen. This may 1) diminish the effect of acetaminophen; and 2) increase the risk of liver damage. *Risk C: Monitor*

Dapsone (Topical): May increase adverse/toxic effects of Methemoglobinemia Associated Agents. *Risk C: Monitor*

Dasatinib: Acetaminophen may increase hepatotoxic effects of Dasatinib. Dasatinib may increase serum concentrations of Acetaminophen. Management: Avoid coadministration of acetaminophen

and dasatinib if possible. If coadministration is unavoidable, monitor for signs/symptoms of hepatotoxicity, particularly in patients with greater acetaminophen exposure. *Risk D: Consider Therapy Modification*

Flucloxacillin: May increase adverse/toxic effects of Acetaminophen. Specifically, the risk for high anion gap metabolic acidosis may be increased. *Risk C: Monitor*

Fosphenytoin-Phenytoin: May decrease serum concentrations of Acetaminophen. Specifically, serum concentrations of acetaminophen may be decreased (leading to decreased efficacy), but the formation of the toxic N-acetyl-p-benzoquinone imine (NAPQI) metabolite may be increased (leading to increased hepatotoxicity). *Risk C: Monitor*

Imatinib: Acetaminophen may increase hepatotoxic effects of Imatinib. *Risk C: Monitor*

Immune Checkpoint Inhibitors (Anti-PD-1, -PD-L1, and -CTLA4 Therapies): Acetaminophen may decrease therapeutic effects of Immune Checkpoint Inhibitors (Anti-PD-1, -PD-L1, and -CTLA4 Therapies). *Risk C: Monitor*

Isoniazid: May increase hepatotoxic effects of Acetaminophen. Isoniazid may increase metabolism of Acetaminophen. Specifically, formation of the hepatotoxic NAPQI metabolite may be increased. *Risk C: Monitor*

Lamotrigine: Acetaminophen may decrease serum concentrations of Lamotrigine. *Risk C: Monitor*

Local Anesthetics: Methemoglobinemia Associated Agents may increase adverse/toxic effects of Local Anesthetics. Specifically, the risk for methemoglobinemia may be increased. *Risk C: Monitor*

Lorlatinib: May decrease serum concentrations of Acetaminophen. *Risk C: Monitor*

Metyrapone: May increase serum concentrations of Acetaminophen. More importantly, by inhibiting the conjugative metabolism of acetaminophen, metyrapone may shift the metabolism towards the oxidative route that produces a hepatotoxic metabolite. *Risk X: Avoid*

Nitric Oxide: May increase adverse/toxic effects of Methemoglobinemia Associated Agents. Combinations of these agents may increase the likelihood of significant methemoglobinemia. *Risk C: Monitor*

Phenobarbital: May increase metabolism of Acetaminophen. Specifically, formation of the hepatotoxic NAPQI metabolite may be increased. *Risk C: Monitor*

Phenylephrine (Systemic): Acetaminophen may increase serum concentrations of Phenylephrine (Systemic). *Risk C: Monitor*

Prilocaine: Methemoglobinemia Associated Agents may increase adverse/toxic effects of Prilocaine. Combinations of these agents may increase the likelihood of significant methemoglobinemia. Management: Monitor for signs of methemoglobinemia when prilocaine is used in combination with other agents associated with development of methemoglobinemia. Avoid use of these agents with prilocaine/lidocaine cream in infants less than 12 months of age. *Risk C: Monitor*

Primaquine: Methemoglobinemia Associated Agents may increase adverse/toxic effects of Primaquine. Specifically, the risk for methemoglobinemia may be increased. Management: Avoid concomitant use of primaquine and other drugs that are associated with methemoglobinemia when possible. If combined, monitor methemoglobin levels closely. *Risk D: Consider Therapy Modification*

Primidone: May increase metabolism of Acetaminophen. Specifically, formation of the hepatotoxic NAPQI metabolite may be increased. *Risk C: Monitor*

Probenecid: May increase serum concentrations of Acetaminophen. Probenecid may also limit the formation of at least one major non-toxic metabolite, possibly increasing the potential for formation of the toxic NAPQI metabolite. *Risk C: Monitor*

Rifampin: May increase hepatotoxic effects of Acetaminophen. Rifampin may decrease serum concentrations of Acetaminophen. *Risk C: Monitor*

Sodium Nitrite: Methemoglobinemia Associated Agents may increase adverse/toxic effects of Sodium Nitrite. Combinations of these agents may increase the likelihood of significant methemoglobinemia. *Risk C: Monitor*

SORafenib: Acetaminophen may increase hepatotoxic effects of SORafenib. SORafenib may increase serum concentrations of Acetaminophen. Management: Avoid coadministration of acetaminophen and sorafenib if possible. If coadministration is unavoidable, monitor for signs/symptoms of hepatotoxicity, particularly in patients with greater acetaminophen exposure. *Risk D: Consider Therapy Modification*

Vaccines: Acetaminophen may decrease therapeutic effects of Vaccines. Management: Consider avoiding routine prophylactic use of acetaminophen before or during vaccine administration when possible. Acetaminophen is still recommended to treat fevers and/or pain that occurs after vaccination. *Risk D: Consider Therapy Modification*

Vitamin K Antagonists: Acetaminophen may increase anticoagulant effects of Vitamin K Antagonists. This appears most likely with daily acetaminophen doses exceeding 1.3 or 2 g/day for multiple consecutive days. *Risk C: Monitor*

Food Interactions

Rate of absorption may be decreased when given with food. Management: Administer without regard to food.

Pregnancy Considerations

Summary: Acetaminophen is considered safe for use in people who are pregnant and is the preferred first-line medication for analgesia and antipyretic use in pregnancy. Use the lowest effective dose for the shortest possible duration (Ref). Currently, the most methodologically rigorous studies do not suggest a causal relationship with adverse effects to the fetus or the newborn as they grow and develop.

Rationale for recommendations:

Infant outcomes: High-quality evidence and expert consensus do not support an association to major adverse outcomes (eg, congenital malformations, miscarriage, stillbirth, neurodevelopment effects (Ref). Less robust data may suggest acetaminophen's association with specific adverse events in offspring but are not supported when compared to more robust evidence (Ref).

Congenital malformations: Based on epidemiological data, an increased risk of major congenital malformations has not been observed following maternal use of acetaminophen during pregnancy (Ref).

Pregnancy loss or stillbirth: The use of acetaminophen in recommended doses during pregnancy has not been associated with an increased risk of miscarriage or still birth (Ref).

Neurodevelopmental effects: Recent evidence does not suggest a causal association of adverse neurodevelopmental effects such as attention-deficit/hyperactivity disorder (ADHD) or autism spectrum disorder (ASD) following *in utero* acetaminophen exposure (Ref). Studies that used in-family siblings as the control group did not find an association between acetaminophen use during pregnancy and either ASD or ADHD (Ref). The FDA has expressed concerns about the use of acetaminophen during pregnancy and the possible small increased risk of neurological conditions (eg, autism, ADHD) in children; however, a causal relationship has not been established (Ref). Other organizations have also stated that causality has not been established (Ref).

Pulmonary effects: Causality of pulmonary conditions such as asthma (presenting in early childhood) have not been established with use of acetaminophen in pregnancy (Ref).

Male reproductive development: Adverse effects on male reproductive development (eg, cryptorchidism) have also been reported; however, causality has not been established (Ref).

Ductus arteriosus constriction: Short-term use of acetaminophen at therapeutic doses during the third trimester has not been associated with fetal prenatal constriction of the ductus arteriosus (Ref).

Hyperbilirubinemia: Avoid acetaminophen use in patients with cholestasis who are pregnant. Acetaminophen use, at typical doses, can impair the placental barrier to bile acids, thereby increasing fetal exposure to bile acids (Ref) in these patients.

Pharmacokinetics:

Acetaminophen crosses the placenta (Ref). Fetal acetaminophen pharmacokinetics parallel that of the mother, suggesting that placental transfer is flow limited. Maternal acetaminophen plasma concentrations can serve as a surrogate for fetal exposure (Ref).

Acetaminophen drug clearance is increased during pregnancy, but dose adjustment is unnecessary (Ref).

Breastfeeding Considerations

Acetaminophen is present in human milk.

Data related to the presence of acetaminophen in human milk are available from multiple sources (Berlin 1980; Bitzén 1981; Hurden 1980; Notarianni 1987).

- Eleven patients 2 to 22 months' postpartum were administered a single oral dose of acetaminophen 650 mg. Plasma concentrations of acetaminophen were measured in 2 women during the first 6 hours after dosing. Acetaminophen appeared in the maternal milk and saliva in similar concentrations within 15 minutes (peak: 10 to 15 mcg/mL at 1 to 2 hours). Acetaminophen was no longer present in human milk 12 hours after dosing. The mean half-life of acetaminophen in human milk was 2.28 hours (range: 1.35 to 3.5 hours). Authors of the study calculated the estimated exposure to the breastfeeding infant to be 0.14% of the maternal dose (range: 0.04% to 0.23%). Acetaminophen was not detected in the urine of breastfeeding infants (Berlin 1980).

- In a study of 3 breastfeeding women given acetaminophen 500 mg orally, peak concentrations were observed 2 hours after the dose in both the milk (4 mcg/mL) and serum (5 to 7 mcg/mL). The mean half-life of acetaminophen was 2.74 hours in the plasma and 2.64 hours in human milk (Bitzén 1981).

- A study evaluated 6 infants (2 to 6 days of age) exposed to acetaminophen via human milk. The maternal dose of acetaminophen was 1 to 2 g taken 2 to 4 hours prior to breastfeeding. Acetaminophen, acetaminophen glucuronide, and acetaminophen sulfate were detected in the urine of all 6 infants. Other metabolites were also found in some samples, and it was assumed all metabolites were synthesized by the infants. Following maternal use of acetaminophen 1 g, the authors estimated exposure to the breastfeeding infant to be 1.85% of the weight-adjusted maternal dose (Notarianni 1987).

- The presence of acetaminophen in human milk was studied in 11 women, 3 to 9 days postpartum given acetaminophen 1 g orally. Multiple paired milk and serum samples were obtained over 4 hours. Human milk concentrations of acetaminophen were 2.1 to 15.9 mcg/mL (n=32) (Hurden 1980). Using a human milk concentration of 15.9 mcg/mL, the estimated exposure of acetaminophen to the breastfed infant would be 2.385 mg/kg/day (relative infant dose [RID]: 3.98% based on a therapeutic infant dose of 60 mg/kg/day).

- In general, breastfeeding is considered acceptable when the RID is <10% (Anderson 2016; Ito 2000).

A rash likely caused by acetaminophen was observed in a 2-month-old fully breastfed infant. Following maternal use of acetaminophen 1 g once daily for 2 days, a maculopapular rash appeared on the baby's upper trunk and face, which resolved within 24 hours. Two weeks later, another maternal dose of acetaminophen 1 g was taken when a similar rash appeared after the infant was breastfed 3 times following the dose (Matheson 1985). A prospective cohort study evaluated the outcomes of breastfed infants whose mothers were taking various medications. Within the study, 43 mother-infant pairs reported acetaminophen exposure (dose, duration, and relationship to breastfeeding not provided). There were no cases of diarrhea, drowsiness, or irritability in the breastfed infants (Ito 1993).

Acetaminophen is the preferred initial treatment for acute migraine headache in lactating patients (ACOG 2022). Nonopioid analgesics are preferred for lactating patients who require pain control

peripartum or for surgery outside of the postpartum period (ABM [Martin 2018]; ABM [Reece-Stremtan 2017]; Sachs 2013). Acetaminophen is one of the preferred non-opioid agents (Sachs 2013) and is considered compatible with breastfeeding when used in usual recommended doses (WHO 2002).

Dietary Considerations

Some products may contain phenylalanine and/or sodium.

Monitoring Parameters

Serum acetaminophen levels: Where acute overdose suspected and with long-term use in patients with hepatic disease; liver enzymes in patients with prolonged use or in populations with reduced hepatic function (eg, neonates); relief of pain or fever.

Mechanism of Action

Although not fully elucidated, the analgesic effects are believed to be due to activation of descending serotonergic inhibitory pathways in the CNS. Interactions with other nociceptive systems may be involved as well (Smith 2009). Antipyresis is produced from inhibition of the hypothalamic heat-regulating center.

Pharmacokinetics (Adult Data Unless Noted)

Onset of action:

Oral: <1 hour.

IV: Analgesia: 5 to 10 minutes; Antipyretic: Within 30 minutes.

Peak effect: IV: Analgesic: 1 hour.

Duration:

IV, Oral: Analgesia: 4 to 6 hours.

IV: Antipyretic: ≥ 6 hours.

Absorption:

Oral: Primarily absorbed in small intestine (rate of absorption dependent upon gastric emptying); minimal absorption from stomach; varies by dosage form. Absorption is delayed in neonates in the first few days of life (Anderson 2002).

Rectal: Delayed and erratic absorption; varies by dosage form and age (Allegaert 2020).

Distribution: V_d :

Neonates and Infants <2 years: V_d : Median range: 0.85 to 0.95 L/kg (Hammer 2020); up to 1.5 L/kg has been reported in extremely premature neonates (Anderson 2002).

Children: V_{ss} : 1.2 ± 0.3 L/kg.

Adolescents: V_{ss} : 1.1 ± 0.3 L/kg.

Adults: V_{ss} : 0.8 ± 0.2 L/kg.

Protein binding: 10% to 25% at therapeutic concentrations; 8% to 43% at toxic concentrations.

Metabolism: At normal therapeutic dosages, primarily hepatic metabolism to sulfate and glucuronide conjugates, while a small amount is metabolized by CYP2E1 to a highly reactive intermediate, N-acetyl-p-benzoquinone imine (NAPQI), which is conjugated rapidly with glutathione and inactivated to nontoxic cysteine and mercapturic acid conjugates. At toxic doses (as little as 4 g daily), glutathione conjugation becomes insufficient to meet the metabolic demand causing an increase in NAPQI concentrations, which may cause hepatic cell necrosis. Neonates (especially preterm neonates) demonstrate higher sulfate metabolites and lower glucuronide metabolites compared to adults (Flint 2017). Oral administration is subject to first-pass metabolism.

Bioavailability:

Oral: Infants and Children <6 years: ~72% (range: 11% to 91%) (Kleiber 2019).

Half-life elimination: Prolonged following toxic doses.

Neonates (Van Lingen 1999):

GA 28 to 32 weeks: 11 ± 5.7 hours (range: 3.5 to 25.2 hours).

GA 32 to 36 weeks: 4.8 ± 1.2 hours (range: 3.6 to 6.8 hours).

Infants and Children <2 years: Median range: 2.4 to 2.8 hours (range: 1.2 to 5.4 hours) (Zuppa 2011).

Children 2 to <12 years: Median range: 2.6 to 2.8 hours (range: 2.2 to 4.9 hours) (Zuppa 2011).

Adolescents: 2.9 ± 0.7 hours (manufacturer's labeling).

Adults: 2.4 ± 0.6 hours (manufacturer's labeling); may be slightly prolonged in severe renal insufficiency (CrCl <30 mL/minute): 2 to 5.3 hours.

Time to peak, serum:

Oral: Immediate release: Adults: 10 to 60 minutes (may be delayed in acute overdoses).

IV:

Infants and Children <2 years: Median range: 0.25 to 0.29 hours (range: 0 to 1.4 hours) (Zuppa 2011).

Children 2 to <12 years: Median range: 0.18 to 0.27 hours (range: 0 to 0.75 hours) (Zuppa 2011).

Children ≥ 12 years and Adolescents: Median range: 0.25 to 0.33 hours (range: 0 to 0.4 hours) (Zuppa 2011).

Adults: 15 minutes.

Rectal:

Neonates (van Lingen 1999):

GA 28 to 32 weeks: Median: 3.9 hours.

GA 32 to 36 weeks: Median: 5.1 hours.

Infants and Children: 2.37 ± 1.1 hours (range: 0.43 to 5.26 hours) (Hahn 2000).

Excretion: Urine (<5% unchanged; 60% to 80% as glucuronide metabolites; 20% to 30% as sulfate metabolites; ~8% cysteine and mercapturic acid metabolites).

Pharmacokinetics: Additional Considerations (Adult Data Unless Noted)

Hepatic function impairment: The half-life may increase 2-fold or more in patients with liver disease.

Patient Counseling Points

(For additional information see "Acetaminophen (paracetamol): Patient drug information")

What is this drug used for?

- It is used to ease pain and fever.

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Nausea
- Vomiting
- Trouble sleeping

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Liver problems like dark urine, fatigue, lack of appetite, nausea, abdominal pain, light-colored stools, vomiting, or yellow skin.
- Unable to pass urine
- Change in amount of urine passed
- Stevens-Johnson syndrome/toxic epidermal necrolysis like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in mouth, throat, nose, or eyes.
- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

Note: This is not a comprehensive list of all side effects. Talk to your doctor if you have questions.

Consumer Information Use and Disclaimer: This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a limited summary of general information about the medicine's uses from the patient education leaflet and is not intended to be comprehensive. This limited summary does NOT include all information available about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not intended to provide medical advice, diagnosis or treatment and does not replace information you receive from the healthcare provider. For a more detailed summary of information about the risks and benefits of using this medicine, please speak with your healthcare provider and review the entire patient education leaflet.

Brand Names: International

International Brand Names by Country

For country code abbreviations (show table)

- (AE) United Arab Emirates: Adol | Adol one | Alceta | Apotel max | Atasol | Calpol | Camdol | Dafalgan | Dolo | Efferalgan | Enelfa | Fanalgic | Fevadol | Omol | Panacare | Panaadol | Panadol actifast | Panadol advance | Panadrex | Panalife | Parafusiv | Paralgin | Parasustain | Perfalgan | Revanin | Tempra | Tylenol;
- (AR) Argentina: Acetagen | Acetolit | Causalon | Cestropan | Custodial | Dirox | Dolex ninos | Fabogesic aliv | Fada paracetamol | Fiebro | Fiebrolex | Fiebrolex forte | Fiebrolyto | Geniol | Geniol forte | Itedal | Mejoral | Mejoral para ninos | Multifebrin | Nodipir | Nodolex | Panadol | Paraz mol cabuchi | Paracetamol b. braun | Paracetamol b.braun | Paracetamol bayer | Paracetamol fecofar | Paracetamol fortbenton | Paracetamol isa forte | Paracetamol kabi | Paracetamol labsa | Paracetamol lafedar | Paracetamol larjan | Paracetamol lazar | Paracetamol monserratt y eclair | Paracetamol northia | Paracetamol puntanos | Paracetamol raffo | Paracetamol raffo ninos gotas | Paracetamol sant gall friburg | Paracetamol schaefer | Paracetamol schaefer | Paracetamol sintesina | Paracetamol teva | Paracetamol vannier | Paracetamol vent 3 | Paracetamol Vent-3 | Paracetamol walker | Paracetamolito | Paragrivdol | Paratral | Paraxamol | Parclen | Plovacal | Predualito | Reanima | Rolfit | Tafirol | Tafirol 1 g | Tafirol forte | Tafirol pediatrico | Tafirolito | Termofren | Tetradox | Tylenol | Vent 3 flash | Viclor | Vivita forte paracetamol;
- (AT) Austria: Ben u ron | Duaneo | Gewadal pamol | Mexalen | Mexaratio | Momentum analgetic | Paracetamol accord | Paracetamol actavis | Paracetamol b. braun | Paracetamol genericon pharma | Paracetamol genfarma | Paracetamol hospira | Paracetamol kabi | Paracetamol roesch u handel | Paracetamol sandoz | Parakapton | Perfalgan;

- (AU) Australia: Aft pharmaceuticals paracetamol | Ah paracetamol xr | Amcal children's paracetamol elixir for 1 to 5 years | Amcal children's paracetamol suspension for 6 to 12 years | Amcal paracetamol | Amcal paralieve | Apo osteo paracetamol | Apo paracetamol | Apo paracetamol xr | Apohealth children's paracetamol | Apohealth children's paracetamol 1 month to 2 years | Apohealth children's paracetamol 6 to 12 years | Apohealth osteo relief paracetamol | Apohealth paracetamol | Apohealth paracetamol pain relief | Avallon paracetamol | Avallon paracetamol for children | Avallon paracetamol for children 1 month to 1 year | B. braun paracetamol | Benylin Sore Throat | Black & gold paracetamol | Blooms the chemist colourfree children's concentrated paracetamol suspension 5-12 years | Blooms the chemist colourfree paracetamol drops infants and children | Blooms the chemist paracetamol | C paracetamol | Ca children's paracetamol 1-5 years | Ca children's paracetamol 6-12 years | Ca-paracetamol | Chemist's own osteo relief paracetamol | Chemists own paracetamol | Chemists own paracetamol pain & fever | Chemists' own children's paracetamol pain and fever suspension 1 to 5 years | Chemists' own children's paracetamol pain and fever suspension 5 to 12 years | Chemists' own osteo relief paracetamol | Chemists' own paracetamol | Chemists' own paracetamol pain & fever drops 1 month - 2 years | Chemists' own paracetamol pain & fever drops 1 month to 2 years | Chemists' own paracetamol pain relief | Chemmart paracetamol | Chemmart pharmacy osteo relief paracetamol | Chemplus osteo relief paracetamol | Chemplus paracetamol | Chlidren's panadol colourfree baby | Cipla australia pain relief para | Cipla pain relief for kids paracetamol | Cipla pain relief for little kids paracetamol | Cipla pain relief paracetamol | Cipla paracetamol | Cm chemadol | Cm paracetamol | Cold & flu | Coles paracetamol | Combolieve day & night pain relief | Comfarol | Cw painex | Discount drug stores paracetamol | Dismol | Duatrol | Dymadon | Dymadon for babies | Dymadon for kids | Dymadon p | Febridol | Febridol suspension for children 6 to 12 years | Febridol syrup for children 1 to 5 years | Fedimol | Genrx paracetamol | Gesimol | Gold cross paracetamol | Hc painex | Hedanol | Hedanol paracetamol | Help@hand for pain relief paracetamol | Herron gold paracetamol | Herron paracetamol | Home brand paracetamol | I paracetamol | Ip paracetamol | Ipca paracetamol | Ipmol | Logicin pain & fever | Logicin pain and fever infant | Medichoice paracetamol | Medihealth paracetamol | Medirelieff paracetamol | Medix paracetamol | Medreich paracetamol | Mendeleev children's 1 month-2 years paracetamol | Neo paracin | Neo paramol paracetamol | Neocombo day & night pain relief | Noumed paracetamol for children 1 to 5 years | Noumed paracetamol for children 5 to 12 years | Nova paracaps | Nova paracetamol | Nyal paracetamol | O/n paracetamol | Osteomol | P paracetamol | Pa painex | Pacimol | Pacimol i | Pacimol paracetamol | Pain & fever | Pain relief | Pain relief osteo paracetamol | Panadol | Panadol back & neck long lasting | Panadol children | Panadol children 1 month - 1 year | Panadol children 1-5 years | Panadol children 5-12 years | Panadol children 5-12 years elixir | Panadol children 6 months - 5 years | Panadol children chewable 3+ years | Panadol children chewable 7+ years | Panadol children suppositories 5 to 12 years | Panadol children suppositories 6 months to 5 years | Panadol Optizorb | Panadol osteo | Panadol rapid | Panamax | Panamax elixir | Parabex osteo | Paracet | Paracetamol aft | Paracetamol alphapharm | Paracetamol apotex | Paracetamol bnm | Paracetamol genpar | Paracetamol gppl | Paracetamol kabi | Paracetamol osteo | Paracetamol pain relief | Paracetamol pfizer | Paracetamol sandoz;
- (AU) Australia: Paracetamol sandoz pharma | Paracetamol watson | Paradyn | Parahexal | Paralgin | Paralgin paracetamol | Paraosteo | Parapaed 1 to 5 years | Parapaed six plus | Parapane | Parmol | Perfalgan | Pharmacist children's paracetamol | Pharmacist formula paracetamol | Pharmaction paracetamol | Pharmacy action osteo relief | Pharmacy action paracetamol | Pharmacy action paracetamol osteo | Pharmacy care paracetamol | Pharmacy choice children's paracetamol 1 month to 2 years infant | Pharmacy choice children's paracetamol suspension 1-5 years | Pharmacy choice children's paracetamol suspension concentrated 5-12 years | Pharmacy choice children's paracetamol suspension concentrated 6-12 years | Pharmacy choice children's paracetamol syrup 1-5 years | Pharmacy choice osteo relief | Pharmacy choice paracetamol | Pharmacy health paracetamol | Pharmacy health paracetamol colour free concentrated suspension 5-12 years | Pharmacy health paracetamol colour free infant | Pharmacy health paracetamol colour free suspension 1-5 years | Pharmacy health paracetamol for children 1 month to 1 year | Pharmacy health paracetamol osteo | Pharmacy health paraosteo | Pharmacy own paracetamol | Priceline paracetamol | Priceline pharmacy paracetamol | Priceline pharmacy paracetamol

- colour free concentrated suspension 5-12 years | Priceline pharmacy paracetamol colour free infant | Priceline pharmacy paracetamol colour free suspension 1-5 years | Priceline pharmacy paracetamol for children 1 month to 1 year | Priceline pharmacy paraosteo | Quimol | Quiphen | Reddycet paracetamol | Sbpa paracetamol | Setamol | Soul pattinson paracetamol | Soul pattinson paracetamol colour free | Soul pattinson paracetamol for children 1 month to 1 year | Soul pattinson paraosteo | Spiro lifecare paracetamol | Tactinal | Tempra | Terry white chemists osteo relief paracetamol | Terry white childrens paracetamol | Terrywhite chemmart para osteo | Terrywhite chemmart paracetamol | Terrywhite chemmart paracetamol colour free concentrated suspension 5-12 years | Terrywhite chemmart paracetamol colour free infant | Terrywhite chemmart paracetamol colour free suspension 1-5 years | Terrywhite chemmart paracetamol for children 1 month to 1 year | Trust for kids paracetamol | Trust for kids paracetamol 1 to 5 years | Trust for kids paracetamol 6 to 12 years | Trust paracetamol xr | Trust pharmadol | Twp paracetamol | Tylenol | Tylenol for babies 1 month - 2 years | Tylenol for children | Tylenol for children 2+ years | Vidalife para original | Wagner health children's paracetamol 1-5 years | Wagner health children's paracetamol 5-12 years | Wagner health liquigesic paracetamol | Wagner health paracetamol | Woolworths essentials paracetamol | Your pharmacy paracetamol;
- (BD) Bangladesh: 2 A | Ace | Ace power | Aceta | Act | Actol | Adcetamol | Alkaparol | Ambeedol | Ampol | Analpain | Analpyrin | Anapol | Antap | Apa | Apedol | Asmol | At98 | Atopen | Atp | Atp xr | Avlopol | Biocetamol | Centamol | Cetadol | Cetajap | Cetajol | Cetal | Cetalgin | Cetam | Cetol | Cipladol | Depol | Depyrin | Disprol | Doliprane | Fap | Fast | Fea | Fendol | Feva | Fevac | Fevimol | Fibi | Fitamol | Flamodol | G-paracetamol | Genamol | Getamol | Getol | Hepa | Ispha | Knock | Lapain | Longpara | M pol | Micetamol | Napa | Napa one | Napa rapid | Navamol | Nebs | Nova | Noval | Orpa | Painil | Pam | Pamidol | Pamix | Pana | Panacin | Para | Para c | Paracid | Paracil | Paracit | Paracitamol | Paracitol | Paraciv | Parafen | Parafof | Paramex | Paramin | Paramol | Parapain | Parapol | Parapyrol | Parasis | Paratem | Paratex | Paraxia | Pol | Preduet | Prefer | Promel | Pyrac | Pyrapol | Pyrin | Qap | Qcet | Remalgin | Renova | Renova max | Reset | Sb opa | Seridol | Servigesic | Silpa | Silpol | Sinapol | Skymol | Tamen | Tamen ultra | Tamino | Tamol | Temfal | Tempac | Tempil | Tempol | Tendamol | Timidal | Tydenol | Tylen | Tynol | Ucet extend | Xcel | Xpa | Zerin;
 - (BE) Belgium: Algostase mono | Ben u ron | Curpol | Dafalgan | Dafalgan Caps Forte | Dafalgan pediatrique | Dolprone | Efferalgan | Kruidvat paracetamol | Lemgrip | Lonarid Mono | Panaadol | Paracaps | Paracetabs | Paracetabs forte | Paracetamol ab | Paracetamol accord | Paracetamol actavis | Paracetamol b. braun | Paracetamol eg | Paracetamol eurogenerics | Paracetamol fresenius kabi | Paracetamol macopharma | Paracetamol mylan | Paracetamol sandoz | Paracetamol teva | Paracetamol teva generics belgium | Paraphar | Pe-tam | Perdolan | Perfusalgan | Sanicopyrine;
 - (BF) Burkina Faso: Apramol | Doliprane | Efferalgan | Efferalgan pediatrique | Efferalgantab | Intralgine | Nesteffeffer | Panadol enfant et nourrisson | Panol | Para denk | Paraceta m | Paracetamol b. braun | Paracetamol tm | Paracetamol ubigen | Parexcel | Paromax | Perfalgan | Philco para | Xykaa rapid;
 - (BG) Bulgaria: Acetophen | Apap | Calpol | Dafalgan | Daleron | Efferalgan | Fervex baby | Panaadol | Panadol baby | Paracedol | Paracerol pf | Paraceta | Paracetamax | Paracetamax express | Paracetamol accord | Paracetamol actavis | Paracetamol danhsen | Paracetamol farma | Paracetamol kabi | Paracetamol panpharma | Perfalgan | Sanidol | Tydol | Tydol 6+;
 - (BR) Brazil: Abidor | Abidor baby | Abidor crianca | Acetam | Acetamil | Acetamol | Acetitol | Acetofen | Aliviol | Ameniflac | Aminofen | Analgisen | Anatyl | Benegrip multi febre e dor | Braim | Bwell paracetamol | Calpol | Cefabrina | Cefalium one | Cetafrin | Cimegripe 77c | Cimegripe bebe | Cimegripe dor e febre | Cristina termus | Cyfenol | Cyfenol baby | Din | Dorfen | Dorfenol | Dorib | Dorico | Dorsanol | Fenalgin | Furp paracetamol | Gripalce uno | Gripotermom | Halaxminophen | Iquego paracetamol | Lafepe paracetamol | Lfm paracetamol | Multigrip dor e febre | Naldecon febre | Naldecon febre bebe | Naldecon febre infantil | Pacemol | Paracen | Paracetamol baby | Paracetamol cinfa | Paracetamol crianca | Paracetrex | Parador | Paraflan | Paralgen | Paramol | Paratermol | Pratum | Resfenax gripe uno | Resfenol thermus | Resfenol thermus bebe | Sonridor dor e febre | Sonridor rapid | Stopain | Termo | Termol | Thylom | Tilekin | Tilemax baby and child | Tilemaxy | Tilemaxy baby | Tilemaxy crianca | Tilenati | Trifen | Tyflen | Tyflen baby | Tyflen bebe | Tyflen crianca | Tyflaflex | Tyflalgin | Tylenax | Tylenax baby | Tylenax crianca | Tylenol | Tylenol

- bebe | Tylenol crianca | Tylephen | Tyolidol | Tylol | Tyramol | Zuplyn;
- (BZ) Belize: Acetaminofen | Acetaminofen bayer | Mencetamol | Panaadol baby | Panaadol children | Panadol extra fuerte;
 - (CH) Switzerland: Acetalgin | Amavita paracetamol | Becetamol | Ben u ron | Contra-schmerz p | Coop vitality paracetamol | Dafalgan | Dafalgan kinder | Doloran | Dolprone | Kafa | Panadol | Panadol extend | Panadol s | Paracetafelan | Paracetamol actavis | Paracetamol apotel | Paracetamol axapharm | Paracetamol b. braun | Paracetamol fresenius | Paracetamol grunenthal | Paracetamol haenseler | Paracetamol mepha | Paracetamol q generics | Paracetamol sandoz | Paracetamol sintetica | Paracetamol Spirig HC | Paracetamol UPSA | Paracetamol zentiva | Paraconica | Perfalgan | Sun store paracetamol | Treuphadol | Tylenol | Tylenol forte | Zolben;
 - (CI) Côte d'Ivoire: Algesic | Algesic 1000 | Apramol | Doliprane | Dolyc | Efferalgantab | Intralgine | Nesteffe | Novadol | Panaadol advance | Panalife | Panol | Paraceta m | Paracetamol kabi | Paracetamol macopharma | Paracetamol winthrop | Paracetawal | Paralgy | Paraveine | Paraven | Parcedol | Parol | Pharmalgan | Pmol | Pyrexal | Xykaa rapid | Zetifen;
 - (CL) Chile: Acamol | Algiafin | Alividol | Bufferin | Bufferin forte | Cotibin analgesico antipiretico | Cotibin infantil | Dolo esan | Dolsom | Dropol | Dropol infantil | Fibrinol | Geniol | Gesidol | Imitinol | Kendol | Kitadol | Kitadol infantil | Panaadol | Panagesic | Pinnacin | Rapidol | Sinflu | Supracalm | Tapsin | Tapsin infantil | Tapsin sin cafeina infantil | Tempra | Winasorb | Xumadol | Zolben;
 - (CN) China: Ai er xing | Ai sen | An yi | Bai fu ning | Bi li tong | Children's Bufferin | Er he yi | Fortolin | Jin fu ning | Panaadol | Pu le er | Tai nuo lin | Tylenol | Yi shang;
 - (CO) Colombia: Acelifen | Aceradox | Acetafen | Acetaminodelt | Acetaminofen | Acetaminofen la sante | Acetaminofen mk | Acetaminofen rapid gel | Acetanovafen | Adorem | Adorem baby | Adorem ninos | Algisil | Ametrex | Aminofen | Analper | Analper forte | Atacadol | Atalmax | Catalac | Cronofen | Dixicol | Dolex | Dolex activgel | Dolex avanzado | Dolex bebes | Dolex ninos | Dolicox | Dolinar | Dolofen | Dolotrin | Dolpirin a | Dolpirin a junior | Dolpirin a ninos | Dristandol | Febrex | Fiemol | Firexifen | Firexyfen | Geniol | Geniol ninos | Gledfen | Kyofen | Mejoral | Mejoralito | Meiorflex | Nadorex | Neofen | Octalgina | Panadol | Paracemiv | Paracetamol denk | Paracetamol kabi | Paragrivdol | Paramol | Parmodol | Pedialab | Prodol | Prodol m | Prodol r | Punzadol | Sanafen | Sevedolito | Supofen | Synfen | Tempalgin | Tempra | Teralgex | Tylenol | Winadol;
 - (CR) Costa Rica: Acetaminofen | Acetaminofen chemo | Acetaminofen forte | Acetaminofen genfar | Acetaminofen mk | Asingesic | Az minofen | Bebetina | Cetamin | Da pharma acetadol | Mencetamol | Nafin | Panadol bebe | Panadol extra fuerte | Panadol ninos | Paracetamol bp | Paracetamol kabi | Paraconica | Tylenol | Tylex | Winasorb | Xypar rapid;
 - (CZ) Czech Republic: Apo acetaminophen | Ben u ron | Dafalgan | Detsky antibol | Efferalgan | Medipyrin | Mexalen | Panadol novum | Paracetamol accord | Paracetamol agmed | Paracetamol apotex | Paracetamol aurovitas | Paracetamol b. braun | Paracetamol basi | Paracetamol dr. max | Paracetamol kabi | Paracetamol zentiva | Paralen sus | Paralgil | Paramax | Paramegal | Parapyrex | Tylenol;
 - (DE) Germany: Abalon | Apiredol | Apogen paracetamol | Apovital apocetamol | Atida paracetamol | Ausbuettels paracetamol | Ben u ron | Curadies paracetamol | Docmorris paracetamol | Docpelin paracetamol | Doloreduct | Duracetamol | Enelfa Dr. Henk | Gib paracetamol | Grippex | Medicon paracetamol | Medpex paracetamol | Mein paracetamol | Mono praecimed | Paedialgon | Paraceta | Paracetalgin | Paracetamol 1 A Pharma | Paracetamol actavis | Paracetamol adgc | Paracetamol aiwa | Paracetamol al | Paracetamol amneal | Paracetamol apogen | Paracetamol axicur | Paracetamol b.braun | Paracetamol bene | Paracetamol dexcel | Paracetamol dura | Paracetamol elac | Paracetamol hormosan | Paracetamol ipa | Paracetamol ipa/apo rot | Paracetamol ipa/ba | Paracetamol IPA/Dr Sandman | Paracetamol ipa/klindwort | Paracetamol ipa/mache | Paracetamol ipa/medibond | Paracetamol ipa/medpex | Paracetamol kabi | Paracetamol l | Paracetamol OPT | Paracetamol rotexmedica | Paracetamol sanavita | Paracetamol sandoz | Paracetamol tad | Paracetawal | Paramol bene | Parapaed | Paraveganio | Pyromed s | Rubiemol | Sinpro n | Tempil paracetamol | Tegal paracetamol | Unsere paracetamol schmerztabletten | Vivimed N;
 - (DK) Denmark: Panodil zapp | Pinemol;
 - (DO) Dominican Republic: Acetagel | Acetamin | Acetaminofen | Acetaminofen Feltrex | Ac-

- etaminofen lam | Acetaminofen mk | Aceteff | Acro temp | Adorem | Alidol | Amolgin | Apamide | Ben u ron | Bioacelifen | Chemifen | Cymatem | Dalgen | Danolux | Defensol junior | Dispamol iqi | Dolenol | Dolfenol | Doloxin | Efferalgan | Eslofem | Febronol | Fidol | Inalgex | Laproter | Melabon infantil | Minofen | Natrafen | Neomol | Notem | Painmax forte | Panadol | Parabay | Piralmin | Pirogesic | Pitirin | Sinedol | Temperal | Tempra | Tylenol | Tylex | Ultrafen | Ultramol | Winasorb;
- (EC) Ecuador: Acestar | Acetagen | Acetamin | Acetaminofen | Acetaminofen genfar | Acetaminofen nifa | Actifen | Alginox | Alividol | Analgan | Analgan rapid | Analpax | Aproxol | Apyral | Axapara | Bephen | Beralfen | Betamol | Calmadol | Cetamihelab | Dolmedica | Dolorfin | Dolorol | Dolotamol | Elbrus | Elbrus flash | Exedol | Febedol flash | Febedol forte | Feverach | Feveril | Fiebrerol | Finalin | Finalin junior | Finalin ninos | Frizac | Frizac pediatricas | Geniol | Ibupaxx | Julphadol | Kabimol | Kro termofin | Kro termofin plus | Mejoral | Milfen | Molzetan | Napafen | Napafen sin alcohol | Neomol | Nexamol | Normotemp | Opemol | Pana  ol | Paracerol | Paracetadol rapid gel | Paracetamol ariston | Paracetamol h g | Paracetamol labovida | Paracetamol mk | Paracetamol nf | Paracetamol nifa | Paraconica | Paralfin | Paralgen | Paramidol | Paranir | Raserflash | Sinalgia | Sodifen | Sodifen ninos | T brax | Tempdol letic | Temperyl | Tempra | Temproxial | Termax | Termax ped | Termax plus | Termifen | Termo out | Termofin | Termofin forte | Termofin plus | Tylenol | Umbral | Umbramil md | Winadol;
 - (EE) Estonia: Calpol | Dafalgan | Efferalgan | Grippostad | Lekadol | Medipyrin | Mexalen | Pamol | Panadol | Panadol Optizorb | Pana  ol rapid | Paracetamol accord | Paracetamol arena | Paracetamol b. braun | Paracetamol kabi | Paracetamol Sopharma | Paracetamol stada | Paracetamol zentiva | Paraflux | Paramax | Perfalgan | Pinex | Rivalgyl | Tachipirina | Tylenol;
 - (EG) Egypt: Abimol | Acetaminophen | Adol | Calpol | Cetal | Cetamol | Doliprane | Eupifalgan | Febrimol | Fevano | Gesicrest | Glasbomol | Injectmol | Novaldol | Paracetamol europe | Paracetamol pharco | Paracpimol | Paragesic | Paragesic baby | Paramol | Pentamol | Perfalgan | Pyral | Pyremol | Pyritrust | Rotapyretic | Stopadol | Targecetal | Tempra | Thera lo | Unicetamol | Vedraphenal;
 - (ES) Spain: Abezen | Actron | Antidol | Antidol infantil | Apiredol | Apiretal | Bandol | Calmanticold | Calthek forte | Cupanol | Cupanol adultos | Dafalgan | Difenatil | Difenatil infantil | Dolgesic | Dolocatil | Dololavi | Dolostop | Dolostop pediatrico | Duorol | Efetamol | Efferalgan | Febrectal | Febrectal adultos | Febrectal ninos | Febrovir | Gelocatil | Gelocatil pediatri | Melabon infantil | Pana  ol | Paracetam mundogen | Paracetamol abamed | Paracetamol accord | Paracetamol actavis | Paracetamol alter | Paracetamol apotex | Paracetamol aristo | Paracetamol aurovitas | Paracetamol b.braun | Paracetamol benel | Paracetamol cinsa | Paracetamol cipla | Paracetamol combino pharm | Paracetamol combix | Paracetamol curaxys | Paracetamol cuve | Paracetamol dermogen | Paracetamol edigen | Paracetamol esteve | Paracetamol farmalid | Paracetamol G.E.S. | Paracetamol galenicum | Paracetamol galenicum vitae | Paracetamol gelos | Paracetamol grifols | Paracetamol interquim | Paracetamol kabi | Paracetamol kern | Paracetamol kern pharma | Paracetamol krka | Paracetamol level | Paracetamol mabo | Paracetamol mabo farma | Paracetamol mundogen | Paracetamol mylan | Paracetamol neuraxpharm | Paracetamol normon | Paracetamol onedose | Paracetamol Pensa | Paracetamol pensavital | Paracetamol pharma combix | Paracetamol pharmagenus | Paracetamol qualigen | Paracetamol qualix | Paracetamol ratio | Paracetamol sandoz | Paracetamol sandoz care | Paracetamol sandoz farmaceutica | Paracetamol serra | Paracetamol serraclinics | Paracetamol stada | Paracetamol tarbis | Paracetamol tarbis farma | Paracetamol tecnigen | Paracetamol tecnimede | Paracetamol teva | Paracetamol Tevagen | Paracetamol vir | Perfalgan | Supofen | Termalgin | Tigetemel;
 - (ET) Ethiopia: Asmol | Cadimol | Febramol | Kelvin | Lotemp | Milzor | Paingo | Pana  ol advance | Pana  ol baby & infant | Para denk | Paracerol | Paramol | Polaminofen | Remedol;
 - (FI) Finland: Arax Junior | Pamol | Pana  ol | Panadol novum | Para | Para caps | Para tabs | Paraceon | Paracetamol accord | Paracetamol actavis | Paracetamol b. braun | Paracetamol fresenius kabi | Paracetamol krka | Paracetamol panpharma | Paracetamol ratiopharm | Paracyl | Paramax | Paramax rap | Paramin | Parasetamoli ya | Perfalgan | Pinex | Polarfen | Pyritol | Rolod | Salpara;
 - (FR) France: Claradol | Dafalgan | Dafalgancaps | Dafalgantabs | Doliprane | Dolipranecaps | Dolipranelib | Dolipraneliquiz | Dolitabs | Dolko | Dolstic | Efferalgan | Efferalganmed | Efferalganmed pediatrique | Efferalgantab | Expandox | Geluprane | Gynospasmine | Libradol | Pana-

- dol | Paracetamol accord | Paracetamol actavis | Paracetamol aguettant | Paracetamol almus | Paracetamol alter | Paracetamol arrow | Paracetamol b.braun | Paracetamol bayer | Paracetamol benta | Paracetamol biogaran | Paracetamol cristers | Paracetamol eg | Paracetamol EGN | Paracetamol evolugen | Paracetamol g gam | Paracetamol gnr | Paracetamol grunenthal | Paracetamol Hexal | Paracetamol irex | Paracetamol ivax | Paracetamol kabi | Paracetamol krka | Paracetamol lucien | Paracetamol macopharma | Paracetamol Merck | Paracetamol qualimed | Paracetamol ranbaxy | Paracetamol ranbaxy medication officinale | Paracetamol ratiopharm | Paracetamol rpg | Paracetamol sandoz | Paracetamol teva | Paracetamol zentiva | Paracetamol zydus | Perfalgan;
- (GB) United Kingdom: Alvedon | Anadin paracetamol | Apamide | Calpahalal | Calpol | Calpol six plus | Care Infant paracetamol | Cupanol | Disprol | Disprol Paracetamol | Disprol Soluble Paracetamol | Fanalgic | Galpamol | Galpharm paracetamol | Hedex | Infadrops | Mandanol | Medinol | Miradol | Mrs cullens | Obimol | Paldesic | Panaadol | Panaadol actifast | Panadol advance | Panadol solid | Panaleve | Panets | Paracetamol accord | Paracetamol almus | Paracetamol arrow | Paracetamol berk | Paracetamol cox | Paracetamol cupal | Paracetamol kent | Paracetamol sfp | Paracets | Paracets plus | Paraclear | Paramin | Parapaed | Paravict | Placidex | Salzone | Sootheze six plus | Tabalgin | Ticegesic | Tixymol;
 - (GH) Ghana: Primadol | Zinel;
 - (GR) Greece: Algocit | Anadin | Apotel | Apotel max | Condalair | Dalminette | Depon | Depon maximum | Dolal | Lonarid aplo | Panaadol | Panaadol actifast | Panadol advance | Paracetamol teva | Paracetamol/kabi | Protalgon;
 - (GT) Guatemala: Acetaminofen | Acetaminofen balaxi | Acetaminofen bayer | Acetaminofen therfam | Acetaminofen therfarm | Avamigran extra forte | Axapara | Cetamin | Dolenfant | Feverach | Nanmol | Temptra | Temptra forte | Tialgin | Zetawin;
 - (HK) Hong Kong: Acetaminophen | Acetamol | Anapar | Anaphen | Apap | Axcel paracetamol | BF Paradac | Biogesic | Calpol | Christamol | Christamol 2 | Dhamol | Dolorol | Endopain | Endopain II | Europain | Fortolin | Kenamose | Konaton | Lycogesic | Madame pearl's mp paracetamol | New eurogesic | Panaadol | Panadol back&neck | Panadol extend | Panaadol joint | Paracetamol aft | Paracetamol b | Paracetamol b braun | Paracetamol cinfa | Paracetamol o | Paramol | Paramolan | Paraway | Parmol | Paromon | Partamol | Perfalgan | Pharmadol | Progesic | Promol | Pymadon | Pyrudon | Qualimol | Serimol | Setamol | Tylenol | Uni-Pamol | Wintamol | Xykaa extend;
 - (HN) Honduras: Acetan;
 - (HR) Croatia: Daleron | Efferalgan | Fibralgin | Imensa | Lekadol | Lekadol Forte | Lupocet | Lupocet BABY | Lupocet forte | Lupocet junior | Lupocet teen | Panaadol | Panaadol baby | Panaadol Optizorb | Paracetamol accord | Paracetamol Jadran | Paracetamol kabi | Paracetamol PharmaS | Perfalgan | Plicet | Plicet rapid;
 - (HU) Hungary: Ben u ron | Doloramol | Febrilin | Mexalen | Panaadol | Panaadol baby | Panadol rapid | Paracetamol accord | Paracetamol actavis | Paracetamol bril | Paracetamol kabi | Paracetamol onkogen | Paracetamol pxgpharma | Paracetamol sandoz | Paramax forte | Perfalgan | Piropill | Roxacet | Rubophen;
 - (ID) Indonesia: Alphamol | Apetic | Betamol | Bodrexin demam | Casetamol | Cetamol | Cetapain | Contrexyn demam | Cupanol | Dapyrin | Dumin | Ekacetol | Erlamol | Etagesic | Eterfix | Farmadol | Fasidol | Fermol | Fevrin | Grafadon | Gunaceta | Hopadol | Hufagesic | Hufagripp demam | Ikacetamol | Itamol | Kamolas | Maganol | Mesamol | Metamol | Mirasic | Moretic | Naprex | Neo rheumacyl aktif | Nipe fever | Nufadol | Obat penurun panas | Ottopan | Pamol | Panaadol | Panadol actifast | Paracetamol phapros | Paracetol | Paraco | Paramol | Pehamol | Piosfen | Pireta | Poro | Praxion | Prodenas | Progesic | Pyrex | Pyrexin | Pyridol | Samconal | Sanmol | Selesmol | Sumagesic | Tamoliv | Temptra | Termagon | Termorex | Trifamol | Tylenol | Unibebi demam | Xepamol | Zetamol;
 - (IE) Ireland: Alvedon | Anadin paracetamol | Calpol | Calpol six plus | Calpol six plus sugar | Calpol sugar free | Disprol | Hedex | Medinol | Panadol | Panadol actifast | Panadol advance | Panaadol original | Panaleve | Paracetamol actavis | Paralief | Paralink | Paralink six plus | Parapaed | Paratabs | Perfalgan | Rimadol | Rowalief;
 - (IL) Israel: Abrol | Acamol | Acamoli | Aldolor | Maccabimol kid | Panadol | Paracet | Rokamol | Tiptipot novimol;
 - (IN) India: 37 C | 99f | Acetaminophen | Aceto | Acton | Acton or | Admol | Afeb | Aladol | Aldomol

- | Algina | Alice | Aminol | Amipar | Amipol | Amol | Anamol | Anemol | Anetol | Arden | Aristomol | Aterpa | Bactpar | Bepamol | Betmol | Biocetamol | Biopyrin | Brikamol | C 37 | Cachpar | Cadipar | Cadpar | Calpol | Calridol | Cemol | Centamol | Cepal | Ceta | Cetafen | Cetamol | Chilmol | Cipmol | Cofamol | Compain | Coolant | Crocin advance | Cropara | Cruzmol | Decomb i.v. | Detemp | Dispar | Disprin Paracetamol | Disprin plus | Docin | Dolcera | Doliprane | Doliza | Dolo | Dolo 650 | Doloan | Dologlan | Dolomark | Dolopar 650 | Eboo | Ecocin | Elpar | End ache | Essmol | Eupyrlic | Espresso | Expres | F norm | Fabrimol | Febrex | Febrimol kid | Febrinil | Fenace | Fepamol | Fepanil | Feudamol | Fev r | Feva | Fevanex | Fevanex rapid | Fevanil | Fevastin | Fevecet | Fever x | Fevridol | Fibromet | Finamac | Flumol | Fnll | Frutimol | Galmol | Genidol | Glitmol | Gratacin | Gufimol | Healthy para | Ibibull | Ifimol | Indamol | Infupar | Iphamol | Ivifen | Jacphen | Jagcin | K 37 forte | Kabimol | Kabipara | Kaldol | Kelvin | Kelvinil ds | Kemcin | Lanol | Larkin | Lit king | Lofe | Lotemp | Lupicin | Lupipara | Macfast | Malidens | Medimol | Medomol | Meprocin | Meridol | Metacin | Metalgin | Metaljin | Metamol | Metaplus | Mino | Mol | Mol plus | Mypar | Mypol | Nacmol | Nemflam | Neomol | Neximol | Neximol ds | Nicetamol | Nimol | Nofiva | Nowaste para | Nu 650 | Opara | Opimol | Optimol | Orangemol | Oromol | Oromol a | Orvomol | P 100 | P 120 suspension | P 125 | P 250 | P 37 | P 500 | P 650 | P 750 | P mol | Pa 12 | Paarmol | Pacemo | Pacetamol | Pacimol | Pacinova | Pactiv | Pamol moxy | Panacip | Para ge | Para para | Parabig | Paraboot | Paracetamol apex | Paracetanal rectal supp | Paracin | Paracip | Paracon | Parafuse | Paraglass | Paraglow | Paragon | Paragreat | Paragull | Parahenz | Paralife | Paraluc | Paramax | Paramet | Paramist | Paramol | Paranir | Parapin | Parapol | Paraprime | Paraquick | Parareach | Pararose | Parasafe | Parasoft | Paratee | Paratop | Paratru | Parawel | Parazen | Parazex | Parazine | Parry | Paxol | Pcbig | Pcm | Pedmol | Peemol | Penodin | Perfalgan | Peridyl | Pfc | Pinex | Pinkbery | Pm | Pmo | Pmson | Polo | Polymol | Primol | Primosul | Profuse | PUC | Pymol | Pyracron | Pyrakem | Pyremol | Pyrexin | Pyrexon | Pyricare | Pyrigesic | Pyriped | Pyrit | Racemol | Rectol | Regamol | Relif | Retherma | Retol | Retopyrin | Rivmol | Rupar | Rupar junior | Rupar kid | Rutuvel | Saincitamol | Salofan | Sdmol 250 | Senmol | Sigmol | Skydol | Soparin | Spictol | Splendid | Starmol | Sumo l | Suppol | Swimol | Synamol | Sysdol | T 98 | Tacinol | Tamin | Tamol | Tamolgan | Tecpara | Tempcon | Temped | Temptal | Tepid | Thermgo | Thermodol | Thermol | Thermowin | Trustmol | Trycin | Tyfy | Tynelol | Tynol | Ultragin | Ultramol rf | Ultrapyrin | Unflem | Unmol | Ushnol pain | Ushnol rapid | Ushnol rapid fever | Vetocin | Vyrumol | Walcin | Wellpar | Welset | Xtpara | Xyfix | Xypar | Xytroy | Zepara | Zypara;
- (IQ) Iraq: Acetamol | Antipyral d | Antipyrol | As paracetamol | Asparamol | Dolamol advance | Dolorok | Dr panadolz | Duantipyro | Hayadol | Kanadain | Paindole advance mdi | Panadain | Panadain advance | Pandol d | Paracedain | Paracedol | Paracetamol mdi | Paracetol | Paradol | Paramol | Parazar | Parazar kid | Piodol | Piodol advance | Safacetol | Safadol | Safadol actifast | Safadol baby | Safadol child | Samadol | Samadol advance;
 - (IT) Italy: Acetamol | Baby rinolo febbre e dolore | Efferalgan | Flogarfev | Minofen | Minofen febbre e dolore bambini | Nirolex febbre e dolore | Normafllu | Panaadol | Paracetamol pensavital | Paracetamolo | Paracetamolo Aurobindo Pharma | Paracetamolo b braun | Paracetamolo baxter | Paracetamolo doc generici | Paracetamolo dr max | Paracetamolo eg | Paracetamolo galenica senese | Paracetamolo marco viti | Paracetamolo mylan | Paracetamolo pensa | Paracetamolo salf | Paracetamolo sandoz | Paracetamolo zentiva | Paracetamolo zentiva italia | Perfalgan | Piros | Puernol | Sanipirina | Tachipirina | Tachiverde | Termadec febbre e dolore;
 - (JO) Jordan: Adol | Calpol | Cetamol | Cetamol advance | Dolocet | Dolorgan | Newdol | Omol | Pamol | Panaadol | Panaadol advance | Panda | Panda Advance | Paraconica | Paramed | Parasustain | Paravin | Perfalgan | Pymol advance | Remedol | Revanin | Stopalgic | Tamin | Tylenol extra strength;
 - (JP) Japan: Acelio | Acetaminophen | Acetaminophen Isei | Acetaminophen maruishi | Acetaminophen sanwa | Acetaminophen tck | Acetaminophen towa | AcetaminophenK | Alpiny | Alpiny s.s. seiyaku | Alpiny sanken | Anhiba | Anyrume | Anyrume s | Aphlogis | Atenemen | Atoanon | BUFFERIN Luna J | Calonal | Calonal a | Calonal o | Calonal showa | Calonal wakodo | Calsil | Cocarl | Colgen kowa aa | Kiofever | Naron | Neosedenal | Pabron kodomo | Paraceta | Ringl | Sarutu | Tylenol a | Uzu genetsu;
 - (KE) Kenya: Abpara | Acedol | Adco napamol | Adol | Arcet | Asmol | Axapara | Betamol | Biomol | Blink | Calpol | Caremol | Cetamol | Cetamol junior | Cosmol | Curamol | Dentamol | Dolomol | Elymol | Febrex | Fevadol | Feverach | Feverex | Feverex toto | Fremol | Gacet | Ifimol | Infulgan |

- Jotonol | Junior parapaed | Junior sonadol | Kaldol | Kamadol | Micromol | Napa | Neladol | Ossmol | Pa 12 | Paarpin | Pacemol | Pacimol | Painil | Panadol | Panadol advance | Panadol baby & infant | Panadol joint | Parace | Paracerol | Paracetamol b.braun | Paraconica | Paradol | Parafast | Parafil | Parafusiv | Paragen | Paralab | Paralief | Paratal | Paravin | Parol | Pharmadol | Pharmalgan | Procet | Pyremol | Rapidol | Regamol | Reset | Rinamol | Sanmol | Sheladol | Six plus parapaed | Skymol | Sonadol | Sonamol | Tamedol | Tamin | Tamoplus | Thermadol | Trumol | Unimol | Vetocin | Vivamol | Woodward's paracetamol | Xykaa extend | Xykaa rapid | Zover;
- (KR) Korea, Republic of: A.a.p | Ace liquid | Acecort | Acenophen | Acenophen 8 hours er | Acepa | Acepin | Aceraphen | Acet | Acet m 8hours er | Aceta | Acetami | Acetaminophen | Acetaminophen boryung | Acetaminophen daejung | Acetaminophen dongyang | Acetaminophen hana | Acetaminophen hangook | Acetaminophen hangook cosmetic | Acetaminophen hyosung | Acetaminophen inwha | Acetaminophen naewoe | Acetaminophen namyang | Acetaminophen samkyoung | Acetaminophen samnam | Acetaminophen samsungfine | Acetaminophen samwoo | Acetaminophen semo | Acetamol | Acetaphen | Acetphen premix | Actetem er | Aizal | Alkapen | Amiceta | Amiset | Anisphen 8 hours er | Anisphen er | Anophen | Arom | Arucen | Arucen bag | Aspen-s | Astafen | Astamol | Astaphen | Atopen | Bioacet | Bioaset-f | Carol kids a | Cetamin | Cetophen | Champ | Children's tylenol | Cleantal | Coldaewon kids pen | Crیتال | Daewoongbio acetaminophen | Denonex | Donga acetaminophen | Dongkwang acetaminophen | Dupen | E dupef | E q pen kids a | Effena | Endapen | Ensid 8 hours er | Ensid er | Erphen er | Ezn 6 ace | Gcwb acetaminophen | Geworin i | GL acetaminophen | GI cetaminophen | Help Angelica | Hitagen | Huteacs aap | Ilyang acetaminophen | Itarenol | Kabi paracetamol | Koko-pedi | Kolon acetaminophen | Litephen | Madiphen | Mahaphen | Mapen | Mardi | Medytox acetaminophen | Nana | Naspen | New aminophen premix | Newaminophen | Newpropa | Pacemol | Painover | Panadol | Para k | Paracare | Paracet | Pedipan | Pentamol | Penzal 8 hours er | Penzal t | Perfalgan | Phenacol | Phenceta | Phnompen | Prefephen | Profa | Q phen | Raperon | Raperon 8 hours er | Raperon er | Reset | Rhinapan | Rootphen | S vil acet | Samik acetaminophen | Samnam acetaminophen | Sanopen | Sericon | Serycon | Setopen | Setopen 8 hours er | Shatphen | Sinbia | Smaten | Soluaphen | Sumediphen 8 hours ER | Suspen | Suspen 8 hours er | Suspen er | Tacenol | Tacenol 8 hours er | Taganopain | Talenil | Taminol | Taminophen | Tamisphen 8 hours er | Tanophen | Tasmen | Taspen | Taspen er | Tdspharm acetaminofen | Timerol er | Tinaphen | Tolupan | Tolupan 8 hours sr | Torophen | Torupan | Tospan | Tramol | Tramol 8 hours sr | Trisfen ER | Trisphen 8hours er | Tylenol | Tylenol er | Tylicol | Tymax | Tynol | Typen | Tyrex | Tyrex-c | Tyxen | Union acetaminophen | Union acetaminophenbek | Yungjin aminophen;
 - (KW) Kuwait: Adol | Calpol | Efferalgan | Emidol | Fevadol | Medamol | Medinol | Neomol | Panacare | Panaadol | Panaadol actifast | Panaadol advance | Panadol childrens | Panadrex | Paraconica | Perfalgan | Revanin | Temptra | Tylenol | Tylenol arthritis | Vonat;
 - (LB) Lebanon: Adol | Apo acetaminophen | Ben u ron | Calpol | Dafalgan | Doliprane | Dolopan | Efferalgan | Infumol Farmoz | Kidol or | Nomal | Painol | Panadol | Panadol advance | Paracetamol arrow | Paracetamol b.braun | Paraconica | Paramol | Parol | Pediatix | Perfalgan | Sicadol | Suldex a | Supofen | Temptra | Tylenol | Tylenol arthritis;
 - (LT) Lithuania: Acamol | Acetophen | Aldolor | Apo acetaminophen | Calpol | Calpol 6 plus | Daleron | Efferalgan | Gripex | Medipyrin | Mexalen | Neoparacetamol | Pamol | Panadol | Panadol baby | Panaadol Optizorb | Paracetamol accord | Paracetamol b.braun | Paracetamol ecopharm | Paracetamol kabi | Paracetamol panpharma | Paracetamol zentiva | Paracetamolo galenica senese | Paralen | Paramol | Sanidol | Tylenol;
 - (LU) Luxembourg: Algostase mono | Ben u ron | Curpol | Dafalgan | Dafalgan pediatrique | Dolprone | Efferalgan | Enelfa | Lemgrip | Lonarid Mono | Panaadol | Paracetamol eg | Paracetamol fresenius kabi | Paracetamol sandoz | Perdolan | Perfusalgan;
 - (LV) Latvia: Acamol | Acetophen | Ben u ron | Calpol | Calpol 6 plus | Cefecon d | Dafalgan | Efferalgan | Febridol | Lekadol | Mapap | Mexalen | Non aspirin | Pamol | Panaadol | Panadol Optizorb | Paracet | Paracetamol accord | Paracetamol actavis | Paracetamol baby phs | Paracetamol kabi | Paracetamol panpharma | Paracetamol zentiva | Paraflux | Paralen | Paramax | Paramax junior | Paramol | Perfalgan | Tailenol | Termalgin | Titralgan | Tylenol;
 - (MA) Morocco: Andol | Apiretil | Apotel | Apyrol | Atamel | Calpol | Cetamyl | Claradol | Dolgesic | Doli Pediatrique | Doliprane | Dolmax | Dolostop | Efferalgan | Geluprane | Nofebril | Oralgan | Panadol | Paracetal | Parantal | Perfalgan;

- (MX) Mexico: 36.5 infantil | Abatem | Ac fast | Acetafen | Acetif | Adinol | Adinol infantil | Alpirex | Amolgen | Analphen | Andox | Antidol | Antidol ito | Axapara | Biofer | Bonamol | Brontonyl | Colderina | Coriver | Coriver infantil | Datril | Dolotemp | Doluvit | Doluvital | Facetol | Febran | Febraxito | Febrim | Filanc | Gunartek | Icetazol | Infalgina | Kabuflen | Lunazol | Magnidol plus | Mejoral | Mejoral actirapido | Mejoralito | Mejoralito junior | Mejoralito pediatrico | Minofen | Moltec | Noblak | Nordinet infantil | Notem | Oxigricol | Pana  ol | Panofen | Paracetamol allen | Paracetamol gi arl | Paracetamol gi ser | Paracetamol Merck | Perfalgan | Pharmacen | Pharmacen m | Pharmacen t | Piralgina | Piralyne | Piremol | Portem | Precifen | Prosedal | Quitadol | Relofofor | Rolafen | Salpifar | Sedalmerck infantil | Sedalmerck pediatrico | Sedalmerck sedalito | Sensifazol | Serflax | Sidus | Sinedol | Soltadol | Son s piral | Son's piral | Sudis | Tafirol | Tasinec | Temperal | Tempire | Tempofin | Tempra | Tempra forte | Temprim | Temprin | Temzzard | Termotrin | Tylenol | Tylex | Ubeia | Ulpafie | Vicdol | Wamindel | Winasorb | Xi dol | Xi dol infantil | Zurecet;
- (MY) Malaysia: Acet | Acetamol | Actimax | Actimol | Aeknil | Arfen | Avadol | Avadol fast | Avamol | Axcel paracetamol | Aximol | Axpain | Calpol | Cetamax | Cetamol | Children's protemp | Coxan | Dhamol | Dyna p | Dyna paracetamol | Dynamol | Fepril | Hoemal | Hovid adamol | Hurix's 1000 fever & painrelief | Idamol | Ifimol | Kamex | Kidz PCM | Medon | Metacin | Mili | Milidon | Milidon-120 | Milidon-250 | Milidon-500 | Milidon-650 | Naprex | Oralcet | Pacimol | Palavo | Pamol | Panadol | Panadol extend | Pana  ol Optizorb | Panamol | Paracap | Paracet | Paracetamol aft | Paracetamol b. braun | Paracil | Parafizz | Paramol | Parasustain | Parmol | Partamol | Poro | Progesic | Rapidol | Redamol | Redon | Remedol | Rexom winamol | Saframol | Selmax | Setamol | Setromol | Tamin | Tempol | U-molin | Unimol | Uphamol | Weng heng | Winamol | Wingsic | Winkuadon | Winmotan | Winpamol;
- (NG) Nigeria: Ac mol | Acepol | Acetaminophen | Agary paracetamol | Agemol syrup | Albemol | Architamol | Ausdol | Avro paracetamol | Avrogin | Barimol | Bioramol | Bonadol | Bond paracetamol | Bradmol paracetamol | Cadmol p | Cannon paracetamol | Daga | De shalom paracetamol | Drugamol | Easadol | Emcap | Emzor para | Emzor paracetamol | Faith paracetamol elixir | Febrilix | Feccox paracetamol syrup | Feridol | Gauze paracetamol | Ivamol | Joyfad paracetamol | Juhel paracetamol | Kp paracetamol | Labcetamol | Lapain | Lapar | Leyjay paracetamol | Lotemp paracetamol | Mantra | Micpol | Mildol paracetamol | Milidon | Molgen | Moxie paracetamol | Nacemol | Nalis paracetamol | Ngc valgin | Nichben paracetamol | Novadex paracetamol | Osworth paracetamol | Painkil | Palmol | Panda | Panpek paracetamol | Paracetamol b.braun | Paraconica | Paraecnu | Paragesic | Parakris | Paraneros | Parasam | Peace paracetamol | Petsow's children starcimol | Richdol paracetamol | Richdol paracetamol syrup | Sam paracetamol | Sebesmol | Sofamol paracetamol | Supofen | Suremol | Theracin | Thermadol | Vadimol | Weldol | Zonagin;
- (NL) Netherlands: Albert Heijn Paracetamol | Apotel | Boots pharmaceuticals paracetamol | C1000 kinderparacetamol | DA Paracetamol | Daro | Daro paracetamol | Etos kinderparacetamol | Hedex | Hema paracetamol | Idyl kinderparacetamol | Idyl paracetamol | Jumbo paracetamol | Kinderparacet | Kinderparacetamol | Kinderparacetamol cf | Kinderparacetamol htp | Kruidvat paracetamol | Leidapharm kinderparacetamol | Leidapharm paracetamol | Livsane kinderparacetamol | Livsane paracetamol | Momentum | Pana  ol | Panadol artrose | Paracetalgin | Paracetamol A | Paracetamol actavis | Paracetamol Alparma | Paracetamol aurobindo | Paracetamol b. braun | Paracetamol fresenius kabi | Paracetamol Hexal | Paracetamol htp | Paracetamol linn | Paracetamol mdq | Paracetamol mylan | Paracetamol noridem | Paracetamol PCH | Paracetamol pharmaclan | Paracetamol rxt | Paracetamol sandoz | Paracetamol teva | Perfalgan | Pinex | Roter paracetamol | Rxt paracetamol | San paracetamol | Sinaspril paracetamol;
- (NO) Norway: Acetamol prima infanzia | Calpol six plus | Mexalen | Pamol | Panodil | Panodil zapp | Paracet | Paracetamol actavis | Paracetamol b. braun | Paracetamol baxter | Paracetamol baxter viaflo | Paracetamol benel | Paracetamol euromedica | Paracetamol fresenius kabi | Paracetamol norfri | Paracetamol orifarm | Paracetamol panpharma | Paracetamol ratiopharm | Paracetamol zentiva | Paramax | Perfalgan | Pinemol | Pinex;
- (NZ) New Zealand: Aft paracetamol | Amcal paracetamol | Avallon paracetamol for children | Biomed paracetamol | Disprol | Ethics paracetamol | Medco paracetamol | Pacimol | Pain clear | Pain relief | Pamol | Panadol | Pana  ol Optizorb | Paracare | Paracetamol apotex | Paracetamol kabi | Paracetamol osteo | Paraosteo | Paratabs | Perfalgan | Pharmacare paracetamol | Pharmacy health childrens paracetamol | Pharmacy health paracetamol | Pharmacycare paracetamol | Your pharmacy paracetamol;

- (OM) Oman: Adol | Omol | Pmol;
- (PA) Panama: Acertar | Acetaminofen | Acetaminofen calox | Acetaminofen mk | Acetaminofen rigar | Adorem | Alidol | Analfen | Apotel max | Apyrol | Asingesic | Atermin | Atermin g | Dolex | Dolex ninos | Feverach | Feveridol | Leniol | Mencetamol | Panaadol | Panaadol bebe | Panaadol bebes | Panaadol extra fuerte | Panadol ninos | Paracetamol b. braun | Paracetamol kabi | Paracetamol sintetica | Paraconica | Sensifazol | Tylenol | Tylex | Winasorb | Xfebril | Zetafen;
- (PE) Peru: Acetalgina | Acetam | Alicam forte | Anmol | Apiret | Aproxol | Atamel | Bonadol | Calmalip | Calmalip 500 | Chao febredol | Colmar | Colmar xtra | Dolfever | Dolfin | Dolmacaf | Dolmacaf infantil | Dolocare | Dolocetamol | Dolofebril | Dolol | Dolorless | Dolox forte | Extindol | Farmadol | Febramol | Fevastin | Fibrimol | Fieber ninos | Frenadol | Gesidol | Kitadol | Leniol | Liviolex | Lusadol | Luzetamol | Mafidol | Mafidol extra forte | Magnadol | Maxicalm | Mebinol | Meconadol | Mejoralito | Mejoralito infantil | Mejorax | Milzor | Moxtafen | Nerupol | Pacidol | Padoliex | Pala fiebre | Paldolor | Panaadol | Panadol para ninos | Panafebril | Paracetal | Paradolo | Paramidol | Parazmol | Parinmol | Pharamol | Pluscalm | Pronadol | Supracalm | Tafirol | Tapsin infantil | Teralgex | Termolin | Tonopan | Tonopan ninos | Treupel | Tylenol | Tylex kids | Zatrín;
- (PH) Philippines: Acet | Acevon | Aeknil | Afebral m | Alvedon | Axemol iv | Biogesic | Biogic | Brinogesic | Calpol | Calpol six plus | Cortalgescic xtra | Dipamol iv | Eurogesic | Febrile free | Fevergan | Flugard | Ifimol | Kiddilets | Kidicef | Medicol | Meyerf paracetamol | Molpraxx | Nahalgescic | Nalgescic | Napran | Naprex | Novamol | Opigesic | Panadol | Para | Para250 | Paragesic | Parasaph | Paraseth | Parvid | Prc | Ranigesic | Rapidol | Riber | Ritemed paracetamol | Sanmol | Selegescic | Servigesic | Sintos | Stalpara | Sumagesic | Tamin | Tempaid | Tempcaire | Temperal | Tempmed | Tempra | TGP Paracetamol | Trimitamol | Tylenol | United home fevertab | Watsons paracetamol | Winadol | Zytamol;
- (PK) Pakistan: Acetadol | Acetamol | Adol | Advil 6 plus | Advil oral suspension | Afeb | Alimol | Anamol | Axamol | Axamol plus | Baxterol 6 plus | Bemol | Bencetamol | Bofalgan | Bramol | Calpol | Calpol 6 plus | Celsius | Cetapol | Chikymol | Chilpol | Colbex | Danidol | Delmol | Delmol forte | Disprol | Dumol | Estadol | Falgan | Febrol | Febrol ds | Fevamol | FEVERAL infusion | Fevercare | Feveren | Feveren 6 plus | Fitamol | Geomol | Karmol | Klebro | Klebro plus | Kymol | Macmol | Maxfever | Maxfever plus | Mismol | Mismol plus | Monadol | Mylenol | Mytomol | Nafpol | Nafpol 6 plus | Napa | Normidol | Nymbex | Nymbex 6 plus | Paedal | Panaadol | Panadol forte | Panam | Para | Para 4 kids | Para 6+ | Paracare | Paraceq | Paracetamol kabi | Paracetol | Paradon | Paralord | Paramax | Paramed | Paramol | Paranol | Parapol | Parasup | Parasyp | Parazol | Parcol | Parkomol | Parokal | Perdolan | Pm | Pracmol | Provas | Pyrol forte | Radimole 6 plus | Rascodol | Reliefal | Reliefal 6 plus | Saptamol | Setamol | Skimol | Sn dol | Stanpol | Supamol | Tempol | Tyladol | Tydol | Viodol | Weladol;
- (PL) Poland: Acenol | Acetaminophen | Aletofar | Amipar | Apap | Apap caps | Apap dla dzieci Forte | Apenal | Ben u ron | Benalpharm | Codipar | Dafalgan | Efferalgan | Etoran | Gemipar | Infaceta-mol | Lekadol | Metafen paracetamol | Metafen paracetamol max | Novo gesic | Omnipap | Panacit | Panaadol baby | Panaadol max | Panaadol sprint | Panadon | Panaprex | Paracemol | Paracenol | Paracerol | Paracetamol 123ratio | Paracetamol accord | Paracetamol actavis | Paracetamol aflo-farm | Paracetamol apteo med | Paracetamol aurovitas | Paracetamol b. braun | Paracetamol doz | Paracetamol dr. max | Paracetamol farmina | Paracetamol filofarm | Paracetamol forte apteo med | Paracetamol hasco | Paracetamol hasco forte | Paracetamol kabi | Paracetamol olimp | Paracetamol panpharma | Paracetamol sandoz | Paracetamol teva | Paracetamol zen-tiva | Paracetamololum farmalider | Paralen | Paramax | Paramax rapid | Parokan | Pedicetamol | Rastamol | Sinebriv | Tazamol | Tylenol;
- (PR) Puerto Rico: Acetaminophen | Acetaminophen extended release | Aminofen | Aminofen max | Apap | Apap elixir | Aphen | Children's Accudial acetaminophen | Children's pain & fever | Childrens mapap | Childrens pain and fever | Comtrex sore throat relief | Curanol | Extra strength pain relief | Infants pain & fever | Infants' tylenol | Kids pain & fever | Little fevers | Little remedies for fevers | M pap | Max relief | Max relief junior | Nortemp | Ofirmev | Pain relief | Panadol | Panadol infants | Pediacare fever reducer pain reliever | Pharbetol | Q-pap | Silapap | Tactinal | Tempra | Tylenol arthritis;
- (PT) Portugal: Anfetol | Anti-gripe asclepius | Atralidon | Ben u ron | Dafalgan | Doloramol | Doloramol xpress | Dolostop | Gelocatil | Holdofeb | Nuridol | Pacimol | Panadol | Panaadol prolong | Panasorbe | Pantadolor | Paracetamol accord | Paracetamol algik | Paracetamol

- alter | Paracetamol basi | Paracetamol Bluepharma | Paracetamol Combino | Paracetamol farmoz | Paracetamol GP | Paracetamol Labesfal | Paracetamol mylan | Paracetamol Nideran | Paracetamol Pensa | Paracetamol tecnigen | Paracetamol toLife | Paracetamol wellsmed | Paramolan | Parsel | Perfalgan | Singrips | Supofen | Takipirina | Tylenol | Vimergol | Zaramol;
- (PY) Paraguay: Acetaminofen caplin point | Acetamol | Acicur p | Algidol | Algidol forte | Aliviol | Antiflam p | Dolpiret | Efermol | Febrin | Febrinol | Fiedol | Flodin | Gesin | Matydol | Maximol | No dol p | Nodolex | Novomol | Pacemol | Paracetamol b.braun | Paracetamol bional | Paracetamol cassanello | Paracetamol chile | Paracetamol dasanti | Paracetamol heisecke | Paracetamol kabi | Paracetamol lasca | Paracetamol medifar | Paracetamol millet | Paracetamol mintlab | Paracetamol vent3 | Paracetamol vivele | Paracetamolito | Paradolanet | Pirenzep | Sindrolen | Tafirol | Tafirol pediatrico | Tercedol | Z mol | Zededol;
 - (QA) Qatar: Acetalgin | Adol | Calpol | Calpol Six Plus | Cetamin | Dolomol | Enelfa | Falgan | Fevadol | Feverol Advance | Leniol | Mapap Children | Omol | Pamol | Pamol Child | Pamol Infant | Pan-Q | Panadol Actifast | Panaadol Advance | Panaadol Advance Optizorb | Panadol Baby | Panadol Baby and Infant | Panaadol Children 5-12 years | Panaadol Children 6+ years | Panadol Cold and Flu Vapour Release | Panaadol Joint | Panadol Junior | Panadrex | Panalex Advance | Panda | Panda Advance | Panda DS | Panorix | Paracerol | Paracetamol Kabi | Paracetamol-N | Parafusiv | Paramed | Paranox S | Paravin | Partemol | Perfalgan | Pirofen | Pmol | Pmol Fizz | Revanin | Tylenol | Tylenol Forte;
 - (RO) Romania: Adol | Apap | Apap forte | Calpol sase plus | Calpol trei plus | Daleron | Defebryl | Dulsifeb | Efferalgan | Efferalgan pediatric | Efferalgan pentru copii | Fervex durere si febra | Pacetral | Panaadol | Panaadol artro | Paracetamol accord | Paracetamol arena | Paracetamol aurobindo | Paracetamol biofarm | Paracetamol farmex | Paracetamol helcor | Paracetamol infant | Paracetamol infomed | Paracetamol lph | Paracetamol mcc | Paracetamol polipharma | Paracetamol polisano | Paracetamol rompharm | Paracetamol santa | Paracetamol sintofarm | Paracetamol terapia | Paramol | Perfalgan | Prosinus;
 - (RU) Russian Federation: Acamol | Acetalgan | Aminadol | Apap | Calpol | Cefecon d | Daleron | Deminofen | Dolo | Efferalgan | Ifimol | Lekadol | Mexalen | Pacimol | Pamol | Panaadol | Panadol baby | Paracet | Paracetamol akos | Paracetamol alium | Paracetamol avexima | Paracetamol children | Paracetamol ECO | Paracetamol ekolab | Paracetamol for children | Paracetamol fortekids | Paracetamol lekt | Paracetamol medisorb | Paracetamol ubf | Paracetamol Velpharm | Perfalgan | Sanidol | Strimol | Tylenol;
 - (SA) Saudi Arabia: Acetaminophen | Actidol | Adol | Amol | Calpol | Dafalgan | Efferalgan | Emidol | Fevadol | Medamol | Neomol | Omol | Panaadol | Panaadol actifast | Panaadol advance | Panaadol baby & infant | Panadol extend | Panadrex | Panamol | Panda | Paracetamol b. braun | Paracetamol pharco b | Paracetol | Paraconica | Parafusiv | Paramol | Pediamol | Perfalgan | Pmol | Revanin | Riagesic | Tempra | Tylenol | Vitopeine | Vonal;
 - (SE) Sweden: Alvedon | Alvedon dos | Alvedon forte | Alvedon novum | Curadon | Oparap | Pamol | Panodil | Panodil zapp | Paracetamol abece | Paracetamol actavis | Paracetamol alternova | Paracetamol apofri | Paracetamol b. braun | Paracetamol baxter | Paracetamol bluefish | Paracetamol eql pharma | Paracetamol evolan | Paracetamol fresenius kabi | Paracetamol krka | Paracetamol medic | Paracetamol NET | Paracetamol nordic | Paracetamol novum apofri | Paracetamol orifarm | Paracetamol panpharma | Paracetamol teva | Paracetamol zentiva | Paracut | Paracut forte | Parapo | Perfalgan | Pinex;
 - (SG) Singapore: Alcetamol | Calpol | Cetamol | Dhamol | Dolo | Fepril | Fever & pain relief | Milidon | Napa | Naprex | Pacemol | Pamol | Panaadol | Panaadol actifast | Panaadol extend | Panadol for children | Panamol | Paracetamol aft | Paracetamol b. braun | Paracil | Paramol | Paratab | Parcemol | Parmol | Poro | Pritamol | Progesic | Redon | Remedol | Setamol | Sp febril | Uphamol | Zerine;
 - (SI) Slovenia: Calpol | Calpol 6 | Daleron | Lekadol | Panaadol | Panaadol baby | Panaadol Optizorb | Paracetamol accord | Paracetamol b.braun | Paracetamol kabi | Paracetamol Lekarna Ljubljana | Perfalgan | Tevitamol;
 - (SK) Slovakia: Calpol | Efferalgan | Lekadol | Medipyrin | Mexalen | Panadol | Panaadol baby | Panadol novum | Paracetamol agmed | Paracetamol b. braun | Paracetamol dr. max | Paracetamol grindeks | Paracetamol k | Paracetamol kabi | Paracetamol noridem | Paracut | Paralen | Parapyrex | Perfalgan | Plicet;

- (SL) Sierra Leone: Excel's paracetamol | Gacet | Mikalgin ap | Miral paracetamol | Parakant | Pekin para | Pyrigen | Welcetamol | Zinol;
- (SR) Suriname: Ace | Algina | Da kinderparacetamol | Idyl kinder paracetamol | Idyl paracetamol | Kinderparacetamol htp | Kinderparacetamol sam | Livsane paracetamol | Painol | Panadol actifast | Para denk | Paracetamol apotex | Paracetamol htp | Paracetamol sam | Paracetamol sandoz | Paracetamol sanias | Paracetamol stada | Paracetamol teva | Paramol | Paranip | Parmol | Placidex | Sinaspril paracetamol | Tylenol children's | Tylenol extra strength | Tylenol infants;
- (SV) El Salvador: Acetaminofen | Tempydol;
- (TH) Thailand: Acenol | Aceta | Acetaminophen | Acetophen | Aeknil | Algogen | Ampol | Angela | Bakamol | Beramol | Beramol green | Beramol white | Calpol | Cemol | Cemol p | Cetamol | Cetapol | Cetophen | Cetta | Children's protemp | Detemp | Faron | Feehamol | Femol | Foramol | G pamol | Godcemol | Infants' Protemp | Infulgan | Kamol | Kit F | Kressmol | Livsamol | Lotemp | Masapara | Mcxy para | Mymol | Mymol m.d. | Mypara | Nasa | Nasa para | Newtamol | Newtol | Newtol kids | O chin | Orecet | Pacemol | Pamol | Panaadol | Panaadol actifast | Panamol | Pancemol | Papol | Para | Para baby | Para G | Para g dek | Para sweet | Parac | Paracap 2 | Paracet | Paracet frx | Paracin | Parahof | Paramac | Paramol | Paratab | Paratemp | Parmol | Pat med | Patblu | Patum | Pecet | Phairac | Pipmon | Pondparac | Protemp | Pycetol | Pycetol F | Pyracon | Pyranar | Qualimol P | Saebegin | Sara | Servigesic | Starmol blue | Starmol orange | Temnol | Tempcox | Tempra | Termalgin | Tumpara | Tylenol | Tylenol arthritis pain | Unimol | Unipol;
- (TN) Tunisia: Adol | Adoline | Algesic | Algidol | Analgan | Apyrex | Cetamol | Coldex | Dafalgan | Dolel | Doliprane | Effadol pediatrique | Efferalgan | Fresidol | Novadol | Panadol | Parac info | Paracetamol G.E.S. | Paracetamol ips | Paracetyl | Paradol | Paralvit | Paramed | Perfalgan | Stopalgic;
- (TR) Turkey: A ferin zero 6 plus | A-per | Asetam | Asomal | Babinoks | Berko Setamol | Calpol | Deminofen | Durapan | Ekosetol | Gripin Bebe | Kataprin | Kidmol | Minafen | Minamol | Minamol plus | Minopar plus | Minoset | Noral | Panaadol | Panadol actifast | Para nox | Paracerol | Paranox | Parasedol | Parcetol | Parokan | Parol | Parsetan | Partemol | Peditus uno | Peditus uno plus | Pharmadol | Pirofen | Polaminofen | Poliaminofen | Polmofen | Rastamol | Sedalon | Seskamol | Setamol | Setapar | Sifenol | Tamifen | Tamol | Tempo | Termacet | Termalgin | Tylol | Vermidon | Volpan | Xfebril | Zaldaks;
- (TW) Taiwan: Acenol | Aceta | Acetal | Acetaminophen | Acetamol | Acetaphene welcan | Anacin-3 | Anliron | Anstafen | Anti phen | Antidol | Antipain chinteng | Antone | Apap | Bocoton | Bubdel | Butong | Cetamol | Chitounen | Cochitong | Dikuton | Fucole Paran | Hecamol | Jindonpin | Lactam | Napa | Napaton | Paceton | Panacon | Panacon acefen | Panadol | Panalife | Panamax | Panatol | Paracetamol stada | Paralgin | Paramol | Paranol | Poro | Punortor | Pyrinazin | Ramol | Reh nar tong | Sediton | Shikenuoh | Soonansin | Tachipirina | Tamdron | Taminol | Thioton | Tinten | Tonlan | Tonphen | Yollowin | Zolben;
- (TZ) Tanzania, United Republic of: Cetamol;
- (UA) Ukraine: Aldolor | Anapiron | Apotel max | Blimol | Calpol | Efferalgan | Elgan | Exopon | Gripomedum | Ifimol | Infulgan | Medipyrin | Mexalen | Milistan | Panadol | Panaadol advance | Paracet | Paracetamol b.braun | Paracetamol baby | Paracetamol euro | Paracetamololum novo | Parafast | Parafusiv | Paramax | Paramol | Piaron | Pride | Rapidol | Sanidol | Tylenol;
- (UG) Uganda: Abamol | Abpara | Ace | Adco napamol | Agomol | Agopar | Althemol | Asmol | Axcel paracetamol | Bramol | Cachpar | Calpol | Cetamol | Cosmol | Curamol | Dentamol | Dolomol | Elymol | Feveral | Feverex | Feverix toto | Fremol | Ifimol | Influgan | Infulgan | Ivamol | K mol | Kamadol | Kamadol junior | Kamadol kids | Kelvimol | Kelvin | Leniol | Medomol | Niko | Pa 12 | Paarpin | Paediatric paracetamol | Panadol advance | Panaadol baby & infant | Para | Para bkrs | Paracetamol fresenius | Paracin | Paraconica | Paradol | Parafil | Paramed | Paratal | Paravin | Pharmalgan | Polymol | Rectol | Regamol | Renedol | Sanmol | Simdol | Supadol | Suredol | Tamin | Thermadol;
- (UY) Uruguay: Acf | Ceta mol | Detial | Dolex | Dolex pediatrico | Dolosedol | Dolosedol pediatrico | Dozic | Kitadol | Mejoral | Panaadol | Paracetamol athena | Paracetamol kabi | Paracetamol Opko | Paracetamol szabo | Paracetamol teva | Parageniol | Pirantil | Sacaprim | Tafirol | Tervalan | Tylenol | Zetamol | Zolben | Zolbinet;
- (VE) Venezuela, Bolivarian Republic of: Acetabiofen | Acetadol | Acetafen forte | Acetaminofen | Aceval | Acevalone | Actron | Adorem | Agurin | Alivax forte | Amitafen forte | Analgan | Ananty |

- Apiret | Apyrene | Atamel | Brexin | Brexin forte | Colprin | Dol kids | Dopagan | Febrip | Infalgan | Ronadol | Tachiforte | Tachipirin | Tempra | Tempra forte | Tylenol | Tylex | Ultrafen | Vestax | Winadol;
- (VN) Viet Nam: Acepron | Agi tyfedol | Amvifeta | Anogin | Bakidol | Banalcine | Bivinadol | Bv gestic | Byralen | Fahado | Glotadol | Hapacol | Korando | Mexcold | Nicnotaxgin | Panactol | Panadol | Pandidol | Parazacol | Parcamol | Partamol | Phaanedol | Pharbacol | Repamax | Savipamol | Sixfitol | Tahero | Tatanol forte | Topsea | Tydol;
 - (ZA) South Africa: Actamol | Austell paracetamol | Calpol | Cetafuse | Cetapon | Cetapon junior paediatric | Dis chem paracetamol | Dolorol | Donapan | Dynadol | Fabu paracetamol | Fevamol | Go pain p | Go pain paracetamol | Grays paracetamol | Intramol | Lennon paracetamol | Liberilix | Napamol | Pacimol | Pacipayn | Painamol | Painblok | Painogesic | Panado | Paracetamol biotech | Paracetamol fresenius | Paraspem | Perfalgan | Pharma q paracetamol | Prolief | Pyrex | Tylenol | Varipan | Zypol;
 - (ZM) Zambia: Agomol | Amidol | Anmol | Ar cetamol | Asmol | Babygesic | Betamol | Biomol | Cachpar | Calpol | Cetamol | Curamol | Ecocin | Fego | Fevastin | Feverlet | Gacet | Kelvin | Lanol | Metamol | Milpol | Milzor | Neladol | Novadol | Pa 12 | Pacimol | Panado | Panado for children | Paradenk | Paracetamol agro | Paracip | Parafil | Parafil 120 | Parakant | Paralget | Parol | Pharmadol | Pmol | Regamol | Sheladol | Ucetamol | Varipan | Xykaa extend | Xykaa rapid;
 - (ZW) Zimbabwe: Antalgic | Antalgic paediatric | Betamol | Calpol | Grays paracetamol | Meldol | Novadol | Novadol asp | Novadol rapid | Panado | Paracemol b. braun | Paracet | Paracetamol fresenius | Strimol | Tamin | Varipan

Use of UpToDate is subject to the Terms of Use.

REFERENCES

1. Acephen (acetaminophen suppositories) [prescribing information]. South Plainfield, NJ: G&W Laboratories; April 2014.
2. Acetaminophen injection for IV [prescribing information]. Bethlehem, PA: B. Braun Medical Inc; February 2021.
3. Acetaminophen Viaflo bag [prescribing information]. Deerfield, IL: Baxter Healthcare Corporation; May 2024.
4. Achterbergh R, Lammers LA, Kuijsten L, Klümpen HJ, Mathôt RAA, Romijn JA. Effects of nutritional status on acetaminophen measurement and exposure. *Clin Toxicol (Phila)*. 2019;57(1):42-49. doi:10.1080/15563650.2018.1487563 [PubMed 29974811]
5. Ahlfors CE. Benzyl alcohol, kernicterus, and unbound bilirubin. *J Pediatr*. 2001;139(2):317-319. [PubMed 11487763]
6. Ahlqvist VH, Sjöqvist H, Dalman C, et al. Acetaminophen use during pregnancy and children's risk of autism, ADHD, and intellectual disability. *JAMA*. 2024;331(14):1205-1214. doi:10.1001/jama.2024.3172 [PubMed 38592388]
7. Alade SL, Brown RE, Paquet A. Polysorbate 80 and E-Ferol toxicity. *Pediatrics*. 1986;77(4):593-597. [PubMed 3960626]
8. Alander SW, Dowd MD, Bratton SL, Kearns GL. Pediatric acetaminophen overdose: risk factors associated with hepatocellular injury. *Arch Pediatr Adolesc Med*. 2000;154(4):346-350. [PubMed 10768670]
9. Allegaert K. A critical review on the relevance of paracetamol for procedural pain management in neonates. *Front Pediatr*. 2020;8:89. doi:10.3389/fped.2020.00089 [PubMed 32257982]
10. Allegaert K, Anderson BJ, Naulaers G, et al. Intravenous paracetamol (propacetamol) pharmacokinetics in term and preterm neonates. *Eur J Clin Pharmacol*. 2004;60(3):191-197. doi:10.1007/s00228-004-0756-x [PubMed 15071761]

11. Allegaert K, Mian P, Lapillonne A, van den Anker JN. Maternal paracetamol intake and fetal ductus arteriosus constriction or closure: a case series analysis. *Br J Clin Pharmacol*. 2019;85(1):245-251. [PubMed 30300944]
12. Allegaert K, Palmer GM, Anderson BJ. The pharmacokinetics of intravenous paracetamol in neonates: size matters most. *Arch Dis Child*. 2011;96(6):575-580. [PubMed 21317433]
13. Amar PJ, Schiff ER. Acetaminophen safety and hepatotoxicity--where do we go from here? *Expert Opin Drug Saf*. 2007;6(4):341-355. doi: 10.1517/14740338.6.4.341 [PubMed 17688378]
14. American Academy of Pediatrics Committee on Drugs. Acetaminophen toxicity in children. *Pediatrics*. 2001;108(4):1020-1024. [PubMed 11581462]
15. American College of Obstetricians and Gynecologists (ACOG). Acetaminophen use in pregnancy and neurodevelopmental outcomes. <https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2025/09/acetaminophen-use-in-pregnancy-and-neurodevelopmental-outcomes>. Updated September 2025. Accessed October 7, 2025.
16. American College of Obstetricians and Gynecologists (ACOG). ACOG committee opinion no. 753: assessment and treatment of pregnant women with suspected or confirmed influenza. *Obstet Gynecol*. 2018;132(4):e169-e173. [PubMed 30247362]
17. American College of Obstetricians and Gynecologists (ACOG). ACOG practice bulletin no. 209: obstetric analgesia and anesthesia. *Obstet Gynecol*. 2019;133(3):e208-e225. [PubMed 30801474]
18. American College of Obstetricians and Gynecologists (ACOG). ACOG response to consensus statement on paracetamol use during pregnancy. Updated September 29, 2021. Accessed September 24, 2025.
19. American College of Obstetricians and Gynecologists (ACOG). Headaches in pregnancy and postpartum: ACOG clinical practice guideline no. 3. *Obstet Gynecol*. 2022;139(5):944-972. doi:10.1097/AOG.0000000000004766 [PubMed 35576364]
20. American Geriatrics Society (AGS). American Geriatrics Society Panel on Pharmacological Management of Persistent Pain in Older Persons. Pharmacological management of persistent pain in older persons. *J Am Geriatr Soc*. 2009;57(8):1331-1346. [PubMed 19573219]
21. Anand KJ, Chair, International Evidence-Based Group for Neonatal Pain, personal correspondence, April 2002.
22. Anand KJ, International Evidence-Based Group for Neonatal Pain. Consensus statement for the prevention and management of pain in the newborn. *Arch Pediatr Adolesc Med*. 2001;155(2):173-180.
23. Anderson BJ, van Lingen RA, Hansen TG, Lin YC, Holford NH. Acetaminophen developmental pharmacokinetics in premature neonates and infants: a pooled population analysis. *Anesthesiology*. 2002;96(6):1336-1345. doi:10.1097/00000542-200206000-00012 [PubMed 12170045]
24. Anderson PO, Sauberman JB. Modeling drug passage into human milk. *Clin Pharmacol Ther*. 2016;100(1):42-52. [PubMed 27060684]
25. Anderson PO, Sauberman JB, Lane JR, et al. Hydrocodone Excretion Into Breast Milk: The First Two Reported Cases. *Breastfeed Med*. 2007;2(1):10-14. [PubMed 17661614]
26. Bagheri MM, Bahman-Bijari B, Torabi-Nejad MH, et al. Is prophylactic parenteral paracetamol effective to diminish incidence of PDA in preterm neonates? A randomized trial. *Iran J Pediatr*. 2018;28(4):e11735. doi:10.5812/ijp.11735
27. Bagheri H, Bernhard NB, Montastruc JL. Potentiation of the Acenocoumarol Anticoagulant Effect by Acetaminophen. *Ann Pharmacother*. 1999;33(4):506. [PubMed 10332548]
28. Bagheri MM, Niknafs P, Sabsevari F, et al. Comparison of oral acetaminophen versus ibuprofen in premature infants with patent ductus arteriosus. *Iran J Pediatr*. 2016;26(4):e3975. doi:10.5812/ijp.3975 [PubMed 27713809]

29. Bannwarth B, Péhourcq F. Pharmacological rationale for the clinical use of paracetamol: pharmacokinetic and pharmacodynamic issues. *Drugs*. 2003;63(2):5-13. [PubMed 14758786]
30. Bartocci M, Lundeborg S. Intravenous paracetamol: the 'Stockholm Protocol' for postoperative analgesia of term and preterm neonates. *Paediatr Anaesth*. 2007;17(11):1120-1121. [PubMed 17897288]
31. Bauerlein DK, Williams AP, John PR. Optimizing acetaminophen use in patients with risk factors for hepatotoxicity: reviewing dosing recommendations in adults. *Pain Med*. 2021;22(7):1469-1472. doi:10.1093/pm/pnaa274 [PubMed 33164103]
32. Benson GD. Acetaminophen in chronic liver disease. *Clin Pharmacol Ther*. 1983;33(1):95-101. doi:10.1038/clpt.1983.14 [PubMed 6848304]
33. Benson GD, Koff RS, Tolman KG. The therapeutic use of acetaminophen in patients with liver disease. *Am J Ther*. 2005;12(2):133-141. doi:10.1097/01.mjt.0000140216.40700.95 [PubMed 15767831]
34. Berg KJ, Djøseland O, Gjellan A, et al. Acute effects of paracetamol on prostaglandin synthesis and renal function in normal man and in patients with renal failure. *Clin Nephrol*. 1990;34(6):255-262. [PubMed 2073768]
35. Berlin CM Jr, Yaffe SJ, Ragni M. Disposition of acetaminophen in milk, saliva, and plasma of lactating women. *Pediatr Pharmacol (New York)*. 1980;1(2):135-141. [PubMed 7202185]
36. Birmingham PK, Tobin MJ, Fisher DM, et al. Initial and Subsequent Dosing of Rectal Acetaminophen in Children: A 24-Hour Pharmacokinetic Study of New Dose Recommendations. *Anesthesiology*. 2001;94(3):385-389. [PubMed 11374595]
37. Bitzén PO, Gustafsson B, Jostell KG, Melander A, Wåhlin-Boll E. Excretion of paracetamol in human breast milk. *Eur J Clin Pharmacol*. 1981;20(2):123-125. doi:10.1007/BF00607148 [PubMed 7262173]
38. Blazquez AG, Briz O, Gonzalez-Sanchez E, Perez MJ, Ghanem CI, Marin JJ. The effect of acetaminophen on the expression of BCRP in trophoblast cells impairs the placental barrier to bile acids during maternal cholestasis. *Toxicol Appl Pharmacol*. 2014;277(1):77-85. doi:10.1016/j.taap.2014.02.019 [PubMed 24631341]
39. Böhler J, Donauer J, Keller F. Pharmacokinetic principles during continuous renal replacement therapy: drugs and dosage. *Kidney Int Suppl*. 1999;(72):S24-S288. [PubMed 10560800]
40. Bosilkovska M, Walder B, Besson M, Daali Y, Desmeules J. Analgesics in patients with hepatic impairment: pharmacology and clinical implications. *Drugs*. 2012;72(12):1645-1669. [PubMed 22867045]
41. Brindle ME, McDiarmid C, Short K, et al. Consensus guidelines for perioperative care in neonatal intestinal surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations. *World J Surg*. 2020;44(8):2482-2492. doi:10.1007/s00268-020-05530-1 [PubMed 32385680]
42. Brookhuis SAM, Allegaert K, Hanff LM, Lub-de Hooge MN, Dallmann A, Mian P. Modelling tools to characterize acetaminophen pharmacokinetics in the pregnant population. *Pharmaceutics*. 2021;13(8):1302. doi:10.3390/pharmaceutics13081302 [PubMed 34452263]
43. Buck ML. Perioperative Use of High-Dose Rectal Acetaminophen. *Pediatr Pharmacol (New York)*. 2001;7(9). http://www.medscape.com/viewarticle/415082_2
44. Burns MJ. Acetaminophen poisoning in adults: Pathophysiology, presentation, and evaluation. Updated April 6, 2020. <https://www.uptodate.com/contents/acetaminophen-paracetamol-poisoning-in-adults-pathophysiology-presentation-and-evaluation?search=acetaminophen%20toxicity&so>
45. Butler JM, Begg EJ. Free drug metabolic clearance in elderly people. *Clin Pharmacokinet*. 2008;47(5):297-321. [PubMed 18399712]

46. Caldeira D, Costa J, Barra M, Pinto FJ, Ferreira JJ. How safe is acetaminophen use in patients treated with vitamin K antagonists? A systematic review and meta-analysis. *Thromb Res*. 2015; 135(1):58-61. [PubMed 25456003]
47. Caparrotta TM, Antoine DJ, Dear JW. Are some people at increased risk of paracetamol-induced liver injury? A critical review of the literature. *Eur J Clin Pharmacol*. 2018;74(2):147-160. doi:10.1007/s00228-017-2356-6 [PubMed 29067481]
48. Capici F, Ingelmo PM, Davidson A, et al. Randomized controlled trial of duration of analgesia following intravenous or rectal acetaminophen after adenotonsillectomy in children. *Br J Anaesth*. 2008;100(2):251-255. [PubMed 18211998]
49. Ceelie I, de Wildt SN, van Dijk M, et al. Effect of intravenous paracetamol on postoperative morphine requirements in neonates and infants undergoing major noncardiac surgery: a randomized controlled trial. *JAMA*. 2013;309(2):149-154. doi:10.1001/jama.2012.148050 [PubMed 23299606]
50. Centers for Disease Control (CDC). Neonatal deaths associated with use of benzyl alcohol—United States. *MMWR Morb Mortal Wkly Rep*. 1982;31(22):290-291. <http://www.cdc.gov/mmwr/preview/mmwrhtml/00000319.htm> [PubMed 6810084]
51. Centers for Disease Control (CDC). Unusual syndrome with fatalities among premature infants: association with a new intravenous vitamin E product. *MMWR Morb Mortal Wkly Rep*. 1984;33(14):198-199. <http://www.cdc.gov/mmwr/preview/mmwrhtml/00000319.htm> [PubMed 6423951]
52. Chandok N, Watt KD. Pain management in the cirrhotic patient: the clinical challenge. *Mayo Clin Proc*. 2010;85(5):451-458. doi:10.4065/mcp.2009.0534 [PubMed 20357277]
53. Chen KF, Chan LN, Senn TD, et al. The impact of proximal Roux-en-Y gastric bypass surgery on acetaminophen absorption and metabolism. *Pharmacotherapy*. 2020;40(3):191-203. doi:10.1002/phar.2368 [PubMed 31960977]
54. Children's Tylenol chewable tablet (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil Consumer Healthcare; January 2018.
55. Children's Tylenol Meltaways chewable tablets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014a.
56. Children's Tylenol oral suspension liquid (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014.
57. Children's Tylenol oral suspension liquid (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil Consumer Healthcare; January 2020.
58. Chou R. Subacute and chronic low back pain: Management. Connor RF, ed. UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed August 12, 2021.
59. D'Antonio F, Flacco ME, Valle LD, et al. Prenatal paracetamol exposure and child neurodevelopment: a systematic review and meta-analysis. *Lancet Obstet Gynaecol*. Published online January 16, 2026. doi:10.1016/S3050-5038(25)00211-0
60. Damkier P, Gram EB, Ceulemans M, et al. Acetaminophen in pregnancy and attention-deficit and hyperactivity disorder and autism spectrum disorder. *Obstet Gynecol*. 2025;145(2):168-176. doi:10.1097/AOG.0000000000005802 [PubMed 39637384]
61. Dang D, Wang D, Zhang C, Zhou W, Zhou Q, Wu H. Comparison of oral paracetamol versus ibuprofen in premature infants with patent ductus arteriosus: a randomized controlled trial. *PLoS One*. 2013;8(11):e77888. doi:10.1371/journal.pone.0077888 [PubMed 24223740]
62. Dart RC, Bailey E. Does therapeutic use of acetaminophen cause acute liver failure? *Pharmacotherapy*. 2007;27(9):1219-1230. doi: 10.1592/phco.27.9.1219. [PubMed 17723075]

63. Dart RC, Erdman AR, Olson KR, et al; American Association of Poison Control Centers. Acetaminophen poisoning: an evidence-based consensus guideline for out-of-hospital management. *Clin Toxicol (Phila)*. 2006;44(1):1-18. doi:10.1080/15563650500394571 [PubMed 16496488]
64. Dart RC, Rumack BH. Intravenous acetaminophen in the United States: iatrogenic dosing errors. *Pediatrics*. 2012;129(2):349-353. doi: 10.1542/peds.2011-2345 [PubMed 22271694]
65. Dash SK, Kabra NS, Avasthi BS, Sharma SR, Padhi P, Ahmed J. Enteral paracetamol or intravenous indomethacin for closure of patent ductus arteriosus in preterm neonates: a randomized controlled trial. *Indian Pediatr*. 2015;52(7):573-578. doi:10.1007/s13312-015-0677-z [PubMed 26244949]
66. Dathe K, Frank J, Padberg S, et al. Negligible risk of prenatal ductus arteriosus closure or fetal renal impairment after third-trimester paracetamol use: evaluation of the German Embryotox cohort. *BJOG*. 2019;126(13):1560-1567. doi:10.1111/1471-0528.15872 [PubMed 31310697]
67. Davison SN, Koncicki H, Brennan F. Pain in chronic kidney disease: a scoping review. *Semin Dial*. 2014;27(2):188-204. doi:10.1111/sdi.12196 [PubMed 24517512]
68. Dreier JW, Andersen AM, Berg-Beckhoff G. Systematic review and meta-analyses: fever in pregnancy and health impacts in the offspring. *Pediatrics*. 2014;133(3):e674-688. doi:10.1542/peds.2013-3205 [PubMed 24567014]
69. Dvořáčková E, Pilková A, Matoulek M, Slanař O, Hartinger JM. Bioavailability of orally administered drugs after bariatric surgery. *Curr Obes Rep*. 2024;13(1):141-153. doi:10.1007/s13679-023-00548-7 [PubMed 38172482]
70. Dwyer JP, Jayasekera C, Nicoll A. Analgesia for the cirrhotic patient: a literature review and recommendations. *J Gastroenterol Hepatol*. 2014;29(7):1356-1360. doi:10.1111/jgh.12560 [PubMed 24548074]
71. El-Mashad AE, El-Mahdy H, El Amrousy D, Elgendy M. Comparative study of the efficacy and safety of paracetamol, ibuprofen, and indomethacin in closure of patent ductus arteriosus in preterm neonates. *Eur J Pediatr*. 2017;176(2):233-240. doi:10.1007/s00431-016-2830-7 [PubMed 28004188]
72. European Medicines Agency (EMA). Use of paracetamol during pregnancy unchanged in the EU. <https://www.ema.europa.eu/en/news/use-paracetamol-during-pregnancy-unchanged-eu#related-documents-79478>. Updated September 23, 2025. Accessed October 7, 2025.
73. Expert opinion. Senior Hepatic Editorial Team: Matt Harris, PharmD, MHS, BCPS, FAST, FCCP; Jeong Park, PharmD, MS, BCTXP, FCCP, FAST; Arun Jesudian, MD; Sasan Sakiani, MD.
74. Expert opinion. Senior Renal Editorial Team: Bruce Mueller, PharmD, FCCP, FASN, FNKF; Jason A. Roberts, PhD, BPharm (Hons), B App Sc, FSHP, FISAC; Michael Heung, MD, MS.
75. FDA Drug Safety Communication: Prescription acetaminophen products to be limited to 325 mg per dosage unit; boxed warning will highlight potential for severe liver failure. <https://www.fda.gov/Drugs/DrugSafety/ucm239821.htm>. Accessed September 29, 2017.
76. Feldkamp ML, Meyer RE, Krikov S, Botto LD. Acetaminophen use in pregnancy and risk of birth defects: findings from the National Birth Defects Prevention Study. *Obstet Gynecol*. 2010;115(1):109-115. doi:10.1097/AOG.0b013e3181c52616 [PubMed 20027042]
77. FEVERALL (acetaminophen) [prescribing information]. Cranford, NJ: Actavis; received August 2015.
78. FEVERALL Children's (acetaminophen) [prescribing information]. Hawthorne, NY: Taro Pharmaceuticals USA, Inc; received July 2019.
79. FEVERALL Infant's (acetaminophen) [prescribing information]. Hawthorne, NY: Taro Pharmaceuticals USA, Inc; received July 2019.

80. Feverall Junior Strength (acetaminophen) [prescribing information]. Hawthorne, NY: Taro Pharmaceuticals USA, Inc; received July 2019.
81. FeverAll suppositories (acetaminophen) [prescribing information]. Lincoln, NC: Actavis; 2014.
82. Fitzcharles MA, Lussier D, Shir Y. Management of chronic arthritis pain in the elderly. *Drugs Aging*. 2010;27(6):471-490. [PubMed 20524707]
83. Flint RB, Roofthoof DW, van Rongen A, et al. Exposure to acetaminophen and all its metabolites upon 10, 15, and 20 mg/kg intravenous acetaminophen in very-preterm infants. *Pediatr Res*. 2017;82(4):678-684. doi:10.1038/pr.2017.129 [PubMed 28553988]
84. Forrest JA, Clements JA, Prescott LF. Clinical pharmacokinetics of paracetamol. *Clin Pharmacokinet*. 1982;7(2):93-107. doi:10.2165/00003088-198207020-00001 [PubMed 7039926]
85. Gadisseur AP, Van Der Meer FJ, Rosendaal FR. Sustained Intake of Paracetamol (Acetaminophen) During Oral Anticoagulant Therapy With Coumarins Does Not Cause Clinically Important INR Changes: A Randomized Double-Blind Clinical Trial. *J Thromb Haemost*. 2003;1(4):714-717. [PubMed 12871405]
86. Gebauer MG, Nyfort-Hansen K, Henschke PJ, et al. Warfarin and Acetaminophen Interaction. *Pharmacotherapy*. 2003;23(1):109-112. [PubMed 12523469]
87. Ghaderian M, Armanian AM, Sabri MR, Montaseri M. Low-dose intravenous acetaminophen versus oral ibuprofen for the closure of patent ductus arteriosus in premature neonates. *J Res Med Sci*. 2019;24:13. doi:10.4103/jrms.JRMS_631_17 [PubMed 30988681]
88. Goday Arno A, Farré M, Rodríguez-Morató J, et al. Pharmacokinetics in morbid obesity: influence of two bariatric surgery techniques on paracetamol and caffeine metabolism. *Obes Surg*. 2017;27(12):3194-3201. doi:10.1007/s11695-017-2745-z [PubMed 28560524]
89. Gou X, Wang Y, Tang Y, et al. Association of maternal prenatal acetaminophen use with the risk of attention deficit/hyperactivity disorder in offspring: a meta-analysis. *Aust N Z J Psychiatry*. 2019;53(3):195-206. doi:10.1177/0004867418823276 [PubMed 30654621]
90. Gover A, Levy PT, Rotschild A, et al. Oral versus intravenous paracetamol for patent ductus arteriosus closure in preterm infants. *Pediatr Res*. 2022;92(4):1146-1152. doi:10.1038/s41390-022-01944-w [PubMed 35087197]
91. Gupta NK, Lewis JH. Review article: the use of potentially hepatotoxic drugs in patients with liver disease. *Aliment Pharmacol Ther*. 2008;28(9):1021-1041. doi:10.1111/j.1365-2036.2008.03822.x [PubMed 18671777]
92. Gustavson K, Ystrom E, Ask H, et al. Acetaminophen use during pregnancy and offspring attention deficit hyperactivity disorder - a longitudinal sibling control study. *JCPP Adv*. 2021;1(2):e12020. doi:10.1002/jcv2.12020 [PubMed 37431475]
93. Hahn TW, Henneberg SW, Holm-Knudsen RJ, et al. Pharmacokinetics of rectal paracetamol after repeated dosing in children. *Br J Anaesth*. 2000;85(4):512-519. [PubMed 11064607]
94. Hamilton JP, Goldberg E, Chopra S. Management of pain in patients with advanced chronic liver disease or cirrhosis. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed November 25, 2019b.
95. Hammer GB, Maxwell LG, Taicher BM, et al. Randomized population pharmacokinetic analysis and safety of intravenous acetaminophen for acute postoperative pain in neonates and infants. *J Clin Pharmacol*. 2020;60(1):16-27. doi:10.1002/jcph.1508 [PubMed 31448420]
96. Härkin P, Härmä A, Aikio O, et al. Paracetamol accelerates closure of the ductus arteriosus after premature birth: a randomized trial. *J Pediatr*. 2016;177:72-77.e2. doi:10.1016/j.jpeds.2016.04.066 [PubMed 27215779]
97. Hauben M, Bai S, Hung E, Lobello K, Tressler C, Zucal VP. Maternal paracetamol intake and fetal ductus arteriosus constriction/closure: comprehensive signal evaluation using the Austin

- Bradford Hill criteria. *Eur J Clin Pharmacol.* 2021;77(7):1019-1028. doi:10.1007/s00228-020-03039-z [PubMed 33410971]
98. Hayward KL, Powell EE, Irvine KM, Martin JH. Can paracetamol (acetaminophen) be administered to patients with liver impairment? *Br J Clin Pharmacol.* 2016;81(2):210-222. doi:10.1111/bcp.12802 [PubMed 26460177]
 99. Heard K, Bui A, Mlynarchek SL, et al. Toxicity from repeated doses of acetaminophen in children: assessment of causality and dose in reported cases. *Am J Ther.* 2014;21(3):174-183. [PubMed 22407198]
 100. Hey P, Gow P. Letter to the editor: rethinking public health measures to reduce liver injury from therapeutic acetaminophen ingestion. *Hepatology.* 2022;75(2):495-496. doi:10.1002/hep.32171 [PubMed 34553389]
 101. Ho MH, Tung JY, Lee TL, Tsoi NS, Lau YL. Anaphylaxis to paracetamol. *J Paediatr Child Health.* 2008;44(12):746-747. doi:10.1111/j.1440-1754.2008.01419.x [PubMed 19166535]
 102. Howard CR, Howard FM, Weitzman ML. Acetaminophen analgesia in neonatal circumcision: the effect on pain. *Pediatrics.* 1994;93(4):641-646. [PubMed 8134222]
 103. Howell TK, Patel D. Plasma paracetamol concentrations after different doses of rectal paracetamol in older children A comparison of 1 g vs. 40 mg x kg(-1). *Anaesthesia.* 2003;58(1):69-73. doi:10.1046/j.1365-2044.2003.02788_5.x [PubMed 12523328]
 104. Hurden EL, Harvey DL, Lewis PJ. Excretion of paracetamol in human breast milk, Neonatal Society Meeting Held on 4 July 1980 in Southampton. *Arch Dis Child.* 1980;55(12):969-972. [PubMed 21032602]
 105. Hutson JR, Lurie A, Eastabrook G, de Vrijer B, Garcia-Bournissen F. Acetaminophen in late pregnancy and potential for in utero closure of the ductus arteriosus-a pharmacokinetic evaluation and critical review of the literature. *Am J Obstet Gynecol MFM.* 2021;3(1):100288. doi:10.1016/j.ajogmf.2020.100288 [PubMed 33451624]
 106. Hylek EM, Heiman H, Skates SJ, et al. Acetaminophen and Other Risk Factors for Excessive Warfarin Anticoagulation. *JAMA.* 1998;279(9):657-662. [PubMed 9496982]
 107. Imani F, Motavaf M, Safari S, Alavian SM. The therapeutic use of analgesics in patients with liver cirrhosis: a literature review and evidence-based recommendations. *Hepat Mon.* 2014;14(10):e23539. [PubMed 25477978]
 108. "Inactive" ingredients in pharmaceutical products: update (subject review). American Academy of Pediatrics (AAP) Committee on Drugs. *Pediatrics.* 1997;99(2):268-278. [PubMed 9024461]
 109. Infants' Tylenol oral suspension (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014.
 110. Institute for Safe Medication Practices (ISMP). ISMP List of High-Alert Medications in Community/Ambulatory Care Settings. ISMP; 2021. https://online.ectri.org/hubfs/ISMP/Resources/ISMP_HighAlert_Co
 111. Institute for Safe Medication Practices (ISMP). IV Acetaminophen and Overdoses in Kids. *ISMP Medication Safety Alert.* April 19, 2012.
 112. Isaksson M, Jansson L. Contact allergy to Tween 80 in an inhalation suspension. *Contact Dermatitis.* 2002;47(5):312-313. [PubMed 12534540]
 113. Ito S. Drug therapy for breast-feeding women. *N Engl J Med.* 2000;343(2):118-126. [PubMed 10891521]
 114. Ito S, Blajchman A, Stephenson M, Eliopoulos C, Koren G. Prospective follow-up of adverse reactions in breast-fed infants exposed to maternal medication. *Am J Obstet Gynecol.* 1993;168(5):1393-1399. [PubMed 8498418]
 115. Jr. Tylenol Meltaways chewable tablets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014b.

116. Khan FY, Kabiraj G, Ahmed MA, et al. A systematic review of the link between autism spectrum disorder and acetaminophen: a mystery to resolve. *Cureus*. 2022;14(7):e26995. doi:10.7759/cureus.26995 [PubMed 35989852]
117. Kleiber N, Calvier E, Mooij MG, et al. Enteral acetaminophen bioavailability in pediatric intensive care patients determined with an oral microtracer and pharmacokinetic modeling to optimize dosing. *Crit Care Med*. 2019;47(12):e975-e983. doi:10.1097/CCM.0000000000004032 [PubMed 31609773]
118. Kliegman RM, Stanton BF, St. Gemell JW, et al, eds. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia, PA: Saunders Elsevier; 2011.
119. Konstantinidou SK, Argyrakopoulou G, Dalamaga M, Kokkinos A. The effects of bariatric surgery on pharmacokinetics of drugs: a review of current evidence. *Curr Nutr Rep*. 2023;12(4):695-708. doi:10.1007/s13668-023-00498-5 [PubMed 37857987]
120. Krenzelok EP, Royal MA. Confusion: acetaminophen dosing changes based on NO evidence in adults. *Drugs R D*. 2012;12(2):45-48. doi: 10.2165/11633010-000000000-00000 [PubMed 22530736]
121. Kulo A, Van Calsteren K, van de Velde M, et al. Weight, pregnancy and oral contraceptives affect intravenous paracetamol clearance in young women. *Eur Rev Med Pharmacol Sci*. 2014;18(5):599-604. [PubMed 24668698]
122. Kumar A, Gosavi RS, Sundaram V, et al. Oral paracetamol vs oral ibuprofen in patent ductus arteriosus: a randomized, controlled, noninferiority trial. *J Pediatr*. 2020;222:79-84.e2. doi:10.1016/j.jpeds.2020.01.058 [PubMed 32336479]
123. Kwan D, Bartle WR, Walker SE. The Effects of Acute and Chronic Acetaminophen Dosing on the Pharmacodynamics and Pharmacokinetics of (R)- and (S)-Warfarin. *Clin Pharmacol Ther*. 1995;57:212.
124. Kwok J, Luedecke E, Hall HA, Murray AL, Auyeung B. Analgesic drug use in pregnancy and neurodevelopment outcomes: an umbrella review. *Neurosci Biobehav Rev*. 2022;136:104607. doi:10.1016/j.neubiorev.2022.104607 [PubMed 35276298]
125. Larson AM. Acetaminophen hepatotoxicity. *Clin Liver Dis*. 2007;11(3):525-548. doi: 10.1016/j.cld.2007.06.006 [PubMed 17723918]
126. Larson AM, Polson J, Fontana RJ, et al; Acute Liver Failure Study Group. Acetaminophen-induced acute liver failure: results of a United States multicenter, prospective study. *Hepatology*. 2005;42(6):1364-1372. doi:10.1002/hep.20948 [PubMed 16317692]
127. Lavonas EJ, Reynolds KM, Dart RC. Therapeutic acetaminophen is not associated with liver injury in children: a systematic review. *Pediatrics*. 2010;126(6):e1430-e1444. [PubMed 21098156]
128. Lee HS, Ti TY, Lye WC, Khoo YM, Tan CC. Paracetamol and its metabolites in saliva and plasma in chronic dialysis patients. *Br J Clin Pharmacol*. 1996;41(1):41-47. [PubMed 8824692]
129. Lee P, Chuang Y, Hu Y, et al. Maternal acetaminophen use and child neurodevelopment. *JAMA Pediatr*. Published online March 9, 2026. doi:10.1001/jamapediatrics.2026.0071 [PubMed 41801232]
130. Lee WM. Acetaminophen (APAP) hepatotoxicity-Isn't it time for APAP to go away? *J Hepatol*. 2017;67(6):1324-1331. doi:10.1016/j.jhep.2017.07.005 [PubMed 28734939]
131. Li DK, Liu L, Odouli R. Exposure to non-steroidal anti-inflammatory drugs during pregnancy and risk of miscarriage: population based cohort study. *BMJ*. 2003;327(7411):368. doi:10.1136/bmj.327.7411.368 [PubMed 12919986]
132. Liew Z, Yuan Y, Meng Q, et al. Prenatal exposure to acetaminophen and childhood asthmatic symptoms in a population-based cohort in Los Angeles, California. *Int J Environ Res Public Health*. 2021;18(19):10107. doi:10.3390/ijerph181910107 [PubMed 34639411]

133. Liukas A, Kuusniemi K, Aantaa R, et al. Pharmacokinetics of intravenous paracetamol in elderly patients. *Clin Pharmacokinet*. 2011;50(2):121-129. [PubMed 21241071]
134. Lourido-Cebreiro T, Salgado FJ, Valdes L, Gonzalez-Barcala FJ. The association between paracetamol and asthma is still under debate. *J Asthma*. 2017;54(1):32-38. doi:10.1080/02770903.2016.1194431 [PubMed 27575940]
135. Louvet A, Ntandja Wandji LC, Lemaître E, et al. Acute liver injury with therapeutic doses of acetaminophen: a prospective study. *Hepatology*. 2021;73(5):1945-1955. doi:10.1002/hep.31678 [PubMed 33306215]
136. Lucente P, Iorizzo M, Pazzaglia M. Contact sensitivity to Tween 80 in a child. *Contact Dermatitis*. 2000;43(3):172. [PubMed 10985636]
137. Luecke CM, Liviskie CJ, Zeller BN, Vesoulis ZA, McPherson C. Acetaminophen for patent ductus arteriosus in extremely low-birth-weight neonates. *J Pediatr Pharmacol Ther*. 2017;22(6):461-466. doi:10.5863/1551-6776-22.6.461 [PubMed 29290747]
138. Makin AJ, Wendon J, Williams R. A 7-year experience of severe acetaminophen-induced hepatotoxicity (1987-1993). *Gastroenterology*. 1995;109(6):1907-1916. doi:10.1016/0016-5085(95)90758-0 [PubMed 7498656]
139. Makris UE, Abrams RC, Gurland B, Reid MC. Management of persistent pain in the older patient: a clinical review. *JAMA*. 2014;312(8):825-836. [PubMed 25157726]
140. Mapap Children's (acetaminophen) [prescribing information]. Livonia, MI: Major Pharmaceuticals; October 2015.
141. Mariano ER. Approach to the management of acute pain in adults. Connor RF, ed. UpToDate. Waltham, MA: UpToDate Inc. <https://www.uptodate.com>. Accessed October 2, 2025.
142. Martin E, Vickers B, Landau R, Reece-Stremtan S. ABM clinical protocol #28, peripartum analgesia and anesthesia for the breastfeeding mother. *Breastfeed Med*. 2018;13(3):164-171. [PubMed 29595994]
143. Martin U, Temple RM, Winney RJ, Prescott LF. The disposition of paracetamol and the accumulation of its glucuronide and sulphate conjugates during multiple dosing in patients with chronic renal failure. *Eur J Clin Pharmacol*. 1991;41(1):43-46. [PubMed 1782975]
144. Martin U, Temple RM, Winney RJ, Prescott LF. The disposition of paracetamol and its conjugates during multiple dosing in patients with end-stage renal failure maintained on haemodialysis. *Eur J Clin Pharmacol*. 1993;45(2):141-145. [PubMed 8223835]
145. Matheson I, Lunde PK, Notarianni L. Infant rash caused by paracetamol in breast milk? *Pediatrics*. 1985; 76(4):651-652. [PubMed 2931668]
146. Mayoral CE, Marino RV, Rosenfeld W, et al. Alternating Antipyretics: Is This an Alternative? *Pediatrics*. 2000;105(5):1009-1012. [PubMed 10790455]
147. McLachlan AJ, Bath S, Naganathan V, et al. Clinical pharmacology of analgesic medicines in older people: impact of frailty and cognitive impairment. *Br J Clin Pharmacol*. 2011;71(3):351-364.
148. McNeil Consumer Healthcare website. Frequently Asked Questions About New Dosing for Extra Strength Tylenol Products. Published 2011. Accessed May 6, 2014.
149. Mehlich DR, Aspley S, Daniels SE, et al. Comparison of the Analgesic Efficacy of Concurrent Ibuprofen and Paracetamol With Ibuprofen or Paracetamol Alone in the Management of Moderate to Severe Acute Postoperative Dental Pain in Adolescents and Adults: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group, Single-Dose, Two-Center, Modified Factorial Study. *Clin Ther*. 2010;32(5):882-895. [PubMed 20685496]
150. Memisoglu A, Alp Ünkar Z, Cetiner N, et al. Ductal closure with intravenous paracetamol: a new approach to patent ductus arteriosus treatment. *J Matern Fetal Neonatal Med*. 2016;29(6):987-990. doi:10.3109/14767058.2015.1029912 [PubMed 25781500]

151. Mireskandari SM, Makarem J. Effect of rectal diclofenac and acetaminophen alone and in combination on postoperative pain after cleft palate repair in children. *J Craniofac Surg*. 2011;22(5):1955-1959. [PubMed 21959480]
152. Myers RP, Shaheen AA, Li B, Dean S, Quan H. Impact of liver disease, alcohol abuse, and unintentional ingestions on the outcomes of acetaminophen overdose. *Clin Gastroenterol Hepatol*. 2008;6(8):918-25; quiz 837. doi:10.1016/j.cgh.2008.02.053 [PubMed 18486561]
153. Naga Rani MA, Joseph T, Narayanan R. Placental transfer of paracetamol. *J Indian Med Assoc*. 1989;87(8):182-183. [PubMed 2621359]
154. National Center for Immunization and Respiratory Diseases, Advisory Committee on Immunization Practices (NCIRD/ACIP). General recommendations on immunization --- recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2011;60(2):1-64. [PubMed 21293327]
155. Nitsche JF, Patil AS, Langman LJ, et al. Transplacental passage of acetaminophen in term pregnancy. *Am J Perinatol*. 2017;34(6):541-543. doi:10.1055/s-0036-1593845 [PubMed 27806383]
156. Notarianni LJ, Oldham HG, Bennett PN. Passage of paracetamol into breast milk and its subsequent metabolism by the neonate. *Br J Clin Pharmacol*. 1987, 24(1):63-67. [PubMed 3620287]
157. Numata T, Fukushi R, Ito T, Tsuboi R, Harada K. Acetaminophen anaphylaxis diagnosed by skin prick test. *Allergol Int*. 2016;65(4):490-491. doi:10.1016/j.alit.2016.04.019 [PubMed 27460168]
158. Ofirmev (acetaminophen) [prescribing information]. Hazelwood, MO: Mallinckrodt; June 2020.
159. Ofirmev (acetaminophen) [prescribing information]. Hazelwood, MO: Mallinckrodt; March 2018.
160. Ohlsson A, Shah PS. Paracetamol (acetaminophen) for prevention or treatment of pain in newborns. *Cochrane Database Syst Rev*. 2020;1(1):CD011219. doi:10.1002/14651858.CD011219.pub4 [PubMed 31985830]
161. Øie S, Lowenthal DT, Briggs WA, Levy G. Effect of hemodialysis on kinetics of acetaminophen elimination by anephric patients. *Clin Pharmacol Ther*. 1975;18(6):680-686. [PubMed 1204276]
162. Okubo Y, Hayakawa I, Sugitate R, Nariai H. Maternal acetaminophen use and offspring's neurodevelopmental outcome: a nationwide birth cohort study. *Paediatr Perinat Epidemiol*. Published online September 2, 2025. doi:10.1111/ppe.70071 [PubMed 40898607]
163. Oncel MY, Yurttutan S, Erdevi O, et al. Oral paracetamol versus oral ibuprofen in the management of patent ductus arteriosus in preterm infants: a randomized controlled trial. *J Pediatr*. 2014;164(3):510-4.e1. doi:10.1016/j.jpeds.2013.11.008 [PubMed 24359938]
164. Pandharipande P, McGrane S. Pain control in the critically ill adult patient. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed December 9, 2019.
165. Patel R, Sushko K, van den Anker J, Samiee-Zafarghandy S. Long-term safety of prenatal and neonatal exposure to paracetamol: a systematic review. *Int J Environ Res Public Health*. 2022;19(4):2128. doi:10.3390/ijerph19042128 [PubMed 35206317]
166. Perzanowski MS, Miller RL, Tang D, et al. Prenatal acetaminophen exposure and risk of wheeze at age 5 years in an urban low-income cohort. *Thorax*. 2010, 65(2):118-23. [PubMed 19850963]
167. Peterson SJB, Weisman SJ. Pediatric pain management. In: Kliegman RM, St. Geme J, eds. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2024:chap. 93.
168. Porat D, Markovic M, Zur M, et al. Increased paracetamol bioavailability after sleeve gastrectomy: a crossover pre- vs. post-operative clinical trial. *J Clin Med*. 2019;8(11):1949. doi:10.3390/jcm8111949 [PubMed 31726725]
169. Prescott LF. Paracetamol overdosage. Pharmacological considerations and clinical management. *Drugs*. 1983;25(3):290-314. doi:10.2165/00003495-198325030-00002 [PubMed 6343056]

170. Prescott LF, Speirs GC, Critchley JA, Temple RM, Winney RJ. Paracetamol disposition and metabolite kinetics in patients with chronic renal failure. *Eur J Clin Pharmacol.* 1989;36(3):291-297. doi:10.1007/BF00558162 [PubMed 2744070]
171. Prins SA, Van Dijk M, Van Leeuwen P, et al. Pharmacokinetics and analgesic effects of intravenous propacetamol vs rectal paracetamol in children after major craniofacial surgery. *Paediatr Anaesth.* 2008;18(7):582-592. [PubMed 18482233]
172. Prymula R, Siegrist CA, Chlibek R, et al. Effect of prophylactic paracetamol administration at time of vaccination on febrile reactions and antibody responses in children: two open-label, randomised controlled trials. *Lancet.* 2009;374(9698):1339-1350. [PubMed 19837254]
173. Ramachandran A, Jaeschke H. Acetaminophen hepatotoxicity. *Semin Liver Dis.* 2019;39(2):221-234. doi:10.1055/s-0039-1679919 [PubMed 30849782]
174. Rastogi R, Meek BD. Management of chronic pain in elderly, frail patients: finding a suitable, personalized method of control. *Clin Interv Aging.* 2013;8:37-46. [PubMed 23355774]
175. Rebordosa C, Kogevinas M, Bech BH, Sørensen HT, Olsen J. Use of acetaminophen during pregnancy and risk of adverse pregnancy outcomes. *Int J Epidemiol.* 2009;38(3):706-714. doi:10.1093/ije/dyp151 [PubMed 19332503]
176. Rebordosa C, Kogevinas M, Sørensen HT, Olsen J. Pre-natal exposure to paracetamol and risk of wheezing and asthma in children: a birth cohort study. *Int J Epidemiol.* 2008;37(3):583-590. doi:10.1093/ije/dyn070 [PubMed 18400839]
177. Reece-Stremtan S, Campos M, Kokajko L; Academy of Breastfeeding Medicine. ABM clinical protocol #15: analgesia and anesthesia for the breastfeeding mother, revised 2017. *Breastfeed Med.* 2017;12(9):500-506. [PubMed 29624435]
178. Refer to manufacturer's labeling.
179. Riad W, Moussa A. Pre-operative analgesia with rectal diclofenac and/or paracetamol in children undergoing inguinal hernia repair. *Anaesthesia.* 2007;62(12):1241-1245. [PubMed 17991260]
180. Riggs BS, Bronstein AC, Kulig K, et al. Acute acetaminophen overdose during pregnancy. *Obstet Gynecol.* 1989;74(2):247-253. [PubMed 2748061]
181. Rogal SS, Hansen L, Patel A, et al. AASLD practice guidance: palliative care and symptom-based management in decompensated cirrhosis. *Hepatology.* 2022;76(3):819-853. doi:10.1002/hep.32378 [PubMed 35103995]
182. Roofthoof DWE, Simons SHP, van Lingen RA, et al. Randomized controlled trial comparing different single doses of intravenous paracetamol for placement of peripherally inserted central catheters in preterm infants. *Neonatology.* 2017;112(2):150-158. doi:10.1159/000468975 [PubMed 28558384]
183. Sachs HC, Committee On Drugs. The transfer of drugs and therapeutics into human breast milk: an update on selected topics. *Pediatrics.* 2013;132(3):e796-809. [PubMed 23979084]
184. Schindler T, Smyth J, Bolisetty S, et al. Early PARacetamol (EPAR) trial: a randomized controlled trial of early paracetamol to promote closure of the ductus arteriosus in preterm infants. *Neonatology.* 2021;118(3):274-281. doi:10.1159/000515415 [PubMed 33845473]
185. Scialli AR, Ang R, Breitmeyer J, Royal MA. Childhood asthma and use during pregnancy of acetaminophen. A critical review. *Reprod Toxicol.* 2010;30(4):508-519. doi:10.1016/j.reprotox.2010.09.005 [PubMed 20933592]
186. Schwenk ES. Nonopioid pharmacotherapy for acute pain in adults. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed December 5, 2019.
187. Scoville BA, Vulaj V, Mueller BA, Annich GM, Wagner DS. Acetaminophen clearance during ex vivo continuous renal replacement therapies. *J Artif Organs.* 2018;21(2):215-219. doi:10.1007/s10047-017-1001-6 [PubMed 29063343]

188. Serper M, Wolf MS, Parikh NA, Tillman H, Lee WM, Ganger DR. Risk factors, Clinical presentation, and outcomes in overdose with acetaminophen alone or with combination products: Results from the acute liver failure study group. *J Clin Gastroenterol.* 2016;50(1):85-91. doi:10.1097/MCG.0000000000000378 [PubMed 26166142]
189. Shastri N. Intravenous acetaminophen use in pediatrics. *Pediatr Emerg Care.* 2015;31(6):444-450. [PubMed 26035501]
190. Shehab N, Lewis CL, Streetman DD, Donn SM. Exposure to the pharmaceutical excipients benzyl alcohol and propylene glycol among critically ill neonates. *Pediatr Crit Care Med.* 2009;10(2):256-259. [PubMed 19188870]
191. Sheikh J, Allotey J, Sobhy S, et al. Maternal paracetamol (acetaminophen) use during pregnancy and risk of autism spectrum disorder and attention deficit/hyperactivity disorder in offspring: umbrella review of systematic reviews. *BMJ.* 2025;391:e088141. doi:10.1136/bmj-2025-088141 [PubMed 41207796]
192. Shelley WB, Talanin N, Shelley ED. Polysorbate 80 hypersensitivity. *Lancet.* 1995;345(8980):1312-1313. [PubMed 7746084]
193. Singh M, Varukolu S, Chauhan A, et al. Paracetamol exposure and asthma: what does the evidence say? An overview of systematic reviews. *Pediatr Pulmonol.* 2021;56(10):3189-3199. doi:10.1002/ppul.25595 [PubMed 34425045]
194. Smith HS. Potential analgesic mechanisms of acetaminophen. *Pain Physician.* 2009;12(1):269-280. [PubMed 19165309]
195. Society for Maternal-Fetal Medicine (SMFM) Publications Committee. Prenatal acetaminophen use and outcomes in children. *Am J Obstet Gynecol.* 2017;216(3):B14-B15. doi:10.1016/j.ajog.2017.01.021 [PubMed 28126365]
196. Society for Maternal-Fetal Medicine (SMFM) Publications Committee. SMFM response to administration announcement on acetaminophen use during pregnancy and autism. <https://www.smfm.org/news/smfm-response-to-administration-announcement-on-acetaminophen-use-during-pregnancy-and-autism>. Updated September 22, 2025. Accessed October 7, 2025.
197. Society of Obstetricians and Gynaecologists of Canada (SOGC). SOGC position statement on the use of acetaminophen for analgesia and fever in pregnancy. <https://sogc.org/common/Uploaded%20files/Pos>. Published September 12, 2025. Accessed November 20, 2025.
198. Sullivan JE, Farrar HC. Fever and antipyretic use in children. *Pediatrics.* 2011;127(3):580-587. [PubMed 21357332]
199. Sznajder KK, Teti DM, Kjerulff KH. Maternal use of acetaminophen during pregnancy and neurobehavioral problems in offspring at 3 years: a prospective cohort study. *PLoS One.* 2022;17(9):e0272593. doi:10.1371/journal.pone.0272593 [PubMed 36170224]
200. Tadokoro-Cuccaro R, Fisher BG, Thankamony A, Ong KK, Hughes IA. Maternal paracetamol intake during pregnancy-impacts on offspring reproductive development. *Front Toxicol.* 2022;4:884704. doi:10.3389/ftox.2022.884704 [PubMed 35499038]
201. Temple AR, Benson GD, Zinsenheim JR, Schweinle JE. Multicenter, randomized, double-blind, active-controlled, parallel-group trial of the long-term (6-12 months) safety of acetaminophen in adult patients with osteoarthritis. *Clin Ther.* 2006;28(2):222-235. doi:10.1016/j.clinthera.2006.02.004 [PubMed 16678643]
202. Terrin G, Conte F, Oncel MY, et al. Paracetamol for the treatment of patent ductus arteriosus in preterm neonates: a systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed.* 2016;101(2):F127-F136. [PubMed 26283668]
203. Thompson G, Bundell C, Lucas M. Paracetamol allergy in clinical practice. *Aust J Gen Pract.* 2019;48(4):216-219. doi:10.31128/AJGP-08-18-4689 [PubMed 31256492]

204. Tofe I, Ruiz-González MD, Cañete MD, et al. Efficacy of paracetamol in closure of ductus arteriosus in infants under 32 weeks of gestation. *Front Pediatr*. 2018;6:25. doi:10.3389/fped.2018.00025 [PubMed 29492399]
205. Towers CV, Shelton S, van Nes J, et al. Preoperative cesarean delivery intravenous acetaminophen treatment for postoperative pain control: a randomized double-blinded placebo control trial. *Am J Obstet Gynecol*. 2018;218(3):353.e1-353.e4. doi:10.1016/j.ajog.2017.12.203 [PubMed 29274831]
206. Tylenol 8 Hour Arthritis Pain (acetaminophen) [prescribing information]. Ft. Washington, PA: McNeil Consumer Healthcare Division; received February 2019.
207. Tylenol 8 HR extended release caplets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014.
208. Tylenol arthritis pain extended release caplets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014.
209. Tylenol Children's (acetaminophen) [prescribing information]. Ft. Washington, PA: McNeil Consumer; received July 2018.
210. Tylenol extra strength caplets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil-PPC, Inc; 2014.
211. *Tylenol for Healthcare Professionals*. Fort Washington, PA: McNeil-PPC Inc; 2014.
212. Tylenol Infant's Drops (acetaminophen) [prescribing information]. Ft. Washington, PA: McNeil Consumer Healthcare Division; received January 2019.
213. Tylenol regular strength tablets (acetaminophen) [prescribing information]. Fort Washington, PA: McNeil Consumer Healthcare Division; received June 2020.
214. US Food and Drug Administration (FDA). Notice to physicians on the use of acetaminophen during pregnancy. <https://www.fda.gov/media/188843/download>. Published September 22, 2025. Accessed September 24, 2025.
215. van den Bemt PM, Geven LM, Kuitert NA, et al. The Potential Interaction Between Oral Anti-coagulants and Acetaminophen in Everyday Practice. *Pharm World Sci*. 2002;24(5):201-204. [PubMed 12426965]
216. van Lingen RA, Deinum HT, Quak CM, et al. Multiple-dose pharmacokinetics of rectally administered acetaminophen in term infants. *Clin Pharmacol Ther*. 1999;66(5):509-515. [PubMed 10579478]
217. Vernon S, Bacon C, Weightman D. Rectal paracetamol in small children with fever. *Arch Dis Child*. 1979;54(6):469-470. [PubMed 475432]
218. Viitanen H, Tuominen N, Vääräniemi H, Nikanne E, Annala P. Analgesic efficacy of rectal acetaminophen and ibuprofen alone or in combination for paediatric day-case adenoidectomy. *Br J Anaesth*. 2003;91(3):363-367. [PubMed 12925475]
219. Vojjala N, Mehtani R, Singh V. Letter to the editor: acute liver injury due to therapeutic doses of acetaminophen-confounders must be ruled out! *Hepatology*. 2021;74(5):2904. doi:10.1002/hep.31774 [PubMed 33636014]
220. Watanabe H, Kamiyama T, Sasaki S, et al. Toxic epidermal necrolysis caused by acetaminophen featuring almost 100% skin detachment: Acetaminophen is associated with a risk of severe cutaneous adverse reactions. *J Dermatol*. 2016;43(3):321-324. doi:10.1111/1346-8138.13073 [PubMed 26362011]
221. Watkins PB, Kaplowitz N, Slattery JT, et al. Aminotransferase elevations in healthy adults receiving 4 grams of acetaminophen daily: a randomized controlled trial. *JAMA*. 2006;296(1):87-93. doi:10.1001/jama.296.1.87 [PubMed 16820551]

222. Whitcomb DC, Block GD. Association of acetaminophen hepatotoxicity with fasting and ethanol use. *JAMA*. 1994;272(23):1845-1850. doi: 10.1001/jama.1994.03520230055038 [PubMed 7990219]
223. Wilson-Smith EM, Morton NS. Survey of I.V. paracetamol (acetaminophen) use in neonates and infants under 1 year of age by UK anesthetists. *Paediatr Anaesth*. 2009;19(4):329-337. [PubMed 19335345]
224. Wong A, Gaudins A. Risk prediction of hepatotoxicity in paracetamol poisoning. *Clin Toxicol (Phila)*. 2017;55(8):879-892. doi:10.1080/15563650.2017.1317349 [PubMed 28447858]
225. World Health Organization (WHO). Breastfeeding and maternal medication, recommendations for drugs in the eleventh WHO model list of essential drugs. 2002. <https://apps.who.int/iris/handle/10665/6209-2025-who-statement-on-autism-related-issues>. Published September 24, 2025. Accessed November 20, 2025.
226. World Health Organization (WHO). WHO statement on autism-related issues. <https://www.who.int/news/item/09-2025-who-statement-on-autism-related-issues>. Published September 24, 2025. Accessed November 20, 2025.
227. Worriax JD, Yates JE, Flake D, Saseen JJ. Clinical inquiries. Alcoholic liver disease: is acetaminophen safe?. *J Fam Pract*. 2007;56(8):673-674. [PubMed 17669296]
228. Wynne HA, Cope LH, Herd B, Rawlins MD, James OF, Woodhouse KW. The association of age and frailty with paracetamol conjugation in man. *Age Ageing*. 1990;19:419-424. [PubMed 2285011]
229. Yang B, Gao X, Ren Y, Wang Y, Zhang Q. Oral paracetamol vs. oral ibuprofen in the treatment of symptomatic patent ductus arteriosus in premature infants: A randomized controlled trial. *Exp Ther Med*. 2016;12(4):2531-2536. doi:10.3892/etm.2016.3676 [PubMed 27698754]
230. Yoon E, Babar A, Choudhary M, Kutner M, Pyrsopoulos N. Acetaminophen-induced hepatotoxicity: A comprehensive update. *J Clin Transl Hepatol*. 2016;4(2):131-142. doi:10.14218/JCTH.2015.00052 [PubMed 27350943]
231. Zar T, Graeber C, Perazella MA. Recognition, treatment, and prevention of propylene glycol toxicity. *Semin Dial*. 2007;20(3):217-219. [PubMed 17555487]
232. Zimmerman HJ, Maddrey WC. Acetaminophen (paracetamol) hepatotoxicity with regular intake of alcohol: analysis of instances of therapeutic misadventure. *Hepatology*. 1995;22(3):767-773. Erratum in: *Hepatology* 1995;22(6):1898. [PubMed 7657281]
233. Zuppa AF, Hammer GB, Barrett JS, et al. Safety and population pharmacokinetic analysis of intravenous acetaminophen in neonates, infants, children, and adolescents with pain or Fever. *J Pediatr Pharmacol Ther*. 2011;16(4):246-261. [PubMed 22768009]

Topic 9242 Version 1174.0

Source: UpToDate. Downloaded 2026-05-26.