

# Levobupivacaine: International drug information (concise) - UpToDate

## Levobupivacaine: International drug information (concise) - UpToDate

Levobupivacaine: International drug information (concise)

2026© UpToDate, Inc. and its affiliates and/or licensors. All Rights Reserved.

Access UpToDate Lexidrug for additional drug information, tools, and databases.

Contributor Disclosures

For abbreviations, symbols, and age group definitions show table

International Brand Names

International Brand Names by Country

For country code abbreviations (show table)

- (AU) Australia: Chirocaine;
- (BE) Belgium: Chirocaine | Levobupivacaine Fresenius Kabi;
- (DE) Germany: Chirocaine;
- (EG) Egypt: Chirocaine;
- (ES) Spain: Levobupivacaina g.e.s | Levobupivacaina normon;
- (FI) Finland: Chirocaine;
- (FR) France: Chirocaine | Levobupivacaine altan | Levobupivacaine kabi;
- (GB) United Kingdom: Levobupivacain;
- (IE) Ireland: Chirocaine | Levobupivacain;
- (IT) Italy: Chirocaine | Levobupivacaina altan | Levobupivacaina bioindustria;
- (JP) Japan: Popscaine;
- (KW) Kuwait: Chirocaine;
- (LB) Lebanon: Chirocaine;
- (NL) Netherlands: Chirocaine | Levobupivacaine Fresenius Kabi;
- (PT) Portugal: Chirocaine;
- (QA) Qatar: Chirocaine;
- (SE) Sweden: Chirocaine;
- (SI) Slovenia: Chirocaine;
- (TR) Turkey: Chirocaine

Index Terms

- (-)-bupivacaine;
- (S)-bupivacaine;
- L(-)-Bupivacaine;
- Levobupivacaine Hydrochloride

Pharmacologic Category

- Local Anesthetic

#### Reported Use

Epidural pain management; surgical anesthesia

#### Dosage Range

**Note:** Dose ranges are not intended to guide prescribing; consult local prescribing information for additional dosage and administration information.

Children >6 months to <12 years: Surgical anesthesia: Local infiltration (Ilioinguinal/iliohypogastric block): 0.5 **mL**/kg/side of 0.25% solution (maximum: 1.25 **mg**/kg/side) or 0.25 **mL**/kg/side of 0.5% solution (maximum: 1.25 **mg**/kg/side).

Adults:

Surgical anesthesia:

Dental: 5 to 10 **mL** of 0.5% to 0.75% solution.

Epidural:

Cesarean section: 15 to 30 **mL** of 0.5% solution; maximum single dose: 150 **mg**.

Surgery, **not** cesarean section: 10 to 20 **mL** of 0.5% to 0.75% solution; maximum single dose: 150 **mg**; prolonged surgeries may require a maximum cumulative dose of 400 **mg**/24 hours.

Intrathecal: 3 **mL** of 0.5% solution.

Local infiltration: 1 to 60 **mL** of 0.25% solution; maximum: 150 **mg**.

Ophthalmic (peribulbar block): 5 to 15 **mL** of 0.75% solution.

Peripheral nerve: 1 to 40 **mL** of 0.25% solution or 1 to 30 **mL** of 0.5% solution; maximum: 150 **mg**.

Pain management:

Labor pain (epidural bolus): 6 to 20 **mL** of 0.25% solution.

Labor pain (epidural infusion): 10 to 15 **mL**/hour of 0.0625% solution or 4 to 10 **mL**/hour of 0.125% solution (maximum: 12.5 **mg**/hour; a maximum cumulative dose of 400 **mg** per 24 hours is recommended).

Peri- and postoperative pain (epidural infusion): 10 to 15 **mL**/hour of 0.0625% solution, or 10 to 15 **mL**/hour of 0.125% solution, or 5 to 7.5 **mL**/hour of 0.25% solution (maximum rate: 18.75 **mg**/hour; maximum cumulative dose: 400 **mg**/24 hours; however, postoperative doses up to 570 **mg**/24 hours have been well tolerated).

#### Product Availability

Product available in various countries; not currently available in the US

#### Dosage Forms

Solution, Infusion/Injection: 2.5 mg/mL, 5 mg/mL, 7.5 mg/mL (10 mL)

Solution, Infusion [premixed]: 0.625 mg/mL, 1.25 mg/mL (100 mL, 200 mL)

Use of UpToDate is subject to the Terms of Use.

Topic 96003 Version 28.0

Source: UpToDate. Downloaded 2026-05-26.