		predictive algorithm reported as endometrial window of implantation (e.g., pre-receptive, receptive, post-receptive)
Treatments to improve uterine/endometrial receptivity		
	97810	Acupuncture, 1 or more needles; without electrical stimulation, initial 15 minutes of personal one-on-one contact with the patient
	97811	Acupuncture, 1 or more needles; without electrical stimulation, each additional 15 minutes of personal one-on-one contact with the patient, with re-insertion of needle(s) (List separately in addition to code for primary procedure)
	97813	Acupuncture, 1 or more needles; with electrical stimulation, initial 15 minutes of personal one-on-one contact with the patient
	97814	Acupuncture, 1 or more needles; with electrical stimulation, each additional 15 minutes of personal one-on-one contact with the patient, with re-insertion of needle(s) (List separately in addition to code for primary procedure)

Scientific References

Agency for Healthcare Research and Quality (AHRQ), U.S. Department of Health and Human Services, National Guideline Clearinghouse | Fertility: assessment and treatment for people with fertility problems. http://guideline.gov/content.aspx?f=rss&id=43841&osrc=12[5/20/2013 9:54:47 AM]

Agency for healthcare research and quality (AHRQ), U.S. Department of Health and Human Services, National Guideline Clearinghouse Comparative Effectiveness Review Number 217 Management of Infertility. AHRQ Publication No. 19-EHC014-EF May 2019.

Alasmari NM, Son WY, Dahan MH. The effect on pregnancy and multiples of transferring 1-3 embryos in women at least 40 years old. J Assist Reprod Genet 2016;33:1195-202.

Albrozi S, Motazedian S, Parsanezhad M, et al. Comparison of the effectiveness of single intrauterine insemination (IUI) versus double IUI per cycle in infertile patients. Fertil Steril 2003. 80:595-99.