- Urothelial (transitional cell) histology[1]; **AND**
- BCG unresponsive defined by one of the following:
  - Persistent or recurrent CIS within 12 months of receiving adequate BCG[2]; **OR**
  - Recurrent high-grade Ta/T1 disease within 6 months of receiving adequate BCG[3]; **OR**
  - T1 high-grade disease at the first evaluation following complete induction BCG course alone (5 of 6 doses of an initial induction course); **AND**
  - Prescribed by, or in consultation with, a urologist or oncologist

Inlexzo (gemcitabine)

- Member is 18 years of age or older; **AND**
- Member has BCG-unresponsive non-muscle invasive bladder cancer with carcinoma in situ (CIS), with or without papillary tumors that has the following characteristics:
  - High-grade; **AND**
  - Urothelial (transitional cell) histology **AND**
  - BCG unresponsive defined by one of the following:
    - Persistent or recurrent CIS within 12 months of receiving adequate BCG[4]; **OR**
    - Recurrent high-grade Ta/T1 disease within 6 months of receiving adequate BCG[5]; **OR**
    - T1 high-grade disease at the first evaluation following complete induction BCG course alone (5 of 6 doses of an initial induction course); **AND**
    - Prescribed by, or in consultation with, a urologist or oncologist

Universal Approval Criteria:

Unless otherwise noted, the review criteria used by OncoHealth to determine medical necessity for anticancer treatments and supportive agents include, but is not limited to:

- National Comprehensive Cancer Network® (NCCN) - use consistent with NCCN recommendations carrying a Category 1 or 2A level of evidence; **OR**
- United States Food and Drug Administration (FDA) labeling - new drugs or regimens (combinations of drugs) consistent with the product labeling; **OR**
- Indications not included in the official FDA labeling or recommended by NCCN (Category 1 or 2A level of evidence) may be considered if determined to be medically necessary per one or more of the following compendia:
  - Clinical Pharmacology (Strong For); **OR**
  - Wolters Kluwer Lexi-Drugs® (Level A); **OR**

[1] Mixed histology tumors are supported if transitional cell histology is predominant
[2] Adequate BCG trial is defined as at least 5 of 6 doses of an initial induction PLUS at least 2 doses as part of maintenance treatment or a second induction
[3] Adequate BCG trial is defined as at least 5 of 6 doses of an initial induction PLUS at least 2 doses as part of maintenance treatment or a second induction