

# geha-coverage-policy-elrex fio.pdf

Searchable OCR derivative produced locally with PaddleOCR PP-StructureV3. Vertical bars delimit recognized table columns. OCR may contain errors; the original GEHA PDF controls.

## Source PDF page 1

### Elrex fio (Elranatamab-bcmm)

Effective Date: 1/1/24

Revision Date(s): 4/1/2024, 9/27/2024; 2/16/2026 Review Date: 2/16/2026

Policy type: Medical Necessity.

Line of Business: All G.E.H.A Commercial Medical Plans

Authorizations are issued for 6 (six) months, unless the ordering physician requests a different timespan or the patients' unique circumstance or condition supports the medical necessity for a different authorization timeframe. Reauthorization requests are reviewed for efficacy, safety and tolerability.

### Indication Specific Criteria:

#### Multiple Myeloma (MM)

Member is 18 years of age or older; AND

Member has received at least four prior therapies and is refractory to at least one proteasome inhibitor (PI), one immunomodulatory agent (IMiD), and one anti-CD38 monoclonal antibody; AND

Member has an Eastern Cooperative Oncology Group (ECOG) score of 0 to 2; AND

Member has not had a stem cell transplant within 12 weeks; AND.

Member has not experienced disease progression while on a bispecific B-cell maturation antigen (BCMA) directed CD3 T-cell engager - containing regimen; AND

Must be prescribed by, or in consultation with, an oncologist

### Universal Approval Criteria:

Unless otherwise noted, the review criteria used by OncoHealth to determine medical necessity for anticancer treatments and supportive agents include, but is not limited to:

National Comprehensive Cancer Network® (NCCN) - use consistent with NCCN recommendations carrying a Category 1 or 2A level of evidence; OR

United States Food and Drug Administration (FDA) labeling - new drugs or regimens (combinations of drugs) consistent with all components of the product labeling; OR.

Indications not included in the official FDA labeling or recommended by NCCN (Category 1 or 2A level of evidence) may be considered if determined to be medically necessary per one or more of the following compendia:

O Clinical Pharmacology (Strong For); OR

O Wolters Kluwer Lexi-Drugso (Level A); OR

Other uses of drugs and biologics may be considered medically necessary if supported as safe and effective according to peer-reviewed articles from one of the following journals:

Source PDF page 2

=G.E.H.A

O American Journal of Medicine; Annals of Internal Medicine; Annals of Oncology; Annals of Surgical Oncology; Biology of Blood and Marrow Transplantation; Blood; Bone Marrow Transplantation; British Journal of Cancer; British Journal of Hematology; British Medical Journal; Cancer; Clinical Cancer Research; Drugs; European Journal of Cancer (formerly the European Journal of Cancer and Clinical Oncology); Gynecologic Oncology; International Journal of Radiation, Oncology, Biology, and Physics; The Journal of the American Medical Association, Journal of Clinical Oncology; Journal of the National Cancer Institute; Journal of the National Comprehensive Cancer Network (NCCN); Journal of Urology; Lancet; Lancet Oncology; Leukemia; The New England Journal of Medicine; Radiation Oncology;

O Meeting abstracts and case reports are excluded from consideration;

Non-standard protocols may be approved based on unique clinical circumstances;

Dose and frequency should be consistent with United States Food and Drug Administration (FDA) labeling, National Comprehensive Cancer Network (NCCN), or indication specific peer-reviewed literature;

In the instance that a request is made for drug(s) that was (were) previously tried (including in the same pharmacologic class or with the same mechanism of action) and such drug(s) was (were) discontinued due to a lack of efficacy the request may be subject to an off-label review for medical necessity unless supported by the NCCN or high quality literature (prospective phase 2 or 3 studies published as full manuscripts in a CMS-supported journal)..

Billing

|           |            |                                   |
|-----------|------------|-----------------------------------|
| Drug Name | HCPCS Code | Description                       |
| Elrexio   | J1323      | Injection, elranatamab-bcmm, 1 mg |

References

1. Elrexio [package insert].Pfizer Inc., New York, NY. Available at:<https://labeling.pfizer.com/ShowLabeling.aspx?id=19669>

2. Elrexio. NCCN Drugs and Biologics Compendium. Available at:[https://www.nccn.org/professionals/drug\\_compendium/content/](https://www.nccn.org/professionals/drug_compendium/content/)

3. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines). National Comprehensive Cancer Network, Inc. 2024. All rights reserved. Accessed February 10, 2026. To view the most recent and complete version of the guideline, go online to NCCN.org.

Disclaimer

Drug Coverage Policies are developed as needed, reviewed and updated at least annually, and are subject to change. Other policies and coverage determination guidelines may apply. Federal and state regulatory requirements and member specific benefit plan documents, if applicable, must be reviewed prior to utilizing this Drug Coverage Policy. This Drug Coverage Policy is for informational purposes only and does not constitute medical advice nor dictate how providers should practice medicine. This policy

## Source PDF page 3

should not be reproduced, stored in a retrieval system, or altered from its original form without written permission from OncoHealth, Inc.

<div style="text-align: center;">For Internal Use ONLY </div>

| Date      | Updates                                                                                                                                                                                                                                                              |
|-----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 10/31/23  | Policy creation                                                                                                                                                                                                                                                      |
| 12/11/23  | Added ECOG footnote; Updated references                                                                                                                                                                                                                              |
| 4/1/2024  | HCPCS updated from J9999 to J1323 with updated description                                                                                                                                                                                                           |
| 9/27/2024 | Annual Review                                                                                                                                                                                                                                                        |
| 6/11/25   | Policy adopted by GEHA; document reviewed for universal accessibility                                                                                                                                                                                                |
| 2/16/2026 | Annual Review OH P&T added -- Member has not experienced disease progression while on. a bispecific B-cell maturation antigen (BCMA) - directed CD3 T-cell engager-containing. regimen progression on BCMA target; added prescriber information; approved by OH P&T. |