

Silber S, Zsolt P, Liu NJ, et al. Conventional in-vitro fertilization versus intracytoplasmic sperm injection for patients requiring microsurgical sperm aspiration. *Human Reproduction*.1994;9(9):1705-09.

Simon C, Vladimirov IK, Castillon Cortes G, et al. Prospective, randomized study of the endometrial receptivity analysis (ERA) test in the infertility work-up to guide personalized embryo transfer versus fresh transfer or deferred embryo transfer. *Fertil Steril* 2016;106: e46–7.

Simon C, Gomez C, Cabanillas S, Vladimirov IK, Castillon G, Giles J, et al. In vitro fertilization with personalized blastocyst transfer versus frozen or fresh blastocyst transfer: a multicenter, randomized clinical trial. *Fertil Steril* 2019;112:e56.

Society for Assisted Reproductive Technology: National Data Summary 2011. Available at: https://www.sartcorsonline.com/rptCSR_PublicMultYear.aspx?reporting year+2017.

Steiner AZ, Prtichard D, Stanczyk FZ, et al. Association between biomarkers of ovarian reserve and infertility among older women of reproductive age. *JAMA* 2017; 318: 1367-1376

Styer AK, Wright DL, Wolkovich AM, Veiga C, Toth TL. Single-blastocyst transfer decreases twingestation without affecting pregnancy outcome. *Fertil Steril* 2008;89:1702–8.

Sunderam S, Kissin DM, Flowers L et al. Assisted reproductive technology surveillance — United States, 2009. *MMWR Surveill Summ* 2012;61(7):1-23.

Takeshima K, Jwa SC, Saito H, et al. Impact of single embryo transfer policy on perinatal outcomes in fresh and frozen cycles — analysis of the Japanese Assisted Reproduction Technology registry between 2007 and 2012. *Fertil Steril* 2016 Feb;105(2):337–46.

Tannus S, Cohen Y, Son WY, et al. Cumulative live birth rate following elective single blastocyst transfer compared with double blastocyst transfer in women aged 40 years and over. *Reproductive Biomedicine online* 2017;35:733-738.

Taylor E, Gomel V. The uterus and fertility. *Fertil Steril* 2008; 89: 1-16. Theodorou E, Jones BP, Cawood S, et al. Adding a low-quality blastocyst to a high-quality blastocyst for a double embryo transfer does not decrease pregnancy and live birth rate. *Acta Obstetricia et Gynecologica Scandinavica*. 2021;100:1124-1131.

Thornhill AR, deDie-Smulders CE, Geraedts JP, Harper JC, Harton GL, et al. ESHRE PGD Consortium ‘Best practice guidelines for clinical preimplantation genetic diagnosis (PGD) and preimplantation genetic screening (PGS)’. *Hum Reprod* 2005;20:35–48.

Thurin A, Hausken J, Hillensjo T, Jablonowska B, Pinborg A, Strandell A, et al. Elective single-embryo transfer versus double-embryo transfer in in vitro fertilization. *N Engl J Med*. 2004;351:2392–2402.